

# Dopaminergic Modulation of Cortical Plasticity in Alzheimer's Disease Patients

Giacomo Koch<sup>\*1,2</sup>, Francesco Di Lorenzo<sup>1,2</sup>, Sonia Bonni<sup>1</sup>, Viola Giacobbe<sup>1</sup>, Marco Bozzali<sup>3</sup>, Carlo Caltagirone<sup>1,2</sup> and Alessandro Martorana<sup>2</sup>

<sup>1</sup>Non Invasive Brain Stimulation Unit, Department of Behavioural and Clinical Neurology, Santa Lucia Foundation IRCCS, Rome, Italy;

<sup>2</sup>Department of Neuroscience, Tor Vergata University, Rome, Italy; <sup>3</sup>Neuroimaging Laboratory, Santa Lucia Foundation, IRCCS, Rome, Italy

In animal models of Alzheimer's disease (AD), mechanisms of cortical plasticity such as long-term potentiation (LTP) and long-term depression (LTD) are impaired. In AD patients, LTP-like cortical plasticity is abolished, whereas LTD seems to be preserved. Dopaminergic transmission has been hypothesized as a new player in ruling mechanisms of cortical plasticity in AD. We aimed at investigating whether administration of the dopamine agonist rotigotine (RTG) could modulate cortical plasticity in AD patients, as measured by theta burst stimulation (TBS) protocols of repetitive transcranial stimulation applied over the primary motor cortex. Thirty mild AD patients were tested in three different groups before and after 4 weeks of treatment with RTG, rivastigmine (RVT), or placebo (PLC). Each patient was evaluated for plasticity induction of LTP/LTD-like effects using respectively intermittent TBS (iTBS) or continuous TBS protocols. Short-latency afferent inhibition (SAI) protocol was performed to indirectly assess central cholinergic activity. A group of age-matched healthy controls was recruited for baseline comparisons. Results showed that at baseline, AD patients were characterized by impaired LTP-like cortical plasticity, as assessed by iTBS. These reduced levels of LTP-like cortical plasticity were increased and normalized after RTG administration. No effect was induced by RVT or PLC on LTP. LTD-like cortical plasticity was not modulated in any condition. Cholinergic activity was increased by both RTG and RVT. Our findings reveal that dopamine agonists may restore the altered mechanisms of LTP-like cortical plasticity in AD patients, thus providing novel implications for therapies based on dopaminergic stimulation.

*Neuropsychopharmacology* (2014) **39**, 2654–2661; doi:10.1038/npp.2014.119; published online 18 June 2014

## INTRODUCTION

In the case of Alzheimer's disease (AD), synaptic loss is the strongest pathophysiological correlate of cognitive decline, indicating that synaptic degeneration has a central role in the development of dementia (Klyubin *et al*, 2008). Experimental animal models showed that accumulation of soluble A $\beta$  oligomers specifically block mechanisms of cortical plasticity such as hippocampal long-term potentiation (LTP) (Shankar *et al*, 2008) which is regarded as an electrophysiological correlate of learning and memory (Palop and Mucke, 2010). In contrast, these oligomers have been shown to electrically facilitate evoked long-term depression (LTD) (Li *et al*, 2009). These events can in turn induce changes in conformation of tau proteins, leading to further detrimental effects on synaptic plasticity and cognition (Boekhoorn *et al*, 2006). Similar mechanisms of cortical plasticity can be investigated *in vivo* and non-

invasively in humans, although the plasticity-induction procedures adopted are not completely identical in humans and animals. In humans, LTP- and LTD-like measures may be obtained by applying repetitive transcranial magnetic stimulation (rTMS) over the primary motor cortex (M1), through protocols of theta burst stimulation (TBS) mimicking those described in animal models (Huang *et al*, 2005). In particular, 600 stimuli of continuous TBS (cTBS) are known to induce reliable LTD-like after-effects, whereas intermittent TBS (iTBS) can be used to obtain the opposite LTP-like effect (Huang *et al*, 2005). In AD patients, LTP-like cortical plasticity has been reported to be severely dampened when investigating M1 (Koch *et al*, 2012). This brain area is not severely affected by AD pathology, at least in its early clinical stages. For this reason, M1 can be used as a reliable model to assess cortical plasticity before the occurrence of diffuse synaptic dysfunction and gray matter loss, which instead are typically observed at later stages of the disease (ie, fully developed dementia) (Suvà *et al*, 1999). Additionally, a strict association has been reported between the levels of Tau protein found in the cerebro-spinal fluid of AD patients and the strength of cortical plasticity (Koch *et al*, 2011).

Dopamine is a key neuromodulator affecting several distinct steps of synaptic transmission including the probability of neurotransmitter release, the postsynaptic sensitivity to neurotransmitter and the membrane excitability of the

\*Correspondence: Dr G Koch, Non Invasive Brain Stimulation Unit, Laboratorio di Neurologia Clinica e Comportamentale, IRCCS Fondazione, S. Lucia, Via Ardeatina, Rome 306-00179, Italy,

Tel: +39 0651501181, Fax: +39 0651501180,

E-mail: giakoch@gmail.com

Received 1 April 2014; revised 6 May 2014; accepted 19 May 2014;

accepted article preview online 26 May 2014

pre- and postsynaptic cells (Tritsch and Sabatini, 2012). Its strong impact on motor cortical plasticity has been documented in both healthy individuals (Kuo *et al*, 2008; Monte-Silva *et al*, 2009) and patients with neurodegenerative disorders such as Parkinson's disease (Huang *et al*, 2011; Kishore *et al*, 2012). There is experimental evidence demonstrating that in the cerebral cortex, as well as in the basal forebrain, dopamine modulates the activity not only of pyramidal cells and GABA interneurons, but also of diffuse cholinergic projections from neurons located in the basal forebrain (Paspalas and Goldman-Rakic, 2005; Zhang *et al*, 2009). Notably, the dysfunction of dopaminergic transmission has been hypothesized as a new player in the pathophysiology of AD (Joyce *et al*, 1998; Kemppainen *et al*, 2003; Kumar and Patel, 2007; Martorana *et al*, 2010). In animal models of AD, dopamine agonists may improve memory and even reduce intraneuronal amyloid deposition (Himeno *et al*, 2011). Localization studies of dopamine receptors in AD brains have shown a preferential reduction of dopamine D2-like receptors in the hippocampus and in the prefrontal cortex (Kemppainen *et al*, 2003; Kumar and Patel, 2007). Moreover, recent TMS studies performed over M1 revealed that in AD patients the impaired cholinergic transmission, measured by short-latency afferent inhibition (SAI) protocol (Di Lazzaro *et al*, 2002), can be transiently restored by the administration of Levodopa (Martorana *et al*, 2009). Similar effects can also be observed with rotigotine (RTG), a dopamine D2/D3 agonist (Martorana *et al*, 2013). Taken altogether, these data impose a careful investigation of the potential role of dopaminergic networks in mediating the synaptic efficiency/dysfunction in AD. In this study, we aimed at investigating the impact of D2/D3 agonist RTG on cortical plasticity in patients suffering from mild AD. We hypothesized that dopaminergic stimulation would increase the altered mechanisms of LTP-like cortical plasticity that characterize AD patients (Koch *et al*, 2012). After recruitment and baseline investigation of cortical plasticity, patients were randomly assigned to three experimental arms: RTG, rivastigmine (RVT) or placebo (PLC). After 4 weeks, patients were reassessed for cortical plasticity.

## MATERIALS AND METHODS

### Subjects

Thirty consecutive patients with a diagnosis of probable AD according to the NINCDS-ADRDA Criteria (Varma *et al*, 1999) and 10 age-, sex- and education-matched healthy subjects (HS) were recruited for this study (Supplementary Table S1). All subjects underwent an extensive clinical investigation, including interviews on their medical history, a full neurological examination, the MMSE, a complete blood screening, neuropsychological assessment, neuropsychiatric evaluation, and magnetic resonance imaging. In the 30 days before entering the study, subjects had to be clear of treatment with any drug inducing modulatory effects on the cerebral cortex excitability, such as antidepressants, benzodiazepines, anti-epileptic drugs, or neuroleptics. All AD patients had to show a moderate level of dementia, as assessed by a neuropsychological evaluation including the MMSE (ranging from 18 to 24) and a standardized

neuropsychological battery (see below). Subjects were excluded if they had either two or more hyperintense lesions with a diameter  $\geq 10$  mm or more than eight hyperintense lesions with a diameter between 5 and 9 mm on dual-echo MR images (Bozzali *et al*, 2011). All participants (or their legal guardian in the case they were incapable) signed a written informed consent after receiving an extensive disclosure of the experimental details. The current study was performed according to the Declaration of Helsinki and was approved by the Ethics Committee of Santa Lucia Foundation IRCSS in Rome (Prot. CE/AG4/PROG.392-08).

### Transcranial Magnetic Stimulation

Motor-evoked potentials (MEPs) were recorded from the right first dorsal interosseous (FDI) muscles using 9-mm diameter, Ag–AgCl surface cup electrodes. Responses were amplified with a Digitimer D360 amplifier (Digitimer, UK) through filters set at 20 Hz and 2 kHz, then recorded by computer using SIGNAL software, at a sampling rate of 5 kHz per channel (Cambridge Electronic Devices, UK). A monophasic Magstim 200 device (Magstim, UK) was used to define the motor hot spot in the left hemisphere and to assess MEP size. The motor hot spot was defined as the location where TMS consistently produced the largest MEP size at 120% of resting motor threshold in the target muscle (Rossini *et al*, 1994). A second coil was connected to a biphasic Super Rapid Magstim stimulator (Magstim) to deliver rTMS. In the cTBS protocol bursts at 80% active motor threshold were repeated at 5 Hz (ie, every 200 ms), whereas each burst consisted of three stimuli repeating at 50 Hz, for 40 s (600 pulses). In the iTBS protocol, a 2 s train of TBS was repeated 20 times, every 10 s for a total of 190 s (600 pulses) (Huang *et al*, 2005). Twenty MEPs were collected and averaged at baseline. Then, over the same hot spot, 20 MEPs were recorded at 1–5, 6–10, 11–15, 16–20, and 21–25 min after rTMS, and averaged. Although the LTP-like effects can be still pronounced after this interval, recording of MEP amplitude was stopped after 25 min in order to account for patients' compliance. SAI was investigated using the technique that has been recently described (Sailer *et al*, 2003), where the conditioning stimulus (CS) is a single pulse (200  $\mu$ s) of electrical stimulation applied through bipolar electrodes to the right median nerve at the wrist (cathode proximal). The intensity of the CS was set at just over motor threshold for evoking a visible twitch of the thenar muscles. The intensity of the cortical magnetic test stimulus (TS) was adjusted to evoke a muscle response in the relaxed right FDI with an amplitude of about 1 mV peak-to-peak. The CS to the peripheral nerve preceded the cortical magnetic TS by different interstimulus intervals (ISI), that ranged from  $-4$  to  $+8$  ms from the N20 component, in steps of 4 ms (Martorana *et al*, 2013). Ten stimuli were delivered at each ISI. Subjects were provided with audiovisual feedback at high gain, in order to assist them in maintaining complete relaxation. The inter-trial interval was set at 5 s ( $\pm 10\%$ ), for a total duration of  $\sim 5$  min. Measurements were made on each individual trial. The mean peak-to-peak amplitude of the conditioned MEP at each ISI was expressed as a percentage of the mean peak-to-peak amplitude of the unconditioned test pulse in that block.

## Cognitive Assessment

All AD patients underwent a neuropsychological examination including a general cognitive index (Mini Mental Examination State), and the following neuropsychological tests for the specific cognitive domains (Carlesimo *et al*, 1996): (1) 15 Rey's Word List (immediate and 15-min delayed recall) to investigate verbal episodic long-term memory; (2) forward and backward Digit span and the forward and backward Corsi Block Tapping task to investigate short-term memory and working memory; (3) frontal assessment battery (FAB) to investigate executive functions (Apollonio *et al*, 2005). Parallel versions were used in the different sessions.

## Drug Administration and Experimental Design

After recruitment and baseline assessments, AD patients were assigned to RTG ( $n=10$ ), RVT ( $n=10$ ) or PLC ( $n=10$ ). Physicians involved in TMS recordings and cognitive assessments were completely blind on which experimental branch each patient was belonging to. All treatments and PLC were administered for 4 weeks with no interruptions. RTG was administered through a 4 mg transdermal patch (Neupro, UCB pharma), after having started with a 2 mg patch for 1 week. We chose this scheme because such a drug has been previously found to be effective in modulating cholinergic activity in AD patients (Martorana *et al*, 2013). We chose to test the effects of RTG, which has an impact only on a subset of dopamine receptors, and not those of levo-dopa, which has a more global DA activation, because RTG can be administered with transdermal patches and it is well tolerated in AD patients (Martorana *et al*, 2013). Domperidone was administered in the case of dizziness occurrence. RVT was administered using Exelon Patch 4.6 mg. PLC was administered using a transdermal patch similar to those medicated. In all cases, each daily patch was maintained for 24 h. The Monday before starting drug/PLC administration, and the Friday at the end of 4-week treatment, AD patients underwent TMS recordings and cognitive assessment. There was an interval of at least 1 h between the different TMS protocols. The order of TMS protocols was pseudorandomized in the different sessions.

For further follow-up evaluation, seven patients in the RTG group were kept on the same drug administration (4 mg transdermal patch) for 8 additional weeks. In these patients, another evaluation was performed on the Monday following the end of treatment, and reassess by TMS and neuropsychological testing the Friday of the 12th week.

## Data Analysis

Data were analyzed using SPSS for Windows version 11.0; we measured the percentage of change of peak-to-peak amplitudes of the mean MEPs at baseline for each subject in each condition. At baseline two-way repeated measure ANOVAs were performed for each protocol (cTBS, iTBS) with TIME (1–5, 6–10, 11–15, 16–20, and 21–25 min after TBS) as within subjects factor and GROUP (RTG, RVT, PLC, HS) as between subjects factor. To test the effects of the drugs, different repeated measure ANOVAs were performed for each protocol (cTBS, iTBS) with SESSION (baseline *vs*

end of treatment), TIME (1–5, 6–10, 11–15, 16–20, and 21–25 min after TBS) as within subjects factors and GROUP (RTG, RVT, PLC) as between subjects factor. For the SAI recordings, the electrophysiological parameters of AD patients were compared at baseline with those of age-matched controls by means of repeated measures ANOVA, with ISI (–4, 0, +4 and +8 ms plus the latency of the N20) as within subject factor and GROUP (RTG, RVT, PLC *vs* HS) as between subjects factor. To test the effects of the drugs, different repeated measure ANOVAs were performed with SESSION (before *vs* end of treatment) and ISI (–4, 0, +4 and +8 ms plus the latency of the N20) as within subjects factors and GROUP (RTG, RVT, PLC) as between subjects factor. The Greenhouse–Geisser correction was used for nonspherical data. When a significant main effect was reached, paired *t*-tests with Bonferroni correction ( $p<0.05$ ) were used to characterize the different effects in all experiments. Mauchly's test examined for sphericity. To compare the effects of RTG, RVT, and PLC on cognitive performance before and after the treatment, paired *t*-tests were performed separately for each test. For all statistical analyses, a  $p$  value  $<0.05$  was considered to be significant.


## RESULTS

The entire procedure was well tolerated in all subjects. The mean (SD) AMT (calculated as a percentage of maximal stimulator output) for TMS was significantly lower in AD patients (RTG: 34.2%; RVT: 33.8%; PLC: 35.2%) compared with HS (38.2%). Baseline MEP amplitudes did not differ between AD subgroups (RTG ( $1.12 \pm 0.41$  mV), RVT ( $1.09 \pm 0.42$  mV), PLC ( $1.21 \pm 0.42$  mV)), and HS ( $1.07 \pm 0.45$  mV).

At baseline, for the cTBS protocol a repeated measures ANOVA was performed on the percentage changes of the mean MEP amplitude: no effect was seen for the GROUP ( $F(3,34) = 0.96$ ;  $p = 0.42$ ) and TIME ( $F(4,136) = 0.52$ ;  $p = 0.71$ ) main factors as was not significant ( $F(12,136) = 0.46$ ;  $p = 0.92$ ) the interaction GROUP  $\times$  TIME (Figure 1a). For the iTBS protocol, there was a significant GROUP ( $F(3,34) = 17.02$ ;  $p = 0.0001$ ; RTG *vs* HS: Cohen's  $d = 0.81$ ; RVT *vs* HS: Cohen's  $d = 0.74$ ; PLC *vs* HS Cohen's  $d = 0.41$ ) but not TIME ( $F(4,136) = 1.26$ ;  $p = 0.28$ ) main factor effect; the interaction GROUP  $\times$  TIME was significant ( $F(12,136) = 2.26$ ;  $p = 0.01$ ), revealing that AD patients showed the expected altered LTP-like cortical plasticity in comparison with HS (Figure 1b). Accordingly, only in the HS group mean MEPs amplitudes were increased in comparison with baseline at 6–10, 11–15; 16–20 and 21–25 min. (all  $p < 0.05$  at paired *t*-test analyses). The ANOVA analysis performed on SAI protocol at baseline showed a significant ISI ( $F(2,102) = 31.3$ ;  $p = 0.029$ ) but not GROUP ( $F(3,34) = 0.60$ ;  $p = 0.61$ ) main factor; the interaction GROUP  $\times$  TIME was not significant ( $F(9,102) = 0.95$ ;  $p = 0.48$ ) (Figure 2).

The different drugs (RTG, RVT, PLC) did not induce any effect on the cTBS protocol as shown by the ANOVA (GROUP main factor:  $F(2,25) = 0.20$ ;  $p = 0.81$ ; SESSION main factor:  $F(1,25) = 65.38$ ;  $p = 0.007$ ; TIME main factor:  $F(4,100) = 0.42$ ;  $p = 0.78$ ; interaction GROUP  $\times$  SESSION  $\times$  TIME:  $F(8,100) = 0.70$ ;  $p = 0.75$ ; Figure 3a). On the other hand for the iTBS protocol, RTG induced an increase of







**Figure 1** After effects of the cTBS (a) and iTBS (b) protocols on MEP amplitude in the different groups of Alzheimer's disease patients at baseline before starting the treatment with RTG, RVT, and PLC. A group of age-matched HS was included for comparison at baseline only. Error bars indicate SEM. Asterisks indicate  $p < 0.05$  when comparing AD patients of the different groups.

LTP-like cortical plasticity in AD patients, as revealed by the ANOVA showing significant GROUP ( $F(2,25) = 51.96$ ;  $p = 0.00004$ ; RTG vs PLC: Cohen's  $d = 0.81$ ; RTG vs RVT: Cohen's  $d = 0.69$ ) and SESSION ( $F(1,25) = 103.73$ ;  $p = 0.000001$ ) main factors and GROUP  $\times$  SESSION ( $F(2,25) = 74.7$ ;  $p = 0.000001$ ) and GROUP  $\times$  TIME ( $F(8,100) = 23.4$ ;  $p = 0.02$ ) interactions. The TIME main factor ( $F(4,100) = 0.24$ ;  $p = 0.88$ ) and the other interactions were not significant (TIME  $\times$  SESSION ( $F(4,100) = 19.2$ ;  $p = 0.11$ ); TIME  $\times$  GROUP  $\times$  SESSION ( $F(8,100) = 12.62$ ;  $p = 0.27$ )). *Post hoc* analysis with Bonferroni correction showed that LTP increased in the RTG group only ( $p < 0.0001$  at all time points in comparison with the other AD groups) (Figure 3b). Accordingly, MEPs amplitudes were increased in comparison with baseline only in the RTG group at 1-5, 6-10, 11-15, 16-20 and 21-25 min. (all  $p < 0.05$  at paired *t*-test analyses).


For the SAI protocol, the ANOVA showed significant ISI main factor ( $F(3,75) = 6.63$ ;  $p = 0.00038$ ), ISI  $\times$  SESSION interaction ( $F(3,75) = 23.70$ ;  $p = 0.0001$ ) and GROUP  $\times$



**Figure 2** After effects of the SAI protocol on MEP amplitude in the different groups of Alzheimer's disease patients at baseline before starting the treatment with RTG, RVT, and PLC. A group of age-matched HS was included for comparison at baseline only. Error bars indicate SEM. Asterisks indicate  $p < 0.05$  when comparing AD patients of the different groups.



**Figure 3** After effects of the cTBS (a) and iTBS (b) protocols on MEP amplitude in Alzheimer's disease patients at baseline (pre) and after 4 weeks (post) of treatment with RTG, RVT, and PLC. Error bars indicate SEM. Asterisks indicate  $p < 0.05$  when comparing AD patients of the different groups.




**Figure 4** After effects of the SAI protocols on MEP amplitude in Alzheimer's disease patients at baseline (pre) and after 4 weeks (post) of treatment with RTG, RVT, and PLC. Error bars indicate SEM. Asterisks indicate  $p < 0.05$  when comparing AD patients of the different groups.

SESSION interaction ( $F(2,25) = 3.27$ ;  $p = 0.04$ ). *Post hoc* analysis showed that SAI was increased in comparison with baseline at ISI = 0 ms in both RTG and RVT groups (Figure 4).

For the seven AD patients retested after 12 weeks, the analysis performed on the cTBS data did not show any effect for both the SESSION ( $F(2,12) = 12.2$ ;  $p = 0.07$ ), and the TIME main factors ( $F(4,24) = 26.31$ ;  $p = 0.06$ ), and the SESSION  $\times$  TIME interaction ( $F(8,48) = 20.3$ ;  $p = 0.11$ ) (Figure 5a). For the iTBS protocol, there was a significant SESSION ( $F(2,12) = 94.3$ ;  $p = 0.0000001$ ) but not TIME ( $F(4,24) = 13.7$ ;  $p = 0.16$ ) main factor effect. The SESSION  $\times$  TIME interaction ( $F(8,48) = 16.86$ ;  $p = 0.012$ ) was significant. *Post hoc* analysis with Bonferroni correction showed that LTP increased in the RTG group in comparison with baseline evaluation at 4 weeks at all time points ( $p < 0.001$ ) and at 12 weeks at 1–5, 6–10, 11–15 and 16–20 min (all  $p < 0.01$ ) (Figure 5b). For the SAI data, ANOVA revealed significant SESSION ( $F(2,12) = 44.08$ ;  $p = 0.0001$ ), ISI ( $F(3,18) = 3.25$ ;  $p = 0.046$ ) and SESSION  $\times$  ISI interaction ( $F(6,36) = 4.46$ ;  $p = 0.017$ ). *Post hoc* analyses showed that SAI was increased in comparison with baseline at ISI = 0 ms after 4 and 12 weeks of treatment with RTG ( $p$  all  $< 0.05$ ) (Figure 5c).

To control for any change in cognitive functions, neuropsychological scores obtained at baseline were compared with those obtained after 4 weeks of treatment. MMSE, RAVLT, Digit span, Corsi span, and FAB scores at baseline did not differ between the three AD subgroups. The repeated measure ANOVAs performed on the mean scores for each different cognitive test before and after treatment between the three groups of AD patients did not reveal any difference. However paired *t*-test analysis comparing within each group, the different cognitive scores before and after treatment showed that in the RTG group there was an improvement of MMSE scores ( $t(9) = 2.65$ ,  $p < 0.05$ ) and an improvement of the FAB scores ( $t(9) = 2.47$ ,  $p < 0.05$ ). There was no correlation between the individual percentage of change in the MMSE and the FAB scores and the individual



**Figure 5** After effects of the cTBS (a) and iTBS (b) protocols on MEP amplitude in the RTG Alzheimer's disease patients group at baseline (pre), after 4 and 12 weeks of treatment with RTG. (c) After effects of the SAI protocols on MEP amplitude in the RTG Alzheimer's disease patients group at baseline (pre), after 4 and 12 weeks of treatment with RTG. The figure shows the results of the subgroup of patients ( $n = 7$ ) involved in the whole 12 weeks course. Error bars indicate SEM. \* $p < 0.05$  between the 4 weeks and the baseline evaluation. ^ $p < 0.05$  between the 12 weeks and the baseline evaluation.

mean percentage change of MEP amplitude for the iTBS protocol before and after RTG treatment (MMSE:  $r = 0.21$ ;  $p = \text{n.s.}$ ; FAB:  $r = -0.28$ ;  $p = \text{n.s.}$ ). In the RVT group there was an improvement in the MMSE scores ( $t(9) = 2.36$ ,  $p < 0.05$ ) (Supplementary Table S2), whereas no changes of MMSE scores ( $t(9) = -0.07$ ,  $p = 0.94$ ) and FAB scores ( $t(9) = -1.93$ ,  $p = 0.09$ ) were observed in the PLC group. For the cognitive performances of seven AD patients of the RGT group that were followed up at 12-week paired  $t$ -tests showed that there were improvements in the MMSE ( $t(6) = 2.71$ ,  $p < 0.05$ ) and FAB ( $t(6) = 2.57$ ;  $p < 0.05$ ) (Supplementary Table S3).

## DISCUSSION

Our study demonstrates that the dopamine agonist RTG has the ability to restore altered mechanisms of LTP-like cortical plasticity in AD patients.

The increase of LTP-like cortical plasticity is in line with the notion that dopamine is a key player in modulating cortical plasticity in healthy individuals (Kuo *et al*, 2008; Monte-Silva *et al*, 2009) and in patients with Parkinson's disease (Huang *et al*, 2011; Kishore *et al*, 2012). It is indeed well established from animal models that dopamine D1-receptor activation prolongs LTP and LTD (Otmakhova and Lisman, 1998). Conversely, positive and negative effects can both be identified on LTP and LTD when stimulating D2/D3 receptors (Manahan-Vaughan and Kulla, 2003) indicating that dopamine has a non uniform excitatory or inhibitory function. In this view, dopamine can rather act as a strong neuromodulator, with the ability of inducing an effect on cortical activity and plasticity by changing the signal-to-noise ratio (Seamans and Yang, 2004). This focusing effect was confirmed in humans by use of neuroplasticity-inducing stimulation paradigms, namely transcranial direct current stimulation and paired associative stimulation (PAS), which elicit time-dependent LTP- and LTD-like effects on cortical excitability (Kuo *et al*, 2008). An appropriate balance of D1 and D2 activity seems therefore necessary to elicit a focusing effect (Nitsche *et al*, 2009). D2-like receptor activation produces an inverted 'U'-shaped dose-response curve on plasticity, supporting the assumption that modulation of D2-like receptor activity exerts dose-dependent inhibitory or facilitatory effects on neuroplasticity in the human motor cortex (Monte-Silva *et al*, 2009). Crucially, a recent study examined effects of D2 receptor blockade sulpiride on the LTP/LTD-like effects of TBS. Administration of sulpiride blocked the effects of both iTBS and cTBS, showing that D2 receptors are strongly involved in the generation of LTP/LTD-like effects mediated by TBS (Monte-Silva *et al*, 2011). Dopaminergic stimulation is also effective in modulating TBS effects in pathological conditions in which dopaminergic depletion occurs, such as Parkinson's disease. These patients may present with normal LTP-like effects when they assume a full dose of Levodopa (Huang *et al*, 2011). In contrast, there is no LTP-like effect when they are administered half a dose of Levodopa (Huang *et al*, 2011). Moreover, these beneficial boosting effects on plasticity convert to a negative effect as disease advances (Kishore *et al*, 2012).

Our data indicate that a similar boosting effect can be detected also in patients with mild AD. Notably, our AD

patients were treated with a RTG dosage of 4 mg, which is considered as a relatively low dose. Further studies are needed to clarify whether an increased dose of RTG could further increase or even reduce the effects on LTP-like cortical plasticity resembling the inverted 'U'-shaped dose-response curve on plasticity described in healthy controls (Monte-Silva *et al*, 2009). One might argue that the dopaminergic effects on cortical plasticity may be ascribed to an interplay with the cholinergic transmission. On this issue, it was recently demonstrated that, in patients with mild AD, the administration of both Levodopa and RTG is able to increase the levels of central cholinergic activity as assessed by SAI (Martorana *et al*, 2009, 2013). In the current study, the main analyses of SAI measurements did not show any difference between RTG and RVT. This is in line to what expected as both drugs are known to increase SAI activity in AD (Di Lazzaro *et al*, 2002; Marotrana *et al*, 2009; Monte-Silva *et al*, 2011; Di Lazzaro *et al*, 2005). This might suggest that the effects on cortical plasticity induced by RTG administration are specifically driven by dopaminergic stimulation.

A role for the dopaminergic system in AD brains has long been sought and it is still debated. *Post mortem* studies revealed marked loss of dopamine receptors in the temporal and frontal lobes of AD brains (Kemppainen *et al*, 2003; Kumar and Patel, 2007; Martorana *et al*, 2010), regions classically involved in cognitive decline. Interestingly, most of the changes regarding the dopamine receptors, particularly the D2 subtype, were found at rostral and mid-levels of the temporal cortex, indicating that regions classically affected by AD pathology are also sensitive to the loss of D2 receptors (Joyce *et al*, 1998; Goldsmith and Joyce, 1996; Joyce *et al*, 1993; Ryoo and Joyce, 1994). These dopamine D2/D3 receptors may have an important role in the reciprocal activity of large groups of neurons in the high-order association cortical regions, and may promote the cognitive and behavioural impairments observed in AD. Using anti-peptide rabbit antibodies for each of the five dopamine receptors (D1-D5), Kumar and Patel (2007) mapped their distribution in the frontal cortex of *post mortem* AD brains and reported a severe reduction of D1 receptor, D3 receptor, and D4 receptor expression. Conversely, the D2 receptor expression was only moderately reduced. This pattern could explain why the administration of the D2 agonist exerted such a powerful effect on cortical plasticity in our sample of AD patients. Despite defective, the activity of D2 receptors is likely to be potentiated by increasing the levels of D2 agonists. In another series of studies, D2 receptors activity was evaluated *in vivo* by using PET. D2 receptor binding was significantly reduced in the striatum of AD patients, even in the absence of overt extra-pyramidal symptoms (Pizzolato *et al*, 1998). Kemppainen *et al* (2003) reported that D2 receptor binding potentials are reduced in the hippocampus by 30% in AD patients as compared with controls. This reduction was found to be associated with both cognitive (Kemppainen *et al*, 2003) and behavioral abnormalities in AD patients (Tanaka *et al*, 2003).

Dopamine is known to have an important role in higher cognitive functions such as memory, learning, attention, and decision making (Backman *et al*, 2006; Cole *et al*, 2011). Interestingly, animal and human studies have consistently demonstrated that an inverted-U curve exists between



dopaminergic signaling and cognition. Although low or high levels of dopamine impair cognition, middle dosages of dopamine appears necessary for optimal performances (Baunez and Robbins, 1999; Boussaoud and Kermadi, 1997; Cools and D'Esposito, 2011; Glickstein et al, 2005). This phenomenon somewhat resembles what has already been observed with cortical plasticity measurements (Monte-Silva et al, 2009). Recent evidence revealed that, in animal models of AD, dopamine agonists may improve memory function of transgenic-AD mice, inducing decreases in intraneuronal amyloid beta and tau protein deposition, strongly suggesting that dopaminergic agents could be a novel drug for AD (Himeno et al, 2011). However, the involvement of dopamine in the cognitive symptoms in patients with AD has been poorly investigated. Data are limited to indirect evidence that AD patients with higher MMSE scores have an increased capacity of striatal dopamine synthesis (Kemppainen et al, 2003; Tanaka et al, 2003). Notably in the RTG group, despite the clear effects on LTP-like cortical plasticity, we only found a trend towards some improvement for the frontal executive functions. Moreover, we did not find any correlation between the percentage of change of LTP-like cortical plasticity and the relative change of the scores obtained in the different cognitive tests after RTG administration. Clearly, randomized controlled trials are needed to determine whether dopamine agonists may have a clinical relevance in improving cognitive functions in AD patients. These drugs could involve the modulation of the dopaminergic mesocortico-limbic pathway that has important roles in reward, motivation, executive functions, learning, and memory (Kurniawan et al, 2011).

This study has some limitations. The fact that dopaminergic enhancement improved plasticity in these patients is not a definite proof of the causal pathophysiological involvement of the dopaminergic system in Alzheimer's disease. It could be argued that dopamine compensates for reduced cholinergic functions without being causally involved in pathophysiology. On the other hand, dopaminergic activity could be diminished secondarily owing to loss of other transmitters/modulators. Moreover, measures of plasticity were obtained in the motor cortex, which is typically affected by AD pathology at late disease stages. Future studies should focus on the association cortex (eg, temporo-parietal junction or dorsolateral prefrontal cortex) using different neurophysiological methods such as TMS in combination with EEG (Veniero et al, 2013; Rajji et al, 2013).

In conclusion, we demonstrated that the administration of dopamine agonists such as RTG is effective in modulating the altered mechanisms of cortical plasticity that characterize AD patients. This could open new perspectives for therapeutic strategies to contrast cortical dysfunction in AD.

## FUNDING AND DISCLOSURE

The authors declare no conflict of interest.

## ACKNOWLEDGEMENTS

This work was supported by grants of the Italian Ministry of Health to GK (239/GR-2009-1591859) and to MB (47/RF-2010-2311484).

## REFERENCES

- Appollonio I, Leone M, Isella V, Piamarta F, Consoli T, Villa ML et al (2005). The Frontal Assessment Battery (FAB): normative values in an Italian population sample. *Neurol Sci* 26: 108–116.
- Backman L, Nyberg L, Lindenberger U, Li SC, Farde L (2006). The correlative triad among aging, dopamine, and cognition: current status and future prospects. *Neurosci Biobehav Rev* 30: 791–807.
- Baunez C, Robbins TW (1999). Effects of dopamine depletion of the dorsal striatum and further interaction with subthalamic nucleus lesions in an attentional task in the rat. *Neuroscience* 92: 1343–1356.
- Boekhoorn K, Terwel D, Biemans B, Borghgraef P, Wiegert O, Ramakers GJ et al (2006). Improved long-term potentiation and memory in young tau-P301L transgenic mice before onset of hyperphosphorylation and tauopathy. *J Neurosci* 26: 3514–3523.
- Boussaoud D, Kermadi I (1997). The primate striatum: neuronal activity in relation to spatial attention versus motor preparation. *Eur J Neurosci* 9: 2152–2168.
- Bozzali M, Padovani A, Caltagirone C, Borroni B (2011). Regional grey matter loss and brain disconnection across Alzheimer disease evolution. *Curr Med Chem* 18: 2452–2458.
- Carlesimo GA, Caltagirone C, Gainotti G (1996). The Mental Deterioration Battery: normative data, diagnostic reliability and qualitative analyses of cognitive impairment. The Group for the Standardization of the Mental Deterioration Battery. *Eur Neurol* 36: 378–384.
- Cole DM, Beckmann CF, Searle GE, Plisson C, Tziortzi AC, Nichols TE et al (2011). Orbitofrontal connectivity with resting-state networks is associated with midbrain dopamine D3 receptor availability. *Cereb Cortex* 22: 2784–2793.
- Cools R, D'Esposito M (2011). Inverted-U-shaped dopamine actions on human working memory and cognitive control. *Biol Psychiatry* 69: 113–125.
- Di Lazzaro V, Oliviero A, Pilato F, Saturno E, Dileone M, Marra C et al (2005). Neurophysiological predictors of long term response to AChE inhibitors in AD patients. *J Neurol Neurosurg Psychiatry* 76: 1064–1069.
- Di Lazzaro V, Oliviero A, Tonali PA, Marra C, Daniele A, Profice P et al (2002). Non invasive in vivo assessment of cholinergic cortical circuits in AD using transcranial magnetic stimulation. *Neurology* 59: 392–397.
- Glickstein SB, Desteno DA, Hof PR, Schmauss C (2005). Mice lacking dopamine D2 and D3 receptors exhibit differential activation of prefrontal cortical neurons during tasks requiring attention. *Cereb Cortex* 15: 1016–1024.
- Goldsmith SK, Joyce JN (1996). Dopamine D2 receptors are organized in bands in normal human temporal cortex. *Neuroscience* 74: 435–451.
- Himeno E, Ohyagi Y, Ma L, Nakamura N, Miyoshi K, Sakae N et al (2011). Apomorphine treatment in Alzheimer mice promoting amyloid- $\beta$  degradation. *Ann Neurol* 69: 248–256.
- Huang YZ, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC (2005). Theta burst stimulation of the human motor cortex. *Neuron* 45: 201–206.
- Huang YZ, Rothwell JC, Lu CS, Chuang WL, Chen RS (2011). Abnormal bidirectional plasticity-like effects in Parkinson's disease. *Brain* 134: 2312–2320.
- Joyce JN, Kaeger C, Ryoo H, Goldsmith S (1993). Dopamine D2 receptors in the hippocampus and amygdala in Alzheimer's disease. *Neurosci Lett* 154: 171–174.
- Joyce JN, Myers AJ, Gurevich E (1998). Dopamine D2 receptor bands in normal human temporal cortex are absent in Alzheimer's disease. *Brain Res* 784: 7–17.
- Kemppainen N, Laine M, Laakso MP, Kaasinen V, Någren K, Vahlberg T et al (2003). Hippocampal dopamine D2 receptors correlate with memory functions in Alzheimer's disease. *Eur J Neurosci* 18: 149–154.

- Kishore A, Popa T, Velayudhan B, Joseph T, Balachandran A, Meunier S (2012). Acute dopamine boost has a negative effect on plasticity of the primary motor cortex in advanced Parkinson's disease. *Brain* 135: 2074–2088.
- Klyubin I, Betts V, Welzel AT, Blennow K, Zetterberg H, Wallin A et al (2008). Amyloid beta protein dimer-containing human CSF disrupts synaptic plasticity: Prevention by systemic passive immunization. *J Neurosci* 28: 4231–4237.
- Koch G, Esposito Z, Kusayanagi H, Monteleone F, Codecà C, Di Lorenzo F et al (2011). CSF Tau levels influence cortical plasticity in Alzheimer's disease patients. *J Alzheimers Dis* 26: 181–186.
- Koch G, Di Lorenzo F, Bonni S, Ponzo V, Caltagirone C, Martorana A (2012). Impaired LTP- but not LTD-like cortical plasticity in Alzheimer's disease patients. *J Alzheimers Dis* 31: 593–599.
- Kumar U, Patel SC (2007). Immunohistochemical localization of dopamine receptor subtypes (D1R–D5R) in Alzheimer's disease brain. *Brain Res* 1131: 187–196.
- Kuo MF, Paulus W, Nitsche MA (2008). Boosting focally-induced brain plasticity by dopamine. *Cereb Cortex* 18: 648–651.
- Kurniawan IT, Guitart-Masip M, Dolan RJ (2011). Dopamine and effort-based decision making. *Front Neurosci* 21: 81.
- Li S, Hong S, Shepardson NE, Walsh DM, Shankar GM, Selkoe D (2009). Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 62: 788–801.
- Manahan-Vaughan D, Kulla A (2003). Regulation of depotentiation and long-term potentiation in the dentate gyrus of freely moving rats by dopamine D2-like receptors. *Cereb Cortex* 13: 123–135.
- Martorana A, Mori F, Esposito Z, Kusayanagi H, Monteleone F, Codecà C et al (2009). Dopamine modulates cholinergic cortical excitability in Alzheimer's disease patients. *Neuropsychopharmacology* 34: 2323–2328.
- Martorana A, Esposito Z, Koch G (2010). Beyond the cholinergic hypothesis: do current drugs work in Alzheimer's disease? *CNS Neurosci Ther* 16: 235–245.
- Martorana A, Di Lorenzo F, Esposito Z, Lo Giudice T, Bernardi G, Caltagirone C et al (2013). Dopamine D<sub>2</sub>-agonist rotigotine effects on cortical excitability and central cholinergic transmission in Alzheimer's disease patients. *Neuropharmacology* 64: 108–113.
- Monte-Silva K, Kuo MF, Thirugnanasambandam N, Liebetanz D, Paulus W, Nitsche MA (2009). Dose-dependent inverted U-shaped effect of dopamine (D2-like) receptor activation on focal and nonfocal plasticity in humans. *J Neurosci* 29: 6124–6131.
- Monte-Silva K, Ruge D, Teo JT, Paulus W, Rothwell JC, Nitsche MA (2011). D2 receptor block abolishes  $\theta$  burst stimulation-induced neuroplasticity in the human motor cortex. *Neuropsychopharmacology* 36: 2097–2102.
- Nitsche MA, Kuo MF, Grosch J, Bergner C, Monte-Silva K, Paulus W (2009). D1-receptor impact on neuroplasticity in humans. *J Neurosci* 29: 2648–2653.
- Otmakhova NA, Lisman JE (1998). D1/D5 dopamine receptors inhibit depotentiation at CA1 synapses via cAMP-dependent mechanism. *J Neurosci* 18: 1270–1279.
- Palop JJ, Mucke L (2010). Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat Neurosci* 13: 812–818.
- Paspalas CD, Goldman-Rakic PS (2005). Presynaptic D1 dopamine receptors in primate prefrontal cortex: target-specific expression in the glutamatergic synapse. *J Neurosci* 25: 1260–1267.
- Pizzolato G, Chierichetti F, Fabbri M, Cagnin A, Dam M, Ferlin G et al (1998). Reduced striatal dopamine receptors in Alzheimer's disease: single photon emission tomography study with the D2 tracer [123I]-IBZM. *Neurology* 47: 1065–1068.
- Rajji TK, Sun Y, Zomorodi-Moghaddam R, Farzan F, Blumberger DM, Mulsant BH et al (2013). PAS-induced potentiation of cortical-evoked activity in the dorsolateral prefrontal cortex. *Neuropsychopharmacol* 38: 2545–2552.
- Rossini PM, Barker AT, Berardelli A, Caramia MD, Caruso G, Cracco RQ et al (1994). Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalogr Clin Neurophysiol* 91: 79–92.
- Ryoo HL, Joyce JN (1994). Loss of dopamine D2 receptors varies along the rostrocaudal axis of the hippocampal complex in Alzheimer's disease. *J Comp Neurol* 348: 94–110.
- Sailer A, Molnar GF, Paradiso G, Gunraj CA, Lang AE, Chen R (2003). Short and long latency afferent inhibition in Parkinson's disease. *Brain* 126: 1883–1894.
- Seamans JK, Yang CR (2004). The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog Neurobiol* 74: 1–58.
- Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I et al (2008). Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 14: 837–842.
- Suvà D, Favre I, Kraftsik R, Esteban M, Lóbrinus A, Miklóssy J (1999). Primary motor cortex involvement in Alzheimer disease. *J Neuropathol Exp Neurol* 58: 1125–1134.
- Tanaka Y, Meguro K, Yamaguchi S, Ishii H, Watanuki S, Funaki Y et al (2003). Decreased striatal D2 receptor density associated with severe behavioral abnormality in Alzheimer's disease. *Ann Nucl Med* 17: 567–573.
- Tritsch NX, Sabatini BL (2012). Dopaminergic modulation of synaptic transmission in cortex and striatum. *Neuron* 76: 33–50.
- Varma AR, Snowden JS, Lloyd JJ, Talbot PR, Mann DM, Neary D. (1999). Evaluation of the NINCDS-ADRDA criteria in the differentiation of Alzheimer's disease and frontotemporal dementia. *J Neurol Neurosurg Psychiatry* 66: 184–188.
- Veniero D, Ponzo V, Koch G (2013). Paired associative stimulation enforces the communication between interconnected areas. *J Neurosci* 33: 13773–13783.
- Zhang L, Doyon WM, Clark JJ, Phillips PE, Dani JA (2009). Controls of tonic and phasic dopamine transmission in the dorsal and ventral striatum. *Mol Pharmacol* 76: 396–404.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)