

ERRATA

Capsular Polysaccharide of *Clostridium perfringens* Hobbs 9

ROBERT CHERNIAK* AND HAROLD M. FREDERICK

Department of Chemistry, Georgia State University, Atlanta Georgia 30303

Volume 15, no. 3, p. 767: Because of a printing error, Table 2 is incorrect as printed. The table should read as follows:

TABLE 2. Purification of the capsular polysaccharide of *C. perfringens* Hobbs 9

| Step | Fraction isolated | Fraction treated (g) | Treatment | Amt re-covered (% by wt) | Composition (% by wt) of isolated fraction | | | |
|------|----------------------|----------------------|---|--------------------------|--|------------------|-----------|------------|
| | | | | | Carbohydrate | Protein | Phosphate | Hexosamine |
| 1 | Crude polysaccharide | 62 (cells) | 0.85% NaCl | 9.0 | 16 | 21 | 24 | 1.9 |
| 2 | Supernatant 0 | 5.6 | (NH ₄) ₂ SO ₄ | 20.4 | 17 | 29 | 21 | 5.2 |
| 3 | Supernatant 0-RP | 1.0 | RNase ^a -Papain | 41 | 21 | 10 | 15 | 8.4 |
| 4 | Supernatant 0-RPD | 0.35 | DEAE-Sephadex A25 (15 cm) | 43 | 24 | 3.8 | 3.2 | 14.2 |
| 5 | Supernatant 0-RPDA | 0.042 | Agarose 1.5M | 83 | 31 | 3.1 | 2.8 | 14.7 |
| 6 | Supernatant 0-RPD2 | 0.116 | DEAE-Sephadex A25 (30 cm) | 45 | | | | |
| | Peak IA | | | | ND ^b | 6.0 ^c | 7.6 | 17.6 |
| | Peak IB | | | | ND | 5.3 | 7.5 | 18.3 |
| | Peak II | | | | ND | 5.6 | 1.6 | 23.2 |

^a Ribonuclease.

^b ND, Not done.

^c Calculated by amino acid composition.

Penicillin Sensitivity and Serum Resistance Are Attributes of Strains of *Neisseria gonorrhoeae* Causing Disseminated Gonococcal Infection

B. I. EISENSTEIN,* T. J. LEE, AND P. F. SPARLING

Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, North Carolina 27514

Volume 15, no. 3, p. 835, 2nd column, line 39: Change "0.25" to read "0.025."