



Published in final edited form as:

J Mol Biol. 2015 March 13; 427(5): 1061–1074. doi:10.1016/j.jmb.2014.04.021.

In vivo roles of BamA, BamB and BamD in the biogenesis of BamA, a core protein of the β -barrel assembly machine of *Escherichia coli*

Rajeev Misra^{*}, Ryan Stikeleather, and Rebecca Gabriele

School of Life Sciences, Arizona State University, Tempe, Arizona 85287, USA

Abstract

Assembly of the β -barrel outer membrane proteins (OMPs) is an essential cellular process in Gram negative bacteria and in the mitochondria and chloroplasts of eukaryotes—two organelles of bacterial origin. Central to this process is the conserved β -barrel OMP that belongs to the Omp85 superfamily. In *Escherichia coli*, BamA is the core β -barrel OMP, and together with four outer membrane lipoproteins, BamBCDE, constitute the β -barrel assembly machine (BAM). In this paper, we investigated the roles of BamD, an essential lipoprotein, and BamB in BamA biogenesis. Depletion of BamD caused impairment in BamA biogenesis and cessation of cell growth. These defects of BamD depletion were partly reversed by single amino acid substitutions mapping within the β -barrel domain of BamA. However, in the absence of BamB, the positive effects of the β -barrel substitutions on BamA biogenesis under BamD depletion conditions were nullified. By employing a BamA protein bearing one such substitution, F494L, it was demonstrated that the mutant BamA protein could not only assemble without BamD, but it could also facilitate the assembly of wild-type BamA expressed *in trans*. Based on these data, we propose a model in which the Bam lipoproteins, which are localized to the outer membrane by the BAM-independent Lol pathway, aid in the creation of new BAM complexes by serving as outer membrane receptors and folding factors for nascent BamA molecules. The newly assembled BAM holocomplex then catalyzes the assembly of substrate OMPs and BamA. These *in vivo* findings are corroborated by recently published *in vitro* data.

Keywords

β -barrel outer membrane protein assembly; BAM complex; genetics

© 2014 Elsevier Ltd. All rights reserved.

^{*}Corresponding author. Mailing address: School of Life Sciences, Arizona State University, 427 East Tyler Mall, Tempe, AZ 85287-4501, U.S.A. Phone: (480) 965-3320 Fax: (480) 965-6899 rajeev.misra@asu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

The two major classes of Gram negative outer membrane proteins (OMPs) are lipoproteins and β -barrels [1]. The localization of lipoprotein (Lol) pathway targets mature, N-terminal lipid-modified, proteins from the periplasmic side of the inner membrane to the periplasmic side of the outer membrane [2]. The Lol pathway is essential because it is involved in the biogenesis of cellular machineries that catalyze the assembly of β -barrel OMPs and lipopolysaccharide (LPS) in the outer membrane [36].

The *Escherichia coli* β -barrel assembly machine (BAM) contains four outer membrane lipoproteins, BamBCDE, and a β -barrel OMP, BamA (for recent reviews, see [5-7]; BamD and BamA are the two indispensable components of the BAM complex [8-11]. BamA belongs to the Omp85 superfamily of proteins that mediate protein transport/assembly functions [12-14]. In contrast, the BamBCDE homologs are found only in Gram negative bacteria [15,16].

Both *in vivo* and *in vitro* studies have shown that the BAM holocomplex can exist as two major subcomplexes, BamAB and BamACDE [11,17,18]. These two subcomplexes carry out partially overlapping functions [15,19]. Of the three non-essential lipoproteins, the absence of BamB produces the most significant OMP biogenesis defect [17,20-22]. The defect of *bamB* is further exacerbated when BamC or BamE is also simultaneously absent [17], with the *bamB bamE* mutant displaying the most drastic growth and OMP biogenesis defects [15].

The high-resolution structures of all Bam proteins have been solved and they have begun to provide mechanistic insights into the BAM activity. All five proteins contain motifs known to facilitate protein-protein interactions. The N-terminal soluble domain of BamA is exposed to the periplasm, while its C-terminal domain is embedded in the outer membrane [12-14]. The soluble N-terminal end of BamA folds into five polypeptide-transport-associated (POTRA) domains [23] whose structure revealed possible interactions with substrate polypeptides via β -augmentation [24-26]. Mutational and pull-down assays have also revealed that the POTRA domains of BamA are critical for interactions with the Bam lipoproteins [24] and a periplasmic chaperone, SurA [27,28]. Thus, the POTRA domains appear to serve as the initial staging ground for substrate interaction and folding. The C-terminal β -barrel domain of BamA is thought to catalyze the final steps of OMP assembly [29]. An important feature of this domain is a large loop, L6, which contains the 'VRGF/Y' motif conserved in all members of the Omp85 superfamily [13,14,30]. The functional importance of the BamA L6 region and its conserved VRGF motif was first determined through the genetic analysis [15] and subsequently by site-directed mutagenesis [29,31]. Exactly how the L6 region participates in the β -barrel OMP assembly remains to be elucidated.

Three-dimensional structures of BamB [32-34], BamD [35-37], BamCD [38], and BamE [35,39] have been solved. However, the roles of these lipoproteins in OMP assembly remained largely speculative. It has been hypothesized that BamB may channel substrate OMPs to BamA [32], but this has not been supported experimentally [34]. The structural

and mutational data suggest that the N-terminus of BamD might interact with substrate OMPs [11,35]. Interestingly, the same region of BamD is also shown to interact with BamC [38], thus prompting a suggestion that BamC may play a role in regulating BamD activity. BamE has been proposed to interact with the cell envelope components to stabilize and/or regulate the activity of the BAM complex. A recent study suggested a role for BamE in modulating BamA's conformation during its functional cycle [40].

In a recent *in vitro* study, Hagan *et al.* [19] showed that BamB and BamD can bind to unfolded BamA and facilitate its assembly and insertion into proteoliposomes. Their study also showed that BamCDE-mediated assembly of unfolded BamA does not require preassembled BamA in proteoliposomes, although the presence of BamA in the BamCDE complex greatly enhanced assembly of the unfolded BamA substrate. These results reinforce a previous suggestion that BamA aids in its own assembly [24,27,31]. It is unclear whether the nascent OMPs enter the BamA channel [41] or stay outside the BamA barrel at the protein-lipid interface. An attractive model outlined by Kim *et al.* [42] and Noinaj *et al.* [29] envisions that the genesis of a substrate β -barrel begins with its β -strands aligning with the first and last β -strands of the transiently open BamA barrel, and culminating in the substrate β -sheet budding off into the lipid phase of the membrane to form an enclosed β -barrel.

In this paper, we investigated the *in vivo* roles of the BamD and BamB lipoproteins in BamA biogenesis, and determined the effects of BamA β -barrel substitutions that were previously shown to improve BamA biogenesis in a *bamB bamE* double mutant background [15]. Because BamA itself has a β -barrel domain, it is logical to consider that the pre-assembled BAM complex would also catalyze the assembly of nascent BamA molecules. Consequently, a defective BAM complex should affect the assembly of all β -barrel OMPs, including BamA [27,31,43]. We show here that BamA misfolds when BamD is depleted from the cell and this defect is reversed by the β -barrel substitutions in BamA, but only in the presence of BamB. Moreover, the mutant BamA, which assembles without BamD, in turn facilitates the assembly of wild-type BamA expressed *in trans* under acute BamD depletion conditions. These data reveal that BamD and BamB play crucial roles in BamA folding and biogenesis *in vivo* and give physiological credence to the recent *in vitro* study by Hagan *et al.* [19].

Results

***bamA* suppressor alleles allow growth under BamD depletion conditions**

We have previously reported the isolation of six novel *bamA* suppressor alleles from a strain simultaneously lacking the two non-essential lipoproteins of the BAM complex, BamB and BamE, [15]. The double mutant, expressing wild-type BamA (BamA_{WT}), failed to grow on rich medium, grew poorly on minimal medium, and displayed a severe OMP biogenesis defect [15]. Moreover, the data revealed that without BamB and BamE, BamA could not fold correctly and was present in reduced levels presumably due to proteolysis of misfolded BamA. These data suggested a role for the Bam lipoproteins in BamA biogenesis. The presence of the suppressor mutations in *bamA* in a *bamB bamE* background improved bacterial growth, OMP biogenesis, and restored BamA levels and folding to various degrees [15]. Since all six suppressor alterations map in the BamA β -barrel domain, it stands to

reason that BamB and BamE influence folding of the BamA β -barrel domain, and thus the OMP biogenesis defect stems from defects caused by misfolded BamA and the loss of BamB and BamE activities.

BamB and BamE form two BAM subcomplexes, BamAB and BamACDE [11,17,18]. Whereas BamB interacts with BamA directly [8,22], BamE does so indirectly via BamD [17]. Since BamD is the essential lipoprotein and directly interacts with BamA [11], we reasoned that it is likely to play a more critical role in BamA biogenesis than the non-essential BamB or BamE protein individually. Therefore, we investigated BamA biogenesis under BamD depletion conditions and tested the effects of the six *bamA* suppressor alleles on bacterial growth, BamA levels and folding, and general OMP biogenesis.

The parental *E. coli* strain used in this study had its chromosomal *bamD* deleted and expressed *bamD* from an arabinose-inducible allele inserted at the lambda attachment site [11]. Consequently, growth of the parental strain is dependent on arabinose but is inhibited by glucose, which lowers the level of cAMP required for the activity of the *araBAD* promoter [11]. However, despite this arabinose dependence, the parental strain can grow normally for several generations in a minimal medium containing glycerol, where BamD levels are depleted at a much slower rate than that in the minimal medium supplemented with glucose. Therefore, bacterial cultures were first grown overnight in glycerol minimal medium at 37°C to initiate depletion of BamD, followed by growth next day in glucose medium to completely deplete BamD. The cell densities of all overnight grown cultures were nearly identical. Next day, cultures were diluted 1:100 in glucose minimal medium and grown at 37°C. After 3.5 hours, when OD₆₀₀ reached around 1.0 (Fig. 1), cultures were diluted 1:50 in glucose minimal medium and incubated at 37°C for 4 hours. At the end of this incubation period, cultures were diluted 1:50 one final time in glucose minimal medium and grown for 4 hours. After the third (final) dilution, the two cultures that failed to grow expressed either BamA_{WT} or BamA_{E780D} (Fig. 1). In contrast, the remaining cultures expressing five different BamA suppressor variants grew at rates very similar to that seen after the second dilution. Interestingly, BamA_{E780D} was also the weakest among the six BamA suppressors in terms of improving the growth of the *bamB bamE* mutant [15]. Growth after the fourth dilution was sporadic with some cultures occasionally growing weakly (data not shown). It is worth mentioning that BamD levels under these growth conditions became undetectable at the end of the first dilution in glucose minimal medium (e.g. see Fig. 3). These data show that the expression of the BamA suppressor variants, except that bearing an E780D substitution, afforded bacterial growth even under acute BamD depletion conditions. Note that BamA residue numbers are relative to the mature BamA sequence.

Suppressors stabilize BamA under BamD depletion conditions

BamD has been known to stabilize the BAM complex [11], but its effect on BamA biogenesis is not known. BamA levels were examined by Western blot analysis from cultures grown overnight in glycerol minimal medium and after dilutions in glucose minimal medium. Figure 2 shows data from BamA_{WT} and three BamA mutants; the data from the entire set of BamA mutants are presented in Fig. S1. As mentioned earlier, BamD was only

partially depleted when cultures were grown overnight in glycerol minimal medium, but depleted rapidly during growth in glucose minimal medium (Figs. 3 and 5). BamA levels were very similar in all cultures grown overnight and after the first dilution (Fig. 2), which is consistent with their similar overnight cell densities and first dilution growth rates. However, after the second dilution, BamA levels in cultures expressing BamA_{WT} and BamA_{E780D} dropped more than two fold compared to those in cultures expressing BamA_{F474L} and BamA_{G649A} (Figs. 2 and S1). The drop in BamA levels in these two strains coincided with their growth decline during the same period (Fig. 1). It is important to mention that the decrease in BamA_{WT} and BamA_{E780D} levels (Figs. 2 and S1) is an indication of significant defects in BamA assembly since envelope stress should increase under BamD depletion conditions, elevating *bamA* expression by the σ^E -regulated *bamA* promoter [27,44,45].

A prominent BamA antibody-reacting protein band, migrating at the 35 kDa position, consistently appeared in our work (Figs. 2 and S1). Although the 35 kDa band appeared in the overnight cultures, its relative intensity increased substantially after the first dilution, when *bamD* expression was inhibited by glucose, and stayed high after the second and third dilutions during continual BamD depletion (Figs. 2 and S1). Interestingly, a similar 35 kDa BamA fragment was also observed by Ricci *et al.* [43] in a strain expressing a mutant BamA protein defective in its ability to interact with BamD, but not in a strain expressing BamA_{WT}. In that study and ours the 35 kDa fragment appears to be a proteolysis product of BamA resulting from the loss of BamA-BamD interactions due to a defective BamA protein [43] or BamD depletion (this work). Consistent with Ricci *et al.* [43], we found that the 35 kDa fragment is derived from an N-terminal POTRA domain of BamA, since it interacts with the antibodies raised against a N-terminal derived mature BamA peptide (A21-Q35; [24]) (data not shown). For this reason, the 35 kDa is labeled as BamA(N) in Figs. 2 and S1.

In summary, these data showed that with the exception of BamA_{WT} and BamA_{E780D}, the remaining five BamA variants assumed conformations that resisted *in vivo* destabilization under BamD depletion conditions. Since these five BamA variants also improved bacterial growth without BamD, they are likely to restore some OMP biogenesis function in cells. It is important to mention that despite these positive effects of the BamA suppressor variants under BamD depletion conditions, cell growth deteriorated dramatically after the fourth dilution and we were unable to obtain stable *bamD* transductants in these mutant *bamA* backgrounds. Therefore, while the BamA suppressor variants lower the dependence on BamD, they do not completely eliminate it.

Suppressor alteration renders BamA folding independent of BamD, but not OMP biogenesis

Next we examined the effects of BamA suppressors on BamA folding and OMP levels under BamD depletion conditions. Folded BamA can be separated from unfolded, heat-denatured, BamA on a SDS-polyacrylamide gel since the folded more compact form of BamA migrates faster than the unfolded more extended form [15,24,31,46]. We have shown previously that with the exception of F474L and E780D, the other four BamA substitutions altered the heat-modifiability property of BamA [15], thus making it difficult to assess the

folding status of BamA in these four mutants. Also, as observed in the previous section, the strain expressing BamA_{E780D} behaved very similar to that expressing BamA_{WT} under BamD depletion conditions. Given these two observations, we examined BamA folding and OMP biogenesis status from cultures expressing BamA_{F474L} and BamA_{WT}. Envelopes were extracted from cultures expressing BamD (arabinose added) or depleted of it (glucose added) (Fig. 3).

BamA's folding status was determined from cultures grown after first and second dilutions. In cultures expressing BamD (plus arabinose; Fig. 3), both BamA_{WT} and BamA_{F474L} were present at the same levels and displayed an identical heat-modifiability behavior (Fig. 3a). Likewise, levels and folding behaviors of the two proteins were very similar after the first dilution in glucose medium (minus arabinose; Fig. 3). However, after the second dilution in which BamD was further depleted, nearly all of the BamA_{WT} protein from unheated samples migrated at the unfolded/denatured position (Fig. 3a), indicating severe BamA misfolding. Strikingly, under the identical BamD depletion condition, a significant amount of the BamA_{F474L} protein maintained the normal heat-modifiability property (Fig. 3a), indicating normal folding status. Thus, the F474L alteration in the BamA β -barrel domain renders BamA folding independent of BamD, just as it did in the absence of BamB and BamE [15]. We surmise from these results that BamD directly or indirectly, via the assembled BAM complex, assists folding of the BamA β -barrel domain, and that the growth defects under BamD depletion conditions are due in part to misfolding and thus loss of BamA function.

We also analyzed OMP levels from cultures expressing BamD or depleted of it (Fig. 3b-d). When arabinose was added to the growth medium, all cultures showed identical protein profiles from purified envelopes (Fig. 3b) and expressed readily detectable amounts of BamD (Fig. 3c). In contrast, significantly reduced levels of OMPs were present in envelopes isolated from cultures grown in glucose minimal medium after the first dilution (Fig. 3b and d), while BamD could not be detected from purified envelopes (Fig. 3c). Interestingly, cultures expressing either BamA_{WT} or BamA_{F474L} displayed very similar OMP profiles after the first dilution (Fig. 3b and d). This is consistent with the observation that all cultures exhibited almost identical growth rates after the first dilution in glucose minimal medium, thus presumably similar OMP biogenesis status (Fig. 1). However, there was an appreciable difference in the OMP profile from cultures isolated after the second dilution in glucose minimal medium (Fig. 3b and d) where cells expressing BamA_{F474L} grew better than those expressing BamA_{WT} (Fig. 1). The most noticeable difference was in the levels of OmpA and OmpC, which were not significantly affected in a culture expressing BamA_{F474L} compared to that expressing BamA_{WT} (Fig. 3b and d). In contrast, OmpF levels were severely reduced in both cultures (Fig. 3b and d).

These data indicated that even though the F474L alteration can render BamA folding and biogenesis independent of BamD, these changes do not make the activity of the BAM complex fully independent of BamD. In other words, in addition to assisting BamA folding, BamD must perform some indispensable function in OMP biogenesis.

BamA_{F474L} assists assembly of BamA_{WT} under BamD depletion conditions

BamD-independent assembly of BamA_{F474L} allowed us to ask a crucial question of whether the assembled BamA_{F474L} molecules are functionally active by examining their abilities to assemble BamA_{WT}. To answer this, we built two partially diploid strains expressing either BamA_{WT} or BamA_{F474L} from the chromosome and a His-tagged copy of wild-type BamA (BamA_{WT-His}) from a low copy number plasmid, pZS21 [24,27]. A null *recA* allele was introduced to prevent recombination between the wild-type and mutant alleles of *bamA*. Use of His-Probe allowed for the detection of only the plasmid-expressed BamA_{WT-His} and not the chromosomally expressed BamA.

Under BamD depletion conditions, the strain expressing only the wild-type versions of tagged and untagged BamA proteins are expected to stop growing after the second dilution, as observed when using the haploid strain (Fig. 1). Similarly, BamA_{WT} folding is expected to become acutely defective after the second dilution, as observed in Fig. 3. In contrast, the strain expressing BamA_{F474L} from the chromosome and BamA_{WT-His} from the plasmid may grow past the second dilution and correctly fold BamA_{WT-His}, if folded BamA_{F474L} is functionally active. The data from these experiments are presented in Fig. 4.

As expected, the partially diploid strain expressing BamA_{F474L} from the chromosome grew significantly better than that expressing BamA_{WT} after the second and third dilutions in glucose minimal medium to deplete BamD (Fig. 4a). However, growth rates after these dilutions were noticeably lower than those observed with the haploid strain (compare Figs. 1 and 4a), presumably due to the extra burden imposed by the presence of a plasmid and a null *recA* allele in the partial diploid strains. The strain expressing tagged and untagged versions of wild-type BamA failed to grow after the third dilution (Fig. 4a).

We analyzed the levels of major OMPs (Fig. 4b and c) from samples collected during the growth experiments shown in Fig. 4a. The OMP profiles of the two partially diploid strains from cultures obtained after overnight growth and after the first dilution were almost identical. However, after the second dilution, although the major OMP levels deteriorated further, their levels were higher in the strain expressing BamA_{F474L} than that expressing BamA_{WT} (Fig. 4b and c). We then determined BamA_{WT-His} folding from purified envelopes (Fig. 4d-f). In the strain expressing the chromosomally-encoded BamA_{WT} protein, folding of the plasmid-born BamA_{WT-His} deteriorated rapidly after the second dilution, as no folded species was detected (Fig. 4d and f). However, in the strain expressing the chromosomally-encoded BamA_{F474L} protein, almost 50% of the BamA_{WT-His} protein folded correctly after the second dilution (Fig. 4e and f). Even after the third dilution, between 30 and 40% of the BamA_{WT-His} protein folded correctly (Fig. 4e and f). Therefore, while the chromosomally-encoded BamA_{WT} failed to fold and function correctly in the absence of BamD (Figs. 3 and S2), the BamA_{F474L} protein not only folded correctly without BamD (Figs. 3 and S2), it maintained a significant functional activity, as demonstrated by its ability to catalyze folding of the substrate BamA_{WT-His} protein, whose folding is normally dependent on BamD (Fig. 4d and f). These data showed good correlation between the presence of correctly folded BamA species (Fig. 4d-f) and the major OMP levels (Fig. 4b and c). Therefore, BamA_{F474L}

could catalyze the assembly of BamA and major OMPs under acute BamD depletion conditions, albeit not as efficiently as in the presence of BamD.

Role of BamB in BamA biogenesis

In the absence of BamB, BamA can fold efficiently, but the absence of both BamB and BamE results in BamA misfolding and a conditional lethal phenotype [15]. BamB interacts with BamA directly and the stability of this subcomplex remains unaffected under BamD depletion conditions [11]. Even though BamA_{F474L} folds independently of BamD (Fig. 3a), it may still require BamB for correct folding, since BamB and BamD subcomplexes are known to play an overlapping role in BamA_{WT} folding [15]. To test this possibility, we introduced a *bamB*::Km^r allele in *bamD* depletion strains expressing either wild-type or the various suppressor *bamA* alleles. The *bamB* null allele was introduced by P1 transduction in the presence of arabinose to induce *bamD* expression from the chromosome. All strains yielded stable *bamB*::Km^r transductants when *bamD* expression was induced by arabinose. However, none of the transductants were able to form single colonies on a medium lacking arabinose at 30°C or 37°C. These growth defects were much more severe than those observed in a *bamB*⁺ background. A representative growth pattern of strains expressing BamA_{WT} and three BamA mutants is shown in Fig. S3.

We then carried out growth experiments in liquid cultures to assess the optimum time for sample collection and protein analysis. As described earlier, cultures were grown overnight in glycerol minimal medium. Next day, overnight cultures were diluted 1:100 in glucose minimal medium to further deplete BamD levels. However, *bamB* had a dramatic impact on glucose-grown cells, with the *bamB* culture reaching only half the cell density compared to the *bamB*⁺ culture after 3.5 hours of growth (Fig. 5a). At the end of this growth period, cultures were diluted 1:50 in fresh media and grown for another 3.5 hours. Growth of the *bamB* strain in glucose minimal medium ceased at the onset of the second dilution, whilst others grew somewhat slower than during the first dilution period (Fig. 5a). Cultures grown in the glycerol medium supplemented with arabinose displayed similar growth rates, regardless of the presence or absence of *bamB*, but these rates were lower than those observed after the first dilution in medium containing high energy-yielding glucose (data not shown).

The presence of BamB is critical for BamA_{F474L} folding and OMP biogenesis under BamD depletion conditions

The functional consequence of *bamB* under BamD depletion conditions was determined by analyzing the levels of OmpA, OmpC and OmpF, all of which are BAM substrates [27]. Envelopes isolated from cultures grown above (Fig. 5a) were analyzed on SDS(urea)-polyacrylamide gels and protein bands were visualized after Coomassie blue staining (Fig. 5b). When *bamD* expression was fully induced by arabinose, the effect of *bamB* on OMPs were small albeit noticeable (Fig. 5b). However, the effects of *bamB* on OMP levels were salient in cultures grown overnight in glycerol minimal medium without arabinose, and became more pronounced when BamD levels were further depleted after the first dilution in glucose minimal medium (Fig. 5b).

BamA_{F474L} folding status was assessed from envelopes purified from cultures grown overnight and at the end of the first dilution (Fig. 5c and d). When *bamD* expression was induced by arabinose, BamA_{F474L} displayed the typical heat-modifiability pattern, regardless of the presence or absence of BamB or the age of the culture (Fig. 5c and d). However, under BamD-depletion conditions, BamA_{F474L} folding differed depending on the presence or absence of BamB and the culture age. Note, however, that BamD was readily detected in cultures grown overnight in glycerol minimal medium, though its level was significantly lower than that in cultures supplemented with arabinose (Fig. 5c). BamA_{F474L} in *bamB* and *bamB*⁺ backgrounds from glycerol-grown overnight cultures folded similarly to those in arabinose-supplemented cultures (Fig. 5c). After further depletion of BamD in glucose minimal medium, the effect of *bamB* on BamA_{F474L} folding became more apparent, as virtually no heat-modifiable form of BamA_{F474L} could be detected (Fig. 5d). Moreover, BamA_{F474L} was present at a significantly reduced level compared to the *bamB*⁺ background under identical growth conditions (Fig. 5d). In contrast, BamA_{F474L} folded correctly in the *bamB*⁺ background despite BamD levels being undetectable after the first dilution in glucose minimal medium (Fig. 5d). These data showed that BamA_{F474L} stability and folding were severely compromised in the absence of BamB under BamD depletion conditions. Thus, while BamA_{F474L} folding may become largely independent of BamD (Figs. 3 and 4), it is still dependent on BamB in the absence of BamD.

Collectively, these data showed a good correlation between BamA_{F474L} stability/folding and the OMP biogenesis defect, and revealed a critical role of BamB in the biogenesis of BamA and other OMPs when BamD is depleted in the cell.

Discussion

The *in vivo* roles of the BamBCDE lipoproteins are currently unknown. Initial clues of their importance came from the genetic studies showing that *bamD* could not be deleted from the cell [10,11] and that the simultaneous deletion of any two non-essential BamBCE lipoproteins conferred a synthetic and/or conditional lethal phenotype [8,15, 17]. The conditional lethal phenotype of the *bamB bamE* double mutant was shown to be, in part, due to a severe impairment in BamA biogenesis, causing it to misfold and degrade [15]. Because BamE interacts with BamA indirectly via BamD, it was anticipated that when BamD is depleted, this would severely impair BamA biogenesis. Indeed, we found in this work that when BamD is depleted from the cell, the BamA_{WT} protein failed to fold correctly. The reduced BamA_{WT} level under BamD depletion conditions is most likely due to degradation of misfolded nascent and/or pre-inserted BamA and not its reduced synthesis, since the elevated envelope stress created by these conditions should increase *bamA* expression by activating the σ^E regulon [27,44,45].

If the lethal effects of BamD depletion are mainly due to misfolding of BamA, then certain alterations in BamA that make BamA's folding and/or assembly less dependent on BamD should allow for improved bacterial growth and OMP biogenesis under BamD depletion conditions. To test this hypothesis, we introduced six previously characterized BamA alterations that were obtained in a *bamB bamE* background that improved bacterial growth and stabilized BamA by influencing its folding in the absence of the two Bam

lipoproteins [15]. Five of the six BamA alterations significantly improved bacterial growth under BamD depletion conditions (Fig. 1) and stabilized BamA (Figs. 2 and S1). The sole BamA variant that did not improve growth or stabilize BamA carried an E780D alteration and was also the weakest suppressor in the *bamB bamE* background [15]. Notably, while the BamA suppressor alterations improved bacterial growth and/or OMP biogenesis in *bamB*, *bamB bamE* and BamD depletion backgrounds, they produced no negative side effects when the BAM complex contained all four wild-type Bam lipoproteins [15]. Thus, the suppressor alterations' effects on BamA folding and activity were not specific to defective or partially functional BAM complexes.

Because the F474L alteration does not affect the heat-modifiability property of BamA [15], BamA_{F474L} and BamA_{WT} were chosen to test the effect of BamD depletion on BamA folding. Not only was the bacterial growth significantly improved by the F474L alteration, it allowed the mutant BamA protein to maintain the normal folding state, which the BamA_{WT} protein lost when BamD levels dropped below the detection level of Western blot analysis (Fig. 3). Using a partially diploid strain expressing both BamA_{F474L} and BamA_{WT}, we showed the folding and assembly of BamA_{WT} was significantly improved in the presence of BamA_{F474L} under BamD depletion conditions (Fig. 4). These data indicated that the folded BamA_{F474L} molecules without BamD are functionally active and reinforced the idea that BamA assembles BamA [8,15,27]. Curiously, despite allowing normal BamA folding and improving bacterial growth, the positive effect of BamA_{F474L} on OMPs other than BamA_{WT} biogenesis was modest when BamD was depleted (Figs. 3 and 4). One explanation for this observation could be that the F474L substitution specifically influences folding and assembly of the BamA β-barrel to make it less dependent on BamD, but the overall activity of the BAM complex is still BamD dependent. Therefore, a modest improvement in substrate OMP assembly is likely a consequence of improved mutant BamA assembly and activity, which compensated for a compromised BamD-depleted BAM complex. There could be any number of indispensable roles BamD may play in OMP assembly, including serving as an OMP receptor and directly or indirectly assisting in OMP β-barrel formation and insertion into the outer membrane via BamA. It is worth mentioning that unlike in *E. coli*, BamD in *Salmonella* was shown to be dispensable for growth, although the *bamD* null mutant grew poorly [47]. This could be due to the presence of permissive substitutions in the BamA β-barrel sequence of *Salmonella*. Interestingly, both bacteria contain identical residues at the six BamA suppressor sites identified in *E. coli*.

When BamD is depleted from the cell, the whole BamACDE subcomplex should be effectively disabled because BamCE interact with BamA via BamD [11,17]. However, the effects of BamD depletion on cell viability and OMP biogenesis are by far more catastrophic, as shown in this work, than that caused by the simultaneous absence of BamCE [40]. Thus, while the whole BamACDE subcomplex is rendered inoperative without BamD, it is the absence of the essential BamD protein alone that inflicts the most devastating effects on BamA and OMP biogenesis. As mentioned earlier, the BamA suppressor alterations were isolated for their abilities to allow a *bamB bamE* mutant to grow on a rich medium [15]. In this background, the BamA suppressors also significantly improved BamA and OMP biogenesis. The fact that the absence of BamE and BamB individually does not affect BamA

folding but their simultaneous absence does [15] suggests that both BamB and BamCDE subcomplexes play overlapping roles in BamA biogenesis, with the latter playing a more critical role than the former. This same conclusion was reached in this study while examining BamA_{F474L} folding under BamD depletion conditions in a *bamB* background. Several observations made in our *in vivo* work here are consistent with a recent *in vitro* analysis carried out by Hagan *et al.* [19] which showed that BamB and BamCDE subcomplexes are individually sufficient to facilitate BamA assembly into proteoliposomes. Their analysis also showed that the presence of pre-assembled BamA in the BamCDE subcomplex allowed more efficient assembly of a FLAG-tagged substrate BamA protein than the BamCDE subcomplex alone, again indicating a role for BamA in its own assembly.

Both BamB and BamD make direct contacts with BamA [8,11,17]. BamA-BamB interaction sites, which were identified by mutagenesis and pull-down assays [22], are clustered closely in the folded structure [32-34]. It remains to be determined whether or not these sites of BamB participate in BamA assembly. The essential POTRA 5 domain of BamA is shown to be critical for BamA-BamD interactions [8,43]. Interestingly, the BamCDE subcomplex was shown *in vitro* to efficiently assemble a truncated BamA substrate lacking the POTRA 5 domain, indicating that the site of BamD-BamA interactions during BamA assembly may be different than those that occur in the fully assembled BAM holocomplex. A similar possibility was previously raised based on the structural data for BamD [35,36,38]. Of the five tetratricopeptide repeats (TPR; [48]) of BamD, the first three N-terminal TPR domains have been implicated in the formation of a structural scaffold for substrate peptide binding [35,36], while a mutational analysis revealed the involvement of the C-terminus of BamD in BamA binding [11]. The last β -strand of BamA contains residues that are highly conserved in almost all bacterial β -barrel OMPs [49]. These conserved sequences of OMPs are suggested to form a common motif recognized by the N-terminus substrate binding scaffold of BamD [35,36]. Thus, the last β -strand of nascent BamA molecules may transiently interact with the N-terminal TPR domains of BamD during assembly.

While the folding and insertion of BamA_{WT} are dependent on the Bam lipoproteins, the reverse is not true [24,43]. This is because lipoproteins rely on a well-defined, Lol-dependent sorting and targeting pathway that does not involve any β -barrel proteins [2]. Consequently, the outer membrane lipoproteins, including all four lipoproteins of the BAM complex, will be stably localized and anchored to the outer membrane by the Lol system, even in the absence of the BamA protein (Fig. 6a, top panel). Once localized in the outer membrane by the Lol system, the BamB/CDE lipoproteins in the outer membrane could then potentially act as receptors for the nascent BamA molecules in the periplasm to guide their folding and insertion into the outer membrane (Fig. 6a, middle panel). In turn, the newly assembled BAM holocomplex can catalyze the assembly of all β -barrel OMPs, including BamA (Fig. 6a, bottom panel). When BamD is depleted from the cell, the remaining BamAB subcomplex is not adequate to support BamA and OMP assembly (Fig. 6b), except in the presence of suppressor alterations within the BamA β -barrel domain (Fig. 6c). However, even in the presence of BamA β -barrel suppressor alterations, growth and OMP assembly are not restored to wild-type levels, reflecting an indispensable role of BamD in *E.*

coli BamA and OMP assembly. Perhaps distinct and/or multiple BamA alterations might be required to render BamD dispensable in *E. coli*, like in *Salmonella* [47].

The SAM complex, which catalyzes β -barrel OMPs assembly in yeast mitochondria, also contains two essential proteins, Sam50 and Sam35, which are analogous to BamA and BamD in bacteria ([6]; and references therein). Both BamA and Sam50 belong to the Omp85 superfamily of proteins [14,50]. Just as BamD is required for proper BamA assembly, Sam50 assembly has been reported to be dependent on Sam35 [51]. Interestingly, however, unlike BamA and BamD, Sam50 and Sam35 are interdependent for their assembly and stability [51]. This is because Sam35 is a peripheral mitochondrial OMP [52,53] whose outer membrane localization appears to be catalyzed primarily by direct protein-protein interactions with Sam50. Therefore, when Sam50 is depleted, Sam35 cannot be stabilized in the outer membrane from the cytoplasmic side. Conversely, when Sam35 is depleted, the resulting SAM complex cannot assemble nascent Sam50 molecules. Although the bulk of the Sam35 protein is apparently exposed to the cytosolic side of the mitochondrial outer membrane, data suggest that Sam35 acts as a receptor for nascent β -barrel OMPs, including Sam50, by recognizing their β -signal in the intermembrane space [54]. BamD, like Sam35, may also serve as a general receptor for all nascent β -barrel OMPs, including BamA. Additionally, our data and those of Hagan *et al.* [19] suggest that BamD serves to fulfill a special folding/assembly requirement of BamA.

Whereas Sam35 is analogous to BamD, the third core component of the SAM complex, Sam37, does not appear to be the counterpart of BamB. Sam37 is thought to act downstream of Sam35 to assist in the release of substrates bound to Sam35/Sam50 [55]. In contrast, BamB has been suggested to act at an early step of OMP biogenesis [21,22]. The absence of Sam37 destabilizes the SAM complex, as the levels of both Sam35 and Sam50 are reduced [51], while the absence of BamB has no effect on the BamACDE subcomplex [8] or BamA folding or levels [15]. However, BamA does become unstable when BamBE are simultaneously missing [15] or BamD is depleted (this work). One characteristic Sam37 and BamB do share is that they both are nonessential, indicating either their activities are redundant or regulatory in nature.

Materials and methods

Strains, culture conditions and chemicals

Bacterial strains used in the study are listed in Table 1. M63-salt based minimal medium was prepared as described by Silhavy *et al.* [56]. The medium was supplemented with thiamine (0.1 mg ml^{-1}), MgSO_4 (1 mM), and casamino acids (0.1%). Glucose or glycerol (both at 0.2%) was used as a carbon source. When necessary, L-arabinose was added to final concentration of 0.2%. All media ingredients were purchased from Difco. All other chemicals were of analytical grade. All cultures were grown at 37°C unless specified.

Protein methods

Envelopes were isolated by the French press method as described previously (Bennion *et al.*, 2010). Detergent solubilized envelopes were analyzed by mini sodium dodecyl sulfate

(SDS)-polyacrylamide (11%) gel electrophoresis (PAGE). For a better separation of OmpC and OmpF, 4M urea was added to the running gel. Protein bands were visualized by staining gels with Coomassie blue.

For Western blot analysis, samples from whole cell extracts or purified envelopes were analyzed by SDS-PAGE and transferred onto Immobilon-P polyvinylidene difluoride membranes (Millipore). Membranes were blocked overnight in 5% (w/v) non-dairy cream. After overnight blocking, membranes were incubated with primary antibodies for 1.5 hours. Primary antibodies used were raised in rabbits against purified AcrA (1:16,000), BamA (1:5,000), and BamD (1:2,000), OmpF/C/A (1:25,000). BamA and BamD antibodies were gifts from Tom Silhavy, Princeton University. After incubation with primary antibodies, membranes were washed twice for 15 minutes each and incubated for 1 hour in secondary antibodies (goat anti-rabbit horseradish peroxidase-conjugated IgG). Protein bands were visualized with Bio-Rad Molecular Imager ChemiDoc XRS System and quantified using Quantity One Software (Bio-Rad).

Genetic methods

Bacteriophage P1-mediated transductions were carried out using a protocol described by Silhavy *et al.* [56]. The *bamA* alleles were transduced into a *bamD* (λ att-lom)::bla P_{BAD} *bamD* *araC* *degP*::Km^r strain (RAM1823) using a *bamA*-linked *zae*::Tn10 (Tc^r; 4.75'). Tc^r transductants were screened for kanamycin sensitivity to ensure the replacement of *degP*::Km^r (3.90'), which is located upstream of *bamA* (4.27') by the *degP*⁺ and *bamA* alleles. DNA sequence analysis of the *bamA* gene confirmed that all Tc^r Km^s transductants had replaced the recipient *bamA* allele with that linked to the *zae*::Tn10 from the donor P1 lysates. Strains partially diploid for *bamA* were constructed in two steps. First, a *recA*::Cm^r allele was introduced via P1 transduction so as to prevent homologous recombination between the two alleles of *bamA*. These strains were then transformed with pZS21, which expresses a His-tagged BamA protein.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are indebted to Phu Vuong for critically reading the manuscript. We would also like to thank Lauren Lynch and Mackenzie Lynch for their assistance in SDS-PAGE analysis, and Rene Tellez Jr. for his involvement in the early phase of the project. RS was supported by a Research Fellowship from the School of Life Sciences Undergraduate Research program. This work was supported by grants from the National Institutes of Health (GM048167) and the School of Life Sciences, Arizona State University.

References

1. Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. *Cold Spring Harb Prospect Biol.* 2010; 2:a000414.
2. Okuda S, Tokuda H. Lipoprotein sorting in bacteria. *Annu Rev Microbiol.* 2011; 65:239–259. [PubMed: 21663440]
3. Ruiz N, Kahne D, Silhavy TJ. Transport of lipopolysaccharide across the cell envelope, the long road of discovery. *Nature Rev Microbiol.* 2009; 7:677–683. [PubMed: 19633680]

4. Sperandeo P, Dehò G, Polissi A. The lipopolysaccharide transport system of Gram negative bacteria. *Biochim Biophys Acta*. 2009; 1791:594–602. [PubMed: 19416651]
5. Hagan CL, Silhavy TJ, Kahne D. β -barrel membrane protein assembly by the Bam complex. *Annu Rev Biochem*. 2011; 80:189–210. [PubMed: 21370981]
6. Misra R. Assembly of the β -barrel outer membrane proteins in Gram-negative bacteria, mitochondria, and chloroplasts. *ISRN Mol Biol*. 2012; 2012:1–15.
7. Ricci DP, Silhavy TJ. The Bam machine, a molecular cooper. *Biochim Biophys Acta*. 2012; 1818:1067–1084. [PubMed: 21893027]
8. Wu T, Malinverni J, Ruiz N, Kim S, Silhavy TJ, Kahne D. Identification of a multicomponent complex required for outer membrane biogenesis in *Escherichia coli*. *Cell*. 2005; 121:235–245. [PubMed: 15851030]
9. Werner J, Misra R. YaeT (Omp85). affects the assembly of lipid-dependent and lipid-independent outer membrane proteins of *Escherichia coli*. *Mol Microbiol*. 2005; 57:1450–1459. [PubMed: 16102012]
10. Onufryk C, Crouch M-L, Fang FC, Gross CA. Characterization of six lipoproteins in the σ^E regulon. *J Bacteriol*. 2005; 187:4552–4561. [PubMed: 15968066]
11. Malinverni JC, Werner J, Kim S, Sklar JG, Kahne D, Misra R, Silhavy TJ. YfiO stabilizes the YaeT complex and is essential for outer membrane protein assembly in *Escherichia coli*. *Mol Microbiol*. 2006; 61:151–164. [PubMed: 16824102]
12. Voulhoux R, Bos MP, Geurtsen J, Mols M, Tommassen J. Role of a highly conserved bacterial protein in outer membrane protein assembly. *Science*. 2003; 299:262–265. [PubMed: 12522254]
13. Gentle I, Gabriel K, Beech P, Waller R, Lithgow T. The Omp85 family of proteins is essential for outer membrane biogenesis in mitochondria and bacteria. *J Cell Biol*. 2004; 164:19–24. [PubMed: 14699090]
14. Moslavac S, Mirus O, Bredemeier R, Soll J, Haeseler A von, Schleiff E. Conserved pore-forming regions in polypeptide-transporting proteins. *FEBS J*. 2005; 272:1367–1378. [PubMed: 15752354]
15. Tellez R Jr, Misra R. Substitutions in the BamA β -barrel domain overcome the conditional lethal phenotype of a *bamB bamE* strain of *Escherichia coli*. *J Bacteriol*. 2012; 194:317–324. [PubMed: 22037403]
16. Webb CT, Heinz E, Lithgow T. Evolution of the β -barrel assembly machinery. *Trends Microbiol*. 2012; 20:612–620. [PubMed: 22959613]
17. Sklar JG, Wu T, Gronenberg LS, Malinverni JC, Kahne D, Silhavy TJ. Lipoprotein SmpA is a component of the YaeT complex that assembles outer membrane proteins in *Escherichia coli*. *Proc Natl Acad Sci USA*. 2007; 104:6400–6405. [PubMed: 17404237]
18. Hagan CL, Kim S, Khane D. Reconstitution of outer membrane protein assembly from purified components. *Science*. 2010; 328:890–892. [PubMed: 20378773]
19. Hagan CL, Westwood DB, Kahne D. Bam lipoproteins assemble BamA *in vitro*. *Biochem*. 2013; 52:6108–6113. [PubMed: 23919461]
20. Charlson ES, Werner JN, Misra R. Differential effects of *yfgL* mutation on the biogenesis of *Escherichia coli* outer membrane proteins and lipopolysaccharide. *J Bacteriol*. 2006; 188:7186–7194. [PubMed: 17015657]
21. Ureta AR, Endres RG, Wingreen NS, Silhavy TJ. Kinetic analysis of the assembly of the outer membrane protein LamB in *Escherichia coli* mutants each lacking a secretion or targeting factor in a different cellular compartment. *J Bacteriol*. 2007; 189:446–454. [PubMed: 17071751]
22. Vuong P, Bennion D, Mantei J, Frost D, Misra R. Analysis of YfgL and YaeT interactions through bioinformatics, mutagenesis, and biochemistry. *J Bacteriol*. 2008; 190:1507–1517. [PubMed: 18165306]
23. Sánchez-Pulido L, Devos D, Genevrois S, Vicente M, Valencia A. POTRA, a conserved domain in the FtsQ family and a class of β -barrel outer membrane proteins. *Trends Biochem Sci*. 2003; 28:523–526. [PubMed: 14559180]
24. Kim S, Malinverni JC, Sliz P, Silhavy TJ, Harrison SC, Kahne D. Structure and function of an essential component of the outer membrane protein assembly machine. *Science*. 2007; 317:961–964. [PubMed: 17702946]

25. Gatzeva-Topalova PZ, Walton TA, Sousa MC. Crystal structure of YaeT, conformational flexibility and substrate recognition. *Structure*. 2008; 16:1873–1881. [PubMed: 19081063]
26. Knowles TJ, Jeeves M, Bobat S, Dancea F, McClelland D, Palmer T, et al. Fold and function of polypeptide transport-associated domains responsible for delivering unfolded proteins to membranes. *Mol Microbiol*. 2008; 68:1216–1227. [PubMed: 18430136]
27. Bennion D, Charlson ES, Coon E, Misra R. Dissection of β -barrel outer membrane protein assembly pathways through characterizing BamA POTRA 1 mutants of *Escherichia coli*. *Mol Microbiol*. 2010; 77:1153–1171. [PubMed: 20598079]
28. Workman P, Heide K, Giuliano N, Lee N, Mar J, Vuong P, et al. Genetic, biochemical, and molecular characterization of the polypeptide transport-associated domain of *Escherichia coli* BamA. *J Bacteriol*. 2012; 194:3512–3521. [PubMed: 22544271]
29. Noinaj N, Kuszak AJ, Gumbart JC, Lukacik P, Chang H, Easley NC, et al. Structural insight into the biogenesis of β -barrel membrane proteins. *Nature*. 2013; 501:385–392. [PubMed: 23995689]
30. Jacob-Dubuisson F, Villeret V, Clantin B, Delattre A, Saint N. First structural insights into the TpsB/Omp85 superfamily. *Biol Chem*. 2009; 390:675–684. [PubMed: 19558323]
31. Leonard-Rivera M, Misra R. Conserved residues of the putative L6 loop of *Escherichia coli* BamA play a critical role in the assembly of β -barrel outer membrane proteins, including that of BamA itself. *J Bacteriol*. 2012; 194:4662–4668. [PubMed: 22753067]
32. Heuck A, Schleiffer A, Clausen T. Augmenting β -augmentation, structural basis of how BamB binds BamA and may support folding of outer membrane proteins. *J Mol Biol*. 2011; 406:659–666. [PubMed: 21236263]
33. Kim KH, Paetzel M. Crystal structure of *Escherichia coli* BamB, a lipoprotein component of the β -barrel assembly machinery complex. *J Mol Biol*. 2011; 406:667–678. [PubMed: 21168416]
34. Noinaj N, Fairman JW, Buchanan SK. The crystal structure of BamB suggests interactions with BamA and its role within the BAM complex. *J Mol Biol*. 2011; 407:248–260. [PubMed: 21277859]
35. Albrecht R, Zeth K. Structural basis of outer membrane protein biogenesis in bacteria. *J Biol Chem*. 2011; 286:27792–27803. [PubMed: 21586578]
36. Sandoval CM, Baker SL, Jansen K, Metzner SI, Sousa MC. Crystal structure of BamD, an essential component of the β -barrel assembly machinery of Gram-negative bacteria. *J Mol Biol*. 2011; 409:348–357. [PubMed: 21463635]
37. Dong C, Hou H-F, Yang X, Shen Y-Q, Dong Y-H. Structure of *Escherichia coli* BamD and its functional implications in outer membrane protein assembly. *Acta Cryst*. 2012; 68:95–101.
38. Kim KH, Aulakh S, Paetzel M. Crystal structure of β -barrel assembly machinery BamCD protein complex. *J Biol Chem*. 2011; 286:39116–39121. [PubMed: 21937441]
39. Knowles TJ, Browning DF, Jeeves M, Maderbocus V, Rajesh S, Sridhar P, et al. Structure and function of BamE within the outer membrane and the β -barrel assembly machine. *EMBO J*. 2011; 12:123–128.
40. Rigel NW, Schwalm J, Ricci DP, Silhavy TJ. BamE modulates the *Escherichia coli* β -barrel assembly machine component BamA. *J Bacteriol*. 2012; 194:1002–1008. [PubMed: 22178970]
41. Robert V, Volokhina EB, Senf F, Bos MP, Gelder P van, Tommassen J. Assembly factor Omp85 recognizes its outer membrane protein substrates by a species-specific C-terminal motif. *PLoS Biol*. 2006; 11:1984–1995.
42. Kim KH, Aulakh S, Paetzel M. The bacterial outer membrane β -barrel assembly machinery. *Prot Sci*. 2012; 21:751–768.
43. Ricci DP, Hagan CL, Kahne D, Silhavy TJ. Activation of the *Escherichia coli* β -barrel assembly machine (Bam). is required for essential components to interact properly with substrate. *Proc Natl Acad Sci USA*. 2012; 109:3487–3491. [PubMed: 22331884]
44. Rhodius VA, Suh WC, Nonaka G, West J, Gross CA. Conserved and variable functions of the σ^E stress response in related genomes. *PLoS Biol*. 2006; 4:43–59.
45. Gerken H, Leiser OP, Bennion D, Misra R. Involvement and necessity of the Cpx regulon in the event of aberrant β -barrel outer membrane protein assembly. *Mol Microbiol*. 2010; 75:1033–1046. [PubMed: 20487295]

46. Stegmeier JF, Andersen C. Characterization of pores formed by YaeT (Omp85) from *Escherichia coli*. *J Biochem*. 2006; 140:275–283. [PubMed: 16829683]
47. Fardini Y, Trotureau J, Bottreau E, Souchart C, Velge P, Virlogeux-Payant I. Investigation of the role of the BAM complex and SurA chaperone in outer-membrane protein biogenesis and type III secretion system expression in *Salmonella*. *Microbiol*. 2009; 155:1613–1622.
48. D'Andrea LD, Regan L. TPR proteins, the versatile helix. *Trends Biochem Sci*. 2003; 28:655–662. [PubMed: 14659697]
49. Struyvé M, Moons M, Tommassen J. Carboxy-terminal phenylalanine is essential for the correct assembly of a bacterial outer membrane protein. *J Mol Biol*. 1991; 218:141–148. [PubMed: 1848301]
50. Gentle IE, Burri L, Lithgow T. Molecular architecture and function of the Omp85 family of proteins. *Mol Microbiol*. 2005; 58:1216–1225. [PubMed: 16313611]
51. Waizenegger T, Habib SJ, Lech M, Mokranjac D, Paschen SA, Hell K, et al. Tob38, a novel essential component in the biogenesis of β -barrel proteins of mitochondria. *EMBO Reports*. 2004; 5:704–709. [PubMed: 15205677]
52. Milenkovic D, Kozjak V, Wiedemann N, Lohaus C, Meyer HE, Guiard B, et al. Sam35 of the mitochondrial protein sorting and assembly machinery is a peripheral outer membrane protein essential for cell viability. *J Biol Chem*. 2004; 279:22781–22785. [PubMed: 15067005]
53. Ishikawa D, Yamamoto H, Tamura Y, Moritoh K, Endo T. Two novel proteins in the mitochondrial outer membrane mediate β -barrel protein assembly. *J Cell Biol*. 2004; 166:621–627. [PubMed: 15326197]
54. Kutik S, Stojanovski D, Becker L, Becker T, Meinecke M, Krüger V, et al. Dissecting membrane insertion of mitochondrial β -barrel proteins. *Cell*. 2008; 132:1011–1024. [PubMed: 18358813]
55. Chan NC, Lithgow T. The peripheral membrane subunits of the SAM complex function codependently in mitochondrial outer membrane biogenesis. *Mol Biol Cell*. 2008; 19:126–136. [PubMed: 17978093]
56. Silhavy, TJ.; Berman, ML.; Enquist, LW. *Experiments with gene fusions*. CSHL Press; Cold Spring Harbor, NY: 1984.
57. Casadaban MJ. Transposition and fusion of the *lac* genes to select promoters in *Escherichia coli* using bacteriophage lambda and Mu. *J Mol Biol*. 1976; 141:541–555. [PubMed: 781293]

Highlights

- Without essential BamD, the core BamA protein of the BAM complex becomes misfolded.
- Compensatory alterations within BamA β -barrel restore BamA folding without BamD.
- Folded mutant BamA without BamD then facilitates OMP and wild type BamA assembly.







Fig 1. Effects of BamD depletion on cell growth. Overnight grown cultures in glycerol minimal medium were diluted 1:100 in glucose minimal medium and grown for 3.5 hours. At the end of this period, cultures were diluted 1:50 in the same medium and grown for 4 hours, and then diluted 1:50 one last time in fresh glucose minimal medium and grown for another 4 hours. BamD expression is repressed in glucose minimal medium. Cell densities were measured in a spectrophotometer at 600 nm. All cultures were grown at 37°C in baffled flasks in a water bath incubator shaker set at 180 rpm. Labeled are cells expressing different BamA proteins. Residue numbers are relative to the mature BamA protein sequence.

**Fig. 2.**


Effects of BamD depletion on BamA. Cell lysates, prepared from samples collected during the growth experiments in Fig. 1, were analyzed by SDS-PAGE. For clarity, data from only four strains are shown here; data from the entire set, including samples collected after the third dilution, are shown in Fig. S1. BamA was detected by Western blot analysis using BamA-specific antibodies (left panels). After developing with BamA antibodies, the same blot was probed with antibodies against AcrA (right panels). AcrA, an inner membrane lipoprotein, is used as a gel loading control. Each lane contains protein lysate from equal number of cells based on OD₆₀₀. Positions of AcrA, BamA and a major BamA fragment [BamA-(N)] are shown. [BamA] refers to BamA/AcrA ratios. Residue numbers are relative to the mature BamA protein sequence.

**Fig. 3.**


Analyses of OMP levels and BamA folding from BamD-depleted cells. Envelopes were purified from cultures grown in glucose- or arabinose-supplemented glycerol minimal medium to repress or induce *bamD* expression, respectively. (a) BamA folding status was determined from envelopes obtained from cultures after first and second dilutions. Prior to SDS-PAGE analysis, SDS-sample buffer-solubilized samples were either left at room temperature (25°C) or heated (100°C). Detection of BamA was carried out by Western blot analysis using BamA-specific antibodies. Electrophoresis was carried out at room temperature. BamA_U and BamA_F refer to unfolded and folded forms of BamA, respectively. (b) SDS(urea)-PAGE analysis of purified envelopes. Protein bands were visualized after Coomassie blue staining. Positions of the three major OMPs are shown. (c) Western blot analysis to detect BamD. The Coomassie blue-stained gel in (b) was electro-blotted and the membrane blot was probed with BamD-specific antibodies. (d) Western blot analysis to detect OMPs from BamD-depleted envelope samples.

**Fig. 4.**

Effects of BamD depletion on bacterial growth, OMP levels, and BamA folding in strains partially diploid for *bamA*. (a) Overnight grown cultures in glycerol minimal medium were diluted three times and grown in a 37°C water bath as described in Fig. 1. BamD expression is repressed in glucose minimal medium. Cell densities were measured in a spectrophotometer at 600 nm. The untagged BamA proteins were expressed from the chromosome, while BamA_{WT-His} was expressed from pZS21 (Km^r). Medium was supplemented with kanamycin to retain the pZS21 plasmid and with tetracycline to induce the expression of *bamA* from the plasmid replicon. Without tetracycline, the TetR repressor protein from a chromosomally localized *zae::Tn10* inhibits the expression of *bamA* from the pZS21 plasmid. (b) and (c) SDS(urea)-PAGE and Western blot analyses of envelopes purified from glycerol-grown overnight (o/n) cultures and after three dilutions (1st, 2nd, and 3rd) in glucose minimal medium (Fig. 4a). Protein bands were visualized either after Coomassie blue staining (b) or Western blot analysis using OmpF/C/A antibodies (c). BamA compositions of the partially diploid strains are labeled on top. (d) and (e) BamA folding status was determined from envelopes obtained from cultures grown overnight in glycerol minimal medium and after dilutions in glucose minimal medium (Fig. 4a). Prior to SDS-PAGE analysis, SDS-solubilized samples were either left at room temperature (25°C) or heated (100°C). Detection of BamA_{WT-His} was carried out by Western blot analysis using His-Probe. BamA_U and BamA_F refer to unfolded and folded forms of BamA, respectively. (f) Quantification of BamA folding. The graph shows the ratios of folded (BamA_F) to unfolded (BamA_U) species. Data were obtained from two independent experiments.

**Fig 5.**

Effects of the presence or absence of BamB on $BamA_{F474L}$ folding in BamD-depleted cells. (a) Overnight cultures were grown in glycerol minimal medium, supplemented with or without arabinose. Cultures grown without arabinose were diluted 1:100 in glucose minimal medium and grown for 3.5 hours. At the end of this period, cultures were diluted 1:50 in fresh glucose minimal medium and grown for another 3.5 hours. Cell densities were measured at 600 nm. Cell samples were collected from overnight grown cells and at the end of the first dilution for protein analysis. All cultures were grown at 37°C in baffled flasks in a water bath incubator shaker set at 180 rpm. (b) Envelopes obtained from cultures described in Fig. 5a were analyzed by SDS(urea)-PAGE. BamD is expressed in the presence of arabinose in glycerol minimal medium and repressed without arabinose in glucose minimal medium. Protein bands were visualized after Coomassie blue staining. Positions of the three major OMPs are shown. (c) and (d) BamA folding status was determined using envelopes obtained from cultures grown overnight in glycerol minimal medium (c) or after the first dilution in glucose minimal medium (d). Prior to SDS-PAGE analysis, SDS-solubilized samples were either left at room temperature (25°C) or heated (100°C). Detection of BamA was carried out by Western blot analysis using BamA-specific antibodies. The same blot was subsequently probed with BamD antibodies (lower panels). BamD is expressed in the presence of arabinose and repressed in its absence in glucose minimal medium. Electrophoresis was performed at room temperature. BamA_U and BamA_F refer to unfolded and folded forms of BamA, respectively.

**Fig. 6.**

A hypothetical model for the *in vivo* assembly of an active BAM complex and β -barrel OMPs. (a) The outer membrane lipoproteins, including the four lipoproteins of the BAM complex, BamDCDE, are localized to the outer membrane via the localization of lipoprotein (Lol) system [2] (top panel). The BamB and BamDCE subcomplexes facilitate the assembly of the nascent BamA molecules (middle panel). The assembled BAM holocomplex then catalyzes the assembly of all β -barrel OMPs, including BamA, which is more efficiently assembled in the presence of the BAM holocomplex (bottom panel) than in the presence of the Bam lipoproteins alone. (b) When BamD is depleted, the assembly of BamA_{WT} and β -barrel OMPs is severely compromised. (c) Certain alterations within BamA β -barrel, such as F474L, significantly improve assembly of BamA when BamD is depleted (bottom panel). In contrast to BamA_{F474L}, a relatively modest improvement in the assembly of other β -barrel OMPs under BamD depletion conditions may reflect their continued dependence on BamD.

Table 1

Bacterial strains used in this study.

Strains	Relevant characteristics	Reference
MC4100	F- <i>araD139</i> Δ (<i>argF-lac</i>)U139 <i>rspL150 relA1 flbB5301 ptsF25 deoC1 thi-1 rbsR</i>	[57]
RAM1821	JCM290: MC4100 <i>ard^{tr} ΔbamD Δ(λatt-lom)::bla P_{BAD} bamD araC</i>	[11]
RAM1823	RAM1821 <i>degP::Km^r</i>	This study
RAM2253	RAM1821 <i>zae::Tn10 (4.75') BamA_{F474L} degP⁺</i>	This study
RAM2254	RAM1821 <i>zae::Tn10 (4.75') BamA_{N631T} degP</i>	This study
RAM2255	RAM1821 <i>zae::Tn10 (4.75') BamA_{G635A} degP⁺</i>	This study
RAM2256	RAM1821 <i>zae::Tn10 (4.75') BamA_{G649A} degP⁺</i>	This study
RAM2257	RAM1821 <i>zae::Tn10 (4.75') BamA_{A652V} degP⁺</i>	This study
RAM2258	RAM1821 <i>zae::Tn10 (4.75') BamA_{E780D} degP⁺</i>	This study
RAM2259	RAM1821 <i>zae::Tn10 (4.75') BamA_{WT} degP⁺</i>	This study
RAM2268	RAM2253 <i>ΔbamB::Km^r</i>	This study
RAM2269	RAM2254 <i>ΔbamB::Km^r</i>	This study
RAM2270	RAM2255 <i>ΔbamB::Km^r</i>	This study
RAM2271	RAM2256 <i>ΔbamB::Km^r</i>	This study
RAM2272	RAM2257 <i>ΔbamB::Km^r</i>	This study
RAM2273	RAM2258 <i>ΔbamB::Km^r</i>	This study
RAM2274	RAM2259 <i>ΔbamB::Km^r</i>	This study
RAM2308	RAM2259 <i>recA::Cm^r</i>	This study
RAM2309	RAM2253 <i>recA::Cm^r</i>	This study
RAM2310	RAM2308 (pZS21-BamA _{WT} -His)	This study
RAM2311	RAM2309 (pZS21-BamA _{WT} -His)	This study