

Research article

Open Access

## Patterns of linkage disequilibrium and haplotype distribution in disease candidate genes

Ji-Rong Long<sup>1</sup>, Lan-Juan Zhao<sup>1,2</sup>, Peng-Yuan Liu<sup>1</sup>, Yan Lu<sup>1</sup>,  
Volodymyr Dvornyk<sup>1</sup>, Hui Shen<sup>1,2</sup>, Yong-Jun Liu<sup>1,2</sup>, Yuan-Yuan Zhang<sup>1</sup>,  
Dong-Hai Xiong<sup>1,2</sup>, Peng Xiao<sup>1,2</sup> and Hong-Wen Deng\*<sup>1,2,3</sup>

Address: <sup>1</sup>Osteoporosis Research Center, Creighton University, Omaha, NE 68131, USA, <sup>2</sup>Department of Biomedical Sciences, Creighton University, Omaha, NE 68131, USA and <sup>3</sup>Laboratory of Molecular and Statistical Genetics, College of Life Sciences Hunan Normal University, Changsha, Hunan 410081, P. R. China

Email: Ji-Rong Long - jrlongyou@creighton.edu; Lan-Juan Zhao - zhaolanjuan@creighton.edu; Peng-Yuan Liu - pylu@creighton.edu; Yan Lu - yanlu@creighton.edu; Volodymyr Dvornyk - Dvornyk@creighton.edu; Hui Shen - huishen@creighton.edu; Yong-Jun Liu - jun@creighton.edu; Yuan-Yuan Zhang - zhangy2@creighton.edu; Dong-Hai Xiong - donghai@creighton.edu; Peng Xiao - peng@creighton.edu; Hong-Wen Deng\* - deng@creighton.edu

\* Corresponding author

Published: 24 May 2004

BMC Genetics 2004, 5:11

Received: 29 January 2004

Accepted: 24 May 2004

This article is available from: <http://www.biomedcentral.com/1471-2156/5/11>

© 2004 Long et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** The adequacy of association studies for complex diseases depends critically on the existence of linkage disequilibrium (LD) between functional alleles and surrounding SNP markers.

**Results:** We examined the patterns of LD and haplotype distribution in eight candidate genes for osteoporosis and/or obesity using 31 SNPs in 1,873 subjects. These eight genes are apolipoprotein E (APOE), type I collagen  $\alpha 1$  (COL1A1), estrogen receptor- $\alpha$  (ER- $\alpha$ ), leptin receptor (LEPR), parathyroid hormone (PTH)/PTH-related peptide receptor type I (PTHRI), transforming growth factor- $\beta 1$  (TGF- $\beta 1$ ), uncoupling protein 3 (UCP3), and vitamin D (1,25-dihydroxyvitamin D<sub>3</sub>) receptor (VDR). Yin yang haplotypes, two high-frequency haplotypes composed of completely mismatching SNP alleles, were examined. To quantify LD patterns, two common measures of LD, D' and r<sup>2</sup>, were calculated for the SNPs within the genes. The haplotype distribution varied in the different genes. Yin yang haplotypes were observed only in PTHRI and UCP3. D' ranged from 0.020 to 1.000 with the average of 0.475, whereas the average r<sup>2</sup> was 0.158 (ranging from 0.000 to 0.883). A decay of LD was observed as the intermarker distance increased, however, there was a great difference in LD characteristics of different genes or even in different regions within gene.

**Conclusion:** The differences in haplotype distributions and LD patterns among the genes underscore the importance of characterizing genomic regions of interest prior to association studies.

## Background

Considerable attention is currently being focused on the genotyping of single nucleotide polymorphisms (SNPs) to search for complex disease susceptibility alleles. The success to detect association between marker alleles and disease critically depends on the extent of linkage disequilibrium (LD) between functional alleles and surrounding markers. Studies on some chromosomes and genome regions have shown that LD is highly variable across human genome and is structured into discrete blocks of sequences separated by hotspots of recombination and/or LD breakdown [1-6]. The LD heterogeneity across the genome is a crucial aspect for genome-wide association studies, which has been emphasized as a powerful approach for complex disease study [7-9]. Due to practical limitations of the expenses, the sample size used on the whole genome scale may not be sufficiently large. For example, the ongoing HapMap represents one of the most comprehensive and largest efforts ever attempted in biomedical research, in which a total of 270 subjects from African, Caucasian and Japanese/Chinese were recruited. However, sample size is an important factor influencing the accuracy of LD evaluation. Small sample size will bias the accuracy of LD estimation [2,10].

A limited but hypothesis-driven approach for association studies is to focus on the polymorphisms located in or around the candidate genes known to be of potentially functional importance for complex traits (based on their cellular and molecular studies). LD patterns in specific candidate genes need be empirically assessed to facilitate efficient experimental designs and execution of association studies [1,11]. So far, only limited information is available about the LD patterns and haplotype characteristics in candidate genes for complex diseases [12,13].

The purpose of this study was to determine the LD and haplotype characteristics in eight candidate genes for the two common complex disorders, osteoporosis and obesity. These eight genes are apolipoprotein E (APOE), type I collagen  $\alpha 1$  (COL1A1), estrogen receptor- $\alpha$  (ER- $\alpha$ ), leptin receptor (LEPR), parathyroid hormone (PTH)/PTH-related peptide receptor type 1 (PTH1R), transforming growth factor- $\beta 1$  (TGF- $\beta 1$ ), uncoupling protein 3 (UCP3), and vitamin D (1,25-dihydroxyvitamin D<sub>3</sub>) receptor (VDR).

## Results

### Haplotype analysis

Allele frequencies of the 31 SNPs are shown in Table 1. The information for the main haplotypes whose frequencies were higher than 1% [14], is presented in Table 2. It is clear that the number of observed haplotypes was smaller than that of haplotypes expected. The number of observed haplotypes varied at different loci. In both the

PTH1R gene and the VDR gene, four SNPs were studied. However, seven and twelve haplotypes, respectively, were observed as compared to the expected 16 haplotypes in each gene.

The yin yang haplotypes are defined as two high-frequency haplotypes composed of completely mismatching SNP alleles, i.e., nucleotides differ at every SNP in the haplotype pair [15]. We detected such haplotype phenomenon in two of the eight studied genes (Table 2). In the PTH1R gene, one pair of yin yang haplotypes, GTAA and ACGG (in the order of SNP20-21-22-23), was observed with frequencies of 34.2% and 52.6%, respectively. A pair of yin yang haplotypes, TCT and CTC (in the order of SNP28-30-31), was also observed in the UCP3 gene with frequencies of 23.5% and 40.5%, respectively.

### LD analysis

In this study, we used two most common LD measures,  $D'$  and  $r^2$ , to determine the extent of LD in the genes (Figure 1). The average of  $D'$  values within the eight genes was  $0.475 \pm 0.325$  ( $\pm SD$ ) with the range of 0.020–1.000, whereas the  $r^2$  was  $0.158 \pm 0.255$  ( $\pm SD$ ), which ranged from 0.000 to 0.883. Until now, there was no unified standard for meaningful LD between a marker and a functional mutation. Although we do not know whether any of our studied SNPs is functional for osteoporosis and/or obesity, the knowledge of LD patterns revealed through them is useful in inferring effective intermarker distance. When a criterion of  $D' > 0.33$  [4,16] was used as a meaningful LD, 58.8% (30/51) of the SNP pairs met this criterion with the intermarker physical distance ranging from 138 bp (SNP3-SNP4) to ~122 kb (SNP12-SNP14). If we considered  $D' = 0.7$  as a limit for useful LD [17], only 33.3% (17/51 SNP pairs) were "useful". If a value of  $r^2 > 0.1$  was applied as a criterion for meaningful LD [16,18], 31.4% (16/51 SNP pairs) would be considered in "useful" LD. When we used both  $D' > 0.33$  and  $r^2 > 0.1$  as a criterion for meaningful LD [4,16,18], these 16 pairs still fell below this threshold. Such "useful" LD was relatively common in a pair-wise physical distance smaller than 8.6 kb, though it was detectable at the distance of 83 kb (SNP13-SNP14).

From Figure 1, the distribution of LD versus physical distance was irregular; however, a decay of LD with increasing distance was still evident. The Spearman correlation coefficient between  $D'$  and the distance was -0.70 ( $P < 0.01$ ). The correlation coefficient was -0.49 ( $P < 0.01$ ) when LD was quantified by  $r^2$ . Similar relationship was observed between LD measures and  $\ln(\text{distance})$ , in which the correlation coefficient was -0.68 and -0.47 for  $D'$  and  $r^2$ , respectively. The  $D'/\ln(\text{distance})$  was calculated for pair-wise SNPs within each gene. The average of  $D'/\ln(\text{distance})$  varied a lot among these eight genes with the

**Table 1: Information about the 37 SNPs in the eight genes**

SNP	Gene	dbSNP accession	polymorphism <sup>a</sup>	Domain	Distance <sup>b</sup>	Frequency (%) <sup>c</sup>	Forward Primer	Reverse Primer
1	APOE	ss12568587	G-C	5' UTR		35.7	TCCCCAGGAGCCGGTG A	CCCCAAGCCCGACCCC A
2	APOE	ss12568609	G-A	Intron 2	1,277	39.9	CCTCAGGTGATCTGCCC GTTTC	ACTCCTGGGCTCAAGTG ATCCTC
3	APOE	ss12568607	T-C	Exon 4	1,497	14.9	CGGGCACGGCTGTCCA A	CGAGCATGGCCTGCAC CTC
4	APOE	ss12568612	C-T	Exon 4	138	8.7	GCTGCGTAAGCGGCTC C	GCGGCCCTGTTCCACC C
5	COL1A1	ss12568606	G-T	5' UTR		15.4	GCACCCTGCCCTAGAC CAC	CCTAGTGCCAGCGACT GCA
6	COL1A1	ss12568597	G-T	Intron 1	3,543	18.8	CCAATCAGCCGCTCCC ATTC	CATCGGGAGGGCAGGC TC
7	COL1A1	ss12568598	G-T	Exon 8	2,365	0.0	GGAAGACTGGGATGAG GGCA	GGCTCGCCAGGCTCAC C
8	COL1A1	ss12568584	G-A	Exon 45	9,862	1.9	CTCAGCCTCCCTGGC CAA	AGCGGAAGTTCATTG GCATC
9	ER- $\alpha$	ss12568579	A-G	Exon 1		48.4	TTGAGCTGCGGACGGTT CA	CGCCGGTTTCTGAGCCT TC
10	ER- $\alpha$	ss12568596	T-C	Intron 1	34,258	44.9	TGGGATTCCAGGCATGA ACCAC	TGGCGTCGATTATCTGA ATTTGGCC
11	ER- $\alpha$	ss12568619	G-A	Intron 3	66,110	25.7	CCCAGAAACAAGTCATC TGCTATTGACA	GTAACAAAAGGTTAACA ATGGTTAGCCC
12	ER- $\alpha$	ss12568618	G-C	Exon 4	36,077	21.8	ACAGCCTGGCCTGTCC C	CAGGTTGGTCAGTAAGC CCATCA
13	ER- $\alpha$	ss12568585	G-A	Intron 4	39,074	9.8	GATCAATGAAGTGGGTC TTGAAAAACCAA	GGTGACAAGCTGGAAAT CTAAGCTTCA
14	ER- $\alpha$	ss12568605	G-A	Intron 6	83,068	11.9	GGAACGGCCCTTGAA ATTGTA	CTGCCTACAGAATACAG TCAGCCA
15	ER- $\alpha$	ss12568617	G-A	Exon 8	32,431	20.3	TCGCATTCCTTGCAAAA GTATTACATCAC	CAAGCAAATGAATGGCC ACTCATCTAGAAA
16	LEPR	ss12568615	A-G	Exon 4		29.0	AGATTTAAGTTGTCTTG CATGCCACC	TTAAGCCCAGCATCCAT TAGCTATTCTTTC
17	LEPR	ss12568604	G-C	Exon 14	3,013	18.2	GAGTAATTGGAGCAATC CAGCCTACA	GCTTCAGCCACTGTACA TCTTAGCTC
18	LEPR	ss12568614	G-A	Exon 20	62,803	35.3	GCCACGCTGATCAGCA ACTC	CCTTTGACTTGTGAGTC AAAAGCAC
19	PTHR1	ss12568589	G-A	Intron 1		0.1	GACTTACATTAGGATTC AAGGTTACTGCCA	GGGACGCAAGCCTGAG TCC
20	PTHR1	ss12568592	A-G	Intron 2	4,250	39.8	GCAGAACCCTAAGGGC TTGTCA	GGCGGGACCCAGGATA CA
21	PTHR1	ss12568591	C-T	Intron 8	5,436	37.4	CGAGCCTCAATTCAGGT GAATCTAACC	CCCGCCCCAAGTGGA CA
22	PTHR1	ss12568588	G-A	Intron 10	1,912	39.7	CCTTGAGCCCTTGTTTT TCCTTTC	GCTCCGGGAACAAAA GTGGATCA
23	PTHR1	ss12568590	G-A	Exon 13	1,246	38.0	CTACAAGGCTCAAATTG CCCCAAA	TTGGCGTCCACTACATT GTCTTCA
24	TGF- $\beta$ 1	ss12568613	C-T	5' UTR		31.1	GGGCCAGTTTCCCTAT CTGTAAA	CTGGGCCACCGTCCTC ATC
25	TGF- $\beta$ 1	ss12568603	+C/-C	Intron 4	12,354	2.1	CCACGCCCCACTTATCT ATCCC	GGAAAGGCCGGTTCAT GCCA
26	TGF- $\beta$ 1	ss12568593	C-T	Exon 5	83	0.8	CAGGCTACAAGGCTCA CCTGAA	GGTTCACTACCGGCCG C
27	TGF- $\beta$ 1	ss12568602	C-T	Intron 5	9,644	27.4	GGCTTGCTTAAGCATT GCGTGAAATTA	GTACAGTGCCGCACG C
28	UCP3	ss12568601	C-T	5' UTR		26.5	CACTGCCCTCACCAGC CA	GTGAGTCTGCCACGG CA
29	UCP3	ss12568600	G-A	Exon 2	2,122	0.0	GCCCTAAAGGGACTGG GCA	GAAAGGTAACGAGGTC AGCAAAAACA

**Table 1: Information about the 37 SNPs in the eight genes (Continued)**

30	UCP3	ss12568595	T-C	Exon 3	809	25.8	TGATTCCCCTAACATCT GGACTTTCATC	CTGCCTAAATCCCCTTA GCAGAAAAAAAAA
31	UCP3	ss12568580	T-C	Exon 5	1,712	44.7	CCTAACAGGAACCTTGC CCAACATCA	TCCACGGAGTTCTGGGT TCC
32	UCP3	ss12568599	C-T	Exon 7	2,996	0.0	CCTAACAGGAACCTTGC CCAACATCA	TCCACGGAGTTCTGGGT TCC
33	VDR	ss12568583	G-A	5' UTR		28.1	CAGCATGCCTGTCCTCA GC	CCAGTACTGCCAGCTC CCA
34	VDR	ss12568581	C-T	Exon 2	4,818	37.3	TGGCCCTGGCACTGAC TC	GGCACGTTCCGGTCAA AGTC
35	VDR	ss12568582	C-T	Exon 4	21,590	0.0	GGACAGTCTGCGGCC A	CCCTACTCCCTGGGCC C
36	VDR	ss12568610	G-A	Intron 8	11,470	41.9	GTGCCCTCACTGCC TTA	CCTCAAATAACAGGAAT GTTGAGCCCA
37	VDR	ss12568608	T-C	Exon 9	1,078	40.8	GGGCCAGGCAGTGGTA TCAC	AGGTCGGCTAGCTTCTG GATCA

a: Minor alleles are given in italic. b: Distance from the previous SNP within each gene in the unit of bp. c: Frequency of minor allele

**Table 2: Number of main haplotypes observed within genes compared to the number of possible haplotypes and yin yang haplotype distribution**

Gene	Chromosome location	No. of SNPs	Kb spanned	SNP density (kb/SNP)	Observed No. of haplotypes <sup>a</sup>	Expected No. of haplotypes	Yin Yang haplotypes <sup>b</sup>
ApoE	19q13.2	4	3	0.75	6	16	GATC, CGTC
COL1A1	17q21.3-q22	3	16	5.33	4	8	GGG, GTG
ER- $\alpha$	6q25.1	7	291	41.57	20	128	TCGCGGG, CTGCGGG
LEPR	1p31.2	3	66	22	7	8	GCC, GGA
PTHRI	3p22-p21.1	4	9	2.25	7	16	GTA, ACGC
TGF- $\beta$ 1	19q13.1	3	22	7.33	4	8	C+C, T+C-
UCP3	11q13.4	3	5	1.67	4	8	TCT, CTC
VDR	12q12-14	4	39	9.75	12	16	GTGT, GTAC

a: Only haplotypes with frequencies more than 1% were included [14] b: For the PTHRI and UCP3, Yin yang haplotypes were presented. For the other six genes, the two most common haplotypes were presented. The haplotypes were reconstructed based on the SNPs in the order from 5' to 3' on the chromosome within each gene.

highest value of 0.085 in PTHRI and the lowest value of 0.001 in TGF- $\beta$ 1. The Spearman correlation coefficient between the two LD measures was 0.81 ( $P < 0.01$ ).


**Discussion**

Determining the haplotype and LD characteristics of candidate genes and/or genomic regions has important implications for strategies of complex diseases gene mapping [19]. In the present study, we illustrated the haplotype distribution and LD patterns in eight candidate genes for two complex diseases, obesity and osteoporosis, in 1873 subjects from 405 Caucasian nuclear families.

In this study, the yin yang haplotype pairs, mismatching at every SNP, were observed in two genes, PTHRI and UCP3, but not in the other six. There are several possible reasons for such differences among genes. Other things being the same, lower LD levels result in smaller yin yang

haplotype size, the genomic span per yin yang haplotype region [15]. The different SNP density and LD patterns among genes may be partly responsible for the discrepancies about yin yang haplotypes among genes. Lower minor allele frequencies in some genes may be another reason since minor allele frequency  $\geq 10\%$  is a threshold for yin yang haplotype analysis [15]. Moreover, yin yang haplotypes were affected by the factors that might be different among genes, such as the mutation and recombination rates [15].

So far, LD was inferred either by using unrelated random sample or family data. The LD values evaluated from these two different data sets were correlated, but in some instances they were quite different. Family subjects contain more information than unrelated random individuals, and the inference of LD using family data is more accurate [4]. Sample size is another important factor



**Figure 1**

An exponential decay function was employed to fit the relationship between LD and the physical distance between SNP markers within each gene (SigmaPlot 2000, SPSS Inc., Chicago, IL). X-axis is the physical distance in the unit of base pair.

influencing the accuracy of LD evaluation. Generally, larger sample size can minimize sampling error and produce more accurate evaluation of LD [20]. In this study, haplotypes were reconstructed based on the nuclear families in a larger sample rather than unrelated individuals as the other studies [12,18,21,22]. This increased the precision of LD evaluation.

In the present study, two most common measurements of LD,  $D'$  and  $r^2$ , were used, both measurements varying between 0 and 1. According to their calculating formula [23], they are positively correlated, as observed in this study, but sometimes they are quite different. For example,  $D'$  value for SNP6-SNP8 was 1.000, while the value

for  $r^2$  was only 0.004. Such a discrepancy between the two measures was also observed by Turet et al [17]. One possible explanation for the discrepancy is that the rare allele frequency of SNP8 was very low, only 1.9%, and  $r^2$  is more sensitive to allele frequencies than  $D'$  [23].

Although a trend of decay of LD with increasing physical distance was observed in the present study, there was a great variation in LD with distance. For example, the physical distances for SNP33-SNP34 in VDR and SNP20-SNP21 in PTHR1 were both about 5 kb, however, the  $D'$  values were 0.062 and 0.964, and the  $r^2$  were 0.004 and 0.826, respectively. In the ER- $\alpha$  gene, the physical distance of SNP14-SNP15 was about 2.6 times that of SNP13-SNP14, while the  $D'$  values were 0.675 and 0.028, and  $r^2$  were 0.599 and 0.019, respectively. Such irregularity has been observed in the other studies [6,17,18,24] and is expected, since LD is also affected by other factors, such as natural selection, the rates of recombination and mutation, and gene conversion [25].

Several limitations of this study should be acknowledged. Compared with some recent analyses [4,14,26], we only selected a limited number of SNPs in each gene, which may bring some difficulty to define the LD characteristics. Though 15-kb chromosome-wide resolution for LD pattern and haplotype structure study is acceptable but not ideal [27]. In addition, the yin yang haplotype analysis might be affected by the limited SNPs and genes examined. The SNPs were chosen based on the information from available literature and public databases rather than evenly distributed within gene. Another limitation is that this study was restricted to Caucasians of European descent, which limited the ability to perform cross-population comparisons for yin yang haplotype coverage and haplotype diversities etc. Since genetic diversity and LD extent differ between populations, studying contrasted populations is informative. However, if the purpose is to provide guidelines for the design of association studies in a given population, information drawn from that population should be preferable [5], especially because Caucasians are more susceptible to osteoporosis and obesity compared with other populations [28,29].

In conclusion, the haplotype distribution was different among the eight studied candidate genes for osteoporosis and/or obesity. The LD characteristics were different between genes or even at different regions within gene, though LD decays with the increase of physical distance. The differences in haplotype distributions and LD patterns among genes underscore the importance of characterizing each locus of interest prior to association studies.

## Methods

### Subjects

The study subjects came from the ongoing genetic studies of osteoporosis and obesity that have been approved by the Creighton University Institutional Review Board. All subjects were Caucasians of European origin. We have recruited 405 nuclear families composed of 1,873 healthy individuals, including 840 parents, 744 daughters and 389 sons. All individuals signed informed-consent documents before entering the project.

### Genotyping

After searching public SNP databases, such as dbSNP <http://www.ncbi.nlm.nih.gov/SNP/>, and available literature, 37 SNPs in the eight candidate genes for obesity and/or osteoporosis were chosen. The selection criteria were as follows: 1) functional relevance and importance; 2) position in or around the gene; and 3) their use in previous genetic studies. Information about the 37 SNPs is presented in Table 1. The genotyping procedure for all SNPs was similar, involving polymerase chain reaction (PCR) and invader assay reaction (Third Wave Technology, Madison, WI). PCR was performed in a 10  $\mu$ l reaction mixture. The sequences of the PCR primers for all 37 SNPs are presented in Table 1. After amplification, an invader reaction was performed in a 7.5  $\mu$ l reaction volume. The genotype for every sample was called according to the ratio of the fluorescence intensity of the two dyes, which was read using a Cytofluor 4000 multi-well plate reader (Applied Biosystems, Foster City, CA). Program PedCheck [http://watson.hgen.pitt.edu/register/soft\\_doc.html](http://watson.hgen.pitt.edu/register/soft_doc.html) was employed to verify Mendelian inheritance of all marker alleles within each family [30].

We initially genotyped all 37 SNPs in a random sample of 190 to 380 subjects. The minor allele frequency of six SNPs was found less than 1.0%. Because the precision and variance of LD estimates suffer from low allele frequencies [6], these six SNPs were excluded from further genotyping and data analysis. Finally, 31 SNPs were analyzed in this study. Among them, 14 were located in exons, 12 in introns, and 5 in untranslated regions. SNP23 in the TGF- $\beta$ 1 gene is an insertion/deletion polymorphism of a cytosine (+/- C) and the other 30 SNPs are nucleotide substitutions, including 25 transitions and 5 transversions. These 31 SNPs totally spanned ~451 kb. The average inter-marker distance was 63 kb (ranging from 138 bp to 291 kb).

### Statistical analysis

Allele frequencies of each SNP were estimated in 1,873 subjects using a maximum likelihood method implemented in the program SOLAR which is available at <http://www.sfbr.org/sfbr/public/software/solar>. The haplotypes of each gene for all subjects were reconstructed

using the program Genehunter version 2.1 <http://www.hgmp.mrc.ac.uk/Registered/Option/genehunter.html>. Genehunter extracts complete multipoint inheritance information to infer maximum likelihood haplotypes for all individuals in nuclear families [31]. Haplotype frequencies were estimated from the unrelated subjects (parents of the nuclear families).

Two normalized measures,  $D'$  and  $r^2$ , were used to characterize the LD patterns within the studied candidate gene. Pair-wise  $D'$  values are computed as:  $D' = D/D_{\max}$ ,  $D_{\max} = \min(f_{1+}f_{+2}, f_{+1}f_{2+})$  when  $D > 0$ , and  $D_{\max} = \min(f_{1+}f_{+1}, f_{+2}f_{2+})$  when  $D < 0$ , where  $f$ s are sample estimates of SNP frequencies [32].  $r^2$  is quantified as  $r^2 = (f_{11}f_{22} - f_{12}f_{21})^2 / [(f_{11} + f_{12})(f_{21} + f_{22})(f_{11} + f_{21})(f_{12} + f_{22})]$ , where  $f$ s are sample estimates of haplotype frequencies [23]. An exponential decay function was employed to fit the relationship between LD (both  $D'$  and  $r^2$ ) and the physical distance between SNP markers within each gene (SigmaPlot 2000, SPSS Inc., Chicago, IL). To quantify the relationship between  $D'$  and  $r^2$  both as index of LD, correlation coefficient between them was calculated using SPSS software.

### Author's contribution

JRL performed SNP genotyping, combined the results of analysis and drafted the manuscript. LJZ performed SNP genotyping and data analysis. PYL and YL performed statistical analysis. VD contributed to the manuscript preparation. HS, YJL, YYZ, DHX and PX performed SNP genotyping. HWD conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

### Acknowledgements

The investigators were partially supported by grants from Health Future Foundation of the USA, the National Institute of Health, the State of Nebraska Cancer and Smoking Related Disease Research Program, the State of Nebraska Tobacco Settlement Fund, US Department of Energy, Chinese National Science Foundation, and the Ministry of Education of P. R. China.

### References

1. Ardlie KG, Seielstad M: **Patterns of linkage disequilibrium in the human genome.** *Nat Rev Genet* 2002, **3**:299-309.
2. Reich DE, Cargill M, Bolk S, Ireland J, Sabeti PC, Richter DJ, Lavery T, Kouyoumjian R, Farhadian SF, Ward R, Lander ES: **Linkage disequilibrium in the human genome.** *Nature* 2001, **411**:199-204.
3. Wall JD, Pritchard JK: **Haplotype blocks and linkage disequilibrium in the human genome.** *Nat Rev Genet* 2003, **4**:587-597.
4. Moffatt MF, Traherne JA, Abecasis GR, Cookson WO: **Single nucleotide polymorphism and linkage disequilibrium within the TCR alpha/delta locus.** *Hum Mol Genet* 2000, **9**:1011-1019.
5. Stephens JC, Schneider JA, Tanguay DA, Choi J, Acharya T, Stanley SE, Jiang R, Messer CJ, Chew A, Han JH, Duan J, Carr JL, Lee MS, Koshy B, Kumar AM, Zhang G, Newell WR, Windemuth A, Xu C, Kalbfleisch TS, Shaner SL, Arnold K, Schulz V, Drysdale CM, Nandabalan K, Judson RS, Ruano G, Vovis GF: **Haplotype variation and linkage disequilibrium in 313 human genes.** *Science* 2001, **293**:489-493.
6. Abecasis GR, Noguchi E, Heinzmann A, Traherne JA, Bhattacharyya S, Leaves NI, Anderson GG, Zhang Y, Lench NJ, Carey A, Cardon LR, Moffatt MF, Cookson WO: **Extent and distribution of linkage**

- disequilibrium in three genomic regions. *Am J Hum Genet* 2001, **68**:191-197.
7. Risch N, Merikangas K: **The future of genetic studies of complex human diseases.** *Science* 1996, **273**:1516-1517.
  8. Weiss KM, Clark AG: **Linkage disequilibrium and the mapping of complex human traits.** *Trends Genet* 2002, **18**:19-24.
  9. Jorde LB: **Linkage disequilibrium and the search for complex disease genes.** *Genome Res* 2000, **10**:1435-1444.
  10. Teare MD, Dunning AM, Durocher F, Rennart G, Easton DF: **Sampling distribution of summary linkage disequilibrium measures.** *Ann Hum Genet* 2002, **66**:223-233.
  11. Stephens M, Smith NJ, Donnelly P: **A new statistical method for haplotype reconstruction from population data.** *Am J Hum Genet* 2001, **68**:978-989.
  12. Bonnen PE, Wang PJ, Kimmel M, Chakraborty R, Nelson DL: **Haplotype and linkage disequilibrium architecture for human cancer-associated genes.** *Genome Res* 2002, **12**:1846-1853.
  13. Zhu X, Yan D, Cooper RS, Luke A, Ikeda MA, Chang YP, Weder A, Chakravarti A: **Linkage disequilibrium and haplotype diversity in the genes of the renin-angiotensin system: findings from the family blood pressure program.** *Genome Res* 2003, **13**:173-181.
  14. Croucher PJ, Mascheretti S, Hampe J, Huse K, Frenzel H, Stoll M, Lu T, Nikolaus S, Yang SK, Krawczak M, Kim WH, Schreiber S: **Haplotype structure and association to Crohn's disease of CARD15 mutations in two ethnically divergent populations.** *Eur J Hum Genet* 2003, **11**:6-16.
  15. Zhang J, Rowe WL, Clark AG, Buetow KH: **Genomewide Distribution of High-Frequency, Completely Mismatching SNP Haplotype Pairs Observed To Be Common across Human Populations.** *Am J Hum Genet* 2003, **73**:1073-1081.
  16. Kruglyak L: **Prospects for whole-genome linkage disequilibrium mapping of common disease genes.** *Nat Genet* 1999, **22**:139-144.
  17. Tiret L, Poirier O, Nicaud V, Barbaux S, Herrmann SM, Perret C, Raoux S, Francomme C, Lebard G, Tregouet D, Cambien F: **Heterogeneity of linkage disequilibrium in human genes has implications for association studies of common diseases.** *Hum Mol Genet* 2002, **11**:419-429.
  18. Nakajima T, Jorde LB, Ishigami T, Umemura S, Emi M, Lalouel JM, Inoue I: **Nucleotide diversity and haplotype structure of the human angiotensinogen gene in two populations.** *Am J Hum Genet* 2002, **70**:108-123.
  19. Weale ME, Depondt C, Macdonald SJ, Smith A, Lai PS, Shorvon SD, Wood NW, Goldstein DB: **Selection and evaluation of tagging SNPs in the neuronal-sodium-channel gene SCN1A: implications for linkage-disequilibrium gene mapping.** *Am J Hum Genet* 2003, **73**:551-565.
  20. Guo SV: **Linkage disequilibrium measures for fine-scale mapping: a comparison.** *Hum Hered* 1997, **47**:301-314.
  21. Peterson RJ, Goldman D, Long JC: **Effects of worldwide population subdivision on ALDH2 linkage disequilibrium.** *Genome Res* 1999, **9**:844-852.
  22. Albagha OM, Tasker PN, McGuigan FE, Reid DM, Ralston SH: **Linkage disequilibrium between polymorphisms in the human TNFRSF1B gene and their association with bone mass in perimenopausal women.** *Hum Mol Genet* 2002, **11**:2289-2295.
  23. Devlin B, Risch N: **A comparison of linkage disequilibrium measures for fine-scale mapping.** *Genomics* 1995, **29**:311-322.
  24. Taillon-Miller P, Bauer-Sardina I, Saccone NL, Putzel J, Laitinen T, Cao A, Kere J, Pilia G, Rice JP, Kwok PY: **Juxtaposed regions of extensive and minimal linkage disequilibrium in human Xq25 and Xq28.** *Nat Genet* 2000, **25**:324-328.
  25. Ardlie K, Liu-Cordero SN, Eberle MA, Daly M, Barrett J, Winchester E, Lander ES, Kruglyak L: **Lower-than-expected linkage disequilibrium between tightly linked markers in humans suggests a role for gene conversion.** *Am J Hum Genet* 2001, **69**:582-589.
  26. Zabetian CP, Buxbaum SG, Elston RC, Kohnke MD, Anderson GM, Gelernter J, Cubells JF: **The structure of linkage disequilibrium at the DBH locus strongly influences the magnitude of association between diallelic markers and plasma dopamine beta-hydroxylase activity.** *Am J Hum Genet* 2003, **72**:1389-1400.
  27. Dawson E, Abecasis GR, Bumpstead S, Chen Y, Hunt S, Beare DM, Pabial J, Dibling T, Tinsley E, Kirby S, Carter D, Papaspyridonos M, Livingston S, Ganske R, Lohmussaar E, Zernant J, Tonissou N, Remm M, Magi R, Puurand T, Vilo J, Kurg A, Rice K, Deloukas P, Mott R, Met-
- spalu A, Bentley DR, Cardon LR, Dunham I: **A first-generation linkage disequilibrium map of human chromosome 22.** *Nature* 2002, **418**:544-548.
28. Liu YZ, Liu YJ, Recker RR, Deng HW: **Molecular studies of identification of genes for osteoporosis: the 2002 update.** *J Endocrinol* 2003, **177**:147-196.
  29. Chagnon YC, Rankinen T, Snyder EE, Weisnagel SJ, Perusse L, Bouchard C: **The human obesity gene map: the 2002 update.** *Obes Res* 2003, **11**:313-367.
  30. O'Connell JR, Weeks DE: **PedCheck: a program for identification of genotype incompatibilities in linkage analysis.** *Am J Hum Genet* 1998, **63**:259-266.
  31. Kruglyak L, Daly MJ, Reeve-Daly MP, Lander ES: **Parametric and nonparametric linkage analysis: a unified multipoint approach.** *Am J Hum Genet* 1996, **58**:1347-1363.
  32. Lewontin RC: **The interaction of selection and linkage. I. General considerations; heterotic models.** *Genetics* 1964, **49**:49-67.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

