



Published in final edited form as:

J Am Coll Health. 2014 ; 62(8): 534–541. doi:10.1080/07448481.2014.933344.

Sleep Quality and Body Mass Index in College Students: The Role of Sleep Disturbances

Perla A. Vargas, PhD^{1,2}, Melissa Flores, BA², and Elias Robles, PhD²

¹Southwest Interdisciplinary Research Center, Arizona State University, Glendale, Arizona

²School of Social and Behavioral Sciences, Arizona State University, Glendale, Arizona

Abstract

Objective—Obesity and its comorbidities have emerged as a leading public health concern. Our aim was to explore the relationship between BMI and sleep patterns, including duration and disturbances.

Methods—A convenience sample of 515 college students completed an online survey consisting of the Pittsburgh Sleep Quality Index (PSQI), and self-reported height and weight to calculate Body Mass Index (BMI). Univariate and multivariate logistic regression analyses were performed using components of the PSQI as predictors of overweight (BMI ≥ 25).

Results—One-third of the participants had BMI ≥ 25 , and 51% were poor-quality sleepers (PSQI > 5). Controlling for age and sex, only sleep disturbances were associated to overweight (OR=1.66, 95% CI: 1.08-2.57).

Conclusions—Sleep disturbances, rather than sleep duration predicted overweight among young adults; this is consistent with the most recent evidence in the literature. These findings support expanding the scope of wellness programs to promote healthy sleep among students.

Keywords

body mass index (BMI); college students; overweight; habits; prevalence; sleep; sleep disturbance; sleep duration; sleep quality

College is a time marked by change, when many students experience autonomy and freedom from direct supervision for the first time, greater academic and social pressures, and erratic schedules. Major changes in lifestyle such as unbalanced diets, alcohol intake, and lack of sleep can be detrimental to the students' well-being.¹⁻³

The prevalence of overweight and obesity, measured as Body Mass Index (BMI ≥ 25), has increased dramatically over the past decade.⁴⁻⁸ For example, in 2008 the prevalence of obesity (i.e., BMI ≥ 30) among U.S. adults 20 years or older was 26.7%, 6.9% greater than the estimate for 2000.⁷ Furthermore, the combined estimated prevalence of overweight and

Note: For comments and further information, address correspondence to Perla A. Vargas, Southwest Interdisciplinary Research Center and School of Social and Behavioral Sciences, Arizona state University, 4701 W Thunderbird Rd, MC 3051, Glendale, AZ 85306-4908, USA (perla.vargas@asu.edu).

The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIMHD or the NIH.

obesity (BMI ≥ 25) for 2000 among adults was 68%.⁷ This increase in BMI has been observed across all ages, with overweight and obesity in children and adolescents reaching 31.8% in 2010.⁸ Similarly elevated rates have been estimated among college students; with over one-third reporting BMI ≥ 25 .⁹⁻¹¹

Concurrent with the increase in overweight and obesity among the U.S. population, there has been a change in sleeping patterns.^{12;13} Insomnia and sleep difficulties have been widely studied in the general population.¹⁴ Research shows that, on average, adults sleep less than the recommended 8 hours per night^{9;15} Furthermore, the percentage of adults sleeping 6 hours or less each night increased from 22% in 1985 to nearly 28% in 2007.¹⁶ Unfortunately, as Lund and colleagues¹⁷ point out, research describing the changes in the sleeping patterns and needs of college-age adolescents is limited. Most of the studies examining sleep patterns among college students have focused on the relationship between sleep habits and academic performance,¹⁸⁻²⁰ and mental health problems (i.e., stress, depression and suicidal ideation).²¹⁻²⁴ All the studies concede that college students experience frequent and sometimes severe sleep problems that negatively impact their health and well-being.¹⁷⁻²⁴ Overall, estimates of sleeping problems/disturbances among college students ranged between 13.8% and 68.6% depending on the definition and measurement used.²⁰⁻²¹ In a 2012 national survey of college students (n = 99,066), 26.4% reported sleep difficulties during the previous 12 months as being “traumatic or very difficult to handle”, and 57.1% of the sample reported having enough sleep to feel rested on fewer than four days a week.⁹ In a study examining sleep/wake patterns over the transition from high school to college, Carskadon and Davis²⁵ found a significant reduction in the number of hours of sleep, as well as delay in onset of nighttime sleep, as students transition to college. In addition to the amount of sleep, the quality of sleep seems to be changing and impacting the well-being of college students.^{17;19;26} Using the Pittsburgh Sleep Quality Index (PSQI), Lund identified 63.9% of college students in her sample (N=1,125) as poor-quality sleepers, those who experience restricted total sleep, erratic sleep schedules, and/or sleep disturbances.¹⁷ Along the same line, Gaultney¹⁹ found that 27% of college students (N = 1,845) were at risk for sleep disorders likely to disturb sleep. When Pilcher and co-investigators²⁶ looked at the impact of sleep quantity and quality on the health of college students, they found sleep quality to be a better predictor than sleep quantity of several measures of health including health complaints, depression, fatigue and sleepiness. Results from these studies reveal an unquestionable problem of insufficient and poor quality sleep among adolescents and young adults, as well as growing levels of sleep dissatisfaction.²⁷ The number of people reporting poor quality sleep seems to be reaching “epidemic” proportions, with 73% of college students reporting occasional sleep problems that include insufficient sleep, difficulty falling asleep, sleep disturbances, and excessive daytime sleepiness.^{20;28}

Although it is well established that an imbalance between caloric intake and physical activity are key factors responsible for the current obesity problem,²⁹⁻³⁰ there is emerging evidence suggesting that other factors may be important contributors to the obesity problem.³⁰⁻³² One of the factors in question is sleep.²⁹⁻⁵⁹

Unlike other mammals, humans have consolidated sleep in periods of 7-9 hours. It has been shown that during normal sleep, glucose utilization and glucose production drop simultaneously during the first part of the night and raise simultaneously during the predawn hours.⁶⁰⁻⁶² Scheen found, however, that when subjects are maintained at rest but sleep-deprived, the pattern of glucose utilization is altered, and glucose levels become constant throughout the night.⁶⁰ Following Scheen's landmark study,⁶⁰ a number of studies have shown that the metabolic changes associated with interference of normal sleep patterns may contribute to the development of obesity, cardiovascular disease, insulin resistance, and diabetes.^{33-41;46;56;58-65} Cross-sectional research evaluating the relationship between obesity and sleeping patterns suggests a negative correlation between overweight/obesity and/or related metabolic markers, and sleep duration among children,⁴²⁻⁴⁹ adolescents,^{40;49-52} and adults.⁵³⁻⁵⁹ Moreover, research studies exploring the association between sleep and BMI among adolescents suggest that the relationship might be multilayered.⁶⁶⁻⁶⁸ For example, in a sample of American adolescents,⁶⁷ researchers found that the likelihood of obesity increased by 80% for each hour of lost sleep. Although, sleep disturbances were not directly related to obesity; they were negatively correlated with level of physical activity, suggesting that sleep disturbances could indirectly impact body weight through a reduction in physical activity. On the other hand, no relationship between level of physical activity and sleep duration was found. Similarly, in a study of European adolescents, investigators found that short sleep duration was associated with obesity, increased food intake, and sedentary habits, but not with level of physical activity.⁶⁸ But not all the emerging evidence supports a relationship between short sleep duration and increased BMI; some studies have found no association between sleep duration and BMI,⁶⁹⁻⁷³ while others have found a U-shaped relationship with the odds rising for those with shorter and longer sleep duration.⁷³⁻⁷⁶ In many studies, however, the potential effects of sleep loss cannot be distinguished from the effects of sleep disturbances;⁷⁷ more recent laboratory studies have solved this problem by measuring metabolic changes while controlling for sleep duration and sleep disturbances.^{36;37;41} For example, Tasali and collaborators⁷¹ found that in young healthy adults, suppression of deep non-rapid eye movement sleep without concomitant changes in total sleep time resulted in decreases in insulin sensitivity, which lead to reduced glucose tolerance and increased risk of diabetes.⁴¹ Additional evidence supporting the hypothesis that sleep fragmentation, independent of sleep duration, can alter glucose homeostasis was obtained by Stamatakis & Punjabi.³⁶ In their study with healthy adults, total sleep time was held within normal range (7-7.5 h) while inducing fragmented sleep with auditory and mechanical stimuli across all stages of sleep. The study showed a reduction in insulin sensitivity and glucose effectiveness associated with sleep fragmentation, supporting the hypothesis regarding the negative effects of sleep disturbances on glucose metabolism. These findings are particularly relevant to college students due to the nature of *college culture*. Although there might be significant variations from campus to campus, college culture is characterized by social arrangements and behavior practices that negatively affect students' sleep, including communal living, frequent parties and drinking, poor time management, and high levels of stress and work.⁷⁸⁻⁸⁰

The present study was designed to evaluate the relationship between parameters of sleep, including sleep duration and disturbance, and BMI. Based on the evidence that sleep

fragmentation alters metabolic functions involved in appetite control, our hypothesis was that sleep disturbances, rather than sleep duration, would be associated with BMI ≥ 25 among college students attending a public university.

Method

Participants

The sample consisted of 536 college students from a large university in the southwestern U.S. who completed the study to satisfy a class requirement. Data were collected between October 22, 2010 and May 6, 2011. Twenty one non-traditional students over 34 years of age were excluded, yielding a sample of 515 participants between 18 to 34 years, mostly females (73.2%). The study was approved by the University Committee on Human Subjects. No personal identifiers were collected and, after reading a description of the study, participants gave passive consent by proceeding to complete the survey.

Materials

The PSQI was utilized to assess sleeping patterns. The PSQI is a standardized, quantitative measure of sleep quality with demonstrated high levels of consistency, reliability, and validity.⁸¹⁻⁸⁴ It is composed of 19 self-reported questions grouped into seven component scores. The seven components were scored following the algorithm proposed by Buysse.⁸¹ Accordingly, each component score is weighted equally on a 0-3 scale with lower scores indicating no problems and higher scores indicating progressively worsening problems as follows: 1) subjective sleep quality (very good to very bad), 2) sleep latency (≤ 15 minutes to > 60 minutes), 3) sleep duration (≥ 7 hours to < 5 hours), 4) sleep efficiency ($\geq 85\%$ to $< 65\%$ hours sleep/hours in bed), 5) sleep disturbances (not during the past month to ≥ 3 times per week), 6) use of sleeping medications (none to ≥ 3 times a week), and 7) daytime dysfunction (not a problem to a very big problem). The seven component scores were added to yield a global score ranging from 0 to 21 that was treated as a continuous variable. The reliability of the PSQI Global score was satisfactory in our sample, yielding a Cronbach's α of 0.702. For clinical use, Buysse and colleagues proposed a cut-off score of ≤ 5 to indicate good quality sleep and > 5 to indicate poor sleep quality.⁸¹ Overweight and obesity were defined as recommended by the Clinical Guidelines for the Identification, Evaluation and Treatment of Overweight and Obesity in adults.⁸⁵ BMI was calculated using self-reported height and weight as: $\text{weight (lb)} / [\text{height (in)}]^2 \times 703$. A dichotomous variable was created using BMI ≥ 25 as cutoff to identify those exceeding the recommended weight (overweight and obese). Self-reported and clinically measured height and weight have been found to be highly correlated; but with an underreporting bias of about one unit.⁸⁶⁻⁸⁷ Although, self-reported BMI is not recommended as a clinical tool; it is considered an important health surveillance tool that is valid for epidemiologic research,⁸⁷ in particular with younger adults.⁸⁸ Specifically, self-reported height and weight are currently used in the National Health and Nutrition Examination Survey to estimate rates of overweight and obesity for the U.S. population.⁸⁸

Procedure

All data were collected online. Participants were directed to complete the College Student's Health Profile Questionnaire. This survey consisted of questions addressing demographics, self-reported height and weight, and several standardized health-related measures including the PSQI. Seventy-three students missed at least one question from the PSQI and were excluded from some analyses. As a result, the number of subjects included in the calculations of the descriptive estimates varied, ranging from 442 in the calculations of the PSQI Global score, and 498 - 506 subjects included in the calculations of all other components of the PSQI. Descriptive statistics, chi square, and logistic regression analyses were performed using IBM SPSS 20 (IBM Corp. Released 2011. IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp). We use the Odds Ratio (OR) and 95% confidence intervals (95% CI) to show the estimated effect of the PSQI global score and each one of the seven components on BMI; ORs greater than one indicating increased risk.

Results

Participants were mostly females (73.2%) with a mean age of 21.68 years (SD=3.49). Most participants identified themselves as Caucasian (63.9%), or Hispanic (18.4%). Over 30% (33.3%) of the participants had a BMI ≥ 25 . Of these participants, 102 were classified as overweight (BMI $25 < 30$) and 67 as obese (BMI ≥ 30). BMI rates varied by sex; with more males reporting BMI ≥ 25 than females (41.9% vs. 30.1%, $p < .05$).

Sleep Pattern

Results of the PSQI are presented in Table 1. The mean global PSQI score was 5.9 ± 2.8 (median 5.78, range .89-16.36) with 51% of the participants experiencing poor sleep quality (> 5), and 19% reporting extremely poor sleep quality (≥ 8). However, when asked to subjectively rate their quality of sleep, only 30.7% rated it as fairly bad or very bad. The most important components contributing to poor sleep quality included: restricted sleep time, long sleep latency, poor sleep efficiency, and daytime dysfunction. Overall, most students reported restricted sleep; only 30.11% of students reported the recommended eight or more hours of sleep, and 40.6% reported six hours or less per night (mean 6.8, SD=1.39). Almost 40% of participants (39.6%) reported taking more than 30 minutes to fall asleep (mean 29.5, SD=42 min); with one-third (33.07%) reporting sleep efficiency below 85% (i.e., percent of the time asleep from the moment they go to bed until they get up). Perhaps as a consequence of the poor quality of sleep, 27.9% of the students reported daytime dysfunction at least once a week. Nearly one-fourth of the students (24.3%) reported taking sleep medication in the previous month, and 13.2% reported taking sleep medication at least once a week. Most participants reported some sleep disturbances (93.0%), and 16.1% were classified as experiencing high levels of sleep disturbances according to the PSQI algorithm (Table 1). Some of the students' responses to specific questions on the PSQI are worth mentioning (Table 1); 55.03% reported *waking up in the middle of night or early morning* at least once a week, 32.7% reported *lacking enthusiasm to get things done* at least once a week, and 28% reported *experiencing bad dreams* at least once a week. Of the students who reported "other" reasons for sleep disturbances at least once a month, the most common was "stress-related" (e.g., stress, racing thoughts, worried).

Relationship between BMI and PSQI

Univariate logistic regression analyses were performed to examine the relationship between increased BMI (≥ 25) and PSQI global score, as well as the seven individual components as predictor variables. Table 2 shows that, of all the possible components entered, only the *sleep disturbance* component was a significant ($p < .05$) predictor of overweight/obesity (BMI ≥ 25). After controlling for age and sex, a multivariate logistic regression analysis revealed a 66% increase in the probability of being overweight as sleep disturbances increased (OR=1.66, 95% CI: 1.08-2.57). However, there was a small interaction effect between age and sleep disturbance (OR=1.03, 95% CI: 1.00-1.05), suggesting that sleep disturbances may become more serious with age (3% increase per year of age), even within the narrow range of the sample.

Comment

Conclusions

Although it is well established that an imbalance between caloric intake and physical activity are key factors responsible for the current obesity problem;²⁹⁻³⁰ there is growing interest in studying the environmental and behavioral factors that may be contributing to the problem.³¹⁻³² Emerging evidence suggests an association between body weight and problems in sleeping patterns, particularly its duration and quality.^{39;42;47-51;54;57;67;68} Furthermore, research studies exploring the association between sleep and BMI among adolescents, suggest the relationship is complex.⁶⁶⁻⁶⁸ As discussed by Pilcher,²⁶ in many studies sleep duration and disturbances are combined to estimate sleep quality, making it difficult to evaluate their independent effects on BMI. We used the PSQI global score and its seven individual components (i.e., subjective quality, latency, duration, efficiency, disturbances, medication, and daytime dysfunction) to examine the sleep patterns of college students, and to evaluate the relationship between the different components of sleep quality and overweight/obesity. Overall, the data revealed a major problem with the sleep pattern of college students. In addition to short sleep duration, college students experienced fragmented sleep, as well as poor sleep quality as measured by the PSQI. Consistent with the hypothesis that sleep fragmentation may be responsible for increased body weight, our results show that after controlling for age and sex, sleep duration was not a significant predictor for BMI; but sleep disturbances were. Probing further, we also found an interaction between sleep disturbances and age suggesting that, as students age, sleep disturbances may have a larger effect on BMI. That is, regardless of the amount of time spent sleeping, sleep disruptions may have a significant influence on weight, and this influence may be augmented by age. These findings are consistent with the hypothesis that sleep restrictions alter the circulating levels of metabolically relevant hormones such as leptin and ghrelin, resulting in alterations of glucose homeostasis and appetite regulation.^{6;33-37;40;41} On the other hand, the interaction effect of age and sleep disturbances is congruent with evidence showing age-related changes in sleep timing and structure.⁸⁹⁻⁹¹ In a meta-analysis conducted to derive normative data on the effects of aging on sleep, Ohayon⁹² found that as we age total sleep time, sleep efficiency, percentage of slow-wave sleep, percentage of REM sleep, and REM latency significantly decrease, while sleep latency, and wake after sleep onset significantly increase.

Our results suggest a need to attend to the quality of sleep, particularly in young adults. Furthermore, there is evidence that sleeping problems for many adult insomniacs begins early in life.⁹³⁻⁹⁵ In addition to effects on grades and daily activities, poor-quality sleep puts college students at risk for health problems of long lasting consequences, including excessive body weight. In turn, excessive body weight brings with it an increased risk for diabetes, cardiovascular disease, and some cancers. For example, in a recent meta-analysis, researchers identified 18 co-morbidities associated with excess body weight. Of these, type 2 diabetes was estimated to have the greatest burden, followed by cardiovascular diseases, and several forms of cancer (e.g., esophageal, colon, endometrial gall bladder and breast cancer).⁹⁶ Excess bodyweight has also been associated to non-fatal disorders that impact the patients' quality of life, such as osteoarthritis and asthma.⁹⁷ In the near future, the increased morbidity associated with excess body weight may have a negative impact on life expectancy, as suggested by Calle.⁹⁸ Calle's study of more than one million American adults found that among people who never smoked, heavier men and women had a higher mortality risk from all causes, while those with a BMI < 25 had the lower mortality risk.⁹⁵ Estimates based on current trends in the age of onset of excess body weight suggest that chronic conditions and disabilities associated to excess weight may increase dramatically in the younger generations, with negative consequences to their quality of life and life expectancy.⁹⁹

A large proportion of college students experience sleep disturbances. This study shows that sleep disturbances are significantly linked to overweight and obesity. Our findings suggest the need to expand efforts of wellness programs and health initiatives to promote healthy sleeping habits among college students. These initiatives ought to also include educational components supporting healthy eating and physical activity among college students, emphasizing their intricate relationship, as well as the health benefits associated with undisturbed sleep.

Limitations

Several limitations should be noted. The current study was designed as an exploratory study based on relatively limited evidence showing that disruption of sleep continuity can produce increased body weight by altering metabolic functions. However, our analysis was based on self-reported data, including height and weight, rather than direct measurements. In addition, we did not measure students' level of physical activity or dietary intake. The characteristics of the study participants, college students, limit the generalizability of our findings to the general population of college-age adults. The cross sectional nature of the study precludes conclusions about the causal relationship between sleep disturbances and BMI. Finally, we cannot rule out the possibility that our findings are due to other extraneous variables not measured, or to chance. Future studies would benefit from collecting multiple data points prospectively, using objective measurements of obesity and sleep, and exploring potential causal mechanisms between these variables.

Acknowledgments

We want to acknowledge the contribution of Dr. Oksana Kozhevnikova as well as Tracie Knight, Michael Duhaime, Donna Fendon, and Mikaela Durfey in the design of the study questionnaire and management of data collection.

Funding: Partial support for Dr. Vargas was provided by the National Institute on Minority Health and Health Disparities of the National Institutes of Health (NIMHD/NIH), award P20 MD002316-07 (F. Marsiglia, P.I.).

References

1. Ruthig J, Haynes T, Stupnisky R, Perry R. Perceived Academic Control: mediating the effects of optimism and social support on college students' psychological health. *Soc Psychol Educ*. 2009; 12(2):233–49.
2. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA. Emerging Adulthood and College-aged Youth: An Overlooked Age for Weight-related Behavior Change. *Obesity Res*. 2008; 16(10):2205–11.
3. Arnett, JJ. Emerging adulthood: Understanding the new way of coming of age. In: Arnett, Jeffrey Jensen; Tanner, Jennifer Lynn, editors. *Emerging adults in America: Coming of age in the 21st century*. Washington, DC: American Psychological Association; 2006. p. 3-19.
4. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among U.S. children and adolescents, 1999-2010. *JAMA*. 2012; 307(5):483–90. [PubMed: 22253364]
5. Ogden CL, Carroll MD, Curtin LR, Lamb MM, Flegal KM. Prevalence of high body mass index in U.S. children and adolescents, 2007-2008. *JAMA*. 2010; 303(3):242–9. [PubMed: 20071470]
6. Fryar, CD.; Carroll, MD.; Ogden, CL. [Accessed June 4, 2013] Prevalence of Overweight Obesity, Extreme Obesity Among Adults: United States, Trends 1960-1962 Through 2009-2010. Centers for Disease Control and Prevention (CDC). Jun 9. 2012 Available at: http://198.246.124.29/nchs/data/hestat/obesity_adult_09_10/obesity_adult_09_10.pdf
7. Flegal KM, arll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among us adults, 1999-2008. *JAMA*. 2010; 303(3):235–41. [PubMed: 20071471]
8. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among us children and adolescents, 1999-2010. *JAMA*. 2012; 307(5):483–90. [PubMed: 22253364]
9. American College Health Association. American College Health Association-National College. Hanover, MD: 2012. Health Assessment II: Reference Group Summary Report Spring 2012. Available at: http://www.acha-ncha.org/docs/ACHA-NCHA-II_ReferenceGroup_DataReport_Spring2012.pdf [Accessed June 4, 2013]
10. Lowry R, Galuska DA, Fulton JE, Wechsler H, Kann L, Collins JL. Physical activity, food choice, and weight management goals and practices among U.S. college students. *Am J Prev Med*. 2000; 18(1):18–27. [PubMed: 10808979]
11. Huang TTK, Harris KJ, Lee RE, Nazir N, Born W, Kaur H. Assessing overweight, obesity, diet, and physical activity in college students. *J Am Coll Health*. 2003; 52(2):83–6. [PubMed: 14765762]
12. Knutson KL, Van Cauter E, Rathouz PJ, DeLeire T, Lauderdale DS. Trends in the prevalence of short sleepers in the USA: 1975 - 2006. *Sleep*. 2010; 33(1):37. [PubMed: 20120619]
13. Matricciani L, Olds T, Petkov J. In search of lost sleep: Secular trends in the sleep time of school-aged children and adolescents. *Sleep Med Rev*. 2012; 16(3):203–11. [PubMed: 21612957]
14. Ohayon MM, Roth T. What are the contributing factors for insomnia in the general population? *J of Psychosom Res*. 2001; 51(6):745–55. [PubMed: 11750297]
15. 2005 Adult Sleep habits and Styles. National Sleep Foundation 2005; Mar 29. 2005 Sleep in America Polls. Available at http://www.sleepfoundation.org/sites/default/files/2005_summary_of_findings.pdf [Accessed June 4, 2013]
16. Schoenborn CA, Adams PE. Health behaviors of adults: United States, 2005-2007. *Vital Health Stat 10*. 2010; 245:1–132. [PubMed: 20669609]

17. Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep Patterns and Predictors of Disturbed Sleep in a Large Population of College Students. *J Adolesc Health*. 2010; 46(2):124–32. [PubMed: 20113918]
18. Trockel MT, Barnes MD, Egget DL. Health-related variables and academic performance among first-year college students: Implications for sleep and other behaviors. *J Am Coll Health*. 2000; 49(3):125–31. [PubMed: 11125640]
19. Gaultney JF. The prevalence of sleep disorders in college students: impact on academic performance. *J Am Coll Health*. 2010; 59(2):91–7. [PubMed: 20864434]
20. Buboltz WC Jr, Brown F, Soper B. Sleep Habits and Patterns of College Students: A Preliminary Study. *J Am Coll Health*. 2001; 50(3):131. [PubMed: 11765249]
21. Sing CY, Wong WS. Prevalence of Insomnia and Its Psychosocial Correlates Among College Students in Hong Kong. *J Am Coll Health*. 2010; 59(3):174–82. [PubMed: 21186447]
22. Kenney SR, LaBrie JW, Hummer JF, Pham AT. Global sleep quality as a moderator of alcohol consumption and consequences in college students. *Addict Behav*. 2012; 37(4):507–12. [PubMed: 22285119]
23. Nyer M, Farabaugh A, Fehling K, Soskin D, Holt D, Papakostas GI, Pedrelli P, Fava M, Pisoni A, Vitolo O. Relationship Between Sleep Disturbance and Depression, Anxiety, and Functioning in College Students. *Depress Anxiety*. 2013; 00:1–8.
24. Nadorff MR, Nazem S, Fiske A. Insomnia symptoms, nightmares, and suicidal ideation in a college student sample. *Sleep*. 2011; 34(1):93. [PubMed: 21203379]
25. Carskadon MA, Davis SS. Sleep-wake patterns in the high-school-to-college transition: Preliminary data. *Sleep Res*. 1989; 18:113.
26. Pilcher JJ, Ginter DR, Sadowsky B. Sleep quality versus sleep quantity: Relationships between sleep and measures of health, well-being and sleepiness in college students. *J Psychosom Res*. 1997; 42:583–96. [PubMed: 9226606]
27. Hicks RA, Fernandez C, Pellegrini RJ. Striking changes in the sleep satisfaction of university students over the last two decades. *Percept Mot Skills*. 2001; 93(3):660. [PubMed: 11806582]
28. Buboltz W, Jenkins SM, Soper B, Woller K, Johnson P, Faes T. Sleep habits and patterns of college students: an expanded study. *J Coll Couns*. 2009; 12(2):113–124.
29. Hill JO, Wyatt HR, Reed GW, Peters JC. Obesity and the Environment: Where Do We Go from Here? *Science*. 2003; 299(5608):853–5. [PubMed: 12574618]
30. Veerman JL, Barendregt JJ, van Beeck EF, Seidell JC, Mackenbach JP. Stemming the Obesity Epidemic: A Tantalizing Prospect. *Obesity Res*. 2007; 15(9):2365–70.
31. Chaput JP, Sjodin AM, Astrup A, Despres JP, Bouchard C, Tremblay A. Risk factors for adult overweight and obesity: the importance of looking beyond the ‘big two’. *Obes Facts*. 2010; 3(5): 320–7. [PubMed: 20975298]
32. Keith SW, Redden DT, Katzmarzyk PT, Boggiano MM, Hanlon EC, Benca RM, Allison DB. Putative contributors to the secular increase in obesity: exploring the roads less traveled. *Int J Obesity*. 2006; 30(11):1585–1594.
33. Al-Disi D, Al-Daghri N, Khanam L, Al-Othman A, Al-Saif M, Sabico S, Chrousos G. Subjective sleep duration and quality influence diet composition and circulating adipocytokines and ghrelin levels in teen-age girls. *Endoc J*. 2010; 57(10):915–23.
34. Shankar A, Syamala S, Kalidindi S. Insufficient rest or sleep and its relation to cardiovascular disease, diabetes and obesity in a national, multiethnic sample. *PLoS One*. 2010; 5(11):e14189. [PubMed: 21152066]
35. Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med*. 2004 Dec.1(3):e62. [PubMed: 15602591]
36. Stamatakis KA, Punjabi NM. Effects of Sleep Fragmentation on Glucose Metabolism in Normal Subjects. *Chest*. 2010; 137(1):95–101. [PubMed: 19542260]
37. Lesser DJ, Bhatia R, Tran WH, Oliveira F, Ortega R, Keens TG, Mittelman SD, Khoo MC, Davidson Ward SL. Sleep fragmentation and intermittent hypoxemia are associated with decreased insulin sensitivity in obese adolescent Latino males. *Pediatr Res*. 2012; 72(3):293–8. [PubMed: 22669298]

38. Lucassen EA, Rother KI, Cizza G. Interacting epidemics? Sleep curtailment, insulin resistance, and obesity. *Ann N Y Acad Sci.* 2012; 1264(1):110–34. [PubMed: 22827862]
39. Chaput JP, Brunet M, Tremblay A. Relationship between short sleeping hours and childhood overweight/obesity: results from the Quebec en Forme Project. *Int J Obes.* 2006; 30(7):1080–5.
40. Gangwisch JE, Malaspina D, Babiss LA, Opler MG, Posner K, Shen S, Turner JB, Zammit GK, Ginsberg HN. Short sleep duration as a risk factor for hypercholesterolemia: analyses of the National Longitudinal Study of Adolescent Health. *Sleep.* 2010; 33(7):956–61. [PubMed: 20614855]
41. Tasali E, Leproult R, Ehrmann DA, Van Cauter E. Slow-wave sleep and the risk of type 2 diabetes in humans. *Proc Natl Acad Sci USA.* 2008; 105(3):1044–9. [PubMed: 18172212]
42. Bawazeer NM, Al-Daghri NM, Valsamakis G, Al-Rubeaan KA, Sabico SL, Huang TT, Mastorakos GP, Kumar S. Sleep duration and quality associated with obesity among Arab children. *Obes.* 2000; 17(12):2251–3.
43. Bell JF, Zimmerman FJ. Shortened nighttime sleep duration in early life and subsequent childhood obesity *Arch Pediatr Adolesc Med.* 2010; 164(9):840–5.
44. Biggs SN, Dollman J. Association between sleep, BMI and waist girth in children and adolescents: a retrospective analysis. *Acta Paediatr.* 2007; 96(12):1839–40. [PubMed: 17953733]
45. Agras WS, Hammer LD, McNicholas F, Kraemer HC. Risk factors for childhood overweight: a prospective study from birth to 9.5 years. *The Journal of pediatrics.* 2004; 145(1):20–25. [PubMed: 15238901]
46. Kong AP, Wing YK, Choi KC, Li AM, Ko GT, Ma RC, Tong PC, Ho CS, Chan MH, Ng MH, Lau J, Chan JC. Associations of sleep duration with obesity and serum lipid profile in children and adolescents. *Sleep Med.* 2011; 12(7):659–65. [PubMed: 21689984]
47. Seegers V, Petit D, Falissard B, Vitaro F, Tremblay RE, Montplaisir J, Touchette E. Short sleep duration and body mass index: a prospective longitudinal study in preadolescence. *Am J Epidemiol.* 2011; 173(6):621–9. [PubMed: 21303806]
48. Shi Z, Taylor AW, Gill TK, Tuckerman J, Adams R, Martin J. Short sleep duration and obesity among Australian children. *BMC Public Health.* 2010; 10:609. [PubMed: 20946684]
49. Snell EK, Adam EK, Duncan GJ. Sleep and the body mass index and overweight status of children and adolescents. *Child Dev.* 2007; 78(1):309–23. [PubMed: 17328707]
50. Lytle LA, Pasch KE, Farbaksh K. The relationship between sleep and weight in a sample of adolescents. *Obes.* 2011; 19(2):324–31.
51. Landis AM, Parker KP. A retrospective examination of the relationship between body mass index and polysomnographic measures of sleep in adolescents. *J Adolesc Health.* 2007; 40(1):89–91. [PubMed: 17185213]
52. Weiss A, Xu F, Storfer-Isser A, Thomas A, Ievers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep.* 2010; 33(9):1201–9. [PubMed: 20857867]
53. Altman NG, Izci-Balserak B, Schopfer E, Jackson N, Rattanaumpawan P, Gehrman PR, Patel NP, Grandner MA. Sleep duration versus sleep insufficiency as predictors of cardiometabolic health outcomes. *Sleep Med.* 2012; 13(10):1261–70. [PubMed: 23141932]
54. Anic GM, Titus-Ernstoff L, Newcomb PA, Trentham-Dietz A, Egan KM. Sleep duration and obesity in a population-based study. *Sleep Med.* 2010; 11(5):447–51. [PubMed: 20363668]
55. Baron KG, Reid KJ, Kern AS, Zee PC. Role of sleep timing in caloric intake and BMI. *Obes.* 2011; 19(7):1374–81.
56. Buxton OM, Cain SW, O'Connor SP, Porter JH, Duffy JF, Wang W, Czeisler CA, Shea SA. Adverse metabolic consequences in humans of prolonged sleep restriction combined with circadian disruption. *Sci Transl Med.* 2012; 4(129):129–43.
57. Chaput JP, Despres JP, Bouchard C, Tremblay A. The association between sleep duration and weight gain in adults: a 6-year prospective study from the Quebec Family Study. *Sleep.* 2008; 31:517. [PubMed: 18457239]
58. Kobayashi D, Takahashi O, Deshpande GA, Shimbo T, Fukui T. Relation between metabolic syndrome and sleep duration in Japan: a large scale cross-sectional study. *Intern Med.* 2011; 50(2): 103–7. [PubMed: 21245632]

59. Markwald RR, Melanson EL, Smith MR, Higgins J, Perreault L, Eckel RH, Wright KP. Impact of insufficient sleep on total daily energy expenditure, food intake, and weight gain. *Proc Natl Acad Sci.* 2013; 110(14):5695–5700. [PubMed: 23479616]
60. Scheen AJ, Byrne MM, Plat L, Leproult R, Van Cauter E. Relationships between sleep quality and glucose regulation in normal humans. *American J Physiol-Endocrinol Metab.* 1996; 34(2):E261.
61. Bolli GB, De Feo P, De Cosmo S, Perriello G, Ventura MM, Calcinaro F, Gerich JE. Demonstration of a dawn phenomenon in normal human volunteers. *Diabetes.* 1984; 33(12):1150–1153. [PubMed: 6389230]
62. Boyle PJ, Scott JC, Krentz AJ, Nagy RJ, Comstock E, Hoffman C. Diminished brain glucose metabolism is a significant determinant for falling rates of systemic glucose utilization during sleep in normal humans. *J Clin Invest.* 1994; 93(2):529. [PubMed: 8113391]
63. Roenneberg T, Allebrandt KV, Meroow M, Vetter C. Social jetlag and obesity. *Curr Biol.* 2012; 22(10):939–43. [PubMed: 22578422]
64. Morris CJ, Yang JN, Scheer FA. The impact of the circadian timing system on cardiovascular and metabolic function. *Prog Brain Res.* 2012; 199:337–58. [PubMed: 22877674]
65. Seicean S, Kirchner HL, Gottlieb DJ, Punjabi NM, Resnick H, Sanders M, Redline S. Sleep-Disordered Breathing and Impaired Glucose Metabolism in Normal-Weight and Overweight/Obese Individuals The Sleep Heart Health Study. *Diabetes Care.* 2008; 31(5):1001–1006. [PubMed: 18268072]
66. Knutson KL. The association between pubertal status and sleep duration and quality among a nationally representative sample of U. S. Adolescents. *Am J Hum Biol.* 2005; 17:418–424. [PubMed: 15981178]
67. Gupta NK, Muelle WH, Chan W, Meininger JC. Is obesity associated with poor sleep quality in adolescents? *Am J Hum Biol.* 2002; 14:762–768. [PubMed: 12400037]
68. Garaulet M, Ortega FB, Ruiz JR, Rey-Lopez JP, Beghin L, Manios Y, et al. Short sleep duration is associated with increased obesity markers in European adolescents: effect of physical activity and dietary habits. The Helena study. *Int J Obes.* 2011; 35:1308–1317.
69. Lauderdale DS, Knutson KL, Yan LL, Rathouz PJ, Hulley SB, Sidney S, et al. Objectively measured sleep characteristics among early-middle-aged adults the CARDIA study. *Am J Epidemiol.* 2006; 164:5–16. [PubMed: 16740591]
70. Hasler G, Buysse DJ, Klaghofer R, Gamma A, Ajdacic V, Eich D, Rossler W, Angst J. The association between short sleep duration and obesity in young adults: a 13-year prospective study. *Sleep.* 2004; 27(4):661. [PubMed: 15283000]
71. Calamaro CJ, Park S, Mason TB, Marcus CL, Weaver TE, Pack A, Ratcliffe SJ. Shortened sleep duration does not predict obesity in adolescents. *J Sleep Res.* 2010; 19(4):559–66. [PubMed: 20545836]
72. Gottlieb DJ, Punjabi NM, Newman AB, Resnick HE, Redline S, Baldwin CM, et al. Association of sleep time with diabetes mellitus and impaired glucose tolerance. *Arch Intern Med.* 2005; 165:863–867. [PubMed: 15851636]
73. Kripke DF, Garfinkel L, Wingard DL, Klauber MR, Marler MR. Mortality associated with sleep duration and insomnia. *Arch Gen Psychiat.* 2002; 59:131–136. [PubMed: 11825133]
74. Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med.* 2004; 1:e62. [PubMed: 15602591]
75. Buxton OM, Marcelli E. Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States. *Soc Sci Med.* 2010; 71(5):1027–36. [PubMed: 20621406]
76. Tamakoshi A, Ohno Y. Self-reported sleep duration as a predictor of all-cause mortality: results from the JACC study, Japan. *Sleep.* 2004; 27:51–54. [PubMed: 14998237]
77. Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep deprivation. *Sleep medicine reviews.* 2007; 11(3):163–178. [PubMed: 17442599]
78. Buboltz WC Jr, Soper B, Brown F, Jenkins S. Treatment approaches for sleep difficulties in college students. *Couns Psychol Quart.* 2002; 15:229–237.

79. Jensen DR. Understanding sleep disorders in a college student population. *J Coll Couns.* 2003; 6:25–34.
80. Wechsler H, Kelley K, Weitzman ER, San Giovanni JP, Seibring M. What Colleges Are Doing About Student Binge Drinking A Survey of College Administrators. *J Am Coll Health.* 2000; 48:219–226. [PubMed: 10778022]
81. Buysse DJ, Reynolds CF III, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res.* 1989; 28(2):193–213. [PubMed: 2748771]
82. Carpenter JS, Andrykowski MA. Psychometric evaluation of the Pittsburgh Sleep Quality Index. *J Psychosom Res.* 1998; 45(1):5–13. [PubMed: 9720850]
83. Backhaus J, Junghanns K, Broocks A, Riemann D, Hohagen F. Test-retest reliability and validity of the Pittsburgh Sleep Quality Index in primary insomnia. *J Psychosom Res.* 2002; 53(3):737–40. [PubMed: 12217446]
84. Fichtenberg NL, Putnam SH, Mann NR, Zafonte RD, Millard AE. Insomnia Screening in Postacute Traumatic Brain Injury: Utility and Validity of the Pittsburgh Sleep Quality Index. *Am J Phys Med Rehabil.* 2001; 80(5)
85. National Institutes of Health. National Heart, Lung and Blood Institute, North American Association for the Study of Obesity: *The practical guide identification, evaluation, and treatment of overweight and obesity in adults.* Bethesda, MD: National Institutes of Health; 2000. NIH publication no. 00-4084
86. Elgar FJ, Roberts C, Tudor-Smith C, Moore L. Validity of self-reported height and weight and predictors of bias in adolescents. *J Adolesc Health.* 2005; 37(5):371–5. [PubMed: 16227121]
87. Spencer EA, Appleby PN, Davey GK, Key TJ. Validity of self-reported height and weight in 4808 EPIC-Oxford participants. *Public Health Nutr.* 2002; 5(04):561–5. [PubMed: 12186665]
88. Kuczmarski MF, Kuczmarski RJ, Matt Najjar. Effects of Age on Validity of Self-Reported Height, Weight, and Body Mass Index: Findings from the Third National Health and Nutrition Examination Survey, 1988-1994. *J Am Diet Assoc.* 2001; 101(1):28–34. [PubMed: 11209581]
89. VanSomeren EJ. More than a marker: interaction between the circadian regulation of temperature and sleep, age-related changes, and treatment possibilities. *Chronobiol Int.* 2000; 17:313–354. [PubMed: 10841209]
90. Van Cauter E, Leproult R, Plat L. Age-related changes in slow wave sleep and REM sleep and relationship with growth hormone and cortisol levels in healthy men. *JAMA.* 2000; 284:861–868. [PubMed: 10938176]
91. Dijk DJ, Duffy JF, Czeisler CA. Contribution of circadian physiology and sleep homeostasis to age-related changes in human sleep. *Chronobiol Int.* 2000; 17:285–311. [PubMed: 10841208]
92. Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep.* 2004; 27:1255–1274. [PubMed: 15586779]
93. Hauri PJ. A cluster analysis of insomnia. *Sleep.* 1983; 6(4):326–38. [PubMed: 6665395]
94. Hauri P, Olmstead E. Childhood-onset insomnia. *Sleep.* 1980; 3(1):59–65. [PubMed: 7466126]
95. Bixler EO, Kales A, Soldatos CR, Kales JD, Healey S. Prevalence of sleep disorders in the Los Angeles metropolitan area. *Am J Psychiatry.* 1979; 136:1257. [PubMed: 314756]
96. Guh D, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis A. The incidence of comorbidities related to obesity and overweight: A systematic review and meta-analysis. *BMC Public Health.* 2009; 9:88. [PubMed: 19320986]
97. Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet.* 2011; 378(9793):815–825. [PubMed: 21872750]
98. Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW. Body-Mass Index and Mortality in a Prospective Cohort of U.S. Adults *N Engl J Med.* 1999; 341:1097–1105.
99. Olshansky SJ, Passaro DJ, Hershow RC, et al. A Potential Decline in Life Expectancy in the United States in the 21st Century. *N Engl J Med.* 2005; 352:1138–1145. [PubMed: 15784668]

Table 1

Frequency distributions of sleep quality and sleep disturbance scores as measured by components and individual questions of the PSQI.

Component	Question/ Component Scores	0	1	2	3
1	Rate over all quality of sleep	Very good 11.4%	Fairly good 57.9%	Fairly bad 27.6%	Very bad 3.1%
2	How long does it take you to fall sleep	<=15 min 24.4%	16-30 min 36.0%	31-60 min 25.0%	>60 min 14.6%
3	Hours do you actually sleep	>=7 hrs 58.6%	6 hrs 25.6%	5 hrs 10.1%	< 5 hrs 4.3%
4	% Time in bed sleeping	>85 % 67.0%	75-84 % 20.8%	65-74 % 7.7%	<65 % 4.6%
5	Overall sleep disturbances score	0 6.8%	1-9 77.1%	10-18 15.1%	19-27 1.0%
6	Taken medicine to aid in sleep?	Not in past month 75.7%	< once a week 11.1%	1-2 times a week 5.3%	3 times a week 7.9%
7	Daytime dysfunction	0 18.3%	1-2 54.9%	3-4 23.7%	5-6 3.2%
Individual Questions					
Q5. How often have you had trouble sleeping because:					
	Q5b. Wake up middle of night/ early AM	Not in past month 19.1%	< once a week 25.8%	1-2 times a week 25.8%	3 times a week 29.2%
	Q5c. Wake up to use the bathroom	36.4%	29.2%	14.8%	19.5%
	Q5d. Cannot breathe comfortably	77.8%	13.0%	5.5%	3.7%
	Q5e. Cough or snore loudly	80.4%	11.4%	4.7%	3.4%
	Q5f. Feel too cold	50.4%	23.0%	18.3%	8.3%
	Q5g. Feel too hot	46.6%	24.6%	19.3%	9.3%
	Q5h. Have bad dreams	43.1%	28.7%	20.7%	7.5%
	Q5i. Have pain	64.1%	17.8%	10.7%	7.5%
	Q5j. Other reasons	68.7%	9.1%	10.2%	12.0%
	Q8. Had trouble staying awake during social activities?	63.4%	22.8%	10.5%	3.2%
	Q9. Problem getting enthusiasm	20.1%	47.2%	23.4%	9.3%

Table 2

Univariate logistic regression with overweight/ obesity (BMI ≥ 25) as outcome variable and Sleep Quality (0/1), PSQI global score (0-21), and the seven components of the PSQI (0-3) as predictor variables.

PSQI Components (4-point scale, range 0 - 3)	OR	95% CI	P value
Poor Sleep Quality (PSQI > 5)	1.106	.740 – 1.654	.623
Global PSQI score (0-21)	1.029	.958 – 1.105	.436
1. Subjective Quality	1.060	.806 -- 1.394	.675
2. Latency	1.000	.830 – 1.205	1.000
3. Duration	1.151	.924 – 1.435	.209
4. Efficiency	.999	.784 – 1.274	.997
5. Disturbances	1.701	1.164 – 2.485	.006
6. Medication	.979	.797 – 1.324	.838
7. Daytime Dysfunction	1.026	.795 – 1.324	.846