

## Poster Sessions – Abstract P291

# The impact of nevirapine- versus protease inhibitor-based regimens on virological markers of HIV-1 persistence during seemingly suppressive ART

Kiselinova, Maja<sup>1</sup>; Anna, Maria<sup>2</sup>; Malatinkova, Eva<sup>1</sup>; Vervish, Karen<sup>1</sup>; Beloukas, Apostolos<sup>2</sup>; Messiaen, Peter<sup>3</sup>; Bonczkowski, Pawel<sup>1</sup>; Trypsteen, Wim<sup>1</sup>; Callens, Steven<sup>1</sup>; Verhofstede, Chris<sup>4</sup>; De Spiegelaere, Ward<sup>1</sup> and Vandekerckhove, Linos<sup>1</sup>

<sup>1</sup>Internal Medicine and Infectious Diseases, Ghent University Hospital, Ghent, Belgium. <sup>2</sup>Clinical Infection, Microbiology and Immunology, University of Liverpool, Liverpool, UK. <sup>3</sup>Infectious Diseases and Immunity, Jessa Hospital, Hasselt, Belgium. <sup>4</sup>AIDS Reference Laboratory, Ghent University Hospital, Ghent, Belgium.

**Introduction:** The source and significance of residual plasma HIV-1 RNA detection during suppressive ART remain controversial. It has been proposed that nevirapine (NVP)-based regimens achieve a greater HIV-1 RNA suppression than regimens containing a protease inhibitor (PI). The aim of this study was to compare the effect of receiving NVP- vs PI-based ART on the virological markers of HIV persistence in peripheral blood.

**Materials and Methods:** The study population comprised 161 HIV-1 infected patients receiving either NVP-based (n = 81) or PI-based (n = 80) ART and showing a HIV-1 RNA load stably suppressed <40 copies/mL for median of 5.2 years (IQR 2.2–8.0). Residual viraemia was detected by real-time PCR with 50% and 95% detection thresholds of 1 and 3 HIV-1 RNA copies/mL, respectively. Cell-associated (CA) unspliced HIV-1 RNA, total HIV-1 DNA and 2 LTR circles were quantified in peripheral blood mononuclear cells (PBMCs) using droplet digital PCR. Groups were compared by standard non-parametric tests; factors associated with HIV-1 detection were analyzed by univariate regression analysis and generalized linear models (SPSS<sup>®</sup> V22 and Rstudio).

**Results:** Plasma HIV-1 RNA was detected in 37/81 (45.7%) and 47/80 (58.8%) subjects on NVP- and PI-based ART, with median (IQR) levels of 5 (3–6) and 5 (3–8) copies/mL, respectively. HIV-1 RNA detection was associated with shorter duration of suppressive ART regardless of treatment arm (p = 0.007), and lower CD4 nadir (p = 0.015). HIV-1 DNA levels were median 282 (120–484) and 213 (87–494) copies/106 PBMCs in the two groups respectively, and were lowest (<100 copies/106 PBMCs) in subjects with lower plasma HIV-1 RNA (p = 0.049), CA unspliced HIV-1 RNA (p = 0.0001), 2 LTR circles (p = 0.005) and pre-ART HIV-1 RNA load (p = 0.0001).

**Conclusions:** In this comprehensive characterization of patients on long-term suppressive ART, we did not observe evidence for a greater suppressive activity of NVP-based over PI-based therapy on plasma and intracellular markers of virus persistence. Overall excellent correlation was observed between the markers, allowing the identification of a subset of treated patients with low HIV-1 expression as an important cohort for future HIV cure studies.

Published 2 November 2014

**Copyright:** © 2014 Kiselinova M et al; licensee International AIDS Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.