

# Prevalence of Osteoarthritis of Knee Among Elderly Persons in Urban Slums Using American College of Rheumatology (ACR) Criteria

ARVIND KUMAR SINGH<sup>1</sup>, MANI KALAIVANI<sup>2</sup>, ANAND KRISHNAN<sup>3</sup>, PRAVEEN KUMAR AGGARWAL<sup>4</sup>, SANJEEV KUMAR GUPTA<sup>5</sup>

## ABSTRACT

**Background:** The prevalence of osteoarthritis among elderly is high and it majorly affects the quality of life. Knee osteoarthritis is the most common form of osteoarthritis. Timely diagnosis using clinical criteria and effective intervention is of utmost importance.

**Aim:** To estimate the prevalence and determinants of osteoarthritis of knee joint among elderly persons residing in an urban slum of Delhi using ACR clinical criteria.

**Materials and Methods:** We did a community-based cross-sectional study among 496 elderly ( $\geq 60$  years) persons residing in urban slum of Delhi, India from December 2009 to February 2010. The American College of Rheumatology (ACR) criteria was used to clinically diagnose osteoarthritis knee.

**Statistical Analysis:** Bivariate analysis using Chi-square test and multivariate analysis was done to identify the determinants.

Sensitivity and specificity of individual factors to diagnose osteoarthritis knee was calculated.

**Results:** The prevalence of osteoarthritis was estimated to be 41.1% (95% C.I., 36.7-45.6). Female sex and age  $\geq 70$  y were found to be independent risk factor for osteoarthritis knee. Among those having knee pain, presence of crepitus and tenderness were the most sensitive factors whereas bone overgrowth and bone warmth were most specific factors.

**Conclusion:** The prevalence of osteoarthritis knee was high among this elderly population and increased with age. Overall, individual factors of ACR criteria were both sensitive and specific in diagnosing osteoarthritis knee. In resource constrained setting of urban India, it can be an effective tool in clinical diagnosis of osteoarthritis knee.

**Keywords:** Aged, Diagnosis, Osteoarthritis

## INTRODUCTION

Osteoarthritis, though having a huge burden among the elderly, is not given the importance it deserves in public health. Osteoarthritis may not directly lead to mortality, but influences the Quality of Life (QoL) largely. Presence of osteoarthritis in older adults was associated with more pain, functional limitations, and lower Quality of Life (physical component) [1]. After achieving high life expectancy for the country, the next responsibility of public health is to provide good quality life without any disability. Osteoarthritis is believed to be a disease of old age with three fold increase in prevalence among elderly when compared to the younger counterpart [2,3]. With India witnessing demographic transition leading to proportionate as well as absolute increase in number of elderly, the magnitude of osteoarthritis is bound to increase [4]. Osteoarthritis of knee joint contributes to nearly 80% of total osteoarthritis burden [5]. Radiological assessment remains the mainstay of diagnosis of osteoarthritis of knee. Most of the cases of osteoarthritis seek treatment very late, only when the condition hampers with the physical activity. Moreover, elderly persons residing in urban slums are likely to have even worse health seeking behaviour. Definitive treatment in form of total knee replacement is costly, and unaffordable in Indian setting. Thus, prevention and early diagnosis remains the most cost effective strategy. At a community level, it is difficult to perform radiography for diagnosis. In Indian setting, where health system is already overburdened with more important public health activities, concern for osteoarthritis takes a backseat. Thus, costly treatment and delayed diagnosis hampers the effective intervention for osteoarthritis of knee joint. In wake of this, clinical assessment for diagnosis of osteoarthritis of knee joint is of utmost importance. American College of Rheumatology (ACR) clinical classification criteria is one of most widely used tool

for clinical diagnosis of osteoarthritis of knee joint in epidemiological studies. In this backdrop, we aimed to estimate the prevalence and determinants of osteoarthritis of knee joint among elderly persons residing in an urban slum of Delhi using ACR criteria as clinical tool to diagnose osteoarthritis knee in community based resource constrained setting of India. This might be useful for concerned policy makers to implement effective strategy for timely intervention in such cases.

## MATERIALS AND METHODS

We conducted a community-based cross-sectional study among elderly persons ( $\geq 60$  y of age) residing in an urban slum in southern region of Delhi during December 2009 to February 2010.

Sample size was calculated based on estimated prevalence of osteoarthritis in a community based study among elderly in Chandigarh in year 1999-2000 using clinical criteria. The prevalence of osteoarthritis knee was reported to be 34%. Sample size was calculated using the formula  $N = \frac{4pq}{d^2}$  where N= Sample size, p= reported prevalence, q= complement of prevalence (1-p) and d= absolute precision which was taken as 4.5% for the purpose of calculation of sample size. The sample size thus estimated was 443. Assuming a non response rate of 15%, the final sample to be achieved was 521. Since, population of elderly residing in this slum was largely unknown, a rapid enumeration of all elderly in the study area was done. Rapid enumeration yielded a total of 545 elderly persons residing in the study area. All 545 persons aged 60 y and above were included in the study if residing for more than six month in the same area. Person who were unable to comprehend and answer the questions, or were physically not capable of getting the physical examination done, were excluded from the study.

Variable	Total n=496 (%)
Age 60- 64 years	283 (57.1)
Females	238 (48.0)
More than 25 years of migration	240 (48.4)
Married	361(72.8)
Living with family	428 (86.3)
Economically dependent	310 (62.5)
Working presently	190 (38.3)
Illiterate	328 (66.1)
Smoker	193 (38.9)
Tobacco chewer	163 (32.9)

**[Table/Fig-1]:** Socio-demographic characteristics to participants (n=496)

Variable	Number	% (95% C.I.)
Osteoarthritis knee (Any knee)	204	41.1 (36.7-45.6)
Osteoarthritis knee	Bilateral	187 37.7 (33.4-42.1)
	Unilateral	17 3.4 (2.0-5.4)

**[Table/Fig-2]:** Prevalence of osteoarthritis knee (n=496)

Variable	Category	Unadjusted OR (95% C.I.)	Adjusted OR* (95% C.I.)	P value
Sex	Male	Referent category		
	Female	3.22 (2.21-4.67)	3.23 (2.03-5.14)	<0.001
Age group (years)	60-64	Referent category		
	65-69	1.27 (0.81-1.99)	1.26 (0.78-2.04)	0.34
	>=70	2.45 (1.55-3.89)	2.77 (1.66-4.61)	<0.001
Living status	With spouse	Referent category		
	With family	0.42 (0.21-0.85)	0.44 (0.21-0.93)	0.03
	Living alone	0.11 (0.04-0.37)	0.11 (0.03-0.38)	<0.001

**[Table/Fig-3]:** Association of osteoarthritis knee (any joint) with various socio-demographic characteristics., \*Results presented only for those factors which were significant on multivariate analysis. Adjusted for all socio-demographic variables and co-morbid conditions like diabetes and hypertension

We did a house to house survey interview of all eligible participants using a semi-structures interview schedule in Hindi. Information regarding basic socio-demographic characteristics, self-reported chronic condition and its treatment history was obtained. Maximum three visits were made in case participant was not interviewed in the first visit. Clinical examination of both the knee joints was done using ACR criteria. All the history taking and examination was done by the first author himself.

American College of Rheumatology (ACR) criteria: ACR criteria for diagnosing osteoarthritis of knee joint is presence of pain in knee joint plus any three of six factors listed below [6].

1. Age more than 50 y
2. Presence of crepitus on active motion
3. Less than 30 min of morning stiffness
4. Bony tenderness
5. Bony overgrowth
6. No palpable warmth of synovium.

Since all the participant were >= 60 y, presence of any two factors out of last five were enough for diagnosis.

First author underwent a training of two days for clinical diagnosis of osteoarthritis knee using ACR criteria under an experienced professor of Orthopedics.

**Ethical Statement:** Ethical clearance was obtained from the Institute Ethics Committee of the All India Institute of Medical Sciences, New Delhi, before commencing the study. Written informed consent was elicited from each participant and participant information sheet in Hindi was provided. All the participants requiring any preliminary

Variable	Right Joint				Left Joint			
	Osteoarthritis present (n=196)		Osteoarthritis absent (n=300)		Osteoarthritis present (n=193)		Osteoarthritis absent (n=303)	
	No.	%	No.	%	No.	%	No.	%
Pain	196	100.0	65	21.7	193	100.0	70	23.1
Crepitus	196	100.0	19	6.3	193	100.0	20	6.6
Tenderness	189	96.4	5	1.7	186	96.4	5	1.7
Morning stiffness	102	52.0	1	0.3	101	52.3	1	0.3
Bone warmth	62	31.6	0	0.0	59	30.6	0	0.0
Bony overgrowth	32	16.3	0	0.0	31	16.1	0	0.0

**[Table/Fig-4]:** Positivity rate of various factors of ACR criteria with respect to presence of osteoarthritis knee (n=496)

	Sensitivity (95% C.I.)	Specificity (95% C.I.)
Pain	1.00 (0.98 - 1.00)	0.78 (0.73 - 0.82)
Crepitus	1.00 (0.98 - 1.00)	0.94 (0.90 - 0.96)
Tenderness	0.96 (0.92 - 0.98)	0.98 (0.96 - 0.99)
Morning stiffness	0.52 (0.45 - 0.59)	0.99 (0.98 - 1.00)
Bone warmth	0.32 (0.25 - 0.38)	1.00 (0.98 - 1.00)
Bony overgrowth	0.16 (0.11 - 0.22)	1.00 (0.99 - 1.00)

**[Table/Fig-5]:** Sensitivity and specificity of various individual factors of ACR criteria for diagnosing osteoarthritis knee (right joint)\*., \*Results presented for one joint (right joint) since results of the both the joints were almost similar

treatment were appropriately dealt with in the field itself whereas those requiring specialist treatment were referred to All India Institute of Medical Sciences, New Delhi.

## STATISTICAL ANALYSIS

Data was entered in Epi-Info software version 3.5.1. and analysis was done using STATA version 9. Wherever applicable, proportion and mean were calculated along with 95% confidence interval and standard deviation respectively. Chi-square t-test was applied for bivariate analysis to find the association between outcome and predictor variables. p-value <0.05 was taken as significant. All the predictor variables significant at the level of p < 0.25 were included as a factor in the multivariate model for logistic regression. Results of bivariate and multivariate analysis are presented as unadjusted Odds' Ratio and adjusted Odds' Ratio respectively. Sensitivity and specificity was calculated for each of the factors in ACR criteria and is reported along with 95% confidence interval.

## RESULTS

During the period of cross-sectional survey, there were 545 (290 males: 255 females) eligible elderly in the area; of these, 496 (91%) participated in the study. Socio-demographic characteristics are presented in [Table/Fig-1]. The prevalence of osteoarthritis knee (any joint) was 41.1% (95% C.I., 36.7-45.6), out of which 37.7% (95% C.I., 33.4-42.1) had bilateral osteoarthritis whereas 3.4% (95% C.I., 2.0-5.4) had unilateral osteoarthritis [Table/Fig-2]. Presence of osteoarthritis knee was found to be significantly higher among females (OR=3.23, 95% CI 2.03-5.14) and participants aged 70 y and above (OR=2.77, 95% CI 1.66-4.61) [Table/Fig-3].

Positivity rate of various factors of ACR criteria is given in [Table/Fig-4]. In presence of pain in knee joint, all the five factors were highly specific in diagnosing osteoarthritis knee. Tenderness and crepitus were the most sensitive of all the five factors and were able to capture all the positive cases of osteoarthritis knee. Morning stiffness, bony overgrowth and bone warmth were not very sensitive though they had a very high specificity [Table/Fig 5].

## DISCUSSION

The elderly population studied here was relatively younger with sex ratio skewed towards male as compared to the national statistics. Since, the entire study population was migratory, this is explainable and similar phenomenon is also noted in various previous studies done in urban slums [4,7].

The prevalence of osteoarthritis knee was estimated to be 41.1 % in the present study. The prevalence was lower when compared to as study done in urban elderly population of Chandigarh where prevalence of osteoarthritis was estimated to be 56.6% [8]. The difference may be explained by the fact that they estimated the prevalence of osteoarthritis of all joint and used different criteria to ascertain the presence of osteoarthritis. The age group in this study was also 65 y and above. Osteoarthritis was considered if patient was suffering from pain, swelling, limitation of movement of a large joint. Other rheumatologic conditions may also present with similar symptoms, which are differentiated from osteoarthritis by absence of crepitus and a longer duration of stiffness. Another study done in similar setting in an urban elderly population of Chandigarh using the clinical criteria for estimation of osteoarthritis knee, reported a similar prevalence of 34% [9]. This study has found a considerably high prevalence of osteoarthritis knee among elderly.

Out of all the predictor variables studied, osteoarthritis knee was found to be significantly associated with female sex and age 70 y and above. Zeng et al., in Taiwan and Sharma et al., in India also reported a significantly higher prevalence of knee osteoarthritis in females and older age group [3,8]. Living arrangement was also found to be a predictor. Person either living alone (OR=0.11 95% CI 0.03, 0.38) or living with family (OR=0.44 95% CI 0.21, 0.93) had lesser risk of osteoarthritis knee than those who were living with spouse only. Person living alone due to lack of family support and person living with family due to more burden of household work may be more active in their household activities and thus may be having a lesser body mass index. There is evidence showing a higher risk of knee osteoarthritis with a higher body mass index [3]. A lower body mass index may put them at a lower risk of getting osteoarthritis knee.

We determined the sensitivity and specificity of various individual factors. Tenderness and crepitus were found to be highly sensitive whereas all the five factors were highly specific in diagnosing osteoarthritis knee. Thus, in this backdrop, ACR criterion seems to be a good tool in community based Indian setting where osteoarthritis is quite prevalent. Diagnosis using ACR criteria required only short training and is not a cumbersome modality. It could be performed by para-medical staff as well as volunteers in community based settings. ACR criteria showed a high inter-observer agreement of 0.94 when performed by experienced hands [10]. However, ACR criteria are not without any limitations. Though, the ACR classification criteria correlate well with articular cartilage damage in patients with OA [11]. In another study, it was found to have low sensitivity and moderate specificity when compared with symptomatic radiographic osteoarthritis (sensitivity 41% and specificity 75%) [12].

Nevertheless, this study has some limitations. Since, the study was done among elderly who have a higher prevalence of osteoarthritis than the general population. The higher sensitivity and specificity may be an overestimation and may not be applicable for the general population. Also, since the test was applied only among those who presented with pain in knee joint, thus increasing the baseline risk of osteoarthritis knee, it is likely that sensitivity and specificity would

have been overestimated. Ideal scenario would have been to apply this criterion to entire study population and compare with any of the gold standard test. So, our result of validity may be applicable in diagnosing osteoarthritis only when presented with pain in knee joint, in elderly population.

## CONCLUSION

With India currently witnessing a demographic transition, the absolute number of elderly is going to increase leading to a higher magnitude of burden of osteoarthritis. The elderly are by themselves a vulnerable section of society and with increasing age they become more prone to chronic conditions, osteoarthritis being one of them. Due to lack of affordable and accessible health care services, there is need for community based efforts to effectively screen, diagnose and treat various prevalent conditions in urban slums, osteoarthritis being one of them. Due to costlier treatment of osteoarthritis, early community based diagnosis using easy tools such as ACR and implementing appropriate intervention so as to halt the course of disease at an earlier stage, seems to be an adequate solution. With the new programme focusing on urban health in the pipeline, the present study may be of use to policy makers for strategic implementation of interventions which may reduce the burden and impact of osteoarthritis by preventing and treating it at a relatively early stage.

## REFERENCES

- [1] Jakobsson U, Hallberg IR. Quality of Life among Older Adults with Osteoarthritis: an explorative study. *J Gerontol Nurs*. 2006; 32:51-60.
- [2] Kim I, Kim HA, Seo YI, Song YW, Jeong JY, Kim DH. The prevalence of knee osteoarthritis in elderly community residents in Korea. *J Korean Med Sci*. 2010; 25(2):293-98.
- [3] Zeng QY, Zang CH, Li XF, Dong HY, Zhang AL, Lin L. Associated risk factors of knee osteoarthritis: a population survey in Taiyuan, China. *Chin Med J (Engl)*. 2006; 119:1522-27.
- [4] Registrar General and Census Commissioner of India, Census of India 2001. Available from <http://www.censusindia.gov.in/> (Accessed on 31st August 2013).
- [5] Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, Levy D. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum*. 1997;40:728-33.
- [6] Johns Hopkins Arthritis Centre ACR clinical classification criteria for OA of Knee. Available from [http://www.hopkins-arthritis.org/physiciancorner/education/acr/acr.html#class\\_knee](http://www.hopkins-arthritis.org/physiciancorner/education/acr/acr.html#class_knee). (Accessed on 31st August 2013).
- [7] Kant S, Misra P, Goswami A. Morbidity among elderly persons residing in a resettlement colony of Delhi. *Indian Journal of Preventive and Social Medicine*. 2004;35:1-9.
- [8] Sharma MK, Swami HM, Bhatia V, Verma A, Bhatia SPS, Kaur G. An epidemiological study of correlates of osteo-arthritis in geriatric population of UT Chandigarh. *Ind J Com Med*. 2007;32(1):77-78.
- [9] Joshi K, Kumar R, Avasthi A. Morbidity profile and its relationship with disability and psychological distress among elderly people in Northern India. *International Journal of Epidemiology*. 2003;32:978-87.
- [10] Bellamy N, Klestov A, Muirden K, Kuhnert P, Do KA, O'Gorman L, et al. Perceptual variation in categorizing individuals according to American College of Rheumatology classification criteria for hand, knee, and hip osteoarthritis (OA): observations based on an Australian Twin Registry study of OA. *J Rheumatol*. 1999;26:2654-58.
- [11] Wu CW, Morrell MR, Heinze E, Concoff AL, Wollaston SJ, Arnold EL, et al. Validation of American College of Rheumatology classification criteria for knee osteoarthritis using arthroscopically defined cartilage damage scores. *Semin Arthritis Rheum*. 2005;35:197-201.
- [12] Peat G, Thomas E, Duncan R, Wood L, Hay E, Croft P. Clinical classification criteria for knee osteoarthritis: performance in the general population and primary care. *Ann Rheum Dis*. 2006;65:1363-67.

### PARTICULARS OF CONTRIBUTORS:

1. Senior Resident, Department of Community Medicine, All India Institute of Medical Sciences, New Delhi, India.
2. Scientist- I, Department of Biostatistics, All India Institute of Medical Sciences, New Delhi, India.
3. Professor, Department of Community Medicine, All India Institute of Medical Sciences, New Delhi, India.
4. Professor and Head, Department of Emergency Medicine, All India Institute of Medical Sciences, New Delhi, India.
5. Professor, Department of Community Medicine, All India Institute of Medical Sciences, New Delhi, India.

### NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Sanjeev Kumar Gupta,  
Professor, Department of Community Medicine, All India Institute of Medical Sciences, Ansari Nagar, New Delhi – 110029, India.  
Phone : 98685-28573, E-mail : [sgupta\\_91@yahoo.co.in](mailto:sgupta_91@yahoo.co.in)

FINANCIAL OR OTHER COMPETING INTERESTS: None.

Date of Submission: **Sep 27, 2013**

Date of Peer Review: **Jul 28, 2014**

Date of Acceptance: **Aug 01, 2014**

Date of Publishing: **Sep 20, 2014**