

RESEARCH

Open Access

Comparative efficacy of a multi-epitope DNA vaccine via intranasal, peroral, and intramuscular delivery against lethal *Toxoplasma gondii* infection in mice

Hua Cong^{1*}, Quan Yuan², Qunli Zhao¹, Lingxiao Zhao¹, Huiquan Yin¹, Huaiyu Zhou¹, Shenyi He¹ and Zhiyu Wang^{3*}

Abstract

Background: *Toxoplasmosis* is an important zoonosis, being a cause of congenital disease and abortion in animals and humans. DNA vaccination as a promising vaccine remains a challenge for an improved delivery system.

Methods: In this study, attenuated *Salmonella typhimurium* BRD509 was used to deliver a DNA vaccine encoding several epitopes, derived from the tachyzoite proteins SAG1, GRA1, ROP2, GRA4 and bradyzoite proteins SAG2C, SAG2X of *Toxoplasma gondii* and A₂/B subunit of *cholera toxin*. The recombinant plasmids were electroporated into attenuated *Salmonella typhimurium*. Humoral and cellular immune responses were evaluated for BALB/c mice administered with this attenuated recombinant *Salmonella* vaccine via the oral and nasal route or by intramuscular injection with DNA plasmid directly.

Results: High IgG levels were present in the mice immunized intramuscularly, while IgA levels were higher in the oral and nasal immunization groups. Furthermore, cellular immunity was activated in oral immunization groups with 60% survival rate following challenge with high virulent RH strain.

Conclusions: The results from this study indicate that a DNA vaccine encoding multi-epitopes of *T. gondii* delivered by attenuated *Salmonella* is promising.

Keywords: *Toxoplasma gondii*, Multi-epitopes, *Salmonella typhimurium*, Vaccine

Background

Toxoplasma gondii is a single-cell obligate intracellular protozoan, which is widely prevalent in humans and animals [1,2]. This parasite is of major medical and veterinary importance, being a cause of congenital disease and abortion [3,4]. Despite many efforts and significant advances in the understanding of the immune responses that occur after infection by *T. gondii*, animal experiments have shown that the effect of monovalent antigen vaccine is not ideal [5-7]. Therefore, the development of a variety of antigen combinations for different stages of life cycle, especially epitopes from tachyzoites and

bradyzoites are likely to be needed for full protection [8-11]. We constructed a DNA vaccine encoding five epitopes of *T. gondii* previously. The vaccination injected intramuscularly into BALB/c resulted in an improvement of the humoral and cellular immune response in immunized mice, but there was only 20% survival rate achieved in vaccinated mice [12]. In order to broaden the immunogenicity, we intend to include more epitopes from various proteins of tachyzoites and bradyzoites in the vaccine construction. Other than epitopes from tachyzoite proteins SAG1, GRA1, ROP2, GRA4, epitopes from bradyzoite proteins SAG2C, SAG2X were also included [13,14].

However, DNA vaccines have been used to inject intramuscularly as naked DNA using a gene gun and by subcutaneous inoculation. After vaccination, only a limited amount of DNA reaches professional APCs

* Correspondence: conghua@sdu.edu.cn; zhiyu.wang@sdu.edu.cn

¹Department of human parasitology, Medical school, Shandong University, No.44 Wenhuxi Road, Jinan, Shandong 250012, P. R. China

³School of Public Health, Shandong University, No.44 Wenhuxi Road, Jinan, Shandong 250012, P. R. China

Full list of author information is available at the end of the article

cells [15]. Mechanisms to induce a more effective immune response and to improve protection from immunized mice remains to be determined. Recombinant *Salmonella* have been used as vaccine vectors to deliver both DNA and protein vaccines from a wide variety of bacterial, viral and parasitic sources [16,17]. Attenuation virulence bacteria, which can penetrate host cells delivering vaccine antigen to APCs, can be used effectively to transport immunogens [18,19].

In this study, a recombinant attenuated *Salmonella* DNA vaccine encoding multi-epitopes of *T. gondii* and A₂/B subunits of *cholera toxin* was constructed. The immunity induced by this attenuated recombinant *Salmonella* vaccine administered orally and nasally in BALB/c mice was compared to immunity induced by a plasmid DNA vaccine injected into mice intramuscularly. Protection against challenge with high virulent RH strain of *T. gondii* was evaluated.

Methods

Parasites and bacterial strain

Tachyzoites of the high virulent RH strain of *T. gondii* were cryopreserved in our laboratory. Parasites were maintained by serial intraperitoneal passage in BALB/c mice. Tachyzoites were harvested from the peritoneal fluid of mice after 72 h, and used to challenge immunized mice.

Salmonella typhimurium strain BRD509 is an aroA⁻ and aroD⁻ mutant of SL 1344 [20]. This was kindly

provided by Dr. Jifeng Bian, the Department of Molecular Biology, Shandong University.

Construction of recombinant *Salmonella*

The oligonucleotides of multi-epitope genes (MEG) were designed based on the *T. gondii* peptide sequences: SAG1-I 59–67 (TCPDKKSTA), SAG1-II 246–255 (ILPKLTENPW), GRA1 176–186(DTMKSMQRDED), ROP2 200–215 (GDV VIEELFNRIPESTS), GRA4 235–243 (SGLTGVKDS), SAG 2C 36–44 (SQFLSLSLL), SAG2X 215–223 (AAGTTAT AV). The construction of pVAX1-MEG-CTXA₂/B was shown in Figure 1.

Attenuated recombinant *Salmonella* carrying pVAX1-MEG-CTXA₂/B was generated by electroporation. pVAX1-MEG-CTXA₂/B plasmid was transferred into 100 μl competent *S. typhimurium* BRD509 in the condition of 2.5 kV, 25 uF, 5 ms. To produce inocula, recombinant *Salmonella* were incubated overnight at 37°C until an OD₆₀₀ of 0.8 was reached, then washed and resuspended in PBS to a final density of approximately 1–5 × 10⁹ cfu.

Vaccinations and challenges

SPF BALB/c female mice (6–8 weeks old) were used in the immunization and parasite challenge. They were purchased from The Laboratory Animal Center of Shandong University. All studies were conducted with Animal Care and Use Committee of Shandong University approval.

The mice were randomly divided into intraoral, intranasal and intramuscular immunization groups (30 mice per group). Table 1 summarizes the treatments performed in




Table 1 Summary of treatments performed in the BALB/c mice

Groups	Route of administration	Treatments ^a	Mice number		
			Total mice	In HI ^b and CMI ^c	In challenge ^d
I	Intramuscular vaccination	100 µl PBS	10	3	7
		100 µg pVAX1	10	3	7
		100 µg pVAX1-MEG-CTXA ₂ /B	10	3	7
II	Intranasal vaccination	200 µl BRD509	10	3	7
		200 µl BRD509/pVAX1	10	3	7
		200 µl BRD509/pVAX1-MEG-CTXA ₂ /B	10	3	7
III	Intraoral vaccination	50 µl BRD509	10	3	7
		50 µl BRD509/pVAX1	10	3	7
		50 µl BRD509/pVAX1-MEG-CTXA ₂ /B	10	3	7

^aThe mice were randomly divided into intramuscular, intranasal and intraoral immunization groups (30 mice per group). For the intramuscular group, the mice were injected 100 µl 1 µg/µl recombinant plasmid or empty plasmid in the quadriceps muscle; For the nasal group, each mouse was given a nasal drip of 50 µl 10⁹ cfu/ml *Salmonella* bacteria with recombinant plasmid or empty plasmid; The oral group was administered 200 µl 10⁹ cfu / ml *Salmonella* bacteria with recombinant plasmid or empty plasmid orally; Each group also had 10 mice which were treated with *Salmonella* without plasmid or saline as negative control. Mice were vaccinated three times at 2 week intervals.

^bHumoral immunity (HI) was tested by collecting sera from three immunized mice in each group.

^cCellular immunity (CMI) was evaluated by splenocytes from three immunized mice per group 2 weeks after the last immunization.

^dFour weeks after the last immunization, immunized mice were challenged intraperitoneally with 1 × 10³ tachyzoites of *T. gondii* RH strain.

the mice. Mice were vaccinated three times at 2 weeks intervals. Blood was collected by orbital plexus puncture and sera were stored at -70 °C for further analysis.

For challenge study, immunized mice were challenged intraperitoneally with 1 × 10³ tachyzoites of RH strain *T. gondii* 4 weeks after the last immunization. The survival time and the survival rate was measured.

Measurement of humoral antibodies response

To measure *T. gondii*-specific total IgG and IgA, plates (Dursley, UK) were coated with 10 µg/ml solution of STAg at 4°C overnight. Sera were diluted in 1% PBST-20 (1:100) and incubated in the plates for 1 h at room temperature (RT). After washing the plates, Horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG, IgA (SouthernBiotech, USA) were further incubated for 1 h at RT. Peroxidase activity was revealed by 3,3',5,5'-tetramethylbenzidine (TMB, 10 mg/ml) and stopped by adding 50 µl of 2 M H₂SO₄. The optical density (OD) was read out at 450 nm in a microplate reader (Bio-TEK, USA).

Proliferation assays

Spleens from three immunized mice per group were removed 2 weeks after the last immunization. The splenocytes were adjusted to a concentration of 5 × 10⁶ cells/ml in DMEM with 10% FCS. The suspension of 100 µl in each well was cultured with STAg (10 µg/ml), purified rMEG protein (50 µg/ml) or concanavalin A (Con A, 5 µg/ml; Sigma). Cell proliferative activity was measured according to instructions described in a previous study [21].

T lymphocyte subsets analysis

Cells were stained with FITC-labeled anti-mouse CD8⁺ monoclonal antibody and PE-labeled anti-mouse CD4⁺ monoclonal antibody, T lymphocyte subsets were measured using flow cytometry (Beckman Coulter, USA).

Cytokine assays

Splenocytes from immunized mice were cultured with rMEG protein as described for lymphocyte proliferation assays. Commercial ELISA kits (mouse IL-4 OptEIA, IL-5 OptEIA, IFN-γ OptEIA, IL-2 OptEIA, Endogen, USA) were used according to the manufacturer's instructions to assay cytokine levels in culture supernatants obtained at 24 h for IL-4, at 36 h for IL-5, at 72 h for IL-2 and at 96 h for IFN-γ.

Statistical analysis

The statistically significant differences between groups were calculated with one-factor analysis of variance (ANOVA). Differences were considered to be significant with p < 0.05.

Results

Construction of recombinant *Salmonella*

The diagram for multi-epitope DNA vaccine of *T. gondii* is shown on Figure 1A. Recombinant *Salmonella* carrying pVAX1-MEG-CTXA₂/B plasmid were identified by enzyme digestion, by PCR (Figure 1B) and further confirmed by sequencing.

Humoral immune response

IgG antibody levels in mouse serum indicates that pVAX1-MEG-CTXA₂/B DNA plasmid inoculated intramuscularly

induced higher levels of IgG than mice immunized with BRD509/pVAX1-MEG-CTXA₂/B orally and intranasally, $p < 0.05$ (Figure 2A). On the other hand, higher levels of anti-*T. gondii* IgA were detected in the serum of mice immunized with BRD509/pVAX1-MEG-CTXA₂/B via the oral and nasal route than in the serum of mice injected intramuscularly with pVAX1-MEG-CTXA₂/B plasmid ($p < 0.05$) (Figure 2B).

Cellular immune response analysis

CD4⁺ and CD8⁺ T lymphocyte subsets in immunized mice were assayed by flow cytometry. Table 2 shows the percentage of CD4⁺ and CD8⁺ T-cell. The percentage of CD8⁺ T cells in mice immunized intramuscularly with pVAX1-MEG-CTXA₂/B plasmid was $28.54 \pm 0.92\%$. A even higher percentage of CD8⁺ T cells, $30.01 \pm 1.78\%$, was achieved in mice immunized intranasally with BRD509/ pVAX1-MEG-CTXA₂/B. The highest percentage of CD8⁺ T-cell, $35.55 \pm 0.70\%$, was achieved in mice immunized orally with BRD509/pVAX1-MEG-CTXA₂/B (Table 2). Culture supernatants from antigen-stimulated splenocytes from mice in these groups were quantified with sandwich ELISA for

Th1 and Th2 cytokines. As shown in Table 3, culture supernatant cytokines from the mice vaccinated intramuscularly with pVAX1-MEG-CTXA₂/B demonstrated preferential production of IFN- γ and IL-2, which were significantly higher than that of control groups ($p < 0.05$). The amounts of IFN- γ and IL-2 were significantly higher in the mice via oral and intranasal immunization with recombinant *Salmonella* BRD509 than the mice immunized intramuscularly ($p = 0.02$). However, the levels of IL-4 and IL-5 produced by immunized mice were low, and no statistically significant differences were present between vaccinated groups and control groups ($p > 0.05$). Furthermore, antigen specific lymphocyte proliferation activity was highly enhanced for BRD509/pVAX1-MEG-CTXA₂/B oral and intranasal immunization groups than for the pVAX1-MEG-CTXA₂/B intramuscular immunization group ($p < 0.05$).

Challenge study

Four weeks after the last immunization, mice were challenged with 100 μ l 1×10^3 tachyzoites of *T. gondii* RH strain intraperitoneally. All the control mice died. Mice treated with saline died within 4 days. Mice immunized




Figure 2 Specific humoral response to native *T. gondii* antigens in mice immunized with multi-epitope DNA vaccine was measured using ELISA 2 weeks after the last immunization. (A) Total Anti-*T. gondii* IgG; (B) Anti-*T. gondii* IgA were tested by sera diluted 1:200 from BALB/c mice via three different immunized routes: i.o.(intraoral); i.n. (intranasal); i.m. (intramuscular).

Table 2 CD4⁺ CD8⁺ subtypes of T cells from immunized mice were measured using flow cytometry

Immunization route ^a	Immunization regimen	CD4 ⁺ (%) ^b	CD8 ⁺ (%) ^b
Intramuscular vaccination	PBS	26.71 ± 1.32	21.67 ± 0.82
	pVAX1	25.45 ± 1.13	22.37 ± 0.98
	pVAX1-MEG-CTXA ₂ /B	24.56 ± 1.25	28.54 ± 0.92
Intranasal vaccination	BRD509	25.71 ± 0.64	22.37 ± 0.83
	BRD509/pVAX1	24.32 ± 1.75	26.34 ± 0.93
	BRD509/pVAX1-MEG-CTXA ₂ /B	24.68 ± 1.43	30.01 ± 1.78
Intraoral vaccination	BRD509	27.45 ± 1.91	24.38 ± 0.98
	BRD509/pVAX1	24.32 ± 1.21	30.28 ± 0.53
	BRD509/pVAX1-MEG-CTXA ₂ /B	25.15 ± 1.55	35.55 ± 0.70

^aMice were immunized by three immunization routes, intramuscular, intraoral, intranasal, on day 0 and day 14 and day 28 with different immunization regimens.

^bThe splenocyte culture supernatants taken from mice (n = 3, each group) 2 weeks after the last immunization were stained with FITC-labeled anti-mouse CD8⁺ monoclonal antibody and PE-labeled anti-mouse CD4⁺ monoclonal antibody, T lymphocyte subsets were measured using flow cytometry.

with empty plasmid intramuscularly died within 5 days. Mice treated orally with empty *Salmonella* died within 8 days. While there was a 20% survival rate (2/10) for the mice immunized with pVAX1-MEG-CTXA₂/B plasmid intramuscularly. The recombinant *Salmonella* BRD509/pVAX1-MEG-CTXA₂/B intranasal immunization group had 40% survival rate (4/10) after 10 days. The highest survival rate of 60% (6/10) was achieved in the mice orally

immunized with recombinant *Salmonella* BRD509/pVAX1-MEG-CTXA₂/B (Figure 3).

Discussion

Recently, considerable efforts have been devoted to the development of nucleic acid vaccine to protect against *T. gondii* [22-25]. However, because of the complexity of the parasite life cycle and the variability of the parasite antigens, multi-epitope vaccines have become an attractive strategy for development of vaccines against this parasite [26-28]. In this study, a DNA vaccine encoding seven epitopes of *T. gondii*, derived from the tachyzoite proteins SAG1, GRA1, ROP2, GRA4 and bradyzoite proteins SAG2C, SAG2X of *T. gondii*, were constructed and linked to A₂/B subunit of *cholera toxin*.

However, DNA vaccine as a promising vaccine remains a challenge for an improved delivery system. Mucosal vaccines can enhance mucosal and systemic humoral immunity especially for intracellular parasites such as *T. gondii* [29,30]. Thus, the best way for a vaccine to induce mucosal immunity is directly by inoculation of the mucosal surface. In this study, attenuated strains of *S. typhimurium* BRD509 were used as vaccine vectors to deliver heterologous antigens to the mucosal surface of the vaccine recipients [31,32]. A recombinant attenuated *S. typhimurium* vaccine encoding a multi-epitope gene of *T. gondii* was constructed. The mice were given inoculations via oral, nasal, or muscular routes. Immune responses induced by oral or nasal immunization with attenuated recombinant *Salmonella* BRD509 vaccines were compared to immune response induced by immunization with the plasmid pVAX1-MEG-CTXA₂/B DNA vaccine intramuscularly.

The results showed intramuscular immunization of mice induced higher levels of IgG antibodies than attenuated

Table 3 Cell proliferative assay and cytokine production in the splenocyte cultures obtained from immunized mice

Immunization route ^a	Immunization regimen	Cytokine production (pg/ml) ^b				
		IL-2	IFN-γ	IL-4	IL-5	SI ^c
Intramuscular vaccination	Saline	12 ± 3	13 ± 8	<10	<10	0.23
	pVAX1	16 ± 4	15 ± 5	11 ± 8	<10	0.51
	pVAX1-MEG-CTXA ₂ /B	93 ± 21	385 ± 64	15 ± 7	13 ± 6	1.61
Intranasal vaccination	BRD509	13 ± 4	14 ± 5	<10	17 ± 5	0.75
	BRD509/pVAX1	14 ± 7	15 ± 6	<10	19 ± 5	1.07
	BRD509/ pVAX1-MEG-CTXA ₂ /B	105 ± 30	412 ± 24	23 ± 6	<10	2.12
Intraoral vaccination	BRD509	12 ± 5	15 ± 4	<10	16 ± 8	0.75
	BRD509/pVAX1	15 ± 7	18 ± 4	<10	16 ± 7	1.45
	BRD509/ pVAX1-MEG-CTXA ₂ /B	121 ± 36	507 ± 16	<10	12 ± 4	2.85

^aMice were immunized by three immunization routes, intramuscular, intraoral, intranasal on day 0 and day 14 and day 28 with different immunization regimens.

^bThe splenocyte culture supernatants taken from mice (n = 3, each group) 2 weeks after the last immunization were examined for cytokine production by sandwich ELISA obtained at 24 h for IL-4, at 36 h for IL-5, at 72 h for IL-2 and 96 h for IFN-γ.

^cThe results of proliferation assays are expressed as the stimulation index (SI), calculated as the ratio between the mean counts per minute (cpm) for triplicate stimulated cultures and the mean counts per min for triplicate unstimulated cultures. SI values 2.5-fold greater than the SI of the control groups were considered as significant.




Figure 3 Survival curves of immunized mice after challenge with 10^3 tachyzoite forms of RH *T. gondii* strain 4 weeks after the last immunization. **(A)** Intramuscular immunization group, the mice were injected with 100 μ l 1 μ g/ μ l recombinant plasmid, empty plasmid or PBS in the quadriceps muscle; **(B)** Nasal immunization group, each mouse was given a nasal drip of 50 μ l 10^9 cfu/ml *Salmonella* bacteria with recombinant plasmid, empty plasmid or empty bacteria; **(C)** Oral immunization group, the mice were administrated orally with 200 μ l 10^9 cfu/ml *Salmonella* bacteria with recombinant plasmid, empty plasmid, or empty bacteria.

recombinant *Salmonella* vaccine immunization via oral and nasal routes. However, the attenuated recombinant *Salmonella* oral and intranasal immunization induced higher levels of IgA antibodies than intramuscular immunization. IgA is important in mucosal immunity to oral infection with toxoplasma cysts. Since antibodies of this isotype are important in preventing re-infection with *T. gondii*, inducing IgA may be a major strategic aspect of vaccine development [33,34].

Major mechanisms by which immunocompetent hosts control *T. gondii* infections is thought to be the strong and persistent cell-mediated immunity elicited by parasites [35,36]. In this study, splenocyte proliferation was enhanced in mice immunized by all three routes. Proliferation activity was significantly enhanced in mice immunized with recombinant attenuated *Salmonella* BRD509 via oral and nasal routes.

Both CD4⁺ and CD8⁺ T cell subsets play a central role in the establishment of protective immunity in hosts. In this study, CD4⁺ and CD8⁺ T lymphocytes subsets from immunized mice were assayed by flow cytometry. There was a marked increase in the percentage of CD8⁺ T-cells in mice immunized with the recombinant attenuated *Salmonella* BRD509 vaccine perorally. As we know, the memory T-cell consists of CD4⁺ and CD8⁺ T-cells, which can rapidly acquire effector functions to kill infected cells and secrete inflammatory cytokines that inhibit replication

of the pathogen. Effector CD4⁺ T-cells enhance CD8⁺ T cells develop and help B-cell responses through the activation of antigen-presenting cells or the secretion of cytokines, such as IFN- γ , IL-2 [37,38]. Compared with the intramuscular immunization DNA vaccine, introduction of attenuated *Salmonella* further enhanced Th1 cell-mediated immunity with higher levels of IFN- γ and IL-2 but low levels of IL-4 and IL-5. These results clearly demonstrate that attenuated *Salmonella* BRD509 can significantly augment Th1-type cellular immune responses in BALB/c mice. Both T-cell subsets are important sources of IFN- γ . Optimal protective CD8⁺ T cell responses depend on the ability of CD4⁺ T-cells to provide the growth factor IL-2.

In this study, a highly virulent RH strain of *T. gondii* was used for challenge study. When challenged with lethal doses of *T. gondii* (1×10^3) all three immunization groups prolonged survival time compared to control mice. Especially, mice immunized perorally had a 60% survival rate. There was somewhat less enhancement of survival (40%) following intranasal immunization, while only a 20% survival rate was achieved following intramuscular immunization. In the future, a cyst-forming PRU strain of *T. gondii* will be used to evaluate the protective potency of this recombinant *Salmonella* vaccine via the mucosal immunization route.

Conclusion

In summary, multi-epitope vaccines delivered by attenuated *Salmonella* are a promising way to elicit protective immune responses. Compared to intramuscular injection, oral and nasal immunizations with recombinant *Salmonella* elicits both systemic and mucosal immune responses, accompanied by a significant increase in survival rates in vaccinated mice after challenge with highly virulent parasites. The results from this study should contribute to development of highly protective mucosal vaccines against *T. gondii*.

Competing interests

The authors declare that they have no competing interests with this publication.

Authors' contributions

HC carried out the vaccine construction and drafted the manuscript. XZ performed the immune response analysis. XZ and LZ performed the animal experiments. QY and ZW participated in the design of the study. QY performed the statistical analysis. YZ and SH made a revision of the manuscript. All authors have read and approved the final manuscript.

Acknowledgments

This study was supported by grants from the National Natural Science Foundation Project of China (Grant No. 81171604, No. 81271857 and No. 31170906) and China Postdoctoral Science Foundation (Grant No. 20110491573).

Author details

¹Department of human parasitology, Medical school, Shandong University, No.44 Wenhua Road, Jinan, Shandong 250012, P. R. China. ²School of Mechanical Engineering, Shandong University, Jinan, Shandong 250061, P. R. China. ³School of Public Health, Shandong University, No.44 Wenhua Road, Jinan, Shandong 250012, P. R. China.

Received: 24 August 2013 Accepted: 21 March 2014

Published: 31 March 2014

References

1. Zhou P, Chen Z, Li HL, Zheng H, He S, Lin R, Zhu XQ: *Toxoplasma gondii* infection in humans in China. *Parasit Vectors* 2011, **4**:165.
2. Mwambe B, Mshana SE, Kidenya BR, Massinde AN, Mazigo HD, Michael D, Majinge C, Groß U: Sero-prevalence and factors associated with *Toxoplasma gondii* infection among pregnant women attending antenatal care in Mwanza, Tanzania. *Parasit Vectors* 2013, **6**:222.
3. McLeod R, Kieffer F, Sautter M, Hosten T, Pelloux H: Why prevent, diagnose and treat congenital toxoplasmosis? *Mem Inst Oswaldo Cruz* 2009, **104**:20–344.
4. Wang L, Cheng HW, Huang KQ, Xu YH, Li YN, Du J, Yu L, Luo QL, Wei W, Jiang L, Shen JL: *Toxoplasma gondii* prevalence in food animals and rodents in different regions of China: isolation, genotyping and mouse pathogenicity. *Parasit Vectors* 2013, **6**:273.
5. Couper KN, Nielsen HV, Petersen E, Roberts F, Roberts CW, Alexander J: DNA vaccination with the immunodominant tachyzoite surface antigen (SAG-1) protects against adult acquired *Toxoplasma gondii* infection but does not prevent maternofetal transmission. *Vaccine* 2003, **21**(21–22):2813–2820.
6. Sun XM, Zou J, Elashram Saeed AA, Yan WC, Liu XY, Suo X, Wang H, Chen QJ: DNA vaccination with a gene encoding *Toxoplasma gondii* GRA6 induces partial protection against toxoplasmosis in BALB/c mice. *Parasit Vectors* 2011, **4**:213.
7. Rashid I, Hedhli D, Moiré N, Pierre J, Debierre-Grockiego F, Dimier-Poisson I, Mévélec MN: Immunological responses induced by a DNA vaccine expressing RON4 and by immunogenic recombinant protein RON4 failed to protect mice against chronic toxoplasmosis. *Vaccine* 2011, **29**(48):8838–8846.
8. Zhang NZ, Chen J, Wang M, Petersen E, Zhu XQ: Vaccines against *Toxoplasma gondii*: new developments and perspectives. *Expert Rev Vaccines* 2013, **12**(11):1287–1299.
9. Yan HK, Yuan ZG, Song HQ, Petersen E, Zhou Y, Ren D, Zhou DH, Li HX, Lin RQ, Yang GL, Zhu XQ: Vaccination with a DNA vaccine coding for perforin-like protein 1 and MIC6 induces significant protective immunity against *Toxoplasma gondii*. *Clin Vaccine Immunol* 2012, **19**(5):684–689.
10. Qu D, Han J, Du A: Evaluation of protective effect of multiantigenic DNA vaccine encoding MIC3 and ROP18 antigen segments of *Toxoplasma gondii* in mice. *Parasitol Res* 2013, **112**(7):2593–2599.
11. Cong H, Zhang M, Xin Q, Wang Z, Li Y, Zhao Q, Zhou H, He S: Compound DNA vaccine encoding SAG1/ SAG3 with A2/B subunit of cholera toxin as a genetic adjuvant protects BALB/c mice against *Toxoplasma gondii*. *Parasit Vectors* 2013, **6**:63.
12. Cong H, Gu QM, Yin HE, Wang JW, Zhao QL, Zhou HY, Li Y, Zhang J: Multi-epitope DNA vaccine linked to the A2/B subunit of cholera toxin protect mice against *Toxoplasma gondii*. *Vaccine* 2008, **26**(31):3913–3921.
13. Cong H, Zhang M, Zhang Q, Gong J, Cong H, Xin Q, He S: Analysis of structures and epitopes of surface antigen glycoproteins expressed in bradyzoites of *Toxoplasma gondii*. *BioMed Res Inter* 2013, **2013**:165342.
14. Zhang M, Zhao L, Song J, Li Y, Zhao Q, He S, Cong H: DNA vaccine encoding the *Toxoplasma gondii* bradyzoite-specific surface antigens SAG2CDX protect BALB/c mice against type II parasite infection. *Vaccine* 2013, **31**(41):4536–4540.
15. Mohamed RM, Aosai F, Chen M, Mun HS, Norose K, Belal US, Piao LX, Yano A: Induction of protective immunity by DNA vaccination with *Toxoplasma gondii* HSP70, HSP30 and SAG1 genes. *Vaccine* 2003, **21**(21–22):2852–2861.
16. Gahan ME, Webster DE, Wijburg OL, Wesselingh SL, Strugnell RA: Impact of prior immunological exposure on vaccine delivery by *Salmonella enterica* serovar *Typhimurium*. *Vaccine* 2008, **26**(49):6212–6220.
17. Daiji A, Lage zur S, Garbe Al, Chakraborty T, Weiss S: Oral delivery of DNA vaccines using attenuated *Salmonella typhimurium* as carrier. *Immunol Med Microbiol* 2000, **27**:341–349.
18. Liu XB, Liang AX, Feng XG, Guo AZ, Ke CY, Zhang SJ, Yang LG: Oral and intranasal administration of somatostatin DNA vaccine mediated by attenuated *Salmonella enterica* serovar *Typhimurium* to promote growth of piglets. *Animal* 2011, **5**(8):1231–1236.
19. Chin'ombe N: Recombinant *Salmonella enterica* serovar *Typhimurium* as a vaccine vector for HIV-1 Gag. *Viruses* 2013, **5**(9):2062–2078.
20. Eom JS, Kim JS, Im Jang J, Kim BH, Yoo SY, Choi JH, Bang IS, Lee IS, Park YK: Enhancement of host immune responses by oral vaccination to *Salmonella enterica* serovar *Typhimurium* harboring both FliC and FljB flagella. *PLoS One* 2013, **8**(9):e74850.
21. Cong H, Gu QM, Jiang Y, He S, Zhou HY, YANG TT, Li Y, Zhao QL: Oral immunization with recombinant attenuated *Salmonella* protect mice against *Toxoplasma gondii*. *Parasit Immunol* 2005, **27**(1–2):29–35.
22. Meng M, He S, Zhao G, Bai Y, Zhou H, Cong H, Lu G, Zhao Q, Zhu XQ: Evaluation of protective immune responses induced by DNA vaccines encoding *Toxoplasma gondii* surface antigen 1 (SAG1) and 14-3-3 protein in BALB/c mice. *Parasit Vectors* 2012, **5**:273.
23. Chen J, Huang SY, Zhou DH, Li ZY, Petersen E, Song HQ, Zhu XQ: DNA immunization with eukaryotic initiation factor-2 α of *Toxoplasma gondii* induces protective immunity against acute and chronic toxoplasmosis in mice. *Vaccine* 2013, **31**(52):6225–6231.
24. Yan HK, Yuan ZG, Song HQ, Petersen E, Zhou Y, Ren D, Zhou DH, Li HX, Lin RQ, Yang GL, Zhu XQ: Vaccination with a DNA vaccine coding Perforin-like Protein 1 (TgPLP1) and MIC6 induces significant protective immunity against *Toxoplasma gondii*. *Clin Vaccine Immunol* 2012, **29**:684–689.
25. Zhao G, Zhou A, Lu G, Meng M, Sun M, Bai Y, Han Y, Wang L, Zhou H, Cong H, Zhao Q, Zhu XQ, He S: Identification and characterization of *Toxoplasma gondii* aspartic protease 1 as a novel vaccine candidate against toxoplasmosis. *Parasit Vectors* 2013, **6**:175.
26. Henriquez FL, Woods S, Cong H, McLeod R, Roberts CW: Immunogenetics of *Toxoplasma gondii* informs vaccine design. *Trends Parasitol* 2010, **26**(11):550–555.
27. Zhao L, Zhang M, Cong H: Advances in the study of HLA-restricted epitope vaccines. *Hum Vaccin Immunother* 2013, **9**(12):1–12.
28. Cong H, Mui EJ, Witola WH, Sidney J, Alexander J, Sette A, Maewal A, McLeod R: Towards an immunosense vaccine to prevent toxoplasmosis: Protective *Toxoplasma gondii* epitopes restricted by HLA-A*0201. *Vaccine* 2011, **29**(4):754–762.

29. Del L, Yácono M, Farran I, Becher ML, Sander V, Sánchez VR, Martín V, Veramendi J, Clemente M: **A chloroplast-derived *Toxoplasma gondii* GRA4 antigen used as an oral vaccine protects against toxoplasmosis in mice.** *Plant Biotechnol J* 2012, **10**(9):1136–1144.
30. Ye L, Zeng R, Bai Y, Roopenian DC, Zhu X: **Efficient mucosal vaccination mediated by the neonatal Fc receptor.** *Nat Biotechnol* 2011, **29**(2):158–163.
31. Abdul-Wahid A, Faubert G: **Mucosal delivery of a transmission-blocking DNA vaccine encoding *Giardia lamblia* CWP2 by *Salmonella typhimurium* bactofection vehicle.** *Vaccine* 2007, **25**(50):8372–8383.
32. Chin'ombe N, Ruhanya V: **Recombinant *Salmonella* Bacteria Vectoring HIV/AIDS Vaccines.** *Open Virol J* 2013, **7**:121–126.
33. Lynch LF, Lynch MI, Ferreira RS, Vasconcelos MS, Melo N, Ferreira S, Malagueño E: **Ocular toxoplasmosis: evaluation of lacrimal-specific secretory IgA levels in both patients with active and inactive phases of the disease.** *Mem Inst Oswaldo Cruz* 2011, **106**(5):625–628.
34. Wang Y, Wang G, Zhang D, Yin H, Wang M: **Screening and identification of novel B cell epitopes of *Toxoplasma gondii* SAG1.** *Parasit Vectors* 2013, **6**:125.
35. Bhadra R, Gigley JP, Khan IA: **The CD8 T-cell road to immunotherapy of toxoplasmosis.** *Immunotherapy* 2011, **3**(6):789–801.
36. Yarovinsky F: **Innate immunity to *Toxoplasma gondii* infection.** *Nat Rev Immunol* 2014, **14**(2):109–121.
37. Jongert E, Lemiere A, Van Ginderachter J, De Craeye S, Huygen K, D'Souza S: **Functional characterization of in vivo effector CD4(+) and CD8(+) T cell responses in acute Toxoplasmosis: an interplay of IFN-gamma and cytolytic T cells.** *Vaccine* 2010, **28**(13):2556–2564.
38. Menard LC, Minns LA, Darce S, Mielcarz DW, Foureau DM, Roos D, Dzierszynski F, Kasper LH, Buzoni-Gatel D: **B cells amplify IFN-gamma production by T cells via a TNF-alpha-mediated mechanism.** *J Immunol* 2007, **79**(7):4857–4866.

doi:10.1186/1756-3305-7-145

Cite this article as: Cong *et al.*: Comparative efficacy of a multi-epitope DNA vaccine via intranasal, peroral, and intramuscular delivery against lethal *Toxoplasma gondii* infection in mice. *Parasites & Vectors* 2014 **7**:145.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

