

RESEARCH

Open Access

Synergistic effects of neck circumference and metabolic risk factors on insulin resistance: the Cardiometabolic Risk in Chinese (CRC) study

Jun Liang^{1*†}, Fei Teng^{1†}, Xuekui Liu¹, Caiyan Zou¹, Yu Wang¹, Lianjun Dou¹, Zilin Sun² and Lu Qi^{3,4}

Abstract

Objectives: Recent studies have associated neck circumference (NC) with insulin resistance (IR). We examined whether such relation was modified by other metabolic risk factors.

Methods: The study samples were from a community-based health examination survey in central China. A total of 2588 apparently healthy Chinese men and women were included.

Results: Plasma levels of total cholesterol (TC), HDL-C, uric acid (UA) and diastolic blood pressure (DBP) were independently associated with NC after adjusted for age, sex, body mass index (BMI), waist circumference (WC) and hip circumference (HC) ($P = 0.009, 0.001, 0.015$ and 0.015 , respectively). We observed significant interactions of NC with triglyceride (TG) and UA (all the p for interaction = 0.001) in relation to HOMA-IR. It appeared that the associations between NC and HOMA-IR were more evident in those with higher UA or TG level.

Conclusions: Our data indicate that in apparently healthy Chinese adults, there were synergistic effects of UA, TG and neck circumference on insulin resistance.

Keywords: Insulin resistance, Neck circumference, Uric acid, Triglyceride, Synergistic effects

Background

Neck circumference is a proxy for upper-body fat and a reliable, simple, time saving screening measure, among numerous others such as BMI and WC, for identification of individuals with excess body fat or its abnormal distribution. In Framingham study, NC was positively associated with insulin resistance [1]. In our recent analysis, we found NC is independently related to IR in Chinese [2].

NC has been also related to various metabolic risk factors such as blood pressure and lipids [3,4], independent of overall adiposity (body mass index) and central obesity (waist circumference or visceral adipose tissue) [1]. It remains unclear whether these risk factors modify the relation between NC and IR [5].

In the present study, we comprehensively analyzed the associations of NC and other metabolic risk factors, and their interactions in relation to insulin resistance in a large cohort of apparently healthy Chinese adults.

Materials and methods

Study population

In the Cardiometabolic Risk in Chinese (CRC) Study, we performed a community-based health examination survey for 6,431 individuals (18–93 y) who were randomly selected from residents living in the urban area of central China, in 2009. The details of this study have been presented elsewhere [6–9]. Written consents were obtained from all the participants. The CRC study was reviewed and approved by the ethics committee of the Central Hospital of Xuzhou, China and sponsored by Jiangsu Health International Exchange Program. NC was not measured in all the CRC participants, for the present study, we included adult men and women (≥ 20 y) who were successfully measured NC, and relevant cardiometabolic markers including fasting glucose, insulin, lipids,

* Correspondence: mwj521@163.com

†Equal contributors

¹Department of Endocrinology, Xuzhou Central Hospital, Xuzhou Clinical School of Xuzhou Medical College; Affiliated Hospital of Southeast University, 199# South Jiefang Road, Xuzhou 221009, Jiangsu, China
Full list of author information is available at the end of the article

blood pressure, BMI, WC. We excluded the potential patients with history of diabetes and those with fasting glucose ≥ 7.0 mmol/L and/or 2 h OGTT ≥ 11.1 mmol/L and/or HbA1c $\geq 6.5\%$, also excluded subjects with goiter and other neck masses and deformity with ultrasound. In total 2588 subjects were included. There was no significant difference in the clinical characteristics between the participants of the present analysis and those who were not included.

Anthropometric measures

NC (cm) was measured with head erect and eyes facing forward, horizontally at the upper margin of the laryngeal prominence with a flexible tape [2]. WC (cm) was measured to the nearest 1 cm in the horizontal plane at the midpoint between the lowest rib and the iliac crest. HC (cm) was measured at the level of maximal protrusion of the gluteal muscles. ALL the human body dimensions have three readings and taken the average. WHR was calculated as WC divided by HC. Height and body weight were measured with participants standing without shoes and heavy outer garments. BMI was calculated as weight (in kilograms) divided by height (in meters) squared. Percentage of body fat was estimated with bioelectrical impedance, using the Omron Body Fat Analyzer HBF-306. Subjects with a body fat percentage measured by bipolar bioelectrical impedance analysis (BF% (IMP)) $< \text{or} = 20.9\%$ were considered normal weight, while subjects with a BF% (IMP) $> \text{or} = 21.0\%$ were considered overweight. Blood pressure (BP) was measured after the subject had rested for at least 5 minutes with a mercury manometer by doctors. Three measurements, 60 seconds apart, were taken. The mean of the three measurements was used for analysis.

Assessment of biomarkers

Venous blood sample was drawn from all subjects after an overnight fast (8–12 h). After blood was drawn, specimens were allowed to clot at room temperature for 1–3 h and serum was separated. Immediately following clotting serum was separated by centrifugation for 15 min at 3,000 rpm. Fasting blood specimens were collected for measurement of uric acid (UA), glucose, total cholesterol (TC), triglyceride (TG), high density lipoprotein cholesterol (HDL-C), and low density lipoprotein cholesterol (LDL-C). All biochemical assays were determined by enzymatically on an autoanalyzer (Type 7600, Hitachi Ltd, Tokyo, Japan). Serum insulin concentration was determined by competitive radioimmunoassay (Roche, E170, Germany).

Insulin Resistance was assessed using the Homeostasis Model Assessment-Insulin Resistance (HOMA-IR) index, derived from plasma glucose and insulin [10]. According to recommendations from the European Group for the

Study of Insulin Resistance, the HOMA-IR index was calculated according to the formula: Fasting plasma glucose (mmol/l) \times Fasting serum insulin (U/ml) /22.5. By convention, individuals in the upper quartile in HOMA –IR index for non-diabetics are defined as insulin resistance [11].

Statistical analyses

Fasting glucose, insulin, TG and TC levels were logarithmically transformed to improve the normality. Linear regression models were used to evaluate associations between NC (in sex-specific quintiles) and metabolic markers, adjusting for covariates. Interactions between NC and metabolic risk factors were tested by introduction of the cross-product terms in the linear regression models. All reported P values were two tailed. Variables with P values of < 0.05 were considered statistically significant. Data management and statistical analysis were conducted using SAS statistical software (SAS Institute Inc., Cary, NC, USA).

Results

The characteristics of the study participants by neck circumference

In total 2588 apparently healthy Chinese adults were included in the present study. The mean age of study population was 44.3 (20–87) years and the mean BMI was 24.4 kg/m². Table 1 shows the characteristics of the study participants according to NC levels (in quintiles). NC was positively related to BMI, WC, HC, and WHR (all $p < 0.0001$).

Association of neck circumference with metabolic risk factors

After adjustment for age and sex (Table 2), TC, TG, HDL-C, LDL-C, UA, SBP and DBP were independently associated with NC (all the $P = 0.0001$). In the models further adjusting for BMI, WC and HC, the associations of NC with TG, HDL-C, UA and DBP remained significant ($P = 0.009, 0.001, 0.015$ and 0.015 , respectively). The difference in these markers between the two extreme groups quintiles 1(Q1) and quintiles 5(Q5) were 1.26 mmol/L, 0.34 mmol/L, 118.56, $\mu\text{mol/L}$ and 12 mmHg respectively for TG, HDL-C, UA and DBP.

Synergistic effects of serum uric acid, TG and NC on insulin resistance

We further tested the interactions between NC (in quintiles) and these metabolic risk factors, and found significant interactions for TG and UA (both in tertiles) in relation to HOMA-IR (all the P for interaction = 0.001). Figure 1 shows the joint effects of NC and UA, as well as NC and TG in relation to HOMA-IR. For both UA and TG, we found that the associations between NC and

Table 1 Characteristics of participants by NC in quintiles

Variables	NC (in quintiles, cm)					P for trend
	Q1(≤33)	Q2(34–35)	Q3(36–37)	Q4(38–39)	Q5(≥40)	
N	801	455	427	517	388	
Age, years	43.1 ± 11.3	45.3 ± 12.1	43.3 ± 12.9	44.5 ± 11.3	43.4 ± 9.5	<0.0001
BMI, kg/m ²	21.9 ± 2.4	23.1 ± 2.9	24.3 ± 2.1	25.8 ± 2.1	27.8 ± 2.6	<0.0001
BF, %	28.8 ± 4.8	26.7 ± 6.5	25.1 ± 4.6	26.1 ± 4.2	27.4 ± 3.9	<0.0001
WC, cm	75.5 ± 6.7	81.5 ± 6.3	86.4 ± 5.5	91.3 ± 5.9	97.0 ± 6.9	<0.0001
HC, cm	92.5 ± 5.1	95.0 ± 5.2	96.9 ± 4.3	99.9 ± 4.6	104.4 ± 5.2	<0.0001
WHR, %	0.82 ± 0.06	0.86 ± 0.05	0.89 ± 0.04	0.91 ± 0.05	0.93 ± 0.05	<0.0001

The values are presented as mean ± standard deviation.

Abbreviations: BMI Body mass index, BF Percent, body fat rate, WC Waist circumference, HC Hip circumference, WHR Waist/hip ratio.

HOMA-IR were more evident among those with higher levels of these markers.

Discussion

In a large cohort of apparently healthy Chinese adults, we found that NC was related to metabolic risk factors including TG, HDL-C, UA and DBP. In our previous study, we found significant association between NC and increasing trend of HOMA-IR, even after adjusting for age and other cardiometabolic risk factors [2]. In this study, we found that plasma levels of UA or TG modified the relation between NC and HOMA-IR.

Our findings are in line with several recent studies in Caucasians and Chinese diabetic patients. Selim et al. found that NC was associated with HDL, FBG, TC, TG and blood pressure in Turkey obese children [4]. In the Framingham study, it was found that NC was associated with insulin resistance and cardiovascular risk factors. After adjustment for visceral adipose tissue, NC was positively associated with SBP, DBP, TC and FBG [1]. Locally acting fat depots may contribute to obesity complications and cardiovascular diseases, through direct paracrine effects, and the cardiovascular risk conferred especially by visceral and upper body adiposity. The

carotid arteries are encased in fat, and total upper-body subcutaneous fat is estimated by NC. The recent study highlight NC has been independently correlated with cardiometabolic risk factors above and beyond that of other adiposity measures [12,13].

Several mechanisms may be underlying the associations of high NC and metabolic risk factors. It has been documented that high NC was a significant predictor of obstructive sleep apnea syndrome (OSAS) [3], which has been associated with aggravates glycemic control, even at the earliest stages of glucose intolerance. In addition, intermittent hypoxemia and sleep fragmentation increases the risk of IR [14]. Our data, together with evidence from other studies, suggest that body fat accumulated in upper body segment may also contribute to the adverse consequence.

The synergistic effects of NC with UA and TG on IR remained unclear. Hyperuricemia has been associated with insulin resistance [15], and excess adiposity [16]. In our previous analysis, we found high UA levels were related to various metabolic disorders [8]. High-sensitivity C-reactive protein (hs-CRP) level is often higher in hyperuricemic patients than in the general population, hs-CRP level was found to be an independent predictor


Table 2 Associations between NC and biochemical risk factors

Variables	NC (in quintiles, cm)					P1	P2
	Q1(≤33)	Q2(34–35)	Q3(36–37)	Q4(38–39)	Q5(≥40)		
TC, mmol/L	4.90 ± 0.86	4.92 ± 0.86	5.04 ± 0.87	5.10 ± 0.86	5.18 ± 0.93	0.0001	0.646
TG, mmol/L	1.03 ± 0.95	1.32 ± 1.06	1.71 ± 1.61	2.05 ± 1.68	2.29 ± 1.99	0.0001	0.009
HDL-C, mmol/L	1.44 ± 0.29	1.31 ± 0.29	1.22 ± 0.27	1.13 ± 0.56	1.10 ± 0.24	0.0001	0.001
LDL-C, mmol/L	2.84 ± 0.69	2.91 ± 0.72	3.02 ± 0.78	3.05 ± 0.75	3.11 ± 0.79	0.0001	0.909
UA, μmol/L	233.97 ± 53.55	277.68 ± 67.89	319.40 ± 65.53	338.75 ± 71.79	352.53 ± 75.39	0.0001	0.015
SBP, mmHg	115 ± 14	121 ± 15	125 ± 14	127 ± 16	131 ± 15	0.0001	0.843
DBP, mmHg	73 ± 10	77 ± 7	80 ± 10	82 ± 11	85 ± 11	0.0001	0.015

The values are presented as mean ± standard deviation.

Abbreviations: TC Total cholesterol, TG Triglyceride, HDL-C High density lipoprotein cholesterol, LDL-C Low density lipoprotein cholesterol, UA Uric acid, SBP Systolic blood pressure, DBP Diastolic blood pressure.

P1 was for models with age and sex adjusted; P2 was for models with age, sex, BMI, WC, and HC adjusted.



of homeostatic model assessment with insulin resistance [17]. In addition, soluble uric acid could increase tissue levels of NADPH oxidase and the generation of reactive oxygen species (ROS) in mature adipose tissue; oxidative-stressed adipose tissue shows decreased sensitivity to insulin as a risk factor of insulin resistance [18]. Uric acid levels have also been found to be positively correlated with the number of obstructive respiratory episodes and oxygen desaturations during sleep [19]. NC has been positively related to blood levels of TG [20]. It was recently found that NC was associated with atherogenic dyslipidaemia beyond BMI and waist circumference in both men and women [21]. Our data were consistent with an additive effect of NC and high levels of UA or TG on insulin resistance.

To our knowledge, study about the interactions of NC and metabolic risk factors on insulin resistance in general population of Chinese adults are lacking. The cross-sectional nature of the study to some extent limits its interpretation as to causality, prospective studies and follow-up data on our participants are warranted to confirm the causal relation. In addition, neck circumference is used to represent upper body subcutaneous fat, we did not perform radiographic measures to quantify this depot of fat directly. Physical fitness is very important on metabolic risk reduction, but in present study, we did not well analyze the interactions between metabolic risk factors and life style intervention. Finally, the study was performed in a Chinese population, further studies in other populations of different ethnicities are warranted to verify our findings.

In conclusion, we found significant associations of high NC with a variety of cardiometabolic risk factors including UA, TG, HDL-C, DBP in apparently healthy Chinese adults. In addition, we found significant synergistic effects of neck circumference with UA and TG on insulin resistance.

Abbreviations

NC: Neck circumference; IR: Insulin resistance; HOMA-IR: Homeostasis Model Assessment-Insulin Resistance; BMI: Body mass index; BF%: Body fat rate; WC: Waist circumference; HC: Hip circumference; WHR: Waist/hip ratio; TC: Total cholesterol; TG: Triglyceride; HDL-C: High density lipoprotein cholesterol; LDL-C: Low density lipoprotein cholesterol; UA: Uric acid; SBP: Systolic blood pressure; DBP: Diastolic blood pressure.

Competing interests

The authors declare that they have no competing of interests.

Authors' contributions

JL and FT contributed to the study design, study conduct/data collection, data analysis, and writing of the manuscript. XL contributed to data analysis and critically re-reading the manuscript. CZ and YW contributed to the study design, study conduct/data collection. LD and HL contributed to study conduct/data collection, data analysis. ZS and LQ contributed to the study design, data analysis, and writing of the manuscript. All authors read and approved the final manuscript.

Acknowledgments

Dr. Qi's research is supported by NIH grants DK091718, HL071981, the American Heart Association Scientist Development Award, and the Boston Obesity Nutrition Research Center (DK46200). Dr. Jun Liang's research is sponsored by Jiangsu Health International Exchange Program and supported by Xuzhou Medical Leading Talent Grant and Xuzhou Science and Technology Grant (XM13B066, XZZD1242). Fei Teng's research is sponsored by Xuzhou Science and Technology Grant (KC14SH072). We thank all subjects for participating in this study.

Author details

¹Department of Endocrinology, Xuzhou Central Hospital, Xuzhou Clinical School of Xuzhou Medical College; Affiliated Hospital of Southeast University, 199# South Jiefang Road, Xuzhou 221009, Jiangsu, China. ²Institute of diabetes, Medical School, Southeast University, Nanjing 210009, Jiangsu, China. ³Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts 02115, USA. ⁴Department of Medicine, Channing Laboratory, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts 02115, USA.

Received: 17 August 2014 Accepted: 23 October 2014

Published: 1 November 2014

References

1. Preis SR, Massaro JM, Hoffmann U, D'Agostino RB Sr, Levy D, Robins SJ, Meigs JB, Vasan RS, O'Donnell CJ, Fox CS: **Neck circumference as a novel measure of cardiometabolic risk: the Framingham Heart study.** *J Clin Endocrinol Metab* 2010, **95**:3701-3710.

2. Liang J, Teng F, Li Y, Liu X, Zou C, Wang Y, Li H, Qi L: **Neck circumference and insulin resistance in Chinese adults: the Cardiometabolic Risk in Chinese (CRC) study.** *Diabetes Care* 2013, **36**(9):e145–e146.
3. Onat A, Hergenç G, Yüksel H, Can G, Ayhan E, Kaya Z, Dursunoğlu D: **Neck circumference as a measure of central obesity: associations with metabolic syndrome and obstructive sleep apnea syndrome beyond waist circumference.** *Clin Nutr* 2009, **28**:46–51.
4. Kurtoglu S, Hatipoglu N, Mazicioglu MM, Kondolot M: **Neck circumference as a novel parameter to determine metabolic risk factors in obese children.** *Eur J Clin Invest* 2012, **42**:623–630.
5. Ahbab S, Ataoğlu HE, Tuna M, Karasulu L, Cetin F, Temiz LU, Yenigün M: **Neck circumference, metabolic syndrome and obstructive sleep apnea syndrome; evaluation of possible linkage.** *Med Sci Monit* 2013, **13**:111–117.
6. Liang J, Zhou N, Teng F, Zou C, Xue Y, Yang M, Song H, Qi L: **Hemoglobin A1c levels and aortic arterial stiffness: the Cardiometabolic Risk in Chinese (CRC) study.** *PLoS One* 2012, **7**:e38485.
7. Liang J, Li Y, Zhou N, Teng F, Zhao J, Zou C, Qi L: **Synergistic effects of serum uric acid and cardiometabolic risk factors on early stage atherosclerosis: the cardiometabolic risk in chinese study.** *PLoS One* 2012, **7**:e51101.
8. Liang J, Xue Y, Zou C, Zhang T, Song H, Qi L: **Serum uric acid and prehypertension among Chinese adults.** *J Hypertens* 2009, **27**:1761–1765.
9. Teng F, Zhu R, Zou C, Xue Y, Yang M, Song H, Liang J: **Interaction between serum uric acid and triglycerides in relation to blood pressure.** *J Hum Hypertens* 2011, **25**:686–691.
10. Wallace TM, Matthews DR: **The assessment of insulin resistance in man.** *Diabet Med* 2002, **19**:527–534.
11. Balkau B, Charles MA: **Comment on the provisional report from the WHO consultation. Europe an Group for the Study of Insulin Resistance (EGIR).** *Diabet Med* 1999, **16**:442–443.
12. Rosenquist KJ, Massaro JM, Pencina KM, D'Agostino RB, Beiser A, O'Connor GT, O'Donnell CJ, Wolf PA, Polak JF, Seshadri S, Fox CS: **Neck circumference, carotid wall intima-media thickness, and incident stroke.** *Diabetes Care* 2013, **36**(9):e153–e154.
13. Fitch KV, Stanley TL, Looby SE, Rope AM, Grinspoon SK: **Relationship between neck circumference and cardio metabolic parameters in HIV-infected and non- HIV-infected adults.** *Diabetes Care* 2011, **34**:1026–1031.
14. Steiropoulos P, Papanas N, Bouros D, Maltezos E: **Obstructive sleep apnea aggravates glycemic control across the continuum of glucose homeostasis.** *Am J Respir Crit Care Med* 2010, **182**:286.
15. Li C, Hsieh MC, Chang SJ: **Metabolic syndrome, diabetes, and hyperuricemia.** *Curr Opin Rheumatol* 2013, **25**(2):210–216.
16. Yue JR, Huang CQ, Dong BR: **Association of serum uric acid with body mass index among long-lived Chinese.** *Exp Gerontol* 2012, **47**(8):595–600.
17. Zhu Y, Hu Y, Huang T, Zhang Y, Li Z, Luo C, Luo Y, Yuan H, Hisatome I, Yamamoto T, Cheng J: **High uric acid directly inhibits insulin signalling and induces insulin resistance.** *Biochem Biophys Res Commun* 2014, **447**(4):707–714.
18. Spalding KL, Arner E, Westermark PO, Bernard S, Buchholz BA, Bergmann O, Blomqvist L, Hoffstedt J, Näslund E, Britton T, Concha H, Hassan M, Rydén M, Frisén J, Arner P: **Dynamics of fat cell turnover in humans.** *Nature* 2008, **453**(7196):783–787.
19. Ruiz García R, Sánchez Armengol Á, Luque Crespo E: **Blood uric acid levels in patients with sleep-disordered breathing.** *Arch Bronconeumol* 2006, **42**(10):492–500.
20. Stabe C, Vasques AC, Lima MM, Tambascia MA, Pareja JC, Yamanaka A, Geloneze B: **Neck circumference as a simple tool for identifying the metabolic syndrome and insulin resistance: results from the Brazilian Metabolic Syndrome Study.** *Clin Endocrinol (Oxf)* 2013, **78**(6):874–881.
21. Vallianou NG, Evangelopoulos AA, Bountziouka V, Vogiatzakis ED, Bonou MS, Barbetseas J, Avgerinos PC, Panagiotakos DB: **Neck circumference is correlated with triglycerides and inversely related with HDL cholesterol beyond BMI and waist circumference.** *Diabetes Metab Res Rev* 2013, **29**(1):90–97.

doi:10.1186/1758-5996-6-116

Cite this article as: Liang et al.: Synergistic effects of neck circumference and metabolic risk factors on insulin resistance: the Cardiometabolic Risk in Chinese (CRC) study. *Diabetology & Metabolic Syndrome* 2014 **6**:116.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

