



## Increased expression of native cytosolic Cu/Zn superoxide dismutase and ascorbate peroxidase improves tolerance to oxidative and chilling stresses in cassava (*Manihot esculenta* Crantz)

Xu *et al.*

RESEARCH ARTICLE

Open Access

# Increased expression of native cytosolic Cu/Zn superoxide dismutase and ascorbate peroxidase improves tolerance to oxidative and chilling stresses in cassava (*Manihot esculenta* Crantz)

Jia Xu<sup>1,2</sup>, Jun Yang<sup>1</sup>, Xiaoguang Duan<sup>2</sup>, Yueming Jiang<sup>3</sup> and Peng Zhang<sup>1,2\*</sup>

## Abstract

**Background:** Cassava (*Manihot esculenta* Crantz) is a tropical root crop, and is therefore, extremely sensitive to low temperature; its antioxidative response is pivotal for its survival under stress. Timely turnover of reactive oxygen species (ROS) in plant cells generated by chilling-induced oxidative damages, and scavenging can be achieved by non-enzymatic and enzymatic reactions in order to maintain ROS homeostasis.

**Results:** Transgenic cassava plants that co-express cytosolic superoxide dismutase (SOD), *MeCu/ZnSOD*, and ascorbate peroxidase (APX), *MeAPX2*, were produced and tested for tolerance against oxidative and chilling stresses. The up-regulation of *MeCu/ZnSOD* and *MeAPX2* expression was confirmed by the quantitative reverse transcriptase-polymerase chain reaction, and enzymatic activity analyses in the leaves of transgenic cassava plant lines with a single-transgene integration site. Upon exposure to ROS-generating agents, 100  $\mu$ M ROS-generating reagent methyl viologen and 0.5 M H<sub>2</sub>O<sub>2</sub>, higher levels of enzymatic activities of SOD and APX were detected in transgenic plants than the wild type. Consequently, the oxidative stress parameters, such as lipid peroxidation, chlorophyll degradation and H<sub>2</sub>O<sub>2</sub> synthesis, were lower in the transgenic lines than the wild type. Tolerance to chilling stress at 4°C for 2 d was greater in transgenic cassava, as observed by the higher levels of SOD, catalase, and ascorbate-glutathione cycle enzymes (e.g., APX, monodehydroascorbate reductase, dehydroascorbate reductase and glutathione reductase) and lower levels of malondialdehyde content.

**Conclusions:** These results suggest that the expression of native cytosolic SOD and APX simultaneously activated the antioxidative defense mechanisms via cyclic ROS scavenging, thereby improving its tolerance to cold stress.

**Keywords:** *Manihot esculenta* Crantz, Cytosolic superoxide dismutase, Cytosolic ascorbate peroxidase, Reactive oxygen species scavenging, Abiotic stress resistance

## Background

Cassava (*Manihot esculenta* Crantz) is an important tropical root crop that plays an important role not only in ensuring food security, but also in various bioindustrial applications such as animal feed, modified starch and

biofuels [1,2]. Because cassava is native to tropical regions, it is extremely cold-sensitive; its growth is drastically affected at temperatures lower than 18°C, and the plant cannot survive for long after exposure to freezing conditions [3-5]. Therefore, low temperatures and freezing conditions are the most important limiting factors for the cultivating location, as well as productivity. The damage of apical shoot seems to be more critical than other parts of the cassava plant [5]. Improving the tolerability of the cassava plant to multiple stresses has therefore, become a major objective of cassava breeders, especially in subtropical regions [4,5]. Under cold conditions, up-regulation of

\* Correspondence: zhangpeng@sibs.ac.cn

<sup>1</sup>Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai Chenshan Botanical Garden, 3888 Chenhua Road, Shanghai 201602, China

<sup>2</sup>National Key Laboratory of Plant Molecular Genetics and National Center for Plant Gene Research (Shanghai), Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200032, China

Full list of author information is available at the end of the article

reactive oxygen species (ROS) turnover and scavenging in cassava has been reported, and therefore, genetic manipulation of intracellular ROS level might be an effective approach in improving tolerance to abiotic stresses in this tropical crop [5-7].

In the ROS scavenging system that is responsible for homeostasis in plant cells, superoxide dismutases (SODs, EC 1.15.1.1), enzymes that catalyze the dismutation of superoxide into oxygen and hydrogen peroxide, provide the first line of defense against ROS in various subcellular compartments, i.e. chloroplast, mitochondria and cytosol [8]. Essentially, there are three types of SODs, each containing either manganese, iron, or copper plus zinc as a prosthetic group [9]. Along with other ROS scavenging mechanisms like catalase (CAT; EC 1.11.1.6), glutathione peroxidases (GPXs) and peroxiredoxin reductases (PrxRs), and the ascorbate–glutathione (ASC-GSH) cycle, the ROS levels are maintained in a homeostatic state. In the ASC-GSH cycle, using ascorbate as an electron donor, ascorbate peroxidase (APX, EC 1.11.1.1) scavenges potentially harmful hydrogen peroxide to water from the chloroplasts and mitochondria, as well as other organelles [10,11]. Therefore, the formation of toxic hydroxyl radicals by superoxide and hydrogen peroxide can be controlled by the combined enzymatic actions of SOD and APX [12].

Transgenic plants that express SOD or APX have shown enhanced tolerance to multiple stresses [13]. For example, over-expression of different SODs (FeSOD, MnSOD or Cu/ZnSOD) in transgenic plants of tomato, rice, poplar, alfalfa, etc., showed increased tolerance to methyl viologen (MV), ozone, high salinity, chilling or other stresses [14-17]. Transgenic plants have also demonstrated an increased tolerance against various abiotic stresses by the expression of either cytosolic- or organelle-targeted cytosolic APX [18-22]. However, some reports suggest no change in response to oxidative or environmental stress with the expression of a single antioxidant enzyme [23,24]. These contradictory findings may be due to the complex network of plant antioxidant defenses, which possibly confer a higher tolerance to oxidative stress by pyramiding or stacking of multiple genes in a single genotype [25]. The gene-stacking approach entails manipulation of two or more desirable enzymes mediating the ROS turnover and scavenging pathways, in improving the abiotic stress tolerance in plants.

Indeed, co-expression of two distinct ROS-scavenging enzymes, such as SOD and other ROS-scavenging enzymes, in the chloroplasts or cytosol in transgenic plants has a synergistic effect in increasing the levels of abiotic stress resistance. For example, coupled expression of Cu/ZnSOD and APX in transgenic plants of *Festuca arundinacea*, potato, tobacco, sweet potato and plum led to increased tolerance to multiple abiotic stresses,

e.g., the herbicide methyl viologen (MV), chilling, high temperature and drought [12,26-29]. Payton et al. [30] showed that co-expression of glutathione reductase (GR, EC 1.6.4.2) and APX in cotton improved antioxidant enzyme activity during moderate chilling at high light intensity, in chloroplasts [30]. Co-expression of the *Suaeda salsa* glutathione S-transferase (GST) and CAT1 in rice also caused tolerance to stresses caused by salt and paraquat [31]. Taken together, these data indicated that the combination of transgenes encoding different ROS-scavenging enzymes in various subcellular compartments might have a synergistic effect in improving stress tolerance.

Lately, plant breeders and biotechnologists have appreciated the molecular insights and advances in cassava abiotic stress resistance, on a global scale. Apart from the various approaches from traditional breeding to field evaluation [32], studies of cassava response to drought or cold stress at the molecular level have reportedly used the “omics” technology, e.g., expressed sequence tags, cDNAs and oligonucleotide microarray [5,33-38]. However, few studies on improved tolerance to environmental stresses using genetic engineering have been reported [7,39]. Senescence-induced expression of the isopentenyl transferase gene in cassava showed increased drought resistance, as observed by the elevated content of cytokinin in mature leaves, and prolonged leaf life [39]. Enhanced ROS scavenging by simultaneous expression of cytosolic *MeCu/ZnSOD* and peroxisomal *MeCAT1* in transgenic cassava also confirmed the improved tolerance towards drought and cold temperatures [7].

In the present study, transgenic cassava plants co-expressing both cytosolic *MeCu/ZnSOD* (Genbank accession No. AY642137) and cytosolic *MeAPX2* (GenBank accession No. AY973622) showed enhanced ROS scavenging capacity, thereby leading to enhanced tolerance to oxidative stresses that was induced by MV, H<sub>2</sub>O<sub>2</sub> as well as chilling. Our results suggest that manipulation of ROS-scavenging enzyme systems by the overexpression of both cytosolic SOD and APX is a worthwhile approach to produce transgenic plants with enhanced tolerance to a wide range of abiotic stresses.

## Results

### Subcellular localization of *MeCu/ZnSOD* and *MeAPX2*

Green fluorescent protein (GFP) was the fusion protein used for subcellular localization of *MeCu/ZnSOD* and *MeAPX2*. The control, CaMV 35S-eGFP construct, exhibited GFP fluorescence in the cytoplasm and nucleus of agroinfiltrated cells of *N. benthamiana* leaves. The *MeCu/ZnSOD* has been reported as a cytosolic SOD isoform in our previous report [7]. The *MeAPX2* was located in the cytosol of the leaf cells, as indicated by the fused GFP fluorescence (Additional file 1) that was




consistent with the signal peptide sequences predicted by Reilly et al. [38]. Therefore, both enzymes used in the study were cytosolic proteins.

#### Molecular characterization of the transgenic plants

Four independent transgenic plant lines (named SA1, SA2, SA4 and SA6) harboring the *MeCu/ZnSOD* and *MeAPX2* gene-expressing cassettes (Figure 1a) were produced by the use of *Agrobacterium*-mediated transformation and subcultured *in vitro* regularly. Confirmation of the single integration event of the pC-P54::MeCu/ZnSOD-35S::MeAPX2 T-DNA in these transgenic lines were carried out by the Southern blotting technique using *Xba*I-digested cassava genomic DNA, which were extracted from leaves of *in vitro* plants and hybridized

with DIG-labeled *HPT* probe (Figure 1b, left panel). No signals were detected in the wild-type (WT) plants. However, when these WT plants were hybridized with the *MeAPX2* probe (Figure 1b, right panel), two bands of ~13 kb and ~6.6 kb were detected, indicating the possibility of two *APX* homologs in the cassava genome. The transgenic SA lines showed an additional band in their hybridization pattern, confirming that the transgenic plants were integrated at a single site (Figure 1b, right panel). All transgenic plants except SA2 line thrived successfully in the greenhouse and field, with a relatively normal leaf phenotype, growth capacity and root tuberization, similar to WT plants (Figure 1c); the SA2 line had a slightly dark-green and curled leaves when cultivated in the field. No significant differences of



**Figure 1** Molecular and phenotypic analysis of SA transgenic cassava coupled expression of cytosolic *MeCu/ZnSOD* and cytosolic *MeAPX2* genes. **(a)** Schematic presentation of the T-DNA region of pC-P54::MeCu/ZnSOD-35S::MeAPX2 with unique *Xba*I site. LB, left border; RB, right border of T-DNA;  $T_{35S}$ , CaMV 35S terminator;  $T_{nos}$ , NOS terminator;  $P_{35S}$ , cauliflower mosaic virus 35S promoter;  $P_{54/1.0}$ , vascular-specific promoter p54/1.0; *HPT*, hygromycin phosphotransferase. **(b)** Southern blot analysis of transgenic and WT cassava plants for transgene integration. Transgene integration patterns in SA lines detected *Xba*I-digested genomic DNA by *HPT* (left panel) and *MeAPX2* (right panel) probes. gAPX, genomic DNA of cassava ascorbate peroxidase; T-APX2, transgene *MeAPX2*; M,  $\lambda$  HindIII DIG-labeled molecular marker; P, plasmid pC-P54::MeCu/ZnSOD-35S::MeAPX2; WT, wild-type control. Numbers indicate different transgenic lines. **(c)** Plant growth status and phenotype evaluation in field. WT, wild-type control; SA1, SA2 and SA4, independent transgenic cassava plant lines.



yield were found between WT and transgenic plant lines (Additional file 2).

The basic transcriptional levels of *MeCu/ZnSOD* and *MeAPX2* in cassava leaves were confirmed by quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR) (Figure 2a). The expressions of *MeCu/ZnSOD* and *MeAPX2* in transgenic lines were higher by up to 20- and 5-fold, respectively, when compared to that of WT plants. SOD expression level was about 5-times higher than APX2 in the SA lines, indicating that the *p54* promoter is stronger than CaMV 35S promoter in cassava, which is in agreement with the previous report [40]. Further, the changes in SOD and APX isoenzyme

activity from leaves on non-denaturing gel were also confirmed (Figure 2b). A characteristic SOD isoenzyme banding pattern was found in all cassava leaves with Cu/ZnSOD, MnSOD and FeSOD [6], but the intensity of SOD bands was higher in all transgenic lines than the WT. Similarly, a stronger APX isoenzyme band was also observed in all transgenic lines than that of WT (Figure 2b). These results indicate that the heightened production of SODs and APXs were achieved because of the transgene expression in transgenic cassava.

#### Higher protoplast viability and mitochondrial integrity of mesophyll cells under H<sub>2</sub>O<sub>2</sub> stress

Cell death and loss of mitochondrial integrity are indicators of stress damage. The viability of extracted transgenic and WT mesophyll protoplasts was above 95%, when stained by fluorescein diacetate (FDA). After 1 M H<sub>2</sub>O<sub>2</sub> treatment, the viability of transgenic protoplasts from SA1, SA2, and SA4 decreased to 77%, 74%, and 80%, respectively, but the WT showed 52% (Figure 3a), indicating significant improvement to stress tolerance. The mitochondrial integrity of mesophyll cells were




observed by rhodamine 123 (Rh 123) staining. Strong Rh 123 fluorescent signals were observed in all cassava cells prior to  $H_2O_2$  treatment, indicating normal mitochondrial activity in these cells. After  $H_2O_2$  treatment, the protoplasts of the WT plants displayed diffuse and much weaker fluorescent signals, but the protoplasts derived from the transgenic plants retained their intense fluorescence activity (Figure 3b). The difference between transgenic lines and WT was evident, which indicated that the cells in transgenic plants displayed higher  $H_2O_2$  tolerance than that in WT plants.

#### Enhanced tolerance to $H_2O_2$ -mediated oxidative stress in transgenic leaves

To evaluate the response of transgenic plants to oxidative stress, leaves showing a basal level of  $H_2O_2$  production, as indicated by 3,3'-diaminobenzidine (DAB) staining (data not shown), were selected and treated with 0.5 M  $H_2O_2$ . After 24 hours,  $H_2O_2$  molecules significantly accumulated in WT leaves (Figure 4a) at a concentration of 20.8 mmol/g, which was 1.6-times higher than that in the untreated leaves (Figure 4b). Leaves of transgenic plants showed much less  $H_2O_2$  accumulation (Figure 4a), as observed by

the elimination of the reaction product, DAB- $H_2O_2$ , from a majority of leaves from the SA4 plants (Figure 4b). For example, the  $H_2O_2$  concentration in the leaves of SA4 was 14.6 mmol/g, which was 11.6% higher than that in the untreated leaves.

The activities of SOD and APX in cassava leaves were assessed under normal and stressful conditions (Figure 4c, d). Enzyme activities were not significantly different between WT and transgenic plants under normal conditions. However, after stress induction, the SOD activity of WT plants decreased to 20% of the baseline value in 24 hours ( $1.7 U mg^{-1}$  protein, Figure 4c). In contrast, the SOD activity of all transgenic plant lines increased in 12 hours; for example, the activity level in SA4 showed a 2-fold increase as compared to 0 hours with  $23.4 U mg^{-1}$  protein. However, their activities decreased at 24 hours; the lowest value was observed in SA1 plants with  $6.6 U mg^{-1}$  protein, which was about 4-times higher than that of WT plants (Figure 4c). A consistent increase in APX activity was detected in all leaves, especially in SA1, with the activity level showing a 3-fold increased at 24 h as compared to 0 h; protein levels increased from  $0.06 \mu mol \cdot min^{-1} \cdot mg^{-1}$  protein to  $0.33 \mu mol \cdot min^{-1} \cdot mg^{-1}$




**Figure 4** Enhanced tolerance to  $H_2O_2$ -mediated oxidative stress in SA transgenic leaves. **(a)**  $H_2O_2$  accumulation in leaves detected by 3,3'-diaminobenzidine (DAB) staining. Scale bar = 0.5 cm. **(b)** Changes in levels of  $H_2O_2$  concentration between WT and transgenic cassava during 0.5 M  $H_2O_2$  treatment. **(c)** and **(d)** Changes in SOD and APX activities between WT and transgenic cassava during  $H_2O_2$  treatment. WT, wild-type control; SA1, SA2 and SA4, independent transgenic plant lines. Values represent the means of three independent experiments  $\pm$  SD. \*\*\*Significant at 1% level from WT by *t*-test.

protein. The WT only showed an increase from  $0.05 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$  protein to  $0.17 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$  protein. At 24 hours, all transgenic plant leaves showed a significant increase in activity compared to the WT plant leaves (Figure 4d).

#### Enhanced tolerance to methyl viologen (MV)-mediated oxidative stress in transgenic leaves

The effect of methyl viologen (MV), a superoxide-generating herbicide, on cassava was determined by

subjecting the leaves to  $100 \mu\text{M}$  MV for 2 days. Compared to WT plants, the extent of chlorophyll loss due to MV was significantly less in transgenic leaves (Figure 5a). As shown in Figure 5b, the chlorophyll content was not significantly different between WT and transgenic plants prior to MV treatment. After MV treatment, the chlorophyll content of WT plant leaves decreased by 75%, whereas in SA1, SA2 and SA4 leaves, the chlorophyll loss was only 21%, 37% and 38%, respectively. MV treatment causes membrane-lipid peroxidation, leading to an





increase in the malondialdehyde (MDA) content. After MV treatment, the MDA content in WT plants increased to a maximum of 45%, whereas in transgenic lines the MDA content increased by approximately 14% (Figure 5c). The average amount of MDA in WT was 11.5 nmol/g Fresh Weight (FW), which had 2.8 nmol/g FW more than SA1 line. These data suggests that transgenic cassava has less lipid peroxidation because of timely ROS scavenging. The impact of ROS scavenging enzymes in transgenic cassava was further confirmed when both SOD and APX activities were significantly enhanced after treatment (Figure 5d and e). The SOD and APX activity was approximately 1.5-times higher in transgenic plants as compared to WT plants. The highest activity of SOD reached 21 U/mg protein in SA4 line; and the highest value of APX was found in SA1 line with  $0.18 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$  protein. These data confirmed that improved performance of transgenic cassava leaves against oxidative stress is due to elevated SOD and APX activities of the ROS scavenging system.


#### Improved cold tolerance of transgenic plants

Two-month-old plants were given a chilling treatment by transferring into a growth chamber at 4°C for 2 days. After the treatment, the WT plants wilted severely, whereas the transgenic lines were slightly affected, with fewer leaves wilting (Figure 6a). Among the three transgenic cassava lines, the SA1 line was the least affected by the chilling treatment. The level of MDA increased by 7%, 8%, and 20% in SA1, SA2, and SA4 lines after chilling stress, respectively. However, WT showed a 40% increase in MDA content, which was significantly higher than that of transgenic lines (Figure 6b).

Other ROS-scavenging enzymes were monitored in both WT and SA transgenic cassava for their response to cold stress. Before treatment, both the WT and all the transgenic lines showed similar level of enzymatic activity. Upon cold treatment, increased activity of SOD, CAT, APX, MDHAR, DHAR and GR were confirmed in the transgenic lines as compared to the WT plants (Figure 7). The APX activity in SA lines showed >2-fold increase than WT (Figure 7c). The increase in SOD and CAT activities in transgenic lines was about 1.5-fold that of WT after treatment (Figure 7a, b). In the ascorbate cycle, MDHAR, DHAR and GR increased up to 43.6%, 30.6% and 28.6%, respectively (Figure 7d, e, f). However, no significant changes in enzyme activity were observed in WT before and after the treatment.

#### Discussion


The cassava plant is generally considered to be cold-sensitive, and abiotic stresses, such as low temperature and salinity, dramatically affect their growth, thereby, leading to reduced productivity [41,42]. Excessive ROS



**Figure 6** Phenotypic changes of foliage (a) and the malondialdehyde (MDA) contents of the first fully expanded leaf (b) in low temperature stressed (4°C) two-month-old cassava after 2 h treatment. WT, wild-type control; SA1, SA2 and SA4, independent transgenic plant lines.

generation, which might result in hypersensitive response and cell death in cassava, has been identified as an important indicator of such conditions [5,36]. Therefore, maintaining ROS homeostasis via ROS production and scavenging mechanisms is critical [36,43]. Optimum regulation of ROS generation and scavenging by the mobilization of various pathways has been proposed as a vital mechanism for managing stress in cassava [5,36], and our previous study had confirmed that increased resistance to abiotic stress could be achieved in transgenic cassava by over-expression of cytosolic Cu/ZnSOD and peroxisomal CAT1 [5,7]. In this study, we showed that transgenic cassava with coupled expression of cytosolic





*MeCu/ZnSOD* and cytosolic *MeAPX2* leads to improved oxidative and chilling stress resistance through elevated ROS scavenging. Our studies confirmed that the response to abiotic stress can be improved in cassava by genetic engineering techniques that combine two different

ROS-scavenging pathways, i.e., SOD/CAT and SOD/ASC-GSH cycle.

Most of ROS scavenging enzymes co-expressed targeted the chloroplast [26-28]. Recently, Faize et al. [29] and Diaz-Vivancos et al. [12] reported that over-expression

of cytosolic SOD and APX in tobacco and plums improved drought and salt tolerance. An interesting question raised is the effectiveness of the stacked ROS-scavenging enzymes with various subcellular targeting strategies in plants, undoubtedly their individual important function in stress defense [43,44]. Since the substrate for APX activity is  $H_2O_2$  that is a by-product of SOD action, their individual functions should be combined to achieve a synergistic effect on stress tolerance. Indeed, upon chilling, the enzymatic activity of MDHAR, DHAR and GR in the ASC-GSH cycle in the transgenic cassava was significantly increased (Figure 7). We noticed that, comparing to the transgenic cassava overexpressed cytosolic SOD and peroxisomal CAT [6], the SA transgenic cassava showed less tolerance to post-harvest physiological deterioration of their storage roots (data not shown). We suppose that the PPD occurrence in cassava storage root is directly linked to the oxidization process in the subcellular organelles of parenchyma cells; the action of in the peroxisome enables the timely scavenging of excess ROS generated *in situ*.

Both  $H_2O_2$  and MV treatment in the presence of light lead to the generation of superoxide radicals and  $H_2O_2$  in chloroplasts and mitochondria of plants. *In vivo* imaging of ROS using the fluorescent probe rhodamine-123, a stain readily sequestered by active mitochondria and mesophyll protoplasts, or DAB staining of cassava leaves showed improved tolerance of transgenic cassava cells to oxidative stress caused by  $H_2O_2$  and MV (Figures 3, 4). At the cellular level, the viability of the mesophyll protoplast and integrity of the mitochondrial were clearly indicated by their tolerance to  $H_2O_2$  in transgenic cassava cells (Figure 3). These results indicate that transgenic plants expressing both cytosolic SOD and cytosolic APX are able to rapidly scavenge superoxide and hydrogen peroxide at the site of generation, as well as prevent the formation of hydroxyl radicals, the most toxic ROS, prior to their interaction with target molecules, as noted in our previous report [6]. The changes were observed not only in cytosolic antioxidant system but also in the chloroplasts and mitochondria, which implied that changes in the cytosolic antioxidant defense impacts the subcellular compartments, consistent with previous reports [7,12,29,45]. Therefore, ROS scavenging system in plant cells is an integrative network for developing an antioxidant machinery through cytosol and subcellular organelle interactions.

Under cold conditions, non-transformed cassava plants showed signs of oxidative stress-induced cellular damage such as wilting, and increased lipid peroxidation of leaves (Figure 6). Cassava is very sensitive to low temperature; cold temperatures disrupt the metabolic balance of plant cells, resulting in enhanced production of ROS, e.g.  $H_2O_2$  [5]. The timely detoxification of ROS is necessary for the maintenance of the Calvin cycle and transpiration [18].

Transcriptome profiling of low temperature-exposed cassava showed an increase in transcripts and enzyme activities of ROS scavenging genes and the accumulation of total soluble sugars [5]. In the SA transgenic lines, we confirmed that the enzyme activity of SOD and APX were significantly higher than that of WT during cold treatment (Figure 7). Increased catalytic activity of APX eliminated the endogenous  $H_2O_2$  via the ASC-GSH cycle, which involves GR, DHAR and MDHAR, to restore the cellular redox state, thus, suggesting that the performance of ascorbate-glutathione cycle is essential for the regulation of an efficient system for scavenging the accumulated ROS in leaves during chilling stress. This response was correlated with the up-regulation of APX activity and by maintenance of ASC-GSH redox pools in cold-acclimated plants [46].

A higher CAT activity was also observed in the SA lines under stress conditions. This increase seemed to be involved in the removal of excess  $H_2O_2$ . Dong et al. [47] reported that peroxisomal metabolism responded to cold regulation through ROS by increasing the  $H_2O_2$  production in the peroxisome [47].  $H_2O_2$  could also diffuse through the peroxisomal membrane into the cytosol [48], thus increasing the risk of oxidative damage in this compartment. It suggests that ROS scavenging system in cassava is a complex network involving multiple components of ROS production, turnover and scavenging as well as their cross-interactions.

## Conclusions

In conclusion, our results show the important role of cytosolic *MeCu/ZnSOD* and cytosolic *MeAPX2* in cassava in improving ROS scavenging, thereby leading to reduced  $H_2O_2$  accumulation and improved abiotic stress resistance. The results also confirm that the transgenic approach is effective in improving the stress resistance in cassava via proper gene stacking of ROS scavenging enzymes.

## Methods

### Plasmid constructions, cassava transformation and phenotype evaluation of transgenic cassava

The cDNA amplification of cassava *Cu/ZnSOD* (GenBank accession no. AY642137) and cassava *APX2* (GenBank accession no. AY973622) was determined by PCR using primers covering the full length of transcripts. The PCR fragment was sequenced and cloned into the binary vector pCambia1301 under the control of vascular-specific promoter p54/1.0 promoter (GenBank accession no. AY217353.1) [40] and the ubiquitous CaMV 35S promoter, respectively to generate pC-P54::MeCu/ZnSOD-35S::MeAPX2 (Figure 1a). The plasmid was mobilized into *Agrobacterium tumefaciens* strain LBA4404 for cassava transformation using friable embryogenic callus of

cultivar TMS60444. Embryogenic callus induction of cassava TMS60444, *Agrobacterium*-mediated genetic transformation and plantlet regeneration were performed by the methods described by Zhang et al. [49].

The uniform stem cuttings of field-grown cassava plants were used for phenotype and yield evaluation. Ten stem cuttings per transgenic line and WT were planted in late spring of 2012 in Wushe Plantation for Transgenic Crops, Shanghai, China, and harvested in early November, 2012. The performance of field plants was recorded regularly till harvest.

#### Subcellular localization

The *MeAPX2* genes tagged with green fluorescent protein (GFP) were cloned into pCAMBIA1301, and the resulting constructs were used to transform *Agrobacterium tumefaciens* LBA4404. To identify the subcellular localization of the gene, *N. benthamiana* leaves were agroinfiltrated with the *Agrobacterium* strains harboring pC-35::MeAPX2-eGFP binary vectors, respectively, by the protocol described previously [50]. At 36 hours post-infiltration, the transiently transformed leaves were observed under a confocal microscope (FluoView FV1000, Olympus, Japan).

#### Southern blot and Real-time RT-PCR analyses of transgenic cassava plants

Southern blot analysis entailed digestion of the genomic DNA (20  $\mu$ g) of cassava leaves of *in vitro* plants with *Xba*I, followed by 0.8% (w/v) agarose gel electrophoresis and subsequent transfer to a positively charged nylon membrane (Roche, Mannheim, Germany). The PCR fragment of *MeAPX2* (1 Kb) was labeled with digoxigenin (DIG) using the PCR DIG Probe Synthesis Kit (Roche, Mannheim, Germany). Hybridization and detection were performed by using the DIG-High Prime DNA Labeling and Detection Starter Kit II (Roche, Mannheim, Germany), according to the manufacturer's instructions.

Gene expression was analyzed by real-time RT-PCR (qRT-PCR) of transgenic plants. Briefly, total RNA was extracted from fresh cassava leaves of *in vitro* plants using the RNA Plant plus Reagent (Tiangen, Beijing, China) essentially as described previously [51]. The RNA samples were digested with DNase I and the first strand of cDNA was synthesized from 5  $\mu$ g total RNA from each sample using M-MLV reverse transcriptase (Toyobo, Osaka, Japan). qRT-PCR was carried out using the Bio-Rad CFX96 thermocycler SYBR Green I Master Mix (Toyobo, Osaka, Japan) according to the manufacturer's protocol, under the following PCR conditions: 95°C for 1 minute, followed by 40 cycles at 95°C for 15 seconds, 60°C for 15 seconds and 72°C for 20 seconds. The qRT-PCR primers were designed using Primer 3 Plus Software (<http://www.primer3plus.com>). The primers were: *MeCu*/

*ZnSOD* (forward 5'- ATGTTTCATGCCCTTGGAGAC -3' and reverse 5'- GATCACCAGCATGACGAATG -3'), *MeAPX2* (forward 5'- CATTGATAAGGCCAGGAGGA -3' and reverse 5'- TTGTTAGCAGCATGACCCTG - 3'), and  $\beta$ -*actin* (forward 5'- TGATGAGTCTGGTCCAT CCA -3' and reverse 5'- CCTCCTACGACCCAATC TCA -3'). Fold changes of RNA transcripts were calculated by the  $2^{-\Delta\Delta C_t}$  method [52] with  $\beta$ -*actin* as an internal control [6]. A detailed method for qRT-PCR analysis is available in Additional file 3.

#### Isolation and viability assay of cassava mesophyll protoplasts

*In vitro*-cultured cassava seedlings were grown at 25°C with light at 200  $\mu$ mol  $\cdot$  m<sup>-2</sup>  $\cdot$  s<sup>-2</sup> and a photoperiod 16 hour/8 hour (light/dark). Protoplasts were isolated from the mature leaves of 30-day-old sterile plants, as described by Anthony et al. [53]. Leaves were sliced into 2 mm to 3 mm thin strips and incubated in digestion mixture (pH 5.8) containing 9% (w/v) mannitol, 1% Cellulose R-10 (Yakult Honsha Co., Tokyo, Japan), 1.5% macerozyme R-10 (Yakult Honsha Co., Ltd., Tokyo, Japan) and 25 mM morpholine ethanesulfonic acid (MES) in cell protoplast washing (CPW) medium (27.2 mg of KH<sub>2</sub>PO<sub>4</sub>, 101 mg of KNO<sub>3</sub>, 1480 mg of CaCl<sub>2</sub>  $\cdot$  2H<sub>2</sub>O, 246 mg of MgSO<sub>4</sub>  $\cdot$  7H<sub>2</sub>O, 0.16 mg of KI, 0.025 mg of CuSO<sub>4</sub>  $\cdot$  5H<sub>2</sub>O per liter). After purification, the protoplasts were resuspended in CPW solution supplemented with 9% mannitol (CPW9M, pH 7.0) to a final concentration of 2  $\times$  10<sup>5</sup> cells/mL.

Protoplast viability was determined by fluorescein diacetate (FDA) staining [54]. The purified mesophyll protoplasts were treated for 5 minutes with CPW 9 M solution (pH 7.0) supplemented with 1 M H<sub>2</sub>O<sub>2</sub> for 5 min, and then stained with FDA at a final concentration of 0.01%. Stained protoplasts were observed for fluorescence under a fluorescent microscope (Nikon TE2000-S, Japan). The viable percentage = (the number of protoplasts with green fluorescence)/(the number of total mesophyll protoplasts)  $\times$  100%.

#### Analysis of mitochondrial integrity

Mitochondrial integrity was measured by rhodamine 123 (Rh 123, Molecular Probes-Invitrogen CA, USA) fluorescence [55] after cassava mesophyll protoplasts were treated with 1 M H<sub>2</sub>O<sub>2</sub> for 5 min. Fluorescence was detected using a confocal laser scanning microscope (FluoView FV1000, Olympus, Japan) with Ex/Em of 488 nm/515 nm. In each sample, 20–25 cells were scanned and viewed.

#### Treatment with MV, H<sub>2</sub>O<sub>2</sub> and cold

Fully expanded, healthy leaves were excised from the one-month-old plants in greenhouse and put in 10 cm-diameter petri dishes with different chemical solutions.



For the MV treatment, leaves were allowed to float on 50 mL of 100  $\mu$ M MV, and for the H<sub>2</sub>O<sub>2</sub> treatment, leaves were exposed to 50 mL of 0.5 M H<sub>2</sub>O<sub>2</sub>. Leaves were incubated at 25°C under light conditions.

Four-week-old seedlings of transgenic and the control plants were transplanted into pots (30 cm in diameter, 45 cm in height) and grown in a greenhouse (16 h days, 30°C days and 22°C nights). Two-month-old plants were subjected to stress conditions. For cold treatment, plants with a uniform growth status were transferred to a chamber and incubated at 4°C for 48 hours under weak light (cool-white fluorescent light at approximately 35  $\mu$ mol  $\cdot$  m<sup>-2</sup>  $\cdot$  s<sup>-1</sup>).

#### **Determination of chlorophyll content and lipid peroxidation**

Chlorophyll was isolated from WT and transgenic leaf segments according to the procedure described by Arnon et al. [56]. One-gram leaf disks from the one-month-old plants were homogenized in 10 mL of absolute ethyl alcohol and the homogenate was centrifuged at 3500  $\times$  *g* for 5 minutes. The supernatant was retained and the absorbance was recorded at 663 nm and 646 nm using a Nano-Drop spectrophotometer (Thermo Scientific, Scientific, Wilmington, DE, USA). Lipid peroxidation in leaf tissues was measured in terms of MDA content in the samples, according to method described by Heath and Packer [57]. One-gram leaves were homogenized in 10 mL of 10% (w/v) trichloroacetate (TCA) and centrifuged at 10,000  $\times$  *g* for 10 minutes. Thereafter, 2 mL of 10% trichloroacetic acid containing 0.67% (w/v) thiobarbituric acid was added to 2 mL of the supernatant. The mixture was boiled for 15 min, quickly cooled on ice, and centrifuged at 10,000  $\times$  *g* for 5 minutes. The absorbance of the supernatant was recorded at 532 nm and corrected for non-specific turbidity by subtracting the absorbance at 600 nm using a Nano-Drop spectrophotometer (Thermo Scientific).

#### **Determination of H<sub>2</sub>O<sub>2</sub> content and 3,3'-diaminobenzidine staining**

After determining the lipid peroxidation as above, the quantitative measurement of H<sub>2</sub>O<sub>2</sub> in the leaves was carried out according to the method reported by Velikova et al. [58]. The 3,3'-diaminobenzidine (DAB) staining method was used to qualitatively detect the H<sub>2</sub>O<sub>2</sub> generation in leaves after treatments, using the method described by Thordal-Christensen et al. [59].

#### **Enzyme assays**

SOD isoenzyme was analyzed by separating the protein extracts in 10% native polyacrylamide gel with a 4% stacking gel in standard tris-glycine buffer (pH 8.3). Samples were electrophoresed at 100 V through the stacking gel for 20 min and 120 V through the

separating gel for 60 min. After electrophoresis, the gel was immersed in 0.1% (w/v) nitroblue tetrazolium (NBT) solution for 15 min, briefly washed in ddH<sub>2</sub>O, then immersed in 100 mM potassium phosphate buffer (pH 7.0) containing 0.028 mM riboflavin and 28 mM TEMED (N,N,N',N'-tetramethyl-ethylenediamine) for another 15 min. Gels were briefly washed in ddH<sub>2</sub>O and illuminated on a light box, with a light intensity of 30 mE  $\cdot$  m<sup>-2</sup>  $\cdot$  s<sup>-1</sup> for 15 min to initiate the photochemical reaction [60].

SOD activity was detected according to the method of Beauchamp and Fridovich [61]. A 3-mL reaction mixture contained 50 mM potassium phosphate buffer (pH 7.8), 13 mM methionine, 75 mM nitroblue tetrazolium (NBT), 2 mM riboflavin, 0.1 mM EDTA, and 100 mL enzyme extract. The reaction was initiated by placing the tubes under light intensity of 5000 lx. The absorbance was measured at 560 nm in a spectrophotometer and one unit of SOD is defined as the amount required to inhibit the photo reduction of NBT by 50%.

To analyze the APX isozymes, the protein extracts were separated in 10% native polyacrylamide gels with a 5% stacking gel in standard tris-glycine buffer with 2 mM ascorbate (pH 8.3). Samples were electrophoresed under conditions similar to the SOD isozyme electrophoresis. After electrophoresis, the gels were immersed in 50 mM potassium phosphate buffer, pH 7.0, containing 2 mM ascorbate for 10 min; then in 50 mM potassium phosphate buffer, pH 7.0, containing 4 mM ascorbate and 1 mM H<sub>2</sub>O<sub>2</sub> for 20 min. After rinsing in water, the gels were stained in 50 mM potassium phosphate buffer, pH 7.8, containing 14 mM TEMED and 2.45 mM NBT for 10–30 min [62].

APX activity was determined as described by Nakano and Asada [63]. The reaction mixture contained 50 mmol/L potassium phosphate, pH 7.0, 1 mmol/L ascorbic acid (AsA), 2.5 mmol/L H<sub>2</sub>O<sub>2</sub> and enzyme source (ca 15  $\mu$ g protein) in a final volume of 2 mL at 25°C. Ascorbate oxidation was measured spectrophotometrically by a decrease of A<sub>290</sub>, using the absorption coefficient of 2.8 mM<sup>-1</sup>  $\cdot$  cm<sup>-1</sup>.

The activities of monodehydroascorbate reductase (MDHAR, EC 1.6.5.4), dehydroascorbate reductase (DHAR, EC 1.8.5.1), and GR were assayed as described previously [64–66]. Protein was estimated according to Bradford [67].

#### **Statistical analyses**

All data were represented as mean  $\pm$  SD from at least three independent experiments with three replicates. Statistical analysis was done using SPSS 15.0 for Windows (SPSS, Chicago, IL), with two-tailed Student's *t*-tests for measuring significance. Double asterisks indicate significant differences between transgenic lines and WT at 1% level, one asterisk stands for at 5% level.

## Additional files

**Additional file 1: Subcellular localization of MeAPX2::GFP fusion protein and GFP control in *N. benthamiana* epidermal cells.** Scale bar = 50  $\mu$ m.

**Additional file 2: Yield of fresh storage roots in field-grown (5 months) wild type (WT) and SA transgenic plant lines.** No significant difference was found by t-test ( $p < 0.05$ ).

**Additional file 3: Supplemental methods for qRT-PCR analysis.**

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

JX carried out the transgenic cassava production, molecular and physiological analysis, and wrote the manuscript. JY, XD and YJ partially participated in the experiment and provided helpful suggestions. PZ was responsible for the overall concept, experimental design, data analysis, and revising this manuscript. All authors read and approved the manuscript.

## Acknowledgements

This work was supported by grants from the National Basic Research Program (2010CB126605), the National Natural Science Foundation of China (31271775), the National High Technology Research and Development Program of China (2012AA101204), the SIPPE-SCBG Joint Fund, the Earmarked Fund for China Agriculture Research System (CARS-12) and Shanghai Municipal Afforestation & City Appearance and Environmental Sanitation Administration (G102410, F132427).

## Author details

<sup>1</sup>Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, Shanghai Chenshan Botanical Garden, 3888 Chenhua Road, Shanghai 201602, China. <sup>2</sup>National Key Laboratory of Plant Molecular Genetics and National Center for Plant Gene Research (Shanghai), Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200032, China. <sup>3</sup>Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China.

Received: 30 May 2014 Accepted: 22 July 2014

Published: 5 August 2014

## References

- Hillocks RJ, Thresh JM, Bellotti AC: *Cassava Biology, Production and Utilization*. New York: CAB; 2002.
- Montagnac JA, Davis CR, Tanumihardjo SA: Nutritional value of cassava for use as a staple food and recent advances for improvement. *Compr Rev Food Sci Food Safety* 2009, **8**(3):181–194.
- El-Sharkawy MA: International research on cassava photosynthesis, productivity, eco-physiology, and responses to environmental stresses in the tropics. *Photosynthetica* 2006, **44**(4):481–512.
- Ceballos H, Ramirez J, Bellotti AC, Jarvis A, Alvarez E: Adaptation of Cassava to Changing Climates. In *Crop Adaptation to Climate Change*. Edited by Yadav SS, Redden RJ, Hatfield JL, Lotze-Campen H, Hall AE. Oxford, UK: Wiley-Blackwell; 2011.
- An D, Yang J, Zhang P: Transcriptome profiling of low temperature-treated cassava apical shoots showed dynamic responses of tropical plant to cold stress. *BMC Genomics* 2012, **13**:64.
- Xu J, Duan X, Yang J, Beeching JR, Zhang P: Enhanced reactive oxygen species scavenging by overproduction of superoxide dismutase and catalase delays postharvest physiological deterioration of cassava storage roots. *Plant Physiol* 2013, **161**(3):1517–1528.
- Xu J, Duan X, Yang J, Beeching JR, Zhang P: Coupled expression of Cu/Zn-superoxide dismutase and catalase in cassava improves tolerance against cold and drought stresses. *Plant Signal Behav* 2013, **8**e24525.
- Raychaudhuri SS, Deng XW: The role of superoxide dismutase in combating oxidative stress in higher plants. *Bot Rev* 2000, **66**:89–98.
- Alscher RG, Erturk N, Heath LS: Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 2002, **53**(372):1331–1341.
- Asada K: Ascorbate peroxidase – a hydrogen peroxide-scavenging enzyme in plants. *Physiol Plant* 1992, **85**(2):235–241.
- Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, Yoshimura K: Regulation and function of ascorbate peroxidase isoenzymes. *J Exp Bot* 2002, **53**(372):1305–1319.
- Diaz-Vivancos P, Faize M, Barba-Espin G, Faize L, Petri C, Hernández JA, Burgos L: Ectopic expression of cytosolic superoxide dismutase and ascorbate peroxidase leads to salt stress tolerance in transgenic plums. *Plant Biotechnol J* 2013, **11**(8):976–985.
- Gill SS, Tuteja N: Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Biochem* 2010, **48**(12):909–930.
- Wang Y, Wisniewski M, Meilan R, Uratsu SL, Cui M, Dandekar A, Fuchigami L: Ectopic expression of Mn-SOD in *Lycopersicon esculentum* leads to enhanced tolerance to salt and oxidative stress. *J Appl Hortic* 2007, **9**:3–8.
- Wang YC, Qu GZ, Li HY, Wu YJ, Wang C, Liu GF, Yang CP: Enhanced salt tolerance of transgenic poplar plants expressing a manganese superoxide dismutase from *Tamarix androssowii*. *Mol Biol Rep* 2010, **37**(2):1119–1124.
- Prashanth SR, Sadhasivam V, Parida A: Over expression of cytosolic copper/zinc superoxide dismutase from a mangrove plant *Avicennia marina* in indica rice var Pusa Basmati-1 confers abiotic stress tolerance. *Transgenic Res* 2008, **17**(2):281–291.
- Rubio MC, Gonzalez EM, Minchin FR, Webb KJ, Arrese-Igor C, Ramos J, Becana M: Effects of water stress on antioxidant enzymes of leaves and nodules of transgenic alfalfa overexpressing superoxide dismutases. *Physiol Plant* 2002, **115**(4):531–540.
- Yabuta Y, Motoki T, Yoshimura K, Takada T, Ishikawa T, Shigeoka S: Thylakoid membrane-bound ascorbate peroxidase is a limiting factor of antioxidative systems under photo-oxidative stress. *Plant J* 2002, **32**:912–925.
- Wang Y, Wisniewski M, Meilan R, Cui M, Webb R, Fuchigami L: Overexpression of cytosolic ascorbate peroxidase in tomato confers tolerance to chilling and salt stress. *J Amer Soc Hortic Sci* 2005, **130**(2):167–173.
- Li YJ, Hai RL, Du XH, Jiang XN, Lu H: Over-expression of a populus peroxisomal ascorbate peroxidase (PpAPX) gene in tobacco plants enhances stress tolerance. *Plant Breed* 2009, **128**(4):404–410.
- Sun WH, Duan M, Shu DF, Yang S, Meng QW: Over-expression of StAPX in tobacco improves seed germination and increases early seedling tolerance to salinity and osmotic stresses. *Plant Cell Rep* 2010, **29**(8):917–926.
- Duan M, Feng HL, Wang LY, Li D, Meng QW: Overexpression of thylakoidal ascorbate peroxidase shows enhanced resistance to chilling stress in tomato. *J Plant Physiol* 2012, **169**(9):867–877.
- Pitcher LH, Brennan E, Hurlley A, Dunsmuir P, Tepperman JM, Zilinskas BA: Overproduction of petunia copper/zinc superoxide dismutase does not confer ozone tolerance in transgenic tobacco. *Plant Physiol* 1991, **97**(1):452–455.
- Payton P, Allen RD, Trolinder N, Holaday AS: Overexpression of chloroplast-targeted Mn superoxide dismutase in cotton (*Gossypium hirsutum* L., cv. Coker 312) does not alter the reduction of photosynthesis after short exposures to low temperature and high light intensity. *Photosynth Res* 1997, **52**(3):233–244.
- Hapelin C: Gene stacking in transgenic plant—the challenge for 21st century plant biotechnology. *Plant Biotechnol J* 2005, **3**(2):141–155.
- Kwon SY, Joeng YJ, Lee HS, Kim JS, Cho KY, Allen RD, Kwak SS: Enhanced tolerance of transgenic tobacco plants expressing both superoxide dismutase and ascorbate peroxidase in chloroplasts against methyl viologen mediated oxidative stress. *Plant Cell Environ* 2002, **25**:873–882.
- Tang L, Kwon SY, Kim SH, Kim JS, Choi JS, Sung CK, Kwak SS, Lee HS: Enhanced tolerance of transgenic potato plants expressing both superoxide dismutase and ascorbate peroxidase in chloroplasts against oxidative stress and high temperature. *Plant Cell Rep* 2006, **25**:1380–1386.
- Lee SH, Ahsan N, Lee KW, Kim DH, Kwak SS, Kwon SY, Kim TH, Lee BH: Simultaneous overexpression of both Cu/Zn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses. *J Plant Physiol* 2007, **164**:1626–1638.
- Faize M, Burqos L, Piqueras A, Nicolas E, Barba-Espin G, Clement-Moreno MJ, Alcobendas R, Artlip T, Hernandez JA: Involvement of cytosolic ascorbate peroxidase and Cu/Zn-superoxide dismutase for improved tolerance against drought stress. *J Exp Bot* 2011, **62**:2599–2613.
- Payton P, Webb R, Korniyev D, Allen R, Holaday AS: Protecting cotton photosynthesis during moderate chilling at high light intensity by

- increasing chloroplastic antioxidant enzyme activity. *J Exp Bot* 2001, **52**(365):2345–2354.
31. Zhao F, Zhang H: Salt and paraquat stress tolerance results from co-expression of the *Suaeda salsa* glutathione S-transferase and catalase in transgenic rice. *Plant Cell Tiss Org Cult* 2006, **86**(3):349–358.
  32. Okogbenin E, Setter TL, Ferguson M, Mutegi R, Ceballos H, Olasanmi B, Fregene M: Phenotypic approaches to drought in cassava: review. *Front Physiol* 2013, **4**:93.
  33. Lopez C, Jorge V, Piegu B, Mba C, Cortes D, Restrepo S, Soto M, Laudie M, Berger C, Cooke R, Delseny M, Tohme J, Verdier V: A unigene catalogue of 5700 expressed genes in cassava. *Plant Mol Biol* 2004, **56**:541–554.
  34. Anderson JV, Delseny M, Fregene MA, Jorge V, Mba C, Lopez C, Restrepo S, Soto M, Piegu B, Verdier V, Cooke R, Tohme J, Horvath DP: An EST resource for cassava and other species of Euphorbiaceae. *Plant Mol Biol* 2004, **56**:527–539.
  35. Lokko Y, Anderson JV, Rudd S, Raji A, Horvath D, Mikel MA, Kim R, Liu L, Hernandez A, Dixon AGO, Ingelbrecht IL: Characterization of an 18,166 EST dataset for cassava (*Manihot esculenta* Crantz) enriched for drought-responsive genes. *Plant Cell Rep* 2001, **26**:1605–1618.
  36. Sakurai T, Plata G, Rodriguez-Zapata F, Seki M, Salcedo A, Toyoda A, Ishiwata A, Tohme J, Sakaki Y, Shinozaki K, Ishitani M: Sequencing analysis of 20,000 full-length cDNA clones from cassava reveals lineage specific expansions in gene families related to stress response. *BMC Plant Biol* 2007, **7**:66.
  37. Li YZ, Pan YH, Sun CB, Dong HT, Luo XL, Wang ZQ, Tang JL, Chen BS: An ordered EST catalogue and gene expression profiles of cassava (*Manihot esculenta*) at key growth stages. *Plant Mol Biol* 2010, **74**:573–590.
  38. Reilly K, Bernal D, Cortes DF, Gomez-Vasquez R, Tohme J, Beeching JR: Towards identifying the full set of genes expressed during cassava post-harvest physiological deterioration. *Plant Mol Biol* 2007, **64**:187–203.
  39. Zhang P, Wang WQ, Zhang GL, Kaminek M, Dobrev P, Xu J, Gruissem W: Senescence-inducible expression of isopentenyl transferase extends leaf life, increases drought stress resistance and alters cytokinin metabolism in cassava. *J Integr Plant Biol* 2010, **52**(7):653–669.
  40. Zhang P, Bohl-Zenger S, Puonti-Kaerlas J, Potrykus I, Gruissem W: Two cassava promoters related to vascular expression and storage root formation. *Planta* 2003, **218**:192–203.
  41. El-Sharkawy MA, Cock JH, Held AA: Photosynthetic responses of cassava cultivars (*Manihot esculenta* Crantz) from different habitats to temperature. *Photosynth Res* 1984, **5**(3):243–250.
  42. Carretero CL, Cantos M, Garcia JL, Troncoso A: In vitro-ex vitro salt (NaCl) tolerance of cassava (*Manihot esculenta* Crantz) plants. *In Vitro Cell Dev Biol Plant* 2007, **43**(4):364–369.
  43. Miller G, Shulaev V, Mittler R: Reactive oxygen signaling and abiotic stress. *Physiol Plant* 2008, **133**(3):481–489.
  44. Apel K, Hirt H: Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 2004, **55**:373–399.
  45. Davletova S, Rizhsky L, Liang H, Shengqiang Z, Oliver DJ, Couto J, Shulaev V, Schlauch K, Mittler R: Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of *Arabidopsis*. *Plant Cell* 2005, **17**(1):268–281.
  46. Selote DS, Khanna-Chopra R: Drought acclimation confers oxidative stress tolerance by inducing co-ordinated antioxidant defense at cellular and subcellular level in leaves of wheat seedlings. *Physiol Plant* 2006, **127**:494–506.
  47. Dong CH, Zolman BK, Bartel B, Lee BH, Stevenson B, Agarwal M, Zhu JK: Disruption of *Arabidopsis* CHY1 reveals an important role of metabolic status in plant cold stress signaling. *Mol Plant* 2009, **2**(1):59–72.
  48. Del Rio LA, Pastori GM, Palma JM, Sandalio LM, Sevilla F, Corpas FJ, Jimenez A, Lopez-Huertas E, Hernandez JA: The activated oxygen role of peroxisomes in senescence. *Plant Physiol* 1998, **116**(4):1195–1200.
  49. Zhang P, Potrykus I, Puonti-Kaerlas J: Efficient production of transgenic cassava using negative and positive selection. *Transgenic Res* 2000, **9**:405–415.
  50. Wydro M, Kozubek E, Lehman P: Optimization of transient *Agrobacterium*-mediated gene expression system in leaves of *Nicotiana benthamiana*. *Acta Biochim Pol* 2006, **53**(2):289–298.
  51. Xu J, Aileni M, Abbagani S, Zhang P: A reliable and efficient method for total RNA isolation from various members of spurge family (Euphorbiaceae). *Phytochem Anal* 2010, **21**(5):395–398.
  52. Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔC<sub>t</sub></sup> method. *Methods* 2001, **25**(4):402–408.
  53. Anthony P, Davey MR, Power JB, Lowe KC: An improved protocol for the culture of cassava leaf protoplasts. *Plant Cell Tiss Org Cult* 1995, **42**:229–302.
  54. Larkin PJ: Purification and viability determinations of plant protoplasts. *Planta* 1976, **128**:213–216.
  55. Wu FS: Localization of mitochondria in plant cells by vital staining with rhodamine 123. *Planta* 1987, **171**(3):346–357.
  56. Arnon DI: Copper enzymes in isolated chloroplasts polyphenoloxidase in *Beta vulgaris*. *Plant Physiol* 1949, **24**:1–15.
  57. Heath R, Packer L: Photoperoxidation in isolated chloroplasts. I Kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Biophys* 1968, **125**(1):189–198.
  58. Velikova V, Yordanov I, Edreva A: Oxidative stress and some antioxidant system in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Sci* 2000, **151**:59–66.
  59. Thordal-Christensen H, Zhang Z, Wei Y, Collinge DB: Subcellular localization of H<sub>2</sub>O<sub>2</sub> in plants: H<sub>2</sub>O<sub>2</sub> accumulation in papillae and hypersensitive response during the barley-powdery mildew interaction. *Plant J* 1997, **11**:1187–1194.
  60. Chen CN, Pan SM: Assay of superoxide dismutase activity by combining electrophoresis and densitometry. *Bot Bull Acad Sin* 1996, **37**:107–111.
  61. Beauchamp C, Fridovich I: Superoxide dismutase: improved assays and an assay applicable to acrylamide gel. *Anal Biochem* 1971, **44**:276–287.
  62. Zimmermann P, Heinlein C, Orendi G, Zentgraf U: Senescence-specific regulation of catalases in *Arabidopsis thaliana* (L.) Heynh. *Plant Cell Environ* 2006, **29**:1049–1060.
  63. Nakano Y, Asada K: Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 1981, **22**(5):867–880.
  64. Miyake C, Asada K: Thylakoid-bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product monodehydroascorbate radicals in thylakoids. *Plant Cell Physiol* 1992, **33**(5):541–553.
  65. Dalton DA, Russell SA, Hanus FJ, Pascoe GA, Evans HJ: Enzymatic reactions of ascorbate and glutathione that prevent peroxide damage in soybean root nodules. *Proc Natl Acad Sci U S A* 1986, **83**(11):3811–3815.
  66. Grace SC, Logan BA: Acclimation of foliar antioxidant systems to growth irradiance in three broad-leaved evergreen species. *Plant Physiol* 1996, **112**(4):1631–1640.
  67. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976, **72**:248–254.

doi:10.1186/s12870-014-0208-4

Cite this article as: Xu et al.: Increased expression of native cytosolic Cu/Zn superoxide dismutase and ascorbate peroxidase improves tolerance to oxidative and chilling stresses in cassava (*Manihot esculenta* Crantz). *BMC Plant Biology* 2014 **14**:208.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

