

RESEARCH

Open Access

Self-reported attitudes versus actual practice of oxygen therapy by ICU physicians and nurses

Hendrik JF Helmerhorst^{1,2*}, Marcus J Schultz^{2,3}, Peter HJ van der Voort⁴, Robert J Bosman⁴, Nicole P Juffermans^{2,3}, Evert de Jonge¹ and David J van Westerloo¹

Abstract

Background: High inspiratory oxygen concentrations are frequently administered in ventilated patients in the intensive care unit (ICU) but may induce lung injury and systemic toxicity. We compared beliefs and actual clinical practice regarding oxygen therapy in critically ill patients.

Methods: In three large teaching hospitals in the Netherlands, ICU physicians and nurses were invited to complete a questionnaire about oxygen therapy. Furthermore, arterial blood gas (ABG) analysis data and ventilator settings were retrieved to assess actual oxygen practice in the same hospitals 1 year prior to the survey.

Results: In total, 59% of the 215 respondents believed that oxygen-induced lung injury is a concern. The majority of physicians and nurses stated that minimal acceptable oxygen saturation and partial arterial oxygen pressure (PaO₂) ranges were 85% to 95% and 7 to 10 kPa (52.5 to 75 mmHg), respectively. Analysis of 107,888 ABG results with concurrent ventilator settings, derived from 5,565 patient admissions, showed a median (interquartile range (IQR)) PaO₂ of 11.7 kPa (9.9 to 14.3) [87.8 mmHg], median fractions of inspired oxygen (FiO₂) of 0.4 (0.4 to 0.5), and median positive end-expiratory pressure (PEEP) of 5 (5 to 8) cm H₂O. Of all PaO₂ values, 73% were higher than the upper limit of the commonly self-reported acceptable range, and in 58% of these cases, neither FiO₂ nor PEEP levels were lowered until the next ABG sample was taken.

Conclusions: Most ICU clinicians acknowledge the potential adverse effects of prolonged exposure to hyperoxia and report a low tolerance for high oxygen levels. However, in actual clinical practice, a large proportion of their ICU patients was exposed to higher arterial oxygen levels than self-reported target ranges.

Keywords: Oxygen; Hyperoxia; Mechanical ventilation; Lung injury; Intensive care medicine; Questionnaire

Background

Oxygen supply during mechanical ventilation is a highly effective and uniformly used intervention to support oxygenation of mechanically ventilated patients in the intensive care unit (ICU). Although oxygen administration is a lifesaving strategy in the management of patients with respiratory insufficiency, the clinical implications of hyperoxia remain an important subject of debate [1]. The controversies are triggered by a considerable number of studies showing beneficial, harmful, and/or insignificant effects of oxygen therapy on outcome in different

subgroups [2-15]. However, morbidity and mortality may be substantially impacted by the used threshold and depend on degree, duration, and susceptibility for hyperoxia.

The emerging laboratory evidence for the double-edged nature of oxygen (lifesaving but also potentially harmful) is compelling [16-22], but robust clinical studies and evidence-based guidelines in critically ill patients are still limited [23-26]. Consequently, the attitudes and beliefs in the management of oxygen administration vary considerably in clinical practice [27,28]. In general, physicians are inclined to treat hypoxemia aggressively in order to achieve satisfactory tissue oxygenation [23,29,30]. However, hyperoxemia is often considered acceptable, especially when applied fractions of inspired oxygen (FiO₂) are relatively low [31-33].

* Correspondence: H.J.F.Helmerhorst@lumc.nl

¹Department of Intensive Care Medicine, Leiden University Medical Center, Post Box 9600, Leiden 2300, RC, The Netherlands

²Laboratory of Experimental Intensive Care and Anesthesiology, Academic Medical Center, Amsterdam 1105, AZ, The Netherlands
Full list of author information is available at the end of the article

Given the lack of established guidelines on oxygen therapy in ICU patients, our study was designed to investigate the common beliefs and self-reported attitudes of ICU physicians and nurses on oxygenation targets and to compare this with actual treatment of ICU patients in the same hospitals. We hypothesized that the potential harmful effects of oxygen are well known and generally acknowledged, but that in real clinical practice, hyperoxia is not a major concern for ICU professionals.

Methods

Questionnaire

An anonymous online survey was performed between June and August 2012 to elicit the self-reported behavior of ICU personnel with respect to oxygen therapy. The questionnaire was a modified and comprehensive version of previously used questionnaires from Canada and Australia/New Zealand [27,28] and comprised multiple choice questions (see Additional file 1). The target population consisted of physicians and nurses, working in closed format, mixed medical and surgical, tertiary care ICUs of three participating hospitals, including two academic and one large teaching hospital in the Netherlands. Participants were invited by email to complete the online questionnaire. A reminder was sent once to all participants.

Patient data

Analyses were performed on data recorded between 1 April 2011 and 31 March 2012 for all patients admitted to the ICU departments of the same hospitals that

participated in the questionnaire study. Anonymous encrypted data were collected from the patient data management system (PDMS) database (MetaVision, iMDsoft, Leiden, The Netherlands). According to the Dutch Medical Research Involving Human Subjects Act, there was no need for informed patient consent or approval by ethical committees, as only registries without patient identifying information were used.

Arterial blood gas (ABG) analyses and concurrent ventilator settings were extracted to retrospectively assess actual practices regarding oxygen therapy. Data from ICU admission to dismissal or death were included for analysis. Data with a partial arterial oxygen pressure (PaO₂) of ≤ 4.5 kPa or 33.8 mmHg ($n = 209$) were excluded to prevent confounding by venous blood gas samples. Further exclusion of samples with PaO₂ of 4.5 to 5.0 kPa ($n = 146$) or PaO₂ of 5.0 to 6.0 kPa ($n = 356$) did not materially change our observations (data not shown). Every set of ABG data and ventilator settings was compared with the following set, as described previously [32]. Prone positioning, recruitment maneuvers, and other efforts that may improve oxygenation could not be explored in this database. Clinicians' responses to ABG results were explored by analyzing the FiO₂ and positive end-expiratory pressure (PEEP) adjustments in a subsample of mechanically ventilated patients when more than two ABG samples and ventilator settings were recorded. PaO₂ values were categorized according to the commonly self-reported acceptable range (7 to 10 kPa or 52.5 to 75 mmHg) extracted from the survey results (Figure 1). This range is roughly consistent with




Figure 1 Self-reported tolerance limits for short-term (15 min, open bars) and longer term (24 to 48 h, closed bars) oxygenation. Bars represent percentage of respondents ($n = 200$). The presented case is a young to middle-aged ARDS patient in the ICU requiring mechanical ventilation. Ventilator settings (e.g., PEEP, airway pressures, I/E ratio, flow ratio) are optimized with respect to the PaO₂/FiO₂ ratio and hemodynamic indices. Lung injury due to high FiO₂ and/or ventilator settings is minimized. There is no evidence to indicate end-organ ischemia, and hemodynamics are stable.

oxygenation goals (7.3 to 10.7 kPa or 55 to 80 mmHg) previously suggested by the Acute Respiratory Distress Syndrome (ARDS) network [30,34].

Mechanical ventilation protocol

Local guidelines, applicable during the survey and collection of ABG and ventilation data, instructed for lowest acceptable FiO₂ levels. FiO₂ levels higher than 60% should be avoided by increasing PEEP levels, instituting inverse-ratio ventilation or prone positioning. No explicit target ranges for PaO₂, FiO₂, or saturation were specified in the participating hospitals during this period.

Statistical analysis

All data are expressed as percentages of the total number of respondents for the particular questions, unless otherwise specified. Data are presented as means (\pm SD) or medians (interquartile range (IQR)) depending on data distributions, unless stated otherwise. For assessing differences between physicians and nurses, ICU personnel were grouped according to their respective profession and data were analyzed using Fisher's exact tests. ICU physicians, fellows, and residents were classified as physicians; ICU nurses and ICU nurses in training were classified as nurses.

Statistical analyses were conducted using STATA/SE 10.1 (StataCorp LP, College Station, TX, USA).

Results

Data derived from questionnaire

Respondent characteristics

Approximately 500 potential participants were invited to complete the online questionnaire. Full or partial responses were received from 215 ICU physicians and nurses with a mean age of 40.4 \pm 10.0 years (range 24 to 62). In total, 171 (80%) respondents fully completed all questions. The group of respondents consisted of 28 (13%) critical care physicians, 15 (7%) fellows, 15 (7%) residents, 141 (66%) ICU nurses, 11 (5%) ICU nurses in training, and 5 (2%) ICU clinicians with another type of practice.

Opinions towards oxygen toxicity

The responses are listed in Table 1. Overall, 126 (59%) respondents considered oxygen-induced lung injury in mechanically ventilated patients a major concern. However, the vast majority of respondents (81%) considered high tidal volumes and high inspiratory pressures as the greatest risk for lung injury in mechanically ventilated patients. No differences between physicians and nurses were detected.

Self-reported acceptance of hyperoxemia and hypoxemia

The percentages of respondents accepting various oxygenation ranges in a young to middle-aged mechanically ventilated patient with ARDS are shown in Figure 1.

Table 1 Questionnaire responses regarding risks assessment and management in oxygen therapy

Question	Responses (% of total)	Physicians vs. nurses
Is oxygen induced lung injury a concern when placing a patient on mechanical ventilation?		NS
YES, a major concern	126 (59%)	
due to the <i>high incidence</i> of injury	13 (6%)	
due to the <i>severity</i> of injury	63 (29%)	
due to the <i>high incidence and severity</i> of injury	50 (23%)	
YES, but not a <i>major</i> concern	80 (37%)	
NO, it is not a concern	9 (4%)	
In your opinion, which one of the following two situations poses a greater threat of lung injury for mechanically ventilated patients?		NS
High FiO ₂	35 (16%)	
High tidal volumes and high ventilator pressures	173 (81%)	
Don't know	7 (3%)	
In situations when maximal SaO ₂ achievable is low (\pm 85%) or when FiO ₂ requirements are high, do you assess indices of tissue oxygenation?		P = 0.05
NO	91 (43%)	
YES, lactate	88 (42%)	
YES, microcirculation with OPS/SDF imaging	4 (2%)	
YES, a combination of indices	20 (9%)	
YES, SvO ₂	6 (3%)	
YES, other	2 (1%)	

NS, not significant; FiO₂, fractions of inspired oxygen; OPS, orthogonal polarization spectral; SDF, sidestream dark field; SaO₂, arterial oxygen saturation; SvO₂, mixed venous oxygen saturation.

For both short and longer lasting periods, the vast majority of respondents choose 7 to 10 kPa (52.5 to 75 mmHg) as the lowest acceptable PaO₂ range. Physicians were more tolerant towards lower PaO₂ limits for short duration than nurses ($P < 0.001$).

Presented with a patient whose arterial oxygen saturation (SaO₂) levels are low (<85%) or FiO₂ requirements are high, indices of tissue oxygenation were frequently assessed (Table 1). Differences between physicians and nurses approached statistical significance ($P = 0.05$), with physicians favoring lactate assessment and nurses being less likely to demand some assessment of tissue oxygenation. Nurses in training more often favored lactate assessment than senior ICU nurses ($P = 0.01$).

Adjustment of FiO₂ in specific clinical cases

The proportions of ICU clinicians adjusting FiO₂ levels in specified clinical cases are listed in Table 2. Observed differences by profession were mainly restricted to questions regarding patients with untreatable anemia, where physicians generally favored higher FiO₂ levels than nurses. Only minor differences within the clustered categories of physicians (comparison between physicians, fellows, residents) and nurses (ICU nurses, nurses in training) were observed.

Data derived from ABG measurements and ventilator settings

Descriptive data

A total of 107,888 ABG results with concurrent ventilator settings, covering 5,565 patient admissions, were retrieved and included for analysis over a 1-year period prior to the survey in three hospitals. Median interval between two consecutive ABG samples was 214 min (IQR 130 to 331), and the median number of ABG samples per patient was 7 (IQR 4 to 19). Mean PaO₂ was 12.9 kPa (SD 5.1) or 96.8 mmHg, and median PaO₂ was 11.7 kPa (IQR 9.9 to 14.3) or 87.8 mmHg. Overall, in 25.3% of ABG results, PaO₂ was in the self-reported range (7 to 10 kPa), 1.2% was lower and 73.4% was higher than the predefined range.

Mechanical ventilation settings showed a mean PEEP of 6.1 cm H₂O (SD 4.3), and median PEEP was 5 (IQR 5 to 8). Mean FiO₂ was 0.45 (SD 0.14), and median FiO₂ was 0.40 (IQR 0.40 to 0.50). Only small differences were observed between the three participating hospitals.

Recorded FiO₂ adjustments following ABG analysis

After exclusion of spontaneously breathing and non-invasively ventilated patients, 62,875 ABG records derived from 4,264 mechanically ventilated patients were included for analysis of FiO₂ adjustments in response to ABG samples.

Analyzing every first registered ABG sample in the ICU, 62,222 PaO₂ measurements covering 3,791 patients

were retrospectively categorized in predefined ranges and were followed by a recorded FiO₂ adjustment. The subsequently registered PaO₂ measurement was compared with the first registered PaO₂ (Table 3).

Table 4 shows that quantity, direction, and magnitude of ventilator adjustments in response to high arterial oxygen levels are considerably influenced by the level of FiO₂. In 58.3% of cases with PaO₂ higher than the upper level of the commonly self-reported acceptable oxygenation (10 kPa or 75 mmHg), neither FiO₂ nor PEEP levels had been lowered when the next ABG sample was taken.

Discussion

In accordance with accumulating laboratory evidence for the toxic effects of oxygen in pulmonary injury [1,35-38], the majority of surveyed ICU physicians and nurses consider prolonged hyperoxic exposure to be associated with an increased risk for lung injury, although a lower risk than high tidal volumes and inspiratory pressures. In contrast, in actual clinical practice, the majority of PaO₂ values recorded in ICU patients are higher than recommended targets under comparable conditions and are generally accepted by ICU physicians and nurses without adjustment of ventilator settings.

Compared with previous studies from other countries, more respondents identified oxygen toxicity as a major threat to lung injury in ICU patients (59% compared with 26% and 51% in studies from Australia and Canada, respectively) [27,28]. Similar proportions of respondents considered high inspiratory oxygen concentrations a more important risk than high tidal volumes or inspiratory pressures, and a similar heterogeneity in self-reported attitudes regarding oxygen therapy was observed [27,28,39]. The current results show generally higher minimum allowable SaO₂ ranges than data from Canadian intensivists [28], which may indicate that clinicians' beliefs have changed over the last decade or it may merely reflect geographical differences in oxygen therapy.

It appears that clinicians' opinions regarding optimal oxygen therapy are more variable in case SaO₂ is presented compared to PaO₂ as marker of oxygenation. For PaO₂, the vast majority of clinicians choose 7 to 10 kPa (52.5 to 75 mmHg), whereas the preferred targets for saturation varied between 85% and 95%. Assuming that oxygenation targets should be in line with the best evidence in available guidelines, these preferred ranges may be triggered by recommendations and protocols providing comparable PaO₂ targets [30,40]. However, caution is urged when interpreting pulse oximetry to differentiate between hyperoxemia and normoxia. Saturation levels above 95% require special attention, since the corresponding PaO₂ levels usually cover a wide range and may substantially exceed target levels [24,41].

Table 2 Percentages of respondents adjusting FiO₂ levels in specified clinical conditions with presented levels of arterial oxygenation

	ARDS			Cardiac ischemia			Cerebral ischemia			Sepsis			Untreatable anemia		
	FiO ₂ response			FiO ₂ response			FiO ₂ response			FiO ₂ response			FiO ₂ response		
	Higher	Unchanged	Lower	Higher	Unchanged	Lower	Higher	Unchanged	Lower	Higher	Unchanged	Lower	Higher	Unchanged	Lower
SaO ₂															
80% to 85%	97.4	2.6	0.0	100	0.0	0.0	98.9	1.1	0.0	99.5	0.5	0.0	92.4	7.1	0.5
85% to 90%	61.5	38.5	0.0	96.4	3.6	0.0	91.4	8.0	0.5	92.5	7.0	0.5	85.3	13.6	1.1
90% to 95%	4.1	78.5	17.4	42.6	54.9	2.6	23.0	71.7	5.3	19.3	75.9	4.8	57.6	36.4	6.0
95% to 100%	0.0	17.4	82.6	2.6	61.0	36.4	0.5	41.2	58.3	0.5	40.1	59.4	13.6	58.7	27.7
PaO ₂															
6 kPa	96.6	3.4	0.0	98.3	1.7	0.0	98.3	1.7	0.0	98.3	1.7	0.0	96.5	3.5	0.0
9 kPa	9.0	85.3	5.6	60.1	38.2	1.7	40.6	58.9	0.6	37.7	60.6	1.7	66.7	31.6	1.8
12 kPa	0.6	27.1	72.3	5.6	60.7	33.7	2.3	53.7	44.0	2.9	47.4	49.7	22.2	52.6	25.1
16 kPa	0.0	2.3	97.7	2.2	10.1	87.6	1.1	4.6	94.3	0.6	7.4	92.0	6.4	25.7	67.8

All clinical situations represent patients in the ICU, who have been invasively mechanically ventilated for at least 5 days, with FiO₂ set at 50%. ARDS = patient with acute respiratory distress syndrome and pneumonia; cardiac ischemia = patient with signs of cardiac ischemia (ST-depressions in the anterior leads [max 3 mm]) and pneumonia; cerebral ischemia = patient with recent cerebral ischemia and one-sided hemiplegia; sepsis = patient with a liver abscess and sepsis; untreatable anemia = Jehovah's Witness with stable hemoglobin of 1.8 mmol/L after gastric bleeding; higher = increase FiO₂, higher than current 50%; unchanged = maintain FiO₂ at current 50%; lower = decrease FiO₂, lower than current 50%.

Table 3 FiO₂ adjustment following ABG analysis and its effects on oxygenation measured in the next ABG

PaO ₂ (%) [n = 107,888]	Adjustment of FiO ₂ (%) [n = 62,222]	Successive PaO ₂ [n = 61,073]		
		Higher (Delta PaO ₂)	Unchanged	Lower (Delta PaO ₂)
<7 kPa (1.2)	Higher (34.7)	96.6% (+5.3)	0.4%	3.0% (-0.5)
	Unchanged (46.9)	87.4% (+5.6)	3.2%	9.4% (-0.6)
	Lower (18.4)	95.1% (+7.9)	2.4%	2.5% (-0.9)
7 to 10 kPa (25.3)	Higher (27.9)	76.3% (+3.1)	2.7%	21.0% (-0.9)
	Unchanged (56.0)	66.3% (+2.0)	4.3%	29.4% (-0.8)
	Lower (16.1)	61.3% (+2.6)	3.6%	35.1% (-1.0)
>10 kPa (73.4)	Higher (10.8)	48.6% (+4.6)	2.1%	49.3% (-3.1)
	Unchanged (62.0)	44.7% (+2.1)	3.1%	52.2% (-2.3)
	Lower (27.2)	23.5% (+2.4)	1.7%	74.8% (-4.6)
Total (100)	-	46.3% (+2.6)	2.7%	51.0% (-2.9)

Data presented as percentages of total and irrespective of adjustment of other ventilator settings (e.g., PEEP, I/E ratio). PaO₂ = any recorded PaO₂ stratified by self-reported ranges; successive PaO₂ = PaO₂ from successively registered ABG sample; Delta = mean difference between two successive ABG samples; Higher = increased FiO₂ (column 2) or PaO₂ (column 3), higher than previous level; unchanged = FiO₂ or PaO₂ equal to previous level; lower = decreased FiO₂ or PaO₂, lower than previous level. A total of 62,222 PaO₂ measurements from 3,791 patients (57.7% of all 107,888 ABG samples) in the database is followed by an adjustment of ventilator settings, and 98.2% of PO₂ measurements is followed by a successive PO₂ measurement (n = 61,073) in the same patient when adjustment of FiO₂ is also measured.

According to the results from our questionnaire, the vast majority of respondents stated they would lower FiO₂ levels if PaO₂ was higher than 12 kPa (90 mmHg) or SaO₂ was higher than 95% in ARDS patients. The proportion of respondents that would lower FiO₂ is much lower if patients were presented with sepsis, cardiac and cerebral ischemia, or untreatable anemia. Unfortunately, we do not know whether respondents believe that oxygen is specifically harmful in patients with pre-existing acute lung injury or that higher oxygen levels are considered desirable in patients with ischemia or anemia. The latter hypothesis appears plausible, even though hyperoxemia has been reported to induce important vasoconstriction, which may lead to a paradoxical decrease in oxygen delivery [4,42].

The self-reported low tolerance for higher PaO₂ or SaO₂ than target levels appears to be in contrast with actual treatment of patients in the same three ICUs where the survey was conducted. Neither FiO₂ nor PEEP was changed in the majority of cases when PaO₂ was higher than 15 kPa (112.5 mmHg) and FiO₂ was 40% or lower. In cases when FiO₂ was 40% to 100%, ventilator settings were adjusted more often, but even in these circumstances, hyperoxemia was accepted in approximately 20%. Considering the absence of definitive guidelines and robust controlled clinical evidence, this behavior in itself may still be justifiable [43]. However, the contrast between self-reported attitudes towards oxygen therapy on the one hand and actual treatment by the same healthcare workers on the other hand is striking.

The findings about oxygenation in ICU patients are consistent with previous findings from single center studies,

showing that hyperoxemia was frequently present in mechanically ventilated patients and seldom led to adjustment of ventilator settings [31,32]. Clinicians may have specific reasons not to adjust ventilator settings when PaO₂ levels are higher than the target range. Indeed, we identified a considerable number of cases in which a presumed inadequate adjustment ultimately proved reasonable in the subsequent ABG sample (e.g., high PaO₂ followed by an increase in FiO₂, but resulting in a lower PaO₂). These cases may reflect scenarios in which clinicians anticipate deterioration in oxygenation or otherwise consider PaO₂ values as erroneous (e.g., arguably high PaO₂ - see Table 4, row PaO₂ > 30) or not representative for the current situation of a patient. Alternatively, it may be argued that hypoxemia harbors greater inherent hazards than hyperoxemia [3].

The differences between self-reported attitudes and actual treatment of patients should be interpreted with caution. First, the cases presented to the respondents included only limited details and do not reflect the complexity of clinical situations in daily practice. Further, we presented SaO₂ and PaO₂ categorized in ranges that were arbitrarily chosen. This may have influenced interpretation of the hypothetical cases. Second, ICU clinicians may have given more favorable responses in the online questionnaire due to social desirability and attention bias, although this is less likely as anonymous evaluation was secured. Third, the respondents were asked for the minimum allowable range in a specific ARDS case vignette, which may not reflect their beliefs regarding oxygen therapy in general. In the analysis of actually achieved

Table 4 Adjustment of mechanical ventilation settings following ABG analysis

	FiO ₂ 25% to 40% (n = 37,172)			FiO ₂ 40% to 60% (n = 23,466)			FiO ₂ 60% to 80% (n = 4,318)			FiO ₂ 80% to 100% (n = 2,041)			Total (n = 68,222)		
PaO ₂	13.4 (4.1)			12.6 (5.2)			12.3 (6.6)			16.0 (13.7)			13.1 (5.2)		
SpO ₂	98.0 (2.9)			96.5 (3.8)			94.3 (6.3)			89.7 (12.5)			96.9 (4.6)		
PEEP	5.6 (3.4)			7.3 (4.4)			9.1 (5.6)			9.0 (6.3)			6.5 (4.2)		
PaO ₂ range (kPa)	Delta FiO ₂ (%)	Delta PEEP (cm H ₂ O)	No decrease in FiO ₂ or PEEP (%)	Delta FiO ₂ (%)	Delta PEEP (cm H ₂ O)	No decrease in FiO ₂ or PEEP (%)	Delta FiO ₂ (%)	Delta PEEP (cm H ₂ O)	No decrease in FiO ₂ or PEEP (%)	Delta FiO ₂ (%)	Delta PEEP (cm H ₂ O)	No decrease in FiO ₂ or PEEP (%)	Delta FiO ₂ (%)	Delta PEEP (cm H ₂ O)	No decrease in FiO ₂ or PEEP (%)
<7	+6.1 (13.8)	+0.3 (3.2)	87.6	+5.4 (13.2)	+0.7 (3.7)	81.3	+2.0 (14.5)	+0.6 (6.0)	86.3	-7.0 (16.8)	+1.0 (6.7)	76.9	+3.0 (14.8)	+0.7 (4.3)	83.3
7 to 10	+2.6 (6.7)	+0.1 (3.1)	81.9	+1.6 (7.3)	+0.2 (4.0)	80.3	-0.2 (9.4)	+0.3 (5.2)	80.8	-6.7 (14.8)	+0.1 (5.9)	76.5	+1.3 (8.3)	+0.2 (3.8)	80.8
10 to 15	+0.6 (4.9)	-0.2 (2.7)	72.9	-1.5 (6.6)	-0.2 (3.8)	48.2	-5.2 (9.2)	+0.1 (4.9)	33.7	-15.6 (19.2)	+0.4 (5.7)	31.1	-0.6 (6.7)	-0.1 (3.2)	62.3
15 to 20	-0.3 (5.1)	-0.3 (2.7)	65.0	-4.8 (7.2)	-0.3 (3.3)	25.1	-11.2 (11.4)	+0.1 (4.7)	13.5	-20.4 (20.5)	+0.0 (4.7)	23.6	-2.0 (7.3)	-0.2 (3.0)	52.8
20 to 25	-1.2 (6.4)	-0.2 (2.4)	54.2	-6.1 (9.0)	-0.4 (3.0)	22.1	-12.7 (12.4)	-0.4 (5.3)	11.8	-25.9 (21.7)	-0.1 (4.8)	21.3	-4.2 (10.1)	-0.3 (2.9)	40.4
25 to 30	-1.6 (7.0)	-0.4 (2.8)	51.6	-7.2 (9.6)	-0.3 (2.7)	18.8	-19.0 (12.1)	-0.4 (3.8)	9.2	-25.3 (21.6)	-0.1 (4.8)	22.9	-7.1 (12.2)	-0.3 (2.9)	32.0
>30	-1.0 (6.6)	-0.4 (4.0)	69.8	-3.5 (9.6)	-0.4 (4.5)	43.0	-15.4 (15.8)	+0.3 (5.7)	24.8	-33.6 (23.2)	-0.4 (4.0)	14.3	-5.7 (15.9)	-0.3 (4.2)	51.6

Data are means (±SD), unless stated otherwise. Delta = difference between two successive ABG samples.

oxygenation, we studied *all* patients independent of admission diagnosis. Also, response rates for the survey were relatively modest. However, the profession distribution in the group of respondents closely reflects a typical staff constitution in a general ICU in the Netherlands, which reduces the chance of sampling bias. In the Dutch clinical setting where respiratory therapists are not available, it is often the bedside nurse that responds first to changes in oxygenation. Therefore, the opinions of ICU nurses about oxygen therapy are important in the actual care of critically ill patients [39]. Finally, some ABG samples, taken shortly after ICU arrival, may reflect oxygen therapy initiated on the operating room and influenced by anesthesiological ventilation strategies. However, successive ventilator adjustments were all recorded on the ICU and were supervised by critical care physicians. Therefore, high PaO₂ values in the direct postoperative period are not a plausible explanation for the low proportion of hyperoxic ABG samples not followed by adaptation of the ventilator settings.

Strengths of this study include the large sample of questionnaire responses and the extensive set of ABG data, derived from the same ICUs where the questionnaire was conducted. This facilitated a comprehensive comparison between self-reported attitudes and actual practices of oxygen therapy for both physicians and nurses. Further, the design of the present questionnaire closely resembles previous surveys from Canada and Australia, thereby exploring geographical patterns and trends in time concerning oxygen therapy. Our study extends these data as we have assessed objective data in our analysis including the successively measured PaO₂ after FiO₂ adjustment. This allows further estimation of the effects of recorded FiO₂ adjustments in comparison with previous data [32].

Conclusions

This study shows that most clinicians acknowledge the potential adverse effects of prolonged exposure to hyperoxia, in accordance with emerging evidence for pulmonary toxicity and increased risk of poor outcome in both humans and animals caused by excessive oxygenation [2,4,6,8,16,18,20,35,44]. However, objective data also suggest that clinicians did not consistently accommodate this conception in actual clinical practice and a large proportion of patients was exposed to arterial oxygen levels higher than self-reported as acceptable by nurses and physicians. Additional education, feedback, and implementation strategies, aimed at careful titration of oxygen, may therefore be an effective approach for strict adherence to oxygenation targets [45]. Studies on the effects of different target ranges for PaO₂ on clinically relevant endpoints are needed to guide ICU professionals on how much oxygen should be administered to their patients.

Additional file

Additional file 1: Questionnaire (translated from Dutch) as sent to all participants. The questionnaire is a modified and comprehensive version of previously used questionnaires [27,28].

Abbreviations

ABG: arterial blood gas; FiO₂: fractions of inspired oxygen; ICU: intensive care unit; PaO₂: partial arterial oxygen pressure; PDMS: patient data management system; PEEP: positive end-expiratory pressure; SaO₂: arterial oxygen saturation.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HJFH drafted the manuscript and participated in the conception and design, and collection, analysis, and interpretation of the data. MJS and PHvdV participated in the conception and design, interpretation of data, and critical revision of the article for important intellectual content. RJB was involved in the collection, assembly and interpretation of data, and critical revision. NPJ participated in the interpretation of data and critical revision. EdJ and DJvW were involved in the conception and design, interpretation and collection of data, and critical revision. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by an unrestricted grant from the Netherlands Organisation for Health Research and Development (ZonMw).

Author details

¹Department of Intensive Care Medicine, Leiden University Medical Center, Post Box 9600, Leiden 2300, RC, The Netherlands. ²Laboratory of Experimental Intensive Care and Anesthesiology, Academic Medical Center, Amsterdam 1105, AZ, The Netherlands. ³Department of Intensive Care Medicine, Academic Medical Center, Amsterdam 1105AZ, The Netherlands. ⁴Department of Intensive Care Medicine, Onze Lieve Vrouwe Gasthuis, Amsterdam 1091, AC, The Netherlands.

Received: 1 April 2014 Accepted: 27 June 2014

Published: 25 July 2014

References

1. Altmeier WA, Sinclair SE: **Hyperoxia in the intensive care unit: why more is not always better.** *Curr Opin Crit Care* 2007, **13**:73–78.
2. de Jonge E, Peelen L, Keijzers PJ, Joore H, de Lange D, van der Voort PH, Bosman RJ, de Waal RA, Wesselink R, de Keizer NF: **Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients.** *Crit Care* 2008, **12**:R156.
3. Eastwood G, Bellomo R, Bailey M, Taori G, Pilcher D, Young P, Beasley R: **Arterial oxygen tension and mortality in mechanically ventilated patients.** *Intensive Care Med* 2012, **38**:91–98.
4. Farquhar H, Weatherall M, Wijesinghe M, Perrin K, Ranchord A, Simmonds M, Beasley R: **Systematic review of studies of the effect of hyperoxia on coronary blood flow.** *Am Heart J* 2009, **158**:371–377.
5. Hovaguimian F, Lysakowski C, Elia N, Tramer MR: **Effect of intraoperative high inspired oxygen fraction on surgical site infection, postoperative nausea and vomiting, and pulmonary function: systematic review and meta-analysis of randomized controlled trials.** *Anesthesiology* 2013, **119**:303–316.
6. Kilgannon JH, Jones AE, Shapiro NI, Angelos MG, Milcarek B, Hunter K, Parrillo JE, Trzeciak S, E EMSRN: **Association between arterial hyperoxia following resuscitation from cardiac arrest and in-hospital mortality.** *JAMA* 2010, **303**:2165–2171.
7. Bellomo R, Bailey M, Eastwood GM, Nichol A, Pilcher D, Hart GK, Reade MC, Egi M, Cooper DJ, Study of Oxygen in Critical Care (SOCC) Group: **Arterial hyperoxia and in-hospital mortality after resuscitation from cardiac arrest.** *Crit Care* 2011, **15**:R90.

8. Janz DR, Hollenbeck RD, Pollock JS, McPherson JA, Rice TW: **Hyperoxia is associated with increased mortality in patients treated with mild therapeutic hypothermia after sudden cardiac arrest.** *Crit Care Med* 2012, **40**:3135–3139.
9. Ihle JF, Bernard S, Bailey MJ, Pilcher DV, Smith K, Scheinkestel CD: **Hyperoxia in the intensive care unit and outcome after out-of-hospital ventricular fibrillation cardiac arrest.** *Crit Care Resusc* 2013, **15**:186–190.
10. Wijesinghe M, Perrin K, Ranchord A, Simmonds M, Weatherall M, Beasley R: **Routine use of oxygen in the treatment of myocardial infarction: systematic review.** *Heart* 2009, **95**:198–202.
11. Young P, Beasley R, Bailey M, Bellomo R, Eastwood GM, Nichol A, Pilcher DV, Yunos NM, Egi M, Hart GK, Reade MC, Cooper DJ, Study of Oxygen in Critical Care (SOCC) Group: **The association between early arterial oxygenation and mortality in ventilated patients with acute ischaemic stroke.** *Crit Care Resusc* 2012, **14**:14–19.
12. Rachmale S, Li G, Wilson G, Malinchoc M, Gajic O: **Practice of excessive FiO₂ and effect on pulmonary outcomes in mechanically ventilated patients with acute lung injury.** *Respir Care* 2012, **57**:1887–1893.
13. Raj R, Bendel S, Reinikainen M, Kivisaari R, Siironen J, Lang M, Skrifvars M: **Hyperoxemia and long-term outcome after traumatic brain injury.** *Crit Care* 2013, **17**:R177.
14. Rincon F, Kang J, Maltenfort M, Vibbert M, Urtecho J, Athar MK, Jallo J, Pineda CC, Tzeng D, McBride W, Bell R: **Association between hyperoxia and mortality after stroke: a multicenter cohort study.** *Crit Care Med* 2013, **42**:387–396.
15. Rincon F, Kang J, Vibbert M, Urtecho J, Athar MK, Jallo J: **Significance of arterial hyperoxia and relationship with case fatality in traumatic brain injury: a multicentre cohort study.** *J Neurol Neurosurg Psychiatry* 2013, **85**:799–805.
16. Baleeiro CE, Wilcoxon SE, Morris SB, Standiford TJ, Paine R 3rd: **Sublethal hyperoxia impairs pulmonary innate immunity.** *J Immunol* 2003, **171**:955–963.
17. Kikuchi Y, Tateda K, Fuse ET, Matsumoto T, Gotoh N, Fukushima J, Takizawa H, Nagase T, Standiford TJ, Yamaguchi K: **Hyperoxia exaggerates bacterial dissemination and lethality in *Pseudomonas aeruginosa* pneumonia.** *Pulm Pharmacol Ther* 2009, **22**:333–339.
18. Li LF, Liao SK, Ko YS, Lee CH, Quinn DA: **Hyperoxia increases ventilator-induced lung injury via mitogen-activated protein kinases: a prospective, controlled animal experiment.** *Crit Care* 2007, **11**:R25.
19. Makena PS, Luellen CL, Balazs L, Ghosh MC, Parthasarathi K, Waters CM, Sinclair SE: **Preexposure to hyperoxia causes increased lung injury and epithelial apoptosis in mice ventilated with high tidal volumes.** *Am J Physiol Lung Cell Mol Physiol* 2010, **299**:L711–L719.
20. Nagato A, Silva FL, Silva AR, Bezerra FS, Oliveira ML, Bello-Klein A, Cristovao Porto L, Santos Valenca S: **Hyperoxia-induced lung injury is dose dependent in Wistar rats.** *Exp Lung Res* 2009, **35**:713–728.
21. Tateda K, Deng JC, Moore TA, Newstead MW, Paine R 3rd, Kobayashi N, Yamaguchi K, Standiford TJ: **Hyperoxia mediates acute lung injury and increased lethality in murine *Legionella* pneumonia: the role of apoptosis.** *J Immunol* 2003, **170**:4209–4216.
22. Winslow RM: **Oxygen: the poison is in the dose.** *Transfusion* 2013, **53**:424–437.
23. MacIntyre NR, Cook DJ, Ely EW Jr, Epstein SK, Fink JB, Heffner JE, Hess D, Hubmayer RD, Scheinhorn DJ, American College of Chest Physicians; American Association for Respiratory Care; American College of Critical Care Medicine: **Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College of Critical Care Medicine.** *Chest* 2001, **120**:375S–395S.
24. Martin DS, Grocott MP: **Oxygen therapy in critical illness: precise control of arterial oxygenation and permissive hypoxemia.** *Crit Care Med* 2013, **41**:423–432.
25. Gilbert-Kawai ET, Mitchell K, Martin D, Carlisle J, Grocott MP: **Permissive hypoxaemia versus normoxaemia for mechanically ventilated critically ill patients.** *Cochrane Database Syst Rev* 2014, **5**:CD009931.
26. O'Driscoll BR, Howard LS, Davison AG, British Thoracic S: **BTS guideline for emergency oxygen use in adult patients.** *Thorax* 2008, **63**(Suppl 6):vi1–vi68.
27. Eastwood GM, Reade MC, Peck L, Jones D, Bellomo R: **Intensivists' opinion and self-reported practice of oxygen therapy.** *Anaesth Intensive Care* 2011, **39**:122–126.
28. Mao C, Wong DT, Slutsky AS, Kavanagh BP: **A quantitative assessment of how Canadian intensivists believe they utilize oxygen in the intensive care unit.** *Crit Care Med* 1999, **27**:2806–2811.
29. Tobin MJ: **Mechanical ventilation.** *N Engl J Med* 1994, **330**:1056–1061.
30. Brower RG, Lanken PN, MacIntyre N, Matthay MA, Morris A, Ancukiewicz M, Schoenfeld D, Thompson BT, National Heart L, Blood Institute ACTN: **Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome.** *N Engl J Med* 2004, **351**:327–336.
31. Suzuki S, Eastwood GM, Peck L, Glassford NJ, Bellomo R: **Current oxygen management in mechanically ventilated patients: a prospective observational cohort study.** *J Crit Care* 2013, **28**:647–654.
32. de Graaff AE, Dongelmans DA, Binnekade JM, de Jonge E: **Clinicians' response to hyperoxia in ventilated patients in a Dutch ICU depends on the level of FIO₂.** *Intensive Care Med* 2011, **37**:46–51.
33. Panwar R, Capellier G, Schmutz N, Davies A, Cooper DJ, Bailey M, Baguley D, Pilcher V, Bellomo R: **Current oxygenation practice in ventilated patients - an observational cohort study.** *Anaesth Intensive Care* 2013, **41**:505–514.
34. Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A, Wiedemann HP, Arroliga AC, Fisher CJ, Komara JJ, Perez-Trepichio P, Parsons PE, Wolkin R, Welsh C, Fulkerson WJ, MacIntyre N, Mallatrat L, Sebastian M, McConnell R, Wilcox C, Govert J, Thompson D, Clemmer T, Davis R, Orme J, Weaver L, Grissom C, Eskelson M, Young M, Gooder V, et al: **Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome.** *N Engl J Med* 2000, **342**:1301–1308.
35. Barazzone C, Horowitz S, Donati YR, Rodriguez I, Piguat PF: **Oxygen toxicity in mouse lung: pathways to cell death.** *Am J Respir Cell Mol Biol* 1998, **19**:573–581.
36. Gore A, Muralidhar M, Espey MG, Degenhardt K, Mantell LL: **Hyperoxia sensing: from molecular mechanisms to significance in disease.** *J Immunotoxicol* 2010, **7**:239–254.
37. Nagato AC, Bezerra FS, Lanzetti M, Lopes AA, Silva MA, Porto LC, Valenca SS: **Time course of inflammation, oxidative stress and tissue damage induced by hyperoxia in mouse lungs.** *Int J Exp Pathol* 2012, **93**:269–278.
38. Sinclair SE, Altemeier WA, Matute-Bello G, Chi E: **Augmented lung injury due to interaction between hyperoxia and mechanical ventilation.** *Crit Care Med* 2004, **32**:2496–2501.
39. Eastwood GM, Reade MC, Peck L, Baldwin I, Considine J, Bellomo R: **Critical care nurses' opinion and self-reported practice of oxygen therapy: a survey.** *Aust Crit Care* 2012, **25**:23–30.
40. Blakeman TC: **Evidence for oxygen use in the hospitalized patient: is more really the enemy of good?** *Respir Care* 2013, **58**:1679–1693.
41. Jubran A, Tobin MJ: **Reliability of pulse oximetry in titrating supplemental oxygen therapy in ventilator-dependent patients.** *Chest* 1990, **97**:1420–1425.
42. Bak Z, Sjoberg F, Rousseau A, Steinvall I, Janerot-Sjoberg B: **Human cardiovascular dose-response to supplemental oxygen.** *Acta Physiol (Oxf)* 2007, **191**:15–24.
43. Budinger GR, Mutlu GM: **Balancing the risks and benefits of oxygen therapy in critically ill adults.** *Chest* 2013, **143**:1151–1162.
44. Davis DP, Meade W, Sise MJ, Kennedy F, Simon F, Tominaga G, Steele J, Coimbra R: **Both hypoxemia and extreme hyperoxemia may be detrimental in patients with severe traumatic brain injury.** *J Neurotrauma* 2009, **26**:2217–2223.
45. Suzuki S, Eastwood GM, Glassford NJ, Peck L, Young H, Garcia-Alvarez M, Schneider AG, Bellomo R: **Conservative oxygen therapy in mechanically ventilated patients: a pilot before-and-after trial.** *Crit Care Med* 2014, **42**:1414–1422.

doi:10.1186/s13613-014-0023-y

Cite this article as: Helmerhorst et al.: Self-reported attitudes versus actual practice of oxygen therapy by ICU physicians and nurses. *Annals of Intensive Care* 2014 **4**:23.