

## STUDIES ON ICTERUS NEONATORUM

### THE PRODUCTION OF ICTERUS IN ANIMALS FOLLOWING PROLONGED ANOXAEMIA<sup>1</sup>

BY ALTON GOLDBLOOM AND RUDOLF GOTTLIEB

*(From the Department of Medicine, McGill University Clinic, Royal Victoria Hospital, Montreal)*

(Received for publication October 9, 1929)

In a previous paper (1) in which we reported our studies on icterus neonatorum, we demonstrated the readiness with which the blood of a newly born infant haemolysed as compared with the blood of an older child or an adult. We showed also that the morphologic pattern of the blood of the new born was considerably different; for instance the presence of a polycythaemia and the increase in the number of nucleated red cells and reticulocytes. We showed too, that reduction of the polycythaemia and of the number of immature cells in the circulation coincided with an increase in the bilirubin in the serum, the development of icterus and the development of a normal resistance of the red cells in varying concentrations of salt solution. We showed also that the diminished resistance of the red cells of the newly born to salt solution is due to the immature cells in the circulation, namely that these cells haemolyse even in normal salt solution, and that the cells remaining after these immature forms have been destroyed, show normal resistance.

In attempting to explain the nature of icterus neonatorum, we reasoned that, in view of the admixture of arterial and venous blood, the right to left shunt existing in ante natal life, the foetus, was comparable to any other individual living in a state of diminished oxygen supply. That immediately after birth, with the cessation of admixture in normal infants, the necessity for polycythaemia no longer

<sup>1</sup> This research was made possible by a grant from the Cooper Fund.

TABLE 1  
*Experimental data*

Number of animal	Age <i>weeks</i>	Date	Weight <i>grams</i>	R.B.C. counts	Reticulated R.B.C. number per H.F.F.	Nucleated R.B.C.	Fragility of R.B.C.	Hemoglobin percent- age (Haldane)	Icteric index (Davis)	Van den Bergh units	Remarks	Atmospheric pressure
1	12	January 24	200	5,050,000	0	0	Normal	90	3	Negative	Animal in apparatus Died. Sections of bone marrow made	50 to 300 mm. below normal
		February 1		5,610,000	2	0	Normal	100	3	Negative		
2	10	January 24	225	5,150,000	0	0	Normal	96	1	Negative	Animal in apparatus	50 to 300 mm. below normal
		January 31		6,230,000	3	0	Normal	112	1	Negative		
		February 7		6,820,000	5	0	Normal	124	1	Negative		
		February 16		6,850,000	5	0	Normal	124	1	Negative		
		February 18		6,280,000	4	0	Normal	120	4	Negative		
		February 19		5,820,000	2	0	Normal	110	9	1 $\frac{1}{2}$		
		February 22		5,050,000	0	0	Normal	92	8	1 $\frac{1}{2}$		
February 23		5,030,000	0	0	Normal	92	6	$\frac{3}{4}$				
February 28		5,110,000	0	0	Normal	96	1	Negative				
3	12	February 2	225	5,220,000	0	0	Normal	94	2	Negative	Animal in apparatus	50 to 300 mm. below normal
		February 6		5,960,000	2	0	Normal	110	2	Negative		
		February 12		6,910,000	5	0	Normal	126	2	Negative		
		February 14		6,890,000	5	0	Normal	126	2	Negative		
		February 27		6,940,000	5	0	Normal	126	2	Negative		
		February 28		6,560,000	3	0	Normal	120	5	Negative		

3	12	March 1	6,150,000	2	0	Normal	110	8	1 ‡	Removed to cage	Normal
		March 3	5,200,000	0	0	Normal	96	5	1		
		March 5	5,130,000	0	0	Normal	92	2	Negative		
		March 24	5,230,000	0	0	Normal	94	2	Negative		
4	11	January 25	5,050,000	0	0	Normal	92	1	Negative	Control animal	Normal
		February 17	5,060,000	0	0	Normal	92	1	Negative		
		March 24	5,060,000	0	0	Normal	92	1	Negative		
		April 1	5,080,000	0	0	Normal	92	1	Negative	Killed for section of bone marrow	
5	14	January 25	5,210,000	0	0	Normal	92	2	Negative	Control animal	Normal
		February 12	5,180,000	0	0	Normal	90	2	Negative		
		February 17	5,220,000	0	0	Normal	92	2	Negative		
		March 24	5,220,000	0	0	Normal	92	2	Negative		
6	14	April 10	5,200,000	0	0	Normal	92			Killed for section of bone marrow	
		February 25	5,180,000	0	0	Normal	94	1	Negative	Animal in apparatus	50 to 300 mm. below normal
		March 6	6,010,000	3	0	Normal	112	1	Negative		
		March 10	6,350,000	4	0	Normal	120	1	Negative		
6	14	March 14	6,670,000	5	0	Normal	124	1	Negative		
		March 24	6,660,000	5	0	Normal	124	1	Negative		
		March 25	6,430,000	5	0	Normal	120	5	Trace	Removed to cage	Normal
		March 26	6,150,000	4	0	Normal	112	6	‡		
6	14	March 30	5,120,000	0	0	Normal	92	8	1 ‡		
		April 5	5,190,000	0	0	Normal	94	3	Trace		
		April 12	5,190,000	0	0	Normal	94	1	Negative		

TABLE 1—Continued

Number of animal	Age weeks	Date	Weight grams	R.B.C. counts	Reticulated R.B.C. number per H.P.F.	Nucleated R.B.C.	Fragility of R.B.C.	Hemoglobin percent- age (Haldane)	Icteric index (Davis)	Van den Berg units	Remarks	Atmospheric pressure	
7	12	February 25	250	5,320,000	0	0	Normal	96	3	Negative	Animal in apparatus	50 to 300 mm. below normal	
		March 6		6,400,000	3	0	Normal	116	3	Negative			
		March 10		6,800,000	4	0	Normal	122	3	Negative			
		March 14		6,810,000	5	0	Normal	124	3	Negative			
		March 25	290	6,620,000	4	0	Normal	122	6	Trace		Removed to cage	Normal
		March 26		6,330,000	3	0	Normal	116	8	1½			
		March 27		6,020,000	2	0	Normal	112	8	1½		Killed for sections of bone marrow	
8	12	February 25	275	5,210,000	0	0	Normal	92	1	Negative	Animal in apparatus	50 to 300 mm. below normal	
		March 6		6,130,000	3	0	Normal	110	1	Negative			
		March 14		6,810,000	5	0	Normal	122	1	Negative			
		March 25	310	6,700,000	4	0	Normal	120	3	Negative		Removed to cage	Normal
		March 27		5,940,000	2	0	Normal	108	9	2			
		March 30		5,270,000	0	0	Normal	92	9	2			
		April 7		5,190,000	0	0	Normal	92	2	Trace			
		April 12		5,215,000	0	0	Normal	92	1	Negative			

9	10	February 25	225					94	2	Negative	Animal in apparatus	50 to 300 mm. below normal
		March 6	5,480,000	0	0	Normal	116	2	Negative			
		March 10	6,610,000	4	0	Normal	120	2	Negative			
		March 14	6,800,000	4	0	Normal	126	2	Negative			
		March 24	7,420,000	6	0	Normal	126	2	Negative			
10	11	March 25	260	7,420,000	6	0	Normal	126	2	Negative	Removed to cage	Normal
		March 25	7,180,000	4	0	Normal	120	5	1			
		March 26	6,910,000	3	0	Normal	114	7	1½			
		March 27	6,510,000	2	0	Normal	108	7	1½			
		February 25	250					90	1	Negative	Animal in apparatus	50 to 300 mm. below normal
11	12	March 6	5,100,000	0	0	Normal	110	1	Negative			
		March 10	6,130,000	3	0	Normal	118	1	Negative			
		March 14	6,510,000	5	0	Normal	122	1	Negative			
		March 24	6,740,000	6	0	Normal	122	1	Negative			
		March 25	6,750,000	6	0	Normal	118	5	Trace			
11	12	March 25	6,540,000	5	0	Normal	114	7	1			
		March 26	6,330,000	4	0	Normal	106	7	1			
		March 27	5,870,000	3	0	Normal	90	6	¾			
		March 31	5,060,000	0	0	Normal	90	3	Trace			
		April 5	5,150,000	0	0	Normal	90	1	Negative			
11	12	April 7	5,100,000	0	0	Normal	90	1	Negative			
		April 12	5,120,000	0	0	Normal	94	2	Negative			
		February 25	250	5,390,000	0	0	Normal	104	2	Negative	Animal in apparatus	50 to 300 mm. below normal
		March 6	5,910,000	3	0	Normal	110	2	Negative			
		March 10	6,210,000	5	0	Normal	114	2	Negative			
11	12	March 14	6,470,000	7	0	Normal	116	2	Negative			
		March 18	6,580,000	6	0	Normal		2	Negative			

TABLE 1—Continued

Number of animal	Age weeks	Date	Weight grams	R.B.C. counts	Reticulated R.B.C. number per H.P.F.	Nucleated R.B.C.	Fragility of R.B.C.	Hemoglobin percent- age (Haldane)	Icteric index (Davis)	Van den Bergh units	Remarks	Atmospheric pressure
11	12	March 20		6,590,000	5	0	Normal	116	2	Negative	Removed to cage	Normal
		March 24	280	6,580,000	5	0	Normal	116	2	Negative		
		March 25		6,230,000	3	0	Normal	108	6	$\frac{2}{4}$		
		March 26		5,910,000	3	0	Normal	104	8	$1\frac{1}{4}$		
		March 27		5,580,000	2	0	Normal	96	8	$1\frac{1}{4}$		
12	12	March 29		5,250,000	0	0	Normal	92	5	1	Control animal	Normal
		March 31		5,350,000	0	0	Normal	94	3	Trace		
		April 5		5,380,000	0	0	Normal	94	1	Negative		
		March 1	210	5,210,000	0	0	Normal	94	1	Negative		
		March 5		5,200,000	0	0	Normal	94	1	Negative		
13	9	March 6		5,170,000	0	0	Normal	92	1	Negative	Killed for section of bone marrow	50 to 300 mm below normal
		March 8		5,200,000	0	0	Normal	94	1	Negative		
		April 11	175	5,280,000	0	0	Normal	96	2	Negative		
		April 16		6,380,000	3	0	Normal	116	2	Negative		
		April 20		7,110,000	5	0	Normal	124	2	Negative		
April 23	180	7,010,000	5	0	Normal	122	3	Negative	Removed to cage	Normal		

13	9	April 24	6,650,000	4	0	Normal	120	4	Trace	Animal in apparatus 50 to 300 mm. below normal
		April 25	6,210,000	3	0	Normal	106	8	1½	
		April 29	5,200,000	0	0	Normal	94	8	1½	
		May 1	5,280,000	0	0	Normal	96	6	1	
		May 3	5,290,000	0	0	Normal	96	2	Negative	
14	9	April 11	5,175,000	0	0	Normal	94	1	Negative	Animal in apparatus 50 to 300 mm. below normal
		April 16	6,790,000	4	0	Normal	120	1	Negative	
		April 20	6,850,000	6	0	Normal	126	1	Negative	
		April 21	6,710,000	6	0	Normal	124	1	Negative	
		April 22	6,320,000	4	0	Normal	116	5	½	
		April 23	5,980,000	3	0	Normal	108	8	1½	
		April 26	5,160,000	0	0	Normal	94	8	1½	
		April 28	5,150,000	0	0	Normal	94	6	1	
		April 30	5,180,000	0	0	Normal	94	2	Trace	
May 1	5,170,000	0	0	Normal	94	1	Negative			
15	9	April 11	5,320,000	0	0	Normal	92	1	Negative	Animal in apparatus 50 to 300 mm. below normal
		April 16	6,300,000	3	0	Normal	116	1	Negative	
		April 20	6,850,000	5	0	Normal	124	1	Negative	
		April 23	6,810,000	5	0	Normal	124	1	Negative	
		April 25	6,010,000	3	0	Normal	108	5	1	
		April 26	5,590,000	2	0	Normal	-400	9	1½	
		April 28	5,240,000	0	0	Normal	96	9	1½	
		April 30	5,330,000	0	0	Normal	96	5	1	
		May 2	5,310,000	0	0	Normal	96	1	Negative	

TABLE 1—Concluded

Number of animal	Age weeks	Date	Weight grams	R.B.C. counts	Reticulated R.B.C. number per H.P.F.	Nucleated R.B.C.	Fragility of R.B.C.	Hemoglobin percent- age (Haldane)	Icteric index (Davis)	Van den Bergh units	Remarks	Atmospheric pressure
16	9	April 11	175	5,080,000	0	0	Normal	90	2	Negative	Animal in apparatus	50 to 300 mm. below normal
		April 16		6,160,000	3	0	Normal	110	2	Negative		
		April 20		6,580,000	5	0	Normal	120	2	Negative		
		April 21	180	6,310,000	4	0	Normal	116	3	Negative		
		April 23		6,740,000	3	0	Normal	104	8	1		
17	9	April 27		5,040,000	0	0	Normal	90	8	1	Removed to cage	Normal
		April 29		5,100,000	0	0	Normal	90	3	Trace		
		April 30		5,090,000	0	0	Normal	90	2	Negative		
		April 11	170	5,120,000	0	0	Normal	92	1	Negative		
		April 16		5,090,000	0	0	Normal	92	1	Negative		
18	9	April 21	185	5,130,000	0	0	Normal	92	1	Negative	Control animal	Normal
		April 11	160	5,230,000	0	0	Normal	96	2	Negative		
		April 16		5,250,000	0	0	Normal	96	2	Negative		
		April 21		5,230,000	0	0	Normal	96	2	Negative		
		April 23	185	5,210,000	0	0	Normal	96	2	Negative		
		May 1		5,250,000	0	0	Normal	96	2	Negative		






FIG. 1. APPARATUS FOR REDUCED ATMOSPHERIC PRESSURE

existed, leading to destruction of the excess of red cells and thus to icterus neonatorum.

We felt therefore that if these theories are correct it should be possible not only to produce a polycythaemia by keeping animals under reduced atmospheric pressure but also to produce some degree of jaundice in these animals shortly after the pressure is restored to normal.

Accordingly we have attempted to produce a polycythaemia in




FIG. 2. BONE MARROW OF A NORMAL GUINEA PIG

guinea pigs by keeping them under reduced atmospheric pressure by means of a specially devised apparatus (see fig. 1). Tube 1 connects a glass jar, which is fixed air tight to a glass plate, with a vacuum pump. Tube 2 connects the jar with a manometer. Tube 3 allows air to enter, the amount of which could be regulated by a stop cock.

Eighteen young guinea pigs were used for this experiment, thirteen of them were kept from 2 to 4 weeks under reduced atmospheric pres-

sure while the other five were kept as controls. One animal (no. 1) died at the end of one week in the apparatus, but had already begun to show an increase in the number of red blood cells. Sections of the bone marrow of this animal were made and showed increased activity. A sixth control animal was killed at the beginning of the experiment for bone marrow studies. The pressure in the apparatus was gradu-




FIG. 3. BONE MARROW OF A GUINEA PIG AFTER 4 WEEKS EXPOSURE TO REDUCED ATMOSPHERIC PRESSURE

ally reduced to 300 mm. below normal atmospheric pressure and then kept constant at this level, which corresponds to an altitude of about 18,000 feet. Food was introduced daily through the top opening; for that reason and on account of the blood studies the pressure had to be increased every day for a few minutes. A total of about 200 minutes of normal atmospheric pressure was necessary during the

whole experiment. The blood of the animals was examined every 48 hours. The examinations were made of the number of red blood corpuscles, the percentage of haemoglobin, the number of reticulated and nucleated red blood cells, the fragility of the red blood cells, the icteric index, and the van den Bergh reaction. The blood for the examination was taken by puncture of the ear, except for the van den Bergh reaction, for which 0.5 cc. was taken by heart puncture.




FIG. 4. SHOWING AVERAGE PERCENTILE CHANGE OF RED BLOOD CELLS AND HEMOGLOBIN, INCREASE IN UNITS OF THE VAN DEN BERGH AND THE PRESSURE AT WHICH THE ANIMALS WERE KEPT, PLOTTED AGAINST DAYS OF THE EXPERIMENTS

*Red blood cell counts and haemoglobin percentage*

The number of the red blood cells and the percentage of haemoglobin increased rapidly, reaching an average maximum increase of about 30 per cent within 10 days. Five to eight days after the animals were removed to normal atmospheric pressure again, the number of red blood cells and the percentage of haemoglobin returned to their original figures. The chart (fig. 4) shows the average percentile rise of the red blood cells and the haemoglobin, which always ran parallel.

*Reticulated red blood cells*

Reticulated red blood cells were not present in the blood of the animals at normal atmospheric pressure. Three to four days after exposure of the animals to reduced atmospheric pressure, reticulated red blood cells began to appear, reaching in one week the maximal number of 5 to 6 per high power field in thick smears. Within one week after the removal of the animals to normal atmospheric pressure the reticulated red blood cells disappeared again.

*Nucleated red blood cells and fragility of the red blood cells*

Nucleated red blood cells were at no time present. Fragility of the red blood cells did not show any change during the experiment.

*The icteric index and the van den Bergh reaction*

Normal guinea pigs showed an icteric index of from 1 to 3, while the van den Bergh reaction was always negative. The icteric index and the van den Bergh reaction remained unchanged as long as the animals were in the chamber under reduced atmospheric pressure. About 48 hours after the animals were removed from the apparatus to normal atmospheric pressure, the icteric index began to rise and the indirect van den Bergh reaction became positive, both reaching their maximum within 24 hours, an icteric index of from 7 to 9, and an indirect van den Bergh reaction of from 1 to 2 units. Both icteric index and the van den Bergh reaction remained high for about 8 days and then came gradually down to normal, reaching normal values in 3 to 5 days after the red blood cell count had returned to normal. The table gives a synopsis of the events during the whole experiment.

## THE BONE MARROW

The sections made from the bone marrow of the confined animals showed greater activity than that of the controls (figs. 2 and 3).

## SUMMARY OF FINDINGS

1. Guinea pigs were kept under reduced atmospheric pressure. A polycythaemia developed rapidly, reaching its maximum in about 10 days. The average increase of the red blood cells as well as the haemoglobin was about 30 per cent above the original values. Return to

normal took place in about 5 to 8 days after the pressure was increased to normal.

2. There was with the polycythaemia a marked increase in reticulated red blood cells, 5 to 6 per high power field. When the pressure was increased to normal the reticulated cells disappeared in about one week.

3. There was an increase in the icteric index and an appearance of positive indirect van den Bergh reactions after the animals were removed from the low atmospheric pressure chamber. The icteric index ranged from 7 to 9 (normal 1 to 3) and the van den Bergh reaction from 1 to 2 units (normal 0).

#### COMMENT

Our previous work demonstrated that the infant develops icterus neonatorum as a result of haemolysis of red blood cells in the course of a period of postnatal re-adjustment, during which the polycythaemia disappears.

The above experiments were designed to produce a polycythaemia in animals through a prolonged diminution of their oxygen supply, and to observe whether icterus occurred during their period of re-adjustment from polycythaemia to the normal red cell count.

Polycythaemia was successfully produced. As in new born infants, the animals released from the environment of lowered atmospheric pressure, very soon showed an increasing icteric index and a positive indirect van den Bergh reaction together with a reduction of the polycythaemia and the disappearance of the reticulated cells.

*Thus an icterus, similar to icterus neonatorum, was produced in animals.*

The period of re-adjustment was of somewhat shorter duration than in the human foetus. No nucleated red blood cells were found in these animals and the fragility of their red cells showed no changes.

It would therefore appear quite reasonable to conclude that icterus neonatorum is a haemolytic icterus which is the result of a postnatal re-adjustment from a condition of oxygen unsaturation to a normal oxygen saturation.

#### BIBLIOGRAPHY

1. Goldbloom, A., and Gottlieb, R., *Am. J. Dis. Child.*, 1929, xxxviii, 57. Icterus Neonatorum.