

# Highway proximity and black carbon from cookstoves as a risk factor for higher blood pressure in rural China

Jill Baumgartner<sup>a,b,1</sup>, Yuanxun Zhang<sup>c</sup>, James J. Schauer<sup>d</sup>, Wei Huang<sup>c</sup>, Yuqin Wang<sup>c</sup>, and Majid Ezzati<sup>e</sup>

<sup>a</sup>Institute for Health and Social Policy and Department of Epidemiology, Biostatistics, and Occupational Health, McGill University, Montreal, QC, Canada H3A 1A3; <sup>b</sup>Institute on the Environment, University of Minnesota, St. Paul, MN 55108; <sup>c</sup>College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China; <sup>d</sup>Environmental Chemistry and Technology Program, Department of Civil and Environmental Engineering, University of Wisconsin-Madison, Madison, WI 53706; and <sup>e</sup>MRC-PHE Centre for Environment and Health, Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London W2 1PG, United Kingdom

Edited by Barry R. Bloom, Harvard School of Public Health, Boston, MA, and approved July 29, 2014 (received for review September 13, 2013)

**Air pollution in China and other parts of Asia poses large health risks and is an important contributor to global climate change. Almost half of Chinese homes use biomass and coal fuels for cooking and heating. China's economic growth and infrastructure development has led to increased emissions from coal-fired power plants and an expanding fleet of motor vehicles. Black carbon (BC) from incomplete biomass and fossil fuel combustion is the most strongly light-absorbing component of particulate matter (PM) air pollution and the second most important climate-forcing human emission. PM composition and sources may also be related to its human health impact. We enrolled 280 women living in a rural area of northwestern Yunnan where biomass fuels are commonly used. We measured their blood pressure, distance from major traffic routes, and daily exposure to BC (pyrolytic biomass combustion), water-soluble organic aerosol (organic aerosol from biomass combustion), and, in a subset, hopane markers (motor vehicle emissions) in winter and summer. BC had the strongest association with systolic blood pressure (SBP) (4.3 mmHg;  $P < 0.001$ ), followed by PM mass and water-soluble organic mass. The effect of BC on SBP was almost three times greater in women living near the highway [6.2 mmHg; 95% confidence interval (CI), 3.6 to 8.9 vs. 2.6 mmHg; 95% CI, 0.1 to 5.2]. Our findings suggest that BC from combustion emissions is more strongly associated with blood pressure than PM mass, and that BC's health effects may be larger among women living near a highway and with greater exposure to motor vehicle emissions.**

cardiovascular disease | household air pollution | solid fuels

**P**articulate matter (PM) air pollution is a leading health risk factor (1) and primary contributor to anthropogenic climate change (2). Air pollution is notoriously high in China and other parts of Asia. China's rising energy demands have led to increased air pollution emissions from coal-fired power plants (3). Its motorized transport growth is the fastest in the world with the number of motor vehicles projected to quadruple in the next two decades, reaching over 380 million by 2030 (4). Meanwhile, nearly half of all Chinese still cook and heat their homes with highly polluting biomass and coal fuels (5). The resulting PM concentrations routinely exceed the World Health Organization's (WHO) annual Air Quality Guideline of  $10 \mu\text{g}/\text{m}^3$  by a factor of 10 or more (6) and are associated with a number of adverse health outcomes, including cardiovascular diseases (1, 7).

PM differs in chemical properties, size, and possibly effects on human health. Black carbon (BC) and organic carbon PM are emitted during incomplete biomass and fossil fuel combustion and seem to have important effects on both climate and human health. BC affects the regional and global climate by absorbing solar radiation and heating the atmosphere and is the second most important climate-forcing human emission, after carbon dioxide (8). Coemitted organic carbon may further influence radiative forcing by acting as cloud condensation nuclei (9). These specific characteristics and sources of PM may also impact its toxicity to humans (10).

We previously found that daily exposure to  $\text{PM} < 2.5$  microns in aerodynamic diameter ( $\text{PM}_{2.5}$ ) was associated with higher blood pressure in older Chinese women cooking with biomass fuels (11). In the current study, we used chemical and optical methods to analyze the  $\text{PM}_{2.5}$  exposure samples for BC and organic components and evaluated their associations with blood pressure, the leading risk factor for cardiovascular diseases, worldwide and in China (1). We enrolled 280 women aged 25–90 y who lived in six villages in the surrounding area along the Yunnan–Tibet Highway in the Himalayan foothills of Yunnan Province, China. These women were subsistence farmers and used biomass fuels (largely wood and crop residues) for cooking and space heating. Details about the study site, household fuel and stove use patterns, and exposure to other PM sources like direct and involuntary tobacco smoking are reported elsewhere (12).

The independent and combined effects of different PM components from various sources on human health are poorly understood. We used measurements of women's daily PM exposure and proximity to a highway to examine how PM composition and sources affect the hazardous effects of blood pressure in this group, using proximity to the highway as a proxy for exposure to motor vehicle emissions using organic tracers.

## Results

We enrolled 280 women (mean age: 51.9 y), 18% of whom were overweight [body mass index (BMI) =  $25\text{--}30 \text{ kg}/\text{m}^2$ ] and 4% obese (BMI  $\geq 30 \text{ kg}/\text{m}^2$ ). Mean systolic and diastolic blood pressure (SBP and DBP) were 120 [95% confidence interval (CI), 118 to 122] and 72 mmHg (95% CI, 71 to 73), respectively.

## Significance

**Air pollution is a leading health risk factor and important contributor to regional climate change in China and other parts of Asia. China's particulate matter (PM) air pollution dramatically exceeds health guidelines and is impacted by industrial emissions, motor vehicles, and household use of biomass and coal fuels. Black carbon (BC) from biomass and fossil fuel burning is a major climate-forcing component of PM. We found that BC exposure from biomass smoke is more strongly associated with blood pressure than total PM mass, and that coexposure to motor vehicle emissions may strengthen BC's impact. Air pollution mitigation efforts focusing on reducing combustion pollution are likely to have major benefits for climate and human health.**

Author contributions: J.B., Y.Z., J.J.S., and M.E. designed research; J.B., Y.Z., W.H., and Y.W. performed research; Y.Z. and J.J.S. contributed new reagents/analytic tools; J.B., W.H., and Y.W. analyzed data; and J.B., J.J.S., and M.E. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. Email: jill.baumgartner@mcgill.ca.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1317176111/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1317176111/-DCSupplemental).

Thirteen percent ( $n = 37$ ) of participants were hypertensive, of whom 17% ( $n = 6$ ) were taking antihypertensive medication. None of the participants reported a previous cardiovascular event, and just two women (<1%) reported physician-diagnosed diabetes (Tables S1 and S2).

**Exposure of Rural Chinese Women to PM<sub>2.5</sub> Mass, BC, Organic Carbon, and Motor Vehicle Emission Tracers.** Women's geometric mean 24-h PM<sub>2.5</sub> exposure in summer was 55  $\mu\text{g}/\text{m}^3$  (range: 9–492) and in winter was 117  $\mu\text{g}/\text{m}^3$  (range: 22–634) (12). These levels greatly exceed the WHO's 24-h Air Quality Guideline of 25  $\mu\text{g}/\text{m}^3$ . Geometric mean BC exposure was 5.2  $\mu\text{g}/\text{m}^3$  (4  $\mu\text{g}/\text{m}^3$  in summer and 6  $\mu\text{g}/\text{m}^3$  in winter; range: 2–44) which is lower than BC exposure in central New Delhi traffic (42  $\mu\text{g}/\text{m}^3$ ) (13) but surpasses daytime ambient BC in Beijing, Mexico City, and several cities in Brazil (range: 1.9–4.8  $\mu\text{g}/\text{m}^3$ ) (14–16). Exposure to water-soluble organic mass (WSOM), a more specific marker of primary biomass smoke, was considerably higher than BC (geometric mean of 12  $\mu\text{g}/\text{m}^3$  in summer and 33  $\mu\text{g}/\text{m}^3$  in winter; range: 1–352) (Table 1), indicating that women's exposure is greatly influenced by biomass combustion (17). Exposure to all pollutants was lower in the summer compared with the winter season; however, the relative contribution of BC and WSOM fractions to PM<sub>2.5</sub> mass was approximately the same across seasons and for different age groups (Fig. S1).

The PM components had a low-to-moderate correlation, with higher correlations in summer than in winter (Table S3). The largest correlation was between PM<sub>2.5</sub> and WSOM ( $r = 0.67$ ), followed by PM<sub>2.5</sub> and BC ( $r = 0.48$ ), and finally BC and WSOM ( $r = 0.44$ ). The PM–BC correlation for rural Chinese women in our study was lower than in outdoor air pollution studies in North America and Europe and in urban Shanghai (PM–BC correlation range = 0.50–0.90) (18, 19). Similarly, BC and hopanes were barely correlated ( $r = 0.18$ ) in our study, even though both are emitted from motor vehicles. These correlations were low possibly because women's exposures were dominated by a single emission source, namely biomass combustion, rather than traffic emissions or an urban mixture of sources.

Among women living relatively close to the highway that passed through our study site (i.e., less than the median distance of 208 m in our sample), daily BC exposure was slightly higher than those living farther away in winter, but the opposite occurred in summer (Table S4). Distance from the highway was not strongly related to BC exposure, with each 100 m from the highway associated with a 0.02  $\ln(\mu\text{g}/\text{m}^3)$  lower BC exposure (95% CI, 0 to 0.02;  $P = 0.30$ ). In contrast, average exposure to hopanes, specific tracers of motor vehicle exhaust in our study setting, was significantly higher among women living in the village closest to the highway (median distance = 76 m) compared with those in the village farthest from the highway (median distance = 548 m) (4.6 vs. 1.1  $\text{ng}/\text{m}^3$ ;  $P < 0.001$  for both seasons; Table S4). In fact, the average near-highway hopane exposures exceeded occupational levels among US trucking terminal workers (4.6 vs. 1.9  $\text{ng}/\text{m}^3$ ) (20).

**Associations with Blood Pressure.** We evaluated the associations of SBP and DBP with exposure to PM<sub>2.5</sub> mass, BC, and WSOM. We express the results as the changes in SBP or DBP associated with a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in pollutant exposure using one and two-pollutant multivariate mixed-effects models because there was evidence of a nonlinear association (21, 22).

BC exposure had the largest independent effect on blood pressure among rural Chinese women (Fig. 1). In models with just one pollutant, a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in BC was associated with 4.3-mmHg higher SBP ( $P < 0.001$ ), followed by PM<sub>2.5</sub> mass (2.2 mmHg;  $P = 0.002$ ) and WSOM (1.2 mmHg;  $P = 0.06$ ). The estimated effect of BC on SBP changed little (7% change) and remained statistically significant after including PM<sub>2.5</sub> mass or WSOM in the model. In contrast, the estimated effect of PM<sub>2.5</sub> mass on SBP decreased by 77% and was no longer statistically significant when BC exposure was added to the model. WSOM had no effect on SBP after other PM components were included in the model (Fig. 1A). We found the same strong and statistically robust relationship between BC exposure and DBP (Fig. 1B). Our conclusions remained the same when evaluating the changes in blood pressure associated with an interquartile range (IQR) increment in log-transformed pollution exposures, although the difference in the estimated effect of BC on blood pressure relative to PM and water-soluble organic carbon (WSOC) was slightly reduced. An IQR increase in  $\ln(\text{BC})$  had the strongest and most robust association with higher SBP (3.6 mmHg; 95% CI, 2.0 to 5.2), followed by  $\ln(\text{PM}_{2.5})$  (2.8 mmHg; 95% CI, 1.0 to 4.6) and  $\ln(\text{WSOM})$  (1.8 mmHg; 95% CI, 0.1 to 3.5) (Fig. S2).

We conducted a separate analysis for younger (25–50 y) vs. older women (>50 y). BC was more strongly associated with blood pressure than PM mass among both younger and older women. A 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in BC exposure was associated with a 1.8-mmHg (95% CI, 0 to 3.6) higher SBP in younger women at the sample average, compared with no effect for PM<sub>2.5</sub>. Among women >50 y old, a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in BC was associated with a 7.4-mmHg (95% CI, 4.0 to 10.8) higher SBP and a 2.9-mmHg (95% CI, 1.1 to 4.7) higher DBP (Fig. 2). Our conclusions did not change when evaluating the changes in blood pressure associated with IQR increases in  $\ln(\text{BC})$  and  $\ln(\text{PM})$  exposures by age. The effect of an IQR increase in  $\ln(\text{PM})$  on blood pressure was similar to models estimating a 1- $\ln(\mu\text{g}/\text{m}^3)$  change in pollution, while the effect of the IQR change in  $\ln(\text{BC})$  was slightly reduced. In older women, an IQR increase in  $\ln(\text{BC})$  was associated with a 6.2-mmHg (95% CI, 3.4 to 9.0) higher SBP and a 2.4-mmHg (95% CI, 0.9 to 3.9) higher DBP. In comparison, an IQR increase in  $\ln(\text{PM})$  was associated with a 5-mmHg high SBP (95% CI, 1.9 to 8.1) and a 2.2-mmHg (95% CI, 0.5 to 3.9) higher DBP (Fig. S3).

The relatively consistent pollution ratios across age groups and seasons (Fig. S1) suggest that stronger blood pressure effects of pollution in older women are not a result of age-specific differences in PM composition. Excluding women taking hypertensive medication from the analysis did not change our estimated

**Table 1. Descriptive statistics for 24-h average personal exposure to PM<sub>2.5</sub> mass, BC, and WSOM ( $\mu\text{g}/\text{m}^3$ ) among Chinese women cooking with biomass fuels, by season**

Pollutant	Summer				Winter			
	<i>n</i> (missing)*	GM (95% CI)	Min–max	IQR	<i>n</i> (missing)	GM (95% CI)	Min–max	IQR
PM <sub>2.5</sub> mass	214 (0)	55 (49 to 62)	9–492	61	262 (0)	117 (107 to 128)	22–634	120
BC	211 (3)	4 (4 to 4)	2–14	2	262 (0)	6 (6 to 7)	2–44	7
WSOM	211 (3)	12 (11 to 14)	1–235	21	261 (1)	33 (30 to 37)	1–352	36

\*The sample size (*n*) refers to the 24-h exposure sample for a woman enrolled in our study. Missing values are for filters that were damaged during optical or chemical analysis (~1% of samples). GM, geometric mean; Min–max, minimum to maximum.




Fig. 1. Associations of personal exposure to PM mass, BC, and WSOM on (A) SBP and (B) DBP using one- and two-pollutant mixed-effects regression models.  $\Delta$ SBP or  $\Delta$ DBP represent the difference in SBP or DBP (with 95% CIs) associated with a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in pollutant exposure.

effects of a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase on SBP (7.3 mmHg; 95% CI, 4.2 to 10.5) or DBP (2.9 mmHg; 95% CI, 1.2 to 4.5) in older women >50 y old.

**Distance from Highway, BC Exposure, and Blood Pressure.** BC exposure had a larger effect on blood pressure among women living closer (i.e., <208 m from the highway) to the highway that passes through our study site than those living farther away. In the former group, a 1- $\ln(\mu\text{g}/\text{m}^3)$  increase in BC exposure was associated with almost threefold higher SBP than in women living away from the highway (6.2 mmHg; 95% CI, 3.6 to 8.9 vs. 2.6 mmHg; 95% CI, 0.1 to 5.2; interaction  $P = 0.04$ ). The effect of BC exposure on DBP was also noticeably larger among women living near the highway (2.6 mmHg; 95% CI, 1.0 to 4.2 vs. 0.3 mmHg; 95% CI, -1.3 to 2.0; interaction  $P = 0.02$ ). The 3.6-mmHg larger effect of 1- $\ln(\text{BC})$  on SBP among women near the highway is similar to the SBP impact of a modest reduction in sodium intake (-4.2 mmHg per 4.4-g daily reduction) (23) and within the range of an SBP decrease with use of antihypertensive medication (-2.5 to 12 mmHg) (24).


Highway proximity was also a predictor of the effect of BC on blood pressure at the village level, with the effect strongest among women living in villages near the highway and absent among those living in the two villages farthest away from the highway (Fig. 3). Distance from the highway was not independently related to blood pressure, nor did the relationship between

PM<sub>2.5</sub> or WSOM exposure and blood pressure differ by distance from the highway (all interaction  $P$  values >0.80).

## Discussion

We found that BC exposure was more strongly associated with blood pressure than PM<sub>2.5</sub> mass and WSOM among Chinese women cooking with biomass, and that the effect was stronger among women living near the highway and with greater exposure to motor vehicle emissions. These findings provide several important directions for future health effects studies and for policies and other mitigation strategies aimed at reducing air pollution emissions and exposures.

First, BC may be a useful indicator of the cardiovascular health and climate benefits of interventions that lower air pollution concentrations and exposures. Air pollution mitigation efforts and guidelines, including those in China, have traditionally focused almost exclusively on PM rather than its components or sources (25). However, different interventions may affect PM components, like BC, by varying amounts. For example, some so-called improved cookstoves have emitted higher BC concentrations than traditional stoves, even if they reduced PM<sub>2.5</sub> mass (26, 27). Not all mitigation options that reduce PM emissions will also reduce total climate forcing or, potentially, benefit health. The inclusion of BC as an outdoor air quality indicator has been proposed (18,



**Fig. 2.** Associations of personal exposure to  $PM_{2.5}$  mass and BC on (A) SBP and DBP (B) using one- and two-pollutant mixed-effects regression models, by age.  $\Delta SBP$  or  $\Delta DBP$  represent the difference in SBP or DBP (with 95% CIs) associated with a  $1\text{-ln}(\mu\text{g}/\text{m}^3)$  increase in pollutant exposure.

28) but not adopted, and current guidelines for evaluating biomass cookstoves focus on  $PM_{2.5}$  and carbon monoxide (29).

Second, BC could be an important exposure assessment tool for future health studies. The larger magnitude of blood pressure response and extension of the health impact to younger women strengthens the importance of our initial finding on  $PM_{2.5}$  exposure and blood pressure. Our results support a recent meta-analysis of studies in US and European cities showing that ambient BC concentrations were more strongly associated with cardiovascular mortality and hospital admissions than PM mass (18). If BC is more strongly linked with health than PM, its measurement will facilitate smaller sample sizes and more accurate estimate health impacts of air pollution, and of interventions and policies. There are several methods to measure BC, including simple and low-cost optical assessment on existing PM filter samples or real-time measurement using a new generation of lightweight personal samplers. The inclusion of BC in studies already measuring  $PM_{2.5}$  requires less additional effort and resources compared with other combustion markers like WSOM or organic hopanes.

Finally, we found an indication that the cardiovascular effect of BC from biomass smoke may be stronger if there is coexposure to motor vehicle emissions. Our results demonstrate that the blood pressure effect of BC observed in the United States and Europe (30, 31) is not limited to high-income countries where BC is primarily from motor vehicles, although residential biomass combustion contributes to winter ambient air pollution in northern climates (32–34). The stronger health effect of BC from roadway exposure or combined roadway–biomass exposure may also be an important environmental risk factor for cardiovascular diseases in developing countries like China where the number of motor vehicles is rapidly increasing and household use of biomass and coal fuels persists (5, 35).

There are several possible reasons for the stronger effect of BC on blood pressure. One is that BC more closely identifies PM from combustion sources than heterogeneous PM mass does, which comprises particles from all sources, or WSOM, which is

both emitted as a primary pollutant and formed as a secondary aerosol. Toxicological studies indicate that PM from incomplete combustion may be more toxic in macrophage and fibroblast cell lines than PM from more complete combustion (36, 37). It is also possible that other components of combustion-related PM contribute to the observed health impacts, with BC acting as a surrogate for their levels. In vitro studies indicate possible toxicity of certain organic constituents in PM from biomass combustion and suggest that BC may be a carrier of these compounds for uptake into macrophages and epithelial cells (36, 38). BC may be operating as an indicator for a larger category of primary combustion particles with varying toxicity to humans, which, in addition to BC, could include metals or polycyclic aromatic hydrocarbons, any of which could act individually or in combination to increase blood pressure (39). Although we cannot determine the single or combination of PM components responsible for the stronger BC effect in our study, our results suggest that a reduction in PM exposure containing BC and other combustion-related particles for which BC is an indicator should lead to a reduction in the adverse health and climate impacts of air pollution.

Our study is limited by its cross-sectional design. However, cooking with biomass is a long-term behavior and all residents have lived in their current homes throughout their adult lifetime. Thus, 24-h PM exposure is a measure that is typical except for seasonal and day-to-day variability; we found little variability in the relative composition of PM exposure by season or daily patterns for women in our study. We considered that factors associated with both blood pressure and highway proximity (e.g., excessive noise or stress from living near the highway) might explain the stronger effect of BC among women closer to the highway. However, highway proximity did not affect the relationship between  $PM_{2.5}$  mass and blood pressure, suggesting that distance from the highway is not a proxy for these potentially confounding variables given that  $PM_{2.5}$  exposure is also associated with both motor vehicle emissions and blood pressure.

Our results are consistent with several intervention studies in Latin America that found a decrease in SBP ( $\sim 3\text{--}6$  mmHg) in older women who switched from a traditional open fire cookstove to a less-polluting chimney stove (40, 41). Although blood pressure is an important risk factor for cardiovascular diseases and overall global burden of disease (1), further research should assess the associations of BC with other health outcomes, including those in vulnerable populations like children. Our study is



**Fig. 3.** Associations of personal black carbon exposure and blood pressure (in mm Hg), by village. Estimates and 95% confidence intervals are the changes in SBP and DBP (in mm Hg) associated with a  $1\text{-ln}(\mu\text{g}/\text{m}^3)$  increase in black carbon exposure, by village. The six villages are outlined in red or blue and the yellow line denotes the Yunnan-Tibet highway. DBP, diastolic blood pressure; SBP, systolic blood pressure.

also limited by proxy measurements, using distance, of motor vehicle exposure in the full sample. Distance from the highway has nonetheless been used as an indicator of roadway air pollution exposure in developed countries (42, 43). Further, women living nearest to the highway had significantly higher exposure to hopanes, direct markers of motor vehicle emissions, than women farthest from the highway in our subsample tracer analysis.

## Conclusion

Our results show that the effect of BC exposure on blood pressure is two or more times larger than that of PM<sub>2.5</sub> mass and WSOM among rural Chinese women using biomass fuels. We also found evidence that BC from biomass smoke is associated with higher blood pressure in the presence of motor vehicle emissions as a coexposure. Our findings suggest that BC has direct relevance as an important environmental risk factor for cardiovascular diseases and support the use of BC as a pollution indicator in future health studies and in the evaluation of air pollution mitigation programs. More broadly, our results may be useful in forming policy aimed at reducing air pollution and improving public health in China and other developing countries. China recently committed to spending US\$275 billion over the next 5 y to reduce air pollution (44), but targets for new vehicle emission standards are absent from recently announced mitigation plans (45). In addition, China's current air pollution targets and programs focus on PM reductions. The BC reduction achieved with any mitigation strategy is not always proportional to the reduction in PM mass, and our results show that BC may be more strongly associated with health outcomes in addition to warming the climate. As motorized transport and subsequent traffic emissions increase throughout China, air pollution policies and mitigation efforts that focus on BC control might have the largest benefits for climate and human health.

## Materials and Methods

We recruited 280 women  $\geq 25$  y old between December 2008 and August 2009. None were previous or current tobacco smokers. Families in this region had similar diets, lifestyles, and socioeconomic backgrounds. The Institutional Review Boards at the University of Wisconsin–Madison and Yunnan Provincial Health Bureau approved this research protocol and obtained informed oral consent was obtained from all participants.

**Personal Air Pollution Exposure Measurements.** We measured the participants' 24-h exposure to PM<sub>2.5</sub> on 1–3 consecutive sampling days in winter and summer. Participants wore a waistpack holding lightweight air samplers that collected PM<sub>2.5</sub> on Teflon filters. They were instructed to perform routine daily activities while wearing the waistpack, but could place it within 1 m while sitting or sleeping and within 2 m while bathing. Field staff monitored compliance through home visits.

PM<sub>2.5</sub> mass was gravimetrically estimated on all exposure samples and blanks using a high-precision microbalance in a temperature and humidity controlled room. They were then analyzed for BC, WSOC, and mobile source tracers at the University of the Chinese Academy of Sciences. We estimated BC concentrations using reflectance analysis with an Optical Transmissometer Data Acquisition System (Model OT21; Magee Scientific) (46).

We used the nonpurgeable organic carbon method described in Timonen et al. (47) to estimate WSOC concentrations. Briefly, filter sections were extracted with high purity water and analyzed with an organic carbon analyzer (Shimadzu Corp.). The resulting WSOC exposures were multiplied by 2.0 to yield WSOM from biomass burning (48, 49). For exposure samples from women in the village either nearest to ( $n = 32$  women) or farthest from the highway ( $n = 53$  women), we analyzed the remaining filter sections for hopanes, nonpolar organic tracers of motor vehicle exhaust, using the extraction-derivatization method with GC/MS described by Zhang et al. (50).

Details about these PM components, their measurement, and related quality assurance and control practices are described in Baumgartner et al. (12) and discussed in *SI Text*.

**Physiological and Health Parameters.** Initial questionnaires evaluated household demographics, socioeconomic status, secondhand-smoking status, and medical history. We conducted in-home blood pressure measurements using an automated device and following standard recommendations (51). We recorded the time of measurement, air temperature, and any caffeine consumption in the previous hour. We measured each participant's height (centimeters), weight (kilograms), waist circumference (centimeters), and salt intake from cooked foods and used a pedometer to assess 24-h physical activity. Details on measurement of blood pressure and the other health and sociodemographic factors related to blood pressure are described elsewhere (11) and provided in *Tables S1* and *S2*.

The location of each participant's home was georeferenced using aerial photographs of the study villages from Google Earth (52). We calculated the shortest distance between each participant's home and the closest highway segment to determine the distance from the highway. Smaller roads within villages are narrow and mainly used for walking.

**Analysis.** We estimated the geometric means and ranges of exposure to PM<sub>2.5</sub> mass, BC, WSOM, and hopanes by season and age group. Spearman correlation analysis was used to assess collinearity between nonnormally distributed pollution components.

We analyzed the differences in blood pressure associated with 1- $\ln(\mu\text{g}/\text{m}^3)$  unit increases in pollution exposure using one- and two-pollutant mixed-effects models to determine if observed associations in one-pollutant models were robust to the inclusion of a second pollutant. We only included the first day of pollution exposure in each season because blood pressure was not measured after subsequent second and third days of PM exposure assessment. The two-pollutant models may also help distinguish between the blood pressure effects of combustion vs. noncombustion pollution (BC vs. PM mass), biomass vs. other combustion pollution (WSOM vs. BC), and biomass vs. other nonbiomass pollution (WSOM vs. PM mass).

We used multivariate regression models from our previous study on PM<sub>2.5</sub> exposure and blood pressure (11) so that any differences in the results could be unambiguously attributed to the difference in pollution variables. The following variables were included in all regression models: age, waist circumference, physical activity (daily number of steps), socioeconomic status, daily salt intake, day of the week and time of day of blood pressure measurement, and ambient air temperature. Passive tobacco smoking, education, caffeine intake, and self-reported health were neither associated with blood pressure at  $P < 0.10$  nor did they change the effect of pollution exposure on blood pressure at  $\geq 10\%$ , and were therefore excluded from the final models.

In a second analysis, we allowed the effect of pollution exposure on blood pressure to vary by village using the following model:

$$y_{ifkj} = \mu + \beta_{ifkj} + \zeta_k(\beta_{kj}) + \gamma X_{ifkj} + \eta Z_{ifk} + a_{ifk} + h_{fk} + v_k + e_{ifkj},$$

where  $y_{ifkj}$  is SBP or DBP for individual  $i$  in household  $f$  in village  $k$  at time  $j$ ,  $\beta_{ifkj}$  is an individual's exposure,  $\zeta_k(\beta_{kj})$  is the random slope at the village level,  $X_{ifkj}$  are other individual-level covariates that vary seasonally (e.g., ambient air temperature);  $Z_{ifk}$  are covariates (e.g., socioeconomic status);  $a_{ifk}$  is a random intercept;  $h_{fk}$  and  $v_k$  represent the random effects to account for correlation at the household and village levels, respectively; and  $e_{ifkj}$  is the residual.

For sensitivity analyses, we used a scale of the IQR on log-transformed exposures to facilitate comparison between pollutants with different concentration ranges. To assess the validity of the distance from the highway as a proxy for exposure to motor vehicle emissions, we conducted two-sample  $t$  tests on geometric mean hopane exposure among women living in the villages nearest to and farthest from the highway. We also conducted a separate analysis excluding women who reported taking antihypertensive medication, all of whom were  $>50$  y old.

All statistical analyses were performed in STATA 11 (StataCorp LP).

**ACKNOWLEDGMENTS.** We thank Arden Pope, Brian Robinson, and Gerard Hoek for valuable comments on early results; our field staff in Yunnan for their hard work; and the Lashihai residents for allowing us into their villages. We are grateful for the support of University of the Chinese Academy of Sciences Hundred Talents of the Chinese Academy of Sciences Grant Y12901FEA2 (to Y.Z.) and the Initiative for Renewable Energy and the Environment at the University of Minnesota's Institute on the Environment (J.B.).

- Lim SS, et al. (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859):2224–2260.
- Hansen J, et al. (2005) Efficacy of climate forcings. *J Geophys Res* 110:D18104.

- Schneider K, Turner JL, Jaffe A, Ivanova N (2010) Choke point China: Confronting water scarcity and energy demand in the world's largest country. *Vt J Envtl Law* 12(3):713–733.
- Huo H, Wang M (2012) Modeling future vehicle sales and stock in China. *Energy Policy* 43:17–29.

5. Bonjour S, et al. (2013) Solid fuel use for household cooking: Country and regional estimates for 1980–2010. *Environ Health Perspect* 121(7):784–790.
6. Zheng M, et al. (2005) Seasonal trends in PM<sub>2.5</sub> source contributions in Beijing, China. *Atmos Environ* 39(22):3967–3976.
7. Pope CA III, et al. (2002) Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 287(9):1132–1141.
8. Bond T, et al. (2013) Bounding the role of black carbon in the climate system: A scientific assessment. *J Geophys Res* 118(11):5380–5552.
9. Saxena P, Hildemann LM, McMurry PH, Seinfeld JH (1995) Organics alter hygroscopic behavior of atmospheric particles. *J Geophys Res* 100(D9):18755–18770.
10. Bell ML; Health Effects Institute Health Review Committee (2012) Assessment of the health impacts of particulate matter characteristics. *Res Rep Health Eff Inst* (161):5–38.
11. Baumgartner J, et al. (2011) Indoor air pollution and blood pressure in adult women living in rural China. *Environ Health Perspect* 119(10):1390–1395.
12. Baumgartner J, et al. (2011) Patterns and predictors of personal exposure to indoor air pollution from biomass combustion among women and children in rural China. *Indoor Air* 21(6):479–488.
13. Apte JS, et al. (2011) Concentrations of fine, ultrafine, and black carbon particles in auto-rickshaws in New Delhi, India. *Atmos Environ* 45(26):4470–4480.
14. Salcedo D, et al. (2006) Characterization of ambient aerosols in Mexico City during the MCMA-2003 campaign with aerosol mass spectrometry: Results from the CENICA supersite. *Atmos Chem Phys* 6(4):925–946.
15. Westerdaal D, Wang X, Pan X, Zhang KM (2009) Characterization of on-road vehicle emission factors and microenvironmental air quality in Beijing, China. *Atmos Environ* 43(3):697–705.
16. de Miranda RM, et al. (2012) Urban air pollution: A representative survey of PM<sub>2.5</sub> mass concentrations in six Brazilian cities. *Air Quality Atmosphere & Health* 5(1):63–77.
17. Graham B, et al. (2002) Water-soluble organic compounds in biomass burning aerosols over Amazonia 1. Characterization by NMR and GC-MS. *J Geophys Res* 107(D20):LBA 14-11–LBA 14-16.
18. Janssen NA, et al. (2011) Black carbon as an additional indicator of the adverse health effects of airborne particles compared with PM<sub>10</sub> and PM<sub>2.5</sub>. *Environ Health Perspect* 119(12):1691–1699.
19. Geng F, et al. (2013) Differentiating the associations of black carbon and fine particle with daily mortality in a Chinese city. *Environ Res* 120:27–32.
20. Sheesley RJ, et al. (2009) Tracking personal exposure to particulate diesel exhaust in a diesel freight terminal using organic tracer analysis. *J Expo Sci Environ Epidemiol* 19(2):172–186.
21. Pope CA III, et al. (2009) Cardiovascular mortality and exposure to airborne fine particulate matter and cigarette smoke: Shape of the exposure-response relationship. *Circulation* 120(11):941–948.
22. Burnett R, et al. (2014) An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ Health Perspect* 122(4):397–403.
23. He FJ, Li J, MacGregor GA (2013) Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ* 346.
24. Law MR, Wald NJ, Morris JK, Jordan RE (2003) Value of low dose combination treatment with blood pressure lowering drugs: Analysis of 354 randomised trials. *BMJ* 326(7404):1427.
25. Ayala A, Brauer M, Mauderly JL, Samet JM (2012) Air pollutants and sources associated with health effects. *Air Quality Atmosphere & Health* 5(2):151–167.
26. Just B, Rogak S, Kandlikar M (2013) Characterization of ultrafine particulate matter from traditional and improved biomass cookstoves. *Environ Sci Technol* 47(7):3506–3512.
27. Johnson M, et al. (2011) *In-Home Emissions of Greenhouse Pollutants from Rocket and Traditional Biomass Cooking Stoves in Uganda* (US Agency for International Development, Washington).
28. Grahame TJ, Schlesinger RB (2010) Cardiovascular health and particulate vehicular emissions: A critical evaluation of the evidence. *Air Quality Atmosphere & Health* 3(1):3–27.
29. International Standards Organization (2012) Guidelines for evaluating cookstove performance. *International Workshop Agreement 11* (International Organization for Standardization, Geneva).
30. Mordukhovich I, et al. (2009) Black carbon exposure, oxidative stress genes, and blood pressure in a repeated-measures study. *Environ Health Perspect* 117(11):1767–1772.
31. Wilker EH, et al. (2010) Black carbon exposures, blood pressure, and interactions with single nucleotide polymorphisms in MicroRNA processing genes. *Environ Health Perspect* 118(7):943–948.
32. Puxbaum H, et al. (2007) Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background. *J Geophys Res* 112(D23).
33. Schauer JJ, Cass GR (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environ Sci Technol* 34(9):1821–1832.
34. Cornell AG, et al. (2012) Domestic airborne black carbon and exhaled nitric oxide in children in NYC. *J Expo Sci Environ Epidemiol* 22(3):258–266.
35. Ng W-S, Schipper L, Chen Y (2010) China motorization trends: New directions for crowded cities. *Journal of Transport and Land Use* 3(3):5–25.
36. Jalava PI, et al. (2010) Effect of combustion condition on cytotoxic and inflammatory activity of residential wood combustion particles. *Atmos Environ* 44(13):1691–1698.
37. Tapanainen M, et al. (2011) In vitro immunotoxic and genotoxic activities of particles emitted from two different small-scale wood combustion appliances. *Atmos Environ* 45(40):7546–7554.
38. Bolling AK, et al. (2012) Wood smoke particles from different combustion phases induce similar pro-inflammatory effects in a co-culture of monocyte and pneumocyte cell lines. *Part Fibre Toxicol* 9(45):45.
39. Jacobs L, et al. (2012) Acute changes in pulse pressure in relation to constituents of particulate air pollution in elderly persons. *Environ Res* 117:60–67.
40. Clark ML, et al. (2013) Impact of a cleaner-burning cookstove intervention on blood pressure in Nicaraguan women. *Indoor Air* 23(2):105–114.
41. McCracken JP, Smith KR, Díaz A, Mittleman MA, Schwartz J (2007) Chimney stove intervention to reduce long-term wood smoke exposure lowers blood pressure among Guatemalan women. *Environ Health Perspect* 115(7):996–1001.
42. Zhu Y, Hinds WC, Shen S, Sioutas C (2004) Seasonal trends of concentration and size distribution of ultrafine particles near major highways in Los Angeles Special Issue of Aerosol Science and Technology on Findings from the Fine Particulate Matter Supersites program. *Aerosol Sci Technol* 38(51):5–13.
43. Spira-Cohen A, Chen LC, Kendall M, Sheesley R, Thurston GD (2010) Personal exposures to traffic-related particle pollution among children with asthma in the South Bronx, NY. *J Expo Sci Environ Epidemiol* 20(5):446–456.
44. Anonymous (August 10, 2013) China and the environment: The East is grey. *The Economist*. Available at [www.economist.com/news/briefing/21583245-china-worlds-worst-polluter-largest-investor-green-energy-its-rise-will-have](http://www.economist.com/news/briefing/21583245-china-worlds-worst-polluter-largest-investor-green-energy-its-rise-will-have). Accessed August 18, 2014.
45. Wong E (September 13, 2013) China's plan to curb air pollution sets limits on coal use and vehicles. *The New York Times*. Available at [www.nytimes.com/2013/09/13/world/asia/china-releases-plan-to-reduce-air-pollution.html?\\_r=0](http://www.nytimes.com/2013/09/13/world/asia/china-releases-plan-to-reduce-air-pollution.html?_r=0). Accessed August 18, 2014.
46. Ahmed T, et al. (2009) Measurement of black carbon (BC) by an optical method and a thermal-optical method: Intercomparison for four sites. *Atmos Environ* 43(40):6305–6311.
47. Timonen HJ, Saarikoski SK, Aurela MA, Saarnio KM, Hillamo REJ (2008) Water-soluble organic carbon in urban aerosol: Concentrations, size distributions and contribution to particulate matter. *Boreal Environ Res* 13(4):335–346.
48. Turpin BJ, Lim H-J (2001) Species contributions to PM<sub>2.5</sub> mass concentrations: Revisiting common assumptions for estimating organic mass. *Aerosol Sci Technol* 35(1):602–610.
49. Bae M-S, Schauer JJ, Turner JR (2006) Estimation of the monthly average ratios of organic mass to organic carbon for fine particulate matter at an urban site. *Aerosol Sci Technol* 40(12):1123–1139.
50. Zhang YX, et al. (2009) Harmonizing molecular marker analyses of organic aerosols. *Aerosol Sci Technol* 43(4):275–283.
51. Pickering TG, et al. (2005) Recommendations for blood pressure measurement in humans and experimental animals: Part 1: Blood pressure measurement in humans: A statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation* 111(5):697–716.
52. Google Earth 7 (February 28, 2010) Digital Globe 2013. Available at [www.earth.google.com](http://www.earth.google.com). Accessed May 13, 2013. [Yunnan, China; 26°51'20.41"N, 100°09'02.79"E; eye altitude 24,161 ft.].