

Identification of Isolated *Salmonella enterica* Serotype *gallinarum* Biotype *Pullorum* and *Gallinarum* by PCR-RFLP

Narges Cheraghchi¹; Pejvak Khaki²; Soheila Moradi Bidhendi^{2,*}; Azar Sabokbar¹

¹Department of Microbiology, College of Basic Sciences, Karaj Branch, Islamic Azad University, Karaj, IR Iran

²Department of Microbiology, Razi Vaccine and Serum Research Institute, Karaj, IR Iran

*Corresponding author: Soheila Moradi Bidhendi, Department of Microbiology, Razi Vaccine and Serum Research Institute, P.C. 3197619751, Karaj, IR Iran. Tel: +98-2634570038/+98-9123182404, Fax: +98-2634552194, E-mail: s.bidhendi@rvsri.ac.ir

Received: April 05, 2014; Revised: May 31, 2014; Accepted: June 09, 2014

Background: *Salmonella* spp. is the major bacterial pathogen in poultry and is responsible for significant economic losses of the poultry industry in many parts of the world. Among *Salmonella* spp., *Salmonella gallinarum* and *Salmonella pullorum* are the most common causative agents of chicken salmonellosis resulting in high mortality and morbidity.

Objectives: The aim of this study was to identify *S. gallinarum* and *S. pullorum* by using the polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method.

Materials and Methods: In this study, 13 samples of *Salmonella*, isolated from local poultry, were obtained from Razi Type Culture Collection (RTCC). For the PCR-RFLP method based on the *fliC* gene, extracted DNA was used as a template for amplifying of the *fliC* gene (197bp) using specific primers. PCR products were subjected to digestion using *Hin*pII restriction endonuclease.

Results: For the PCR, 197 bp *fliC* fragment was amplified from all 13 isolates. Ten out of 13 were *S. gallinarum* and the other three were *S. pullorum*. As part of the PCR-RFLP, two fragments were obtained (82 bp and 115 bp) for all *S. gallinarum*, whereas no digestion was observed in *S. pullorum*, and 197 bp fragment was seen.

Conclusions: PCR-RFLP with *fliC* gene and *Hin*pII endonuclease were successfully applied to differentiate the two biotypes. The results suggested that this technique could be effective in detecting *S. gallinarum* and *S. pullorum*.

Keywords: *Salmonella gallinarum*; *Salmonella pullorum*; RFLP; *Hin*pII Endonuclease

1. Background

Under the new naming system the *Salmonella enterica* serovar *gallinarum* is divided into biovar *gallinarum* (*S. gallinarum*) and *pullorum* (*S. pullorum*), which are identified to cause fowl typhoid and pullorum disease in poultry, respectively. While fowl typhoid is a disease of mature birds, pullorum causes mortality of embryos and chicks. Infection with these pathogens is responsible for considerable economic losses in poultry production (1, 2). Among diseases of poultry, salmonellosis is of great concern and has been responsible for serious economic losses of poultry producers (3). *S. gallinarum* and *S. pullorum* are non-motile, host adapted avian pathogens belonging to *Salmonella* serogroup D (4, 5). *S. gallinarum* and *S. pullorum* are very similar from the point of view of their antigenic structure; however they are responsible for distinctly different diseases in chicken (2, 6).

Some countries are considered free of *S. gallinarum* and *S. pullorum*; however, infections are still sometimes reported, and are a matter of concern for the poultry industry (4, 6). *Salmonella* control efforts are complicated due to their sporadic and uneven distribution (7). Specific characterization of *Salmonella* isolates is therefore extremely important in order to attribute an isolate to a previously known epidemic outbreak (5). Although

conventional Kaufmann White scheme is still the only reliable method for serotyping of *Salmonella*, it cannot differentiate between closely related biotypes, such as *S. gallinarum* and *S. pullorum* (4, 5). Differentiation basically takes into account their biochemical characteristics. However, Biochemical characteristics take approximately five to seven days and are very time-consuming.

Recently, biochemical methods have been complemented by DNA-based molecular techniques, because of their sensitivity and specificity. Such methods include restriction fragment length polymorphism (RFLP), which is sometimes associated with PCR (PCR-RFLP), ribotyping, pulse field gel electrophoresis (PFGE) and variable number tandem repeat (VNTR) (8-10). PCR-RFLP is considered as a rapid test with good reproducibility for molecular typing in bacterial epidemiological studies (11). Many researchers have focused on flagellin genes for *Salmonella* subtyping because most of them possess the two structural genes (*fliC* and *fliB*) that contain a hypervariable central region and a conserved flanking DNA region.

Flagellin genes encode proteins that are related to the serotyping scheme (12). The hypervariable central region of *Salmonella* flagellin genes makes it possible to differentiate the *Salmonella* isolates by the PCR-RFLP tech-

nique. Many studies used part I of the gene that encodes flagellin (*fliC*) to differentiate serotypes. Most *Salmonella* strains have two structural genes (*fliC* and *fliB*) that encode flagellins. Non-motile strains generally exhibit these structural genes, but are unable to build up a functional flagellum (13, 14). *S. gallinarum* and *S. pullorum* have been reported to possess phase 1 flagellin C gene (*fliC*) (13-15). There are no data from epidemiological studies of *S. gallinarum* and *S. pullorum* isolates based on molecular typing in Iran. Thus these data could be helpful in this regard and also for the rapid detection of these bacteria.

2. Objectives

The aim of the present study was to differentiate *S. gallinarum* and *S. pullorum* isolated in Iran, based on the PCR-RFLP method and the *HinpII* enzyme.

3. Materials and Methods

3.1. Bacterial Strains

S. gallinarum (n = 10) and *S. pullorum* (n = 3) isolates used in this study were from clinical samples of chickens kept at the Microbiology Department of Razi Vaccine and Serum Research Institute of Karaj (RVSRI). *S. gallinarum* (ATCC 9184) and *S. pullorum* (ATCC 9120) were used as positive controls and *S. enteritidis* (ATCC 13076) was used as a negative control. All isolates were cultured on MacConkey agar (Merck, Germany) for 24 hours at 37°C.

3.2. DNA Extraction

Bacterial DNA was prepared as described by Paiva et al. (13) with some modifications. Briefly, for each isolate, a loopful of an overnight pure culture of bacteria was transferred into 1 ml of tris-EDTA (TE) buffer 1x, boiled at 95°C for 10 minutes then centrifuged at 17900 × g (10 minutes, 4°C), followed by addition of 1 µL proteinase K (Fermentase, Inc., the Netherlands) to the supernatant. This mixture was directly used for the PCR.

3.3. Amplification of the *fliC* Gene by PCR

Specific primers, *CTGGTGATGACGGTAATGGT* (*fliC*F: 866-885) and *CAGAAAGTTTCGCACTCTCG* (*fliC* R: 1063-1044), were used for the amplification of flagellin gene phase 1 (*fliC*) (13, 14). A reaction mixture containing, 5 µL of ultra-pure water (Gibco, Germany), 9 µL of Primix (Ampliqon, Denmark), 2 µL of DNA and 1 µL of each primer, was prepared. The thermocycler was programmed with 1 cycle of 94 °C for 5 minutes, 35 three-step cycles; denaturation at 94°C for 1 minute, annealing at 58°C for 1 minute, extension at 72°C for 1 minute, and a final cycle at 72°C for 10 minutes. The PCR products were electrophoresed on 1% agarose gel for 1 hour at 60 V (16). The product size was compared with 100 bp DNA ladder (Fermentas, Inc. the Netherlands) after staining with ethidium bromide.

3.4. PCR-RFLP

The digestion solution was prepared with 10 µL of the PCR product, 2 µL of *HinpII* buffer (10X), 1 µL of *HinpII* enzyme (Fermentas Inc. the Netherlands) and 17 µL of ultra-pure water (Gibco, Germany). After incubation at 37°C for 16 hours, RFLPs were determined by electrophoresis of the digested DNA on 2% agarose gel for 2.5 hours at 60 V (16). Sizes of the products were analyzed and compared with the 100 bp plus DNA ladder (Fermentas Inc. the Netherlands).

4. Results


4.1. Amplification of the *fliC* Gene

In this study, the 197 bp fragment of the *fliC* gene was amplified from 10 *S. gallinarum* and three *S. pullorum* and no variation in gene size was detected according to gel electrophoresis (Figure 1).

4.2. PCR-RFLP Analysis


Ten *S. gallinarum* and three *S. pullorum* were analyzed. Digestion of *S. gallinarum* amplicons with *HinpII* yielded two bands, of 82 and 115 bp, while no change in *S. pullorum* amplicons was observed, since no digestion occurred (Figure 2).

Figure 1. Electrophoresis of *fliC* Gene amplicons From *S. gallinarum* and *S. pullorum* Samples



M: 100 bp marker plus DNA ladder (Fermentas Inc.). SP: lanes 1 to 3; SG: lanes 4 to 12, NC: negative control (*S. enteritidis*).

Figure 2. Electrophoretic Analysis of the *fliC* Gene After Enzymatic Treatment With *HinpII* Restriction Enzyme



M: 100 bp marker plus DNA ladder (Fermentas Inc.); lane 1: *S. pullorum* positive control; lane 2: *S. gallinarum* positive control; lanes 3, 4, 5, 7, 8, 10 and 12: *S. gallinarum* isolates; lanes 6, 9, and 11: *S. pullorum* isolates.

5. Discussion

S. gallinarum and *S. pullorum* have been known as important bacterial pathogens in chicken (2, 6). These serovars cannot be distinguished by conventional serological methods and biochemical tests are currently complemented by molecular techniques based on *fliC* and *fliB* genes that encode flagellins (13). There has not been any research on molecular typing of *S. gallinarum* and *S. pullorum* in Iran.

In the study of Hong et al., the PCR-RFLP flagella typing scheme was successfully applied for serotype identification of 112 *Salmonella* isolates obtained from poultry and poultry environments (17). Also, our methods were successful in differentiating these two biotypes by PCR-RFLP. Kwon et al. worked on 41 *S. pullorum* and 52 *S. gallinarum*. They showed that PCR-RFLP with *Hinp1I* was successful in differentiating the two biotypes. These results suggested that the variable regions of *fliC* could be used as a genetic marker and allow differentiation of these biotypes from each other and PCR-RFLP with *Hinp1I* for these biotypes is a valuable tool for identification of non-motile serotypes of *Salmonella* (14). Our study was done on ten *S. gallinarum* and three *S. pullorum* and the same result was obtained using *Hinp1I* enzyme. In another study, practical application of restriction patterns of *fliC* gene using a mixture of endonucleases (TaqI and ScaI) to differentiate *S. gallinarum* and *S. pullorum* was performed. According to their results this method with the used enzymes was not useful to differentiate *S. gallinarum* from *S. pullorum* (15) but in our study a different enzyme, *Hinp1I*, could differentiate these two biotypes. In the Kisiela et al. investigation, ScaI enzyme digested PCR amplicons of *fimH* gene. They could successfully differentiate *S. gallinarum* from *S. pullorum* (18). The results obtained from our research and theirs showed that *fimH* and *fliC* gene with restriction enzyme ScaI and *Hinp1I* could differentiate *S. gallinarum* from *S. pullorum*.

In another investigation done by Paiva et al. PCR amplicons (197 bp) of 14 *S. pullorum* and 22 *S. gallinarum* that had various results on biochemical tests, were digested with the *Hinp1I* enzyme and the same results were obtained. In our research the bacterial isolates had different results for biochemical tests but the same pattern for PCR-RFLP (13). A study done by Menghistu et al. (8) on *S. gallinarum* showed three patterns in 12 *S. gallinarum* isolates by PCR-RFLP using the restriction enzyme *AluI*. They demonstrated that *AluI* could be used in epidemiological investigations. Our results showed that the *Hinp1I* enzyme could be the same as *AluI* in such investigation to identify *S. gallinarum* from *S. pullorum*. In the present study, we were able to demonstrate that the use of *fliC* gene restriction patterns is a useful method for allowing the differen-

tiation between *S. gallinarum* and *S. pullorum* isolated in Iran; including those with atypical biochemical behavior. Therefore, our results support that this method may be adopted to differentiate *S. gallinarum* from *S. pullorum*.

Several sequences of the gene encoding phase 1 flagellin (*fliC*) are available (19, 20). The distal parts of the *fliC* alleles are conserved regions, making this gene in any serotype suitable for easy amplification, whereas the central region of the *fliC* gene is hyper variable, making it a target for differentiation among *Salmonella* serotypes (14, 15). *S. gallinarum* and *S. pullorum* *fliC* gene represent allelic variants and differ only in two codons, including 316 and 339, which shows that the *Hinp1I* enzyme recognizes one cleavage site in *S. gallinarum* (codon 316), but not in *S. pullorum* (14, 21). In the present study by using the applied technique, we were able to discriminate all the *S. gallinarum* and *S. pullorum* isolates. Since *S. gallinarum* and *S. pullorum* are important in industry, thus the accurate identification of them with molecular technique can be effective in this area. Our literature review in Iran showed that there isn't any publication in this field. Also our results demonstrated that PCR-RFLP with *fliC* gene and *Hinp1I* endonuclease could be effective in detecting *S. gallinarum* and *S. pullorum*. The result indicated that there was no limitation in this technique for differentiation of these biotypes.

Acknowledgements

The authors thank all members of the Microbiology Department of Razi Vaccine and Serum Research Institute (Karaj-Iran) for their assistance with this study.

Funding/Support

This work was fully financed with state funds from Razi Vaccine and Serum Research Institute, under grant number 12-18- 18-8901-89003.

References

1. Barrow PA, Freitas Neto OC. Pullorum disease and fowl typhoid - new thoughts on old diseases: a review. *Avian Pathol.* 2011;**40**(1):1-13.
2. Kang MS, Kwon YK, Jung BY, Kim A, Lee KM, An BK, et al. Differential identification of *Salmonella enterica* subsp. *enterica* serovar *Gallinarum* biovars *Gallinarum* and *Pullorum* based on polymorphic regions of *glgC* and *speC* genes. *Vet Microbiol.* 2011;**147**(1-2):181-5.
3. Lutful Kabir SM. Avian colibacillosis and salmonellosis: a closer look at epidemiology, pathogenesis, diagnosis, control and public health concerns. *Int J Environ Res Public Health.* 2010;**7**(1):89-114.
4. OIE . Fowl typhoid and pullorum disease. *Manual of diagnostic tests and vaccines for terrestrial .* 21th ed; 2012.
5. Gast RK. *Salmonella* infections, in *Diseases of Poultry, Iowa State University, Press, Ames.* 1997:81-121.
6. Shivaprasad HL., Pullorum Disease and Fowl Typhoid, in *Diseases of Poultry. Iowa State Univ.* 2003;**Ames**:568-82.
7. Jones FT. A review of practical *Salmonella* control measures in animal feed. *J Applied Poultry Res.* 2011;**20**(1):102-13.
8. Menghistu HT, Rathore R, Dhama K, Agarwal RK. Isolation, Identification and Polymerase Chain Reaction (PCR) Detection of *Salmonella* species from field materials of poultry origin. *Int J Microbiol Res.* 2011;**2**(2):135-42.
9. Gallegos-Robles MA, Morales-Loredo A, Alvarez-Ojeda G, Vega PA,

- Chew MY, Velarde S, et al. Identification of Salmonella serotypes isolated from cantaloupe and chile pepper production systems in Mexico by PCR-restriction fragment length polymorphism. *J Food Prot.* 2008;**71**(11):2217-22.
10. Ghaderi R, Tadayon K, Avagyan S, Khaki P, Moradi Bidhendi S, Forbes KJ, et al. The population structure of Salmonella enterica Enteritidis in Iran analyzed by multiple-locus variable-number tandem repeat analysis. *Trop Anim Health Prod.* 2013;**45**(4):889-94.
 11. Taddele HM, Rathore R, Dhama K, Agarwal RK. Epidemiological Characterization of Salmonella gallinarum Isolates of Poultry Origin in India, Employing Two PCR Based Typing Methods of RAPD-PCR and PCR-RFLP. *Asian J Anim Vet Adv.* 2011;**6**(11).
 12. Jong HY, Su PT, Sanpong P, Wajjwalku W, Sukpuaram T, Amavisit P. PCR-based restriction fragment length polymorphism for subtyping of Salmonella from chicken isolates. *Kasetsart J.* 2010;**44**:79-83.
 13. Paiva JB, Cavallini JS, Silva MD, Almeida MA, Ângela HL, Berchieri Junior A. Molecular differentiation of Salmonella Gallinarum and Salmonella Pullorum by RFLP of fliC gene from Brazilian isolates. *Brazil J Poult Sci.* 2009;**11**(4):271-5.
 14. Kwon HJ, Park KY, Yoo HS, Park JY, Park YH, Kim SJ. Differentiation of Salmonella enterica serotype gallinarum biotype pullorum from biotype gallinarum by analysis of phase 1 flagellin C gene (fliC). *J Microbiol Methods.* 2000;**40**(1):33-8.
 15. Kilger G, Grimont PA. Differentiation of Salmonella phase 1 flagellar antigen types by restriction of the amplified fliC gene. *J Clin Microbiol.* 1993;**31**(5):1108-10.
 16. Sambrook J, Russell DW. *Molecular cloning: a laboratory manual (3-volume set)*.: Cold spring harbor laboratory press Cold Spring Harbor, New York; 2001.
 17. Hong Y, Liu T, Hofacre C, Maier M, White DG, Ayers S, et al. A restriction fragment length polymorphism-based polymerase chain reaction as an alternative to serotyping for identifying Salmonella serotypes. *Avian Dis.* 2003;**47**(2):387-95.
 18. Kisiela D, Sapeta A, Kuczkowski M, Stefaniak T, Wieliczko A, Ugoriski M. Characterization of FimH adhesins expressed by Salmonella enterica serovar Gallinarum biovars Gallinarum and Pullorum: reconstitution of mannose-binding properties by single amino acid substitution. *Infect Immun.* 2005;**73**(9):6187-90.
 19. Chansiripornchai N, Ramasoota P, Bangtrakulnonth A, Sasipreeyajan J, Svenson SB. Application of randomly amplified polymorphic DNA (RAPD) analysis for typing avian Salmonella enterica subsp. enterica. *FEMS Immunol Med Microbiol.* 2000;**29**(3):221-5.
 20. Li J, Smith NH, Nelson K, Crichton PB, Old DC, Whittam TS, et al. Evolutionary origin and radiation of the avian-adapted non-motile salmonellae. *J Med Microbiol.* 1993;**38**(2):129-39.
 21. Li J, Nelson K, McWhorter AC, Whittam TS, Selander RK. Recombinational basis of serovar diversity in Salmonella enterica. *Proc Natl Acad Sci U S A.* 1994;**91**(7):2552-6.