

ALTERED EXPRESSION OF THE VOLTAGE-GATED CALCIUM CHANNEL SUBUNIT $\alpha_2\delta$ -1: A COMPARISON BETWEEN TWO EXPERIMENTAL MODELS OF EPILEPSY AND A SENSORY NERVE LIGATION MODEL OF NEUROPATHIC PAIN

M. NIETO-ROSTRO,^a G. SANDHU,^a C. S. BAUER,^{a†}
P. JIRUSKA,^{b‡} J. G. R. JEFFERYS^b AND A. C. DOLPHIN^{a*}

^a Department of Neuroscience, Physiology and Pharmacology, University College London, London WC1E 6BT, UK

^b Neuronal Networks Group, School of Clinical and Experimental Medicine, University of Birmingham, Birmingham B15 2TT, UK

Abstract—The auxiliary $\alpha_2\delta$ -1 subunit of voltage-gated calcium channels is up-regulated in dorsal root ganglion neurons following peripheral somatosensory nerve damage, in several animal models of neuropathic pain. The $\alpha_2\delta$ -1 protein has a mainly presynaptic localization, where it is associated with the calcium channels involved in neurotransmitter release. Relevant to the present study, $\alpha_2\delta$ -1 has been shown to be the therapeutic target of the gabapentinoid drugs in their alleviation of neuropathic pain. These drugs are also used in the treatment of certain epilepsies. In this study we therefore examined whether the level or distribution of $\alpha_2\delta$ -1 was altered in the hippocampus following experimental induction of epileptic seizures in rats, using both the kainic acid model of human temporal lobe epilepsy, in which status epilepticus is induced, and the tetanus toxin model in which status epilepticus is not involved. The main finding of this study is that we did not identify somatic overexpression of $\alpha_2\delta$ -1 in hippocampal neurons in either of the epilepsy models, unlike the upregulation of $\alpha_2\delta$ -1 that occurs following peripheral nerve damage to both somatosensory and motor neurons. However, we did observe local reorganization of $\alpha_2\delta$ -1 immunostaining in the hippocampus only in the kainic acid model, where it was associated with areas of neuronal cell loss, as indicated by absence of NeuN immunostaining, dendritic loss, as identified by areas where microtubule-associated protein-2 immunostaining was missing, and reactive gliosis, determined by regions of strong OX42 staining.

This article is part of a Special Issue entitled: “Brain compensation. For good”. Crown Copyright © 2014 Published by Elsevier Ltd. on behalf of IBRO. All rights reserved. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

Key words: calcium channel, dorsal root ganglion (DRG), alpha2delta subunit, epilepsy, neuropathic pain, reactive gliosis.

INTRODUCTION

Voltage-gated calcium (Ca_V) channels consist of three subgroups, the Ca_V1 , 2 and 3 classes (Catterall, 2011). Most of these channels, apart from $\text{Ca}_V1.1$ which is a skeletal muscle channel, are involved in neuronal function, with their most prevalent functions being in excitation–transcription coupling ($\text{Ca}_V1.2$), synaptic transmission (Ca_V2 channels), and regulation of neuronal excitability and pacemaker activity (Ca_V3 channels). Because of their key roles in neuronal function, it is not surprising that a number of different calcium channels have been implicated in the pathogenesis of various forms of epilepsy, in both humans and in animal models. These channels include T-type (Ca_V3) channels (Su et al., 2002; Tringham et al., 2012; Cheong and Shin, 2013), P/Q-type ($\text{Ca}_V2.1$) channels (Imbrici et al., 2004; Rajakulendran et al., 2012), and the auxiliary subunits, $\beta 4$ (Escayg et al., 1998) and $\alpha_2\delta$ -2 (Barclay et al., 2001; Edvardson et al., 2013). Furthermore several calcium channels are either actual or potential targets for therapeutic intervention (Cain and Snutch, 2012; Powell et al., 2013).

The Ca_V auxiliary $\alpha_2\delta$ and β subunits, both of which have four isoforms, are associated with the “high voltage activated” Ca_V1 (L-type) and Ca_V2 classes (N, P/Q and R-type) of calcium channel, but are not thought to be associated with Ca_V3 (T-type) calcium channels. Both auxiliary subunits increase plasma membrane expression of the Ca_V1 and Ca_V2 channels, and influence their biophysical properties (Dolphin, 2012b). Of relevance to the potential pathological roles of $\alpha_2\delta$ subunits, the $\alpha_2\delta$ -1 isoform is up-regulated following peripheral somatosensory nerve damage (for review see Bauer et al., 2010), whereas mutations in $\alpha_2\delta$ -2 have been linked to absence epilepsy (Barclay et al., 2001;

*Corresponding author. Address: Department of Neuroscience, Physiology and Pharmacology, Andrew Huxley Building, University College London, Gower Street, London WC1E 6BT, UK.

E-mail address: a.dolphin@ucl.ac.uk (A. C. Dolphin).

† Current address: Department of Biomedical Science, University of Sheffield, Sheffield S10 2TN, UK.

‡ Current address: Department of Developmental Epileptology, Institute of Physiology, Academy of Sciences of Czech Republic, Prague CZ-14220, Czech Republic.

Abbreviations: Ca_V , voltage-gated calcium; DAPI, 4',6-Diamidino-2-Phenylindole; DRG, dorsal root ganglion; EtOH, ethanol; HIV, human immunodeficiency virus; MAP2, microtubule-associated protein-2; PB, phosphate buffer; PBS, phosphate-buffered saline; PSNL, partial sciatic nerve ligation; SNL, spinal nerve ligation.

Dolphin, 2012a; Edvardson et al., 2013). The $\alpha_2\delta$ proteins have also been reported to fulfill other functions independent of calcium channels (Eroglu et al., 2009; Kurshan et al., 2009), and are likely to interact with other binding partners, including thrombospondins (Eroglu et al., 2009).

Both $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2 represent binding sites for the anti-epileptic $\alpha_2\delta$ ligand drugs gabapentin and pregabalin (Brown et al., 1998; Klugbauer et al., 2003). These drugs are used as adjunct therapy in several forms of epilepsy, particularly drug-resistant partial seizures (Marson et al., 2000; Arroyo et al., 2004; Taylor et al., 2007). They are also widely used in the treatment of neuropathic pain resulting from peripheral nerve damage of various origins, such as trauma, trigeminal neuralgia, diabetes-induced nerve damage, and chronic pain following viral infection, including post-herpetic neuralgia (Moore et al., 2009, 2011). They have also been used for alleviation of chronic pain resulting both from human immunodeficiency virus (HIV) infection and as a side effect of some of the anti-HIV drugs (Schutz and Robinson-Papp, 2013). Chronic neuropathic pain resulting from cancer chemotherapeutic drugs, including paclitaxel and cisplatin, is also treated with gabapentinoid drugs (Fallon, 2013).

The mechanism of action of the gabapentinoid drugs in the treatment of epilepsies remains unclear. In this study we wished to examine whether the level or distribution of $\alpha_2\delta$ -1 was altered following experimental induction of epileptic seizures in rats, since a change in $\alpha_2\delta$ -1 level or distribution might contribute to the anti-epileptic mechanism of action of gabapentinoid drugs, in a similar way to their therapeutic action in neuropathic pain (see Discussion). It was not possible to examine changes in distribution of $\alpha_2\delta$ -2 protein in parallel in this study, as no antibodies suitable for immunohistochemistry are currently available.

We chose to use the rat kainic acid model of human temporal lobe epilepsy, in which spontaneous seizures have been found to occur following a latent period after the initial induction by kainic acid of persistent seizures, known as status epilepticus (Buckmaster and Dudek, 1997). In this model, rats first develop status epilepticus, and then consistently develop spontaneous seizures which exhibit a gradual increase in spontaneous frequency in the subsequent weeks (Dudek and Staley, 2012). This model is relevant because gabapentin is known to be effective against seizures induced by this means (Cilio et al., 2001). For comparison, we also used the tetanus toxin model of temporal lobe epilepsy (Jefferys et al., 1995), in which status epilepticus is not induced (Finnerty and Jefferys, 2002).

EXPERIMENTAL PROCEDURES

Kainic acid treatment

Ten adult male Sprague–Dawley rats weighing approximately 250 g were injected with kainic acid (2.5 mg/kg i.p.) to induce status epilepticus. Injections were repeated once per hour until 5–9 Racine stage III/

IV/V seizures per hour have occurred (Luttjohann et al., 2009). After 40–60 min from the onset of status epilepticus (near-continuous motor epileptic activity), diazepam (10 mg/kg, i.p.) was injected repeatedly until continuous motor activity disappeared. Following status epilepticus animals were housed separately. When seizures stopped, diazepam (2.5 mg/kg, i.p.) was continued every 30 min. Subcutaneous administration of warmed sterile saline was given if the animals appeared lethargic and/or a significant drop in weight occurred. Rats were housed in single cages under standard conditions in a room with controlled temperature ($22 \pm 1^\circ\text{C}$) and 12/12-h light/dark cycle. The animals had *ad libitum* access to food and water. Immediately following the status epilepticus, rats were manually fed, if necessary until adequate recovery, and provided with standard food and also mashed food and apple slices. Control animals were treated with an equivalent volume and number of injections of sterile saline.

Unilateral intrahippocampal injection of tetanus toxin

Four rats were injected with tetanus toxin and four rats with saline as controls. Surgical preparation was performed as previously described (Jiruska et al., 2013), under ketamine/xylazine anesthesia. A small trephine opening was drilled over the right hippocampus at coordinates 4.1 mm caudal to bregma and 3.9 mm laterally (Paxinos and Watson, 2005). Using a Hamilton microsyringe and infusion pump (KD Scientific Inc., Holliston, USA) 1 μl of tetanus toxin (Sigma–Aldrich, Poole, UK) solution was injected into the stratum radiatum of the right hippocampal CA3 area (depth 3.9 mm). The tetanus toxin solution contained 25 ng of tetanus toxin in 1 μl of 0.05 M phosphate-buffered saline (PBS; Sigma–Aldrich, UK) and 2% bovine serum albumin (Sigma–Aldrich, UK). It was injected at 200 nl/min. The microsyringe was left in the hippocampus for 5 min after the injection ended to avoid the solution leaking back through the injection track. Control animals were injected with 1 μl of 0.05 M PBS with 2% bovine serum albumin. Following surgery, the rats were housed in single cages and allowed to recover for 2 days. Subsequently they were monitored for spontaneous seizures in video monitoring units to verify the development of spontaneous and recurrent seizures. Videos were recorded using digital infra-red cameras (Y-cam Solutions Ltd., Richmond, UK). Animals were video-monitored for 4 weeks.

All animal procedures were licensed and performed in strict accordance with the Animal Scientific Procedures Act (1986) of the United Kingdom and with Birmingham University Ethical Review.

Sample preparation and immunohistochemistry

Rats were deeply anesthetized with an intraperitoneal injection of (600 mg/kg) pentobarbitone (Euthatal, Merial Animal Health, Harlow, UK), perfused transcardially with saline containing heparin, followed by perfusion with 4% paraformaldehyde in 0.1 M phosphate buffer (PB, pH 7.4). Brains were dissected and the tissue was

post-fixed for 1.5–2 h, washed with PB, cryoprotected by incubation in PB with 15% sucrose, and finally frozen before embedding in optical cutting temperature compound (OCT) and sectioning with a cryostat. Serial coronal sections of 25 μm of the brain region including the hippocampus were collected and placed sequentially on a series of six slides, with 4 sections/slide; the distance between each section and the next on any slide was therefore 150 μm . A total of at least eight such series were collected per animal.

For Cresyl Violet staining, the first slide of each series was consecutively immersed for 5 min in PBS, 50%, and 75% ethanol (EtOH) and then stained in 0.1% Cresyl Violet (Sigma) for 15 min; after washing in H_2O , the slides were briefly immersed in 75% EtOH, 0.3% acetic acid, dehydrated, cleared in Histoclear for 5 min and mounted in DPX, neutral mounting medium (Sigma–Aldrich).

For immunofluorescence labeling to detect $\alpha_2\delta$ -1, sections underwent heat-induced antigen retrieval (10 mM citrate buffer, pH 6.0, 0.05% Tween 20, 98 °C for 10 min) prior to blocking with 10% goat serum in the presence of 0.1% Triton X-100 in PBS for 1 h. Sections were then incubated with the mouse monoclonal anti- α_2 -1 antibody (Sigma, 1:100) in 50% blocking buffer for

2 or 3 days at 4 °C. These sections were also stained for microtubule-associated protein-2 (chicken Ab against MAP2, EnCor Biotechnology, Gainesville, FL, 1:1000). After extensive washing with PBS containing 0.1% Triton X-100, sections were incubated with biotinylated goat anti-mouse IgG (1:500) overnight at 4 °C and Streptavidin–Alexa Fluor 488 (1:500) and goat anti-chicken ab coupled to Alexa Fluor 647 (1:500) overnight at 4 °C. Samples processed for NeuN immunoreactivity (mouse monoclonal NeuN Ab, Millipore, 1:500) and OX42 immunoreactivity (mouse monoclonal OX42 Ab, Abcam, 1:200) were treated in the same way except that they did not require antigen retrieval. Samples were then washed and all were stained with 4',6-Diamidino-2-Phenylindole (DAPI), before mounting in VectaShield (Vector Laboratories, Burlingame, CA, USA).

In the kainic acid study, experiments were performed on a total of 16 rats (10 kainate-treated and six saline-treated, Table 1), of which all were analyzed for histological abnormalities in the hippocampus by staining the first slide of each series with Cresyl Violet. This staining was therefore performed on at least 8 slides per animal. Slides from 10 kainate-treated rats and four saline-treated were stained for $\alpha_2\delta$ -1 and MAP2 immunoreactivity, selecting one slide from each

Table 1. Kainic acid study: summary of animals used and histological analysis

ID	Treatment	Status Epilepticus	Hippocampal damage (Cresyl Violet)	Microglia activation (OX42)	Hippocampal neuronal cell loss (NeuN)	Disruption of $\alpha_2\delta$ -1 staining in areas of neuronal cell loss	Ectopic $\alpha_2\delta$ -1 staining in CA3
1	Kainic acid	Yes	CA1 (bi) + CA3 (bi)	ND	ND	No	No
2	Kainic acid	Yes	CA1 (bi) + CA3 (bi)	Yes	Yes	Yes (CA1)	No
3	Kainic acid	Yes	CA3 (bi)	ND	ND	No	No
4	Kainic acid	No	CA1 (bi) + CA3 (bi)	Yes	Yes	No	++
5	Kainic acid	Yes	CA1 + CA3 (bi)	ND	ND	No	No
9*	Kainic acid	Yes	CA1 + CA3 (bi)	Yes	Yes	Yes (CA1)	++
10	Kainic acid	Yes	CA3 (bi)	ND	ND	No	No
11	Kainic acid	Yes	CA1 + CA3 (bi)	Yes	Yes	Yes (CA1)	No
12	Kainic acid	No	No	No	No	No	No
13	Kainic acid	Yes	CA1 (bi) + CA3 (bi)	Yes	Yes	Yes (CA1)	+
6	Saline	No	No	ND	ND	No	No
7	Saline	No	No	No	No	No	No
8	Saline	No	No	ND	ND	ND	ND
14**	Saline	No	No	ND	ND	ND	ND
15	Saline	No	No	No	No	No	++
16	Saline	No	No	No	No	No	No

Summary of all rats used in the kainic acid study, their treatment and histological analysis. Cresyl Violet staining was performed in the first slide of each series (equivalent to at least 8 slides per animal) for all brains. The presence of neuronal cell death was detected in the pyramidal cell layers of CA1 and/or CA3, as stated. In most cases the damage appeared bilaterally (bi). In the kainic acid-treated rats, sections adjacent to those Cresyl Violet stained sections showing hippocampal damage were also stained for $\alpha_2\delta$ -1, OX42 and NeuN, in a subset of rats. ND = not determined. + : staining present; ++ : strong staining.

* Sample with rostral side of right hippocampus deformed.

** Sample not well-perfused.

Table 2. Tetanus toxin study: summary of animals used and histological analysis

ID	Treatment	Status epilepticus	Spontaneous seizures	Hippocampal damage near injection site (DAPI)	Microglia activation near injection site (OX42)	Microglia activation contralateral to injection site (OX42)	Disruption of $\alpha_2\delta$ -1 staining in areas of neuronal damage
TTX4	Tetanus toxin	No	Yes (early onset)	DG + CA1	CA1	No	No
TTX5	Tetanus toxin	No	Yes (infrequent)	CA1	CA1	No	CA1
TTX7	Tetanus toxin	No	Yes (late onset)	CA1	CA1	+	CA1
TTX8	Tetanus toxin	No	No	DG	CA1 + DG	No	No
Sal1	Saline	No	No	CA1 + DG	CA1	No	No
Sal2	Saline	No	No	CA1	CA1 + DG	No	CA1
Sal3	Saline	No	No	CA1 + DG	CA1 + DG	No	No
Sal4	Saline	No	No	CA1	No	No	No

Summary of rats used in the tetanus toxin study, their treatment and histological analysis.

of 2–3 series adjacent to those showing neuronal cell disruption from the Cresyl Violet staining. Adjacent slides were then processed for NeuN and OX42 staining (one slide each from one representative series per animal; from six kainate-treated and three saline-treated rats). In the tetanus toxin study, four tetanus toxin-injected and four saline-injected rats were processed similarly (Table 2).

Image acquisition and analysis and composite assembly

Cresyl Violet staining was visualized on a Leica MZ7.5 stereomicroscope with a DC300 camera under transmitted light using the Leica IM50 software (Leica, Milton Keynes, UK). To produce the hippocampus composites immunofluorescence images were acquired with Volocity using an inverted fluorescence microscope Axiovert 500 M with a 5 \times objective, converted to JPEGs and mounted using Autostich software (www.autostich.net), or manually if the level of signal was too low for the software. For higher resolution, confocal images were acquired in a Zeiss LSM 500 M with a 10 \times objective (15 μ m optical sections) using Zeiss LSM Image acquisition software.

Spinal nerve ligation (SNL) experiments

These data were obtained during the course of a previous study (Bauer et al., 2009), but the images in Fig. 6 were not included in that study. All the experimental detail is identical to that previously described (Bauer et al., 2009).

RESULTS

Gross morphology and cell loss in the hippocampus following kainic acid treatment

Rats were repeatedly injected with kainic acid (2.5 mg/kg, i.p., $n = 10$), until status epilepticus was observed, which occurred in 8/10 kainic acid-treated rats (Table 1). For the control rats, saline was administered ($n = 6$; Table 1). Rats were then perfused and the brains removed after 5 weeks.

The gross morphology of all the hippocampi was examined with Cresyl Violet staining, in order to determine whether neuronal cell death had occurred. Immunostaining was then performed for $\alpha_2\delta$ -1 and the dendritic marker MAP2 in all 10 kainic acid-treated and four of the saline-treated rats. In six kainate-treated and three saline-treated rats, immunostaining was also performed for the microglial marker OX-42 (Shaw et al., 1990) and the neuronal nuclear marker NeuN.

We were interested in whether kainate-induced seizures and neuronal damage were associated with an altered expression of $\alpha_2\delta$ -1. From an examination of gross morphology using Cresyl Violet staining, we found regions of neuronal cell loss in CA1, CA3 and the hilus of the dentate gyrus in 8/8 kainate-treated animals which had developed status epilepticus (Fig. 1A, arrows; Table 1), and 1/2 kainate-treated rats which did not develop status epilepticus (Table 1), but not in any saline-treated rats (Fig. 1B and Table 1). This neuronal cell loss was confirmed by comparing NeuN immunostaining, which stains neuronal nuclei (Fig. 1C, D), with DAPI staining (Fig. 1E, F), which stains all nuclei.

Expression of OX42 in hippocampus following kainic acid treatment

We found that expression of OX42, a marker of microglial activation (Robinson et al., 1986), was present in the hippocampus of 4/4 kainate-treated rats examined that had experienced status epilepticus (Fig. 1G) and in 1/2 rats that did not develop status epilepticus. Expression of OX42 was seen in both CA1 and CA3 regions (Fig. 1G), as well as elsewhere in the brain (data not shown). The location of neuronal damage, determined by loss of NeuN staining was generally associated with the appearance of OX42 staining (compare Fig. 1C, G). In contrast, no OX42 staining was observed in 3/3 hippocampi examined of saline-treated controls or 1/2 kainate-treated rat that did not develop status epilepticus and did not show any brain damage (Table 1 and Fig. 1H).




Fig. 1. Comparison between hippocampi from kainic acid-treated and control rats. (A, B) Cresyl Violet (cv) staining of a representative hippocampal section from a kainate-treated rat (A, rat #2) and a saline control rat (B, rat #16). Arrows indicate cell loss in A. (C, D) Composite image of NeuN immunostaining of a representative hippocampal section from a kainate-treated rat (C, rat #2) and a saline control rat (D, rat #16). (E, F) Composite image of DAPI staining of a representative hippocampal section from a kainate-treated rat (E, rat #2) and a saline control rat (F, rat #16). (G, H) Composite image of OX42 staining in adjacent hippocampal sections from kainate-treated rat (G, rat #2) or saline control rat (H, #16). Scale bar = 1 mm for all panels.

Expression of $\alpha_2\delta-1$ in hippocampus following kainic acid treatment

The $\alpha_2\delta-1$ subunit has been identified previously as a synaptic protein, present particularly in presynaptic terminals (Taylor and Garrido, 2008; Bauer et al., 2009). However, $\alpha_2\delta-1$ is also associated with calcium channels, including L-type calcium channels, present on dendrites (Schlick et al., 2010). We found $\alpha_2\delta-1$ to be expressed throughout the hippocampus (Fig. 2) although largely absent from the main cell body layers, including the granule cells of the dentate gyrus, and the CA3 and CA1 pyramidal cell layers. The $\alpha_2\delta-1$ protein is particularly strongly expressed in the molecular layer and hilus of the dentate gyrus and in the CA3 stratum lucidum, but also in the stratum oriens and stratum radiatum of the CA1 region (Fig. 2A). This distribution is similar to that found previously using paraffin-embedded rat brain sections and Horseradish Peroxidase staining (Taylor and Garrido, 2008).

Since $\alpha_2\delta-1$ is elevated following peripheral nerve damage in both sensory neurons (Luo et al., 2001; Newton et al., 2001; Bauer et al., 2009) and in motor neurons (Bauer et al., 2009; see also Fig. 6), we examined whether $\alpha_2\delta-1$ expression was altered in the hippocampus of kainate-treated rats. We did not observe any consistent increase of $\alpha_2\delta-1$ staining in any region of the hippocampus of any of the kainate-treated rats; on the contrary there was localized loss of $\alpha_2\delta-1$

immunostaining in the CA1 region, occurring in patches (Fig. 2B, left panels, arrows), at sites correlating with CA1 pyramidal neuronal cell loss, shown by NeuN staining (Fig. 2B, upper right panel, arrow). These patches are also associated with regions of reactive gliosis, indicated by OX42 staining (Fig. 2B, lower right panel, arrow).

At higher power magnification (Fig. 3), the patches associated with loss of $\alpha_2\delta-1$ immunoreactivity are also associated with a reduction in MAP-2 staining, indicative of a loss of pyramidal cell dendrites, and an increase in DAPI staining, likely to be associated with reactive gliosis (starred areas in Fig. 3A). In Fig. 3B, a region from the same hippocampus, 1.2 mm rostral to the damaged area, is shown as a control.

Despite extensive pyramidal neuron loss from the CA3 region, in the tissue from 9/10 kainic acid-treated rats (Figs. 1A, C, E, 2B, 3A and 4A), we found that the strong $\alpha_2\delta-1$ immunostaining in the CA3 stratum lucidum was undiminished in these hippocampi (Fig. 4). In Fig. 4Ai and ii, the CA3 regions from two kainate-treated rats are shown in comparison with the same region from a saline-treated rat (Fig. 4B). The CA3 stratum lucidum showing intense $\alpha_2\delta-1$ staining corresponds to the layer containing mossy fiber terminal synapses. In some sections, expression of $\alpha_2\delta-1$ was also observed to be associated with the stratum oriens of the CA3 region in kainate-treated rats (Fig. 4Ai). This was observed in 3/10 kainate-treated rats (Table 1), but




Fig. 2. $\alpha_2\delta-1$ immunostaining in kainic acid-treated rat hippocampus compared to control hippocampus. (A) Low power composite image of $\alpha_2\delta-1$ immunostaining (upper panel) in saline control (rat #6), compared with DAPI staining (lower panel). Scale bars = 1 mm. (B) Low power composite image of staining in hippocampus of a kainate-treated rat (#11) for $\alpha_2\delta-1$ (left, upper and lower panels are 150 μm apart), NeuN (right upper panel) and OX42 (right lower panel), showing patchy loss of $\alpha_2\delta-1$ immunoreactivity, associated with regions of loss of CA1 pyramidal neurons, and up-regulation of OX42 immunoreactivity (arrows). The sections stained for NeuN and OX42 are consecutive to the upper left section stained for $\alpha_2\delta-1$. Scale bars = 1 mm.

we also observed strong $\alpha_2\delta-1$ staining in the stratum oriens of one of the saline-treated rats (Table 1).

Effect of intrahippocampal tetanus toxin on OX42 and $\alpha_2\delta-1$ staining

For comparison with the kainic acid model, we examined four rats given an intra-hippocampal injection of tetanus toxin, with four rats given saline injection as controls. Three of the tetanus toxin-treated animals developed spontaneous seizures (Table 1). None of the saline-treated rats developed seizures.

Hippocampal sections from all rats were subjected to immunohistochemical analysis of $\alpha_2\delta-1$ and OX42 immunoreactivity, and counterstained with DAPI. The pattern of expression of $\alpha_2\delta-1$ did not show any significant change in any of the samples, except a

localized reduction of immunoreactivity in CA1, coinciding with the injection track in two of the tetanus toxin-treated rats (Fig. 5A, top and middle panels, arrowed), and one saline-treated rat (data not shown). OX42 signal also appeared along the injection track in all the tetanus toxin-treated (Fig. 5A, bottom panel, arrow) and 3/4 saline-treated rats (Fig. 5B left panel). A minor contralateral expression of OX42 in CA1 stratum pyramidale was observed in one of the four tetanus toxin-treated rats that developed late seizures (Fig. 5A, bottom panel, arrowheads). DAPI staining revealed that in all cases, there was some cell loss in the CA1 stratum pyramidale and/or lateral dentate gyrus, associated with the injection track (Fig. 5A middle panel and 5B right panel), and this neuronal cell death coincided with the areas showing OX42 immunoreactivity and reduced $\alpha_2\delta-1$.




Fig. 3. Localized reduction of $\alpha_2\delta$ -1 in stratum radiatum CA1 in field associated with loss of pyramidal neurons in kainic acid-treated rats. (A, B) High power images of $\alpha_2\delta$ -1 (left panel), MAP2 (middle panel), DAPI (right panel) in sections of CA1 and dentate gyrus (DG) associated with neuronal loss. (A) Shows reduced $\alpha_2\delta$ -1 immunoreactivity in localized areas (asterisks) where dendritic branches are absent, shown by loss of MAP2 immunoreactivity, and DAPI staining shows glial proliferation around the site of pyramidal neuron death. (B) Shows a section 1.2 mm rostral to the damaged area, as a control. *Abbreviations:* St.Ori = stratum oriens, St.pyr = stratum pyramidale, St.rad = stratum radiatum, fis = hippocampal fissure, St.ml = molecular layer, St.gl = granule cell layer. Scale bars = 100 μ m.

Up-regulation of $\alpha_2\delta$ -1 following peripheral nerve injury

In contrast to the results described above, following physical insult to peripheral axons of somatosensory nerves, such as SNL (Fig. 6A) and partial sciatic nerve ligation (PSNL), $\alpha_2\delta$ -1 mRNA and protein is rapidly and strongly up-regulated on the injured side in dorsal root ganglion (DRG) neuron somata (Fig. 6B compared to 6C), axons (Fig. 6D compared to 6E) and terminals (Fig. 6F), as previously described (Newton et al., 2001; Li et al., 2004; Bauer et al., 2009; Patel et al., 2013). This finding represents an aspect of the well-studied and poorly understood difference between injury-induced regeneration in peripheral and central neurons (Tedeschi, 2011; Cho et al., 2013). Importantly, we found that $\alpha_2\delta$ -1 was also up-regulated in motor neurons following SNL, likely to be a consequence of their axonal damage caused by the ligation (Bauer et al., 2009) (Fig. 6F, G, H).

DISCUSSION

Altered expression of hippocampal $\alpha_2\delta$ -1 following kainic acid-induced status epilepticus

Many molecular changes have been described in chronic epileptic tissue. Furthermore, several types of channelopathies were identified in neurons from chronic models of acquired epilepsy. They involve alterations of

kinetics of voltage-gated channels, changes in subunit composition, expression of new types of channels or abnormalities of subcellular distribution of specific channels (Poolos and Johnston, 2012). Channelopathies affecting sodium, potassium (Bernard et al., 2004), calcium (Su et al., 2002) or I_h (Shah et al., 2004) channels have been described. These alterations represent molecular, structural and functional changes which occur during epileptogenesis and are responsible for increased cellular excitability, altered functional properties of dendrites and conversion from regular firing neurons to bursting neurons. Furthermore, up-regulation of T-type calcium currents was demonstrated in apical dendrites of CA1 neurons following the pilocarpine status epilepticus (Su et al., 2002).

The present results indicate that damage to the central nervous system by kainic acid treatment, which provokes status epilepticus, followed by delayed development of spontaneous seizures in rats (Buckmaster and Dudek, 1997), does not result in a widespread up-regulation of $\alpha_2\delta$ -1 in the hippocampus, at least not at 5 weeks post-kainic acid. Although no overall up-regulation of mossy fiber $\alpha_2\delta$ -1 was observed at the time of this analysis, a reduction of $\alpha_2\delta$ -1 was observed in regions of CA1 pyramidal cell loss. These results may indicate that in the regions of kainate-induced CA1 pyramidal cell loss, the $\alpha_2\delta$ -1 protein associated with presynaptic Ca_v channels, in the *en passant* synapses from CA3 neurons or Schaffer




Fig. 4. $\alpha_2\delta-1$ staining in the stratum lucidum remains unaffected by neuronal cell loss in CA3 stratum pyramidale. (A, B) High power images of $\alpha_2\delta-1$ (left panel), DAPI (middle panel), and MAP2 (right panel), in two example sections of the CA3 region of hippocampus from two kainate-treated rats (Ai, #4 and Aii, #13) and a saline-treated rat (B, #6). For clarity the stratum lucidum is delimited by a solid line and the stratum pyramidale by a dashed line, or dotted line in areas of neuronal cell loss. *Abbreviations:* St.luc = stratum lucidum, St.pyr = stratum pyramidale, St.ori = stratum oriens. Scale bar = 100 μm .

collaterals onto CA1 pyramidal neurons, is redistributed once the target CA1 cells and their dendrites are lost (see Figs. 2B and 3A). This may therefore represent a local up-regulation of $\alpha_2\delta-1$ at neighboring synapses, although to examine this in the future will require quantitative analysis at the electron microscopic level. This increased presynaptic $\alpha_2\delta-1$ localization could have increased transmitter release as a consequence (Hoppa et al., 2012).

Axonal sprouting of mossy fibers has been found to occur after kainate treatment (McNamara and Routtenberg, 1995). Possible ectopic expression of $\alpha_2\delta-1$ was observed in the stratum oriens of the CA3 region of some kainate-treated rats that could be associated with mossy fiber sprouting (see Fig. 4Ai), but there was no up-regulation of $\alpha_2\delta-1$ protein in cell body regions in the dentate gyrus, unlike the situation following peripheral nerve damage.

OX42 up-regulation provides evidence of microglial activation, and has been shown to be an early response to brain injury, including kainate-induced lesions (Akiyama et al., 1994). In the present study we found widespread bilateral microglial activation following kainic

acid administration, in all the rats examined that showed status epilepticus, and in one of two rats that did not demonstrate status epilepticus.

Comparison of histological changes observed in the kainic acid and tetanus toxin-induced seizure models

The hippocampi of rats injected with tetanus toxin only showed localized lesions that mainly coincided with the injection track, and correspondingly localized development of OX42 staining. However, three out of four of these rats developed spontaneous seizures. When compared with the kainate model and other status epilepticus models, the tetanus toxin model possesses distinct features. Tetanus toxin is known to induce chronic epilepsy with spontaneous and recurrent seizures but without morphological changes, and in particular without initial status epilepticus (Jiruska et al., 2013). It was shown previously that hippocampal sclerosis characterized by major cell loss was present in only 10% of animals and affected mainly the CA1 region (Jefferys, 1992; Vreugdenhil et al., 2002; Jiruska et al., 2010, 2013). Selective loss of somatostatin-positive




Fig. 5. Histological examination of tetanus toxin-injected rat hippocampus, compared to saline-injected control. (A) Low power composite images in the hippocampus of a tetanus toxin-treated rat (TTX7), of $\alpha_2\delta-1$ immunostaining (top panel) showing a localized area of reduced $\alpha_2\delta-1$ staining (arrow), DAPI staining (middle panel, same section as top panel) showing pyramidal cell layer disruption due to injection track (arrow), and OX42 staining (bottom panel) from a section 175 μm away from that shown in upper and middle panels, showing microglial activation around injection site (arrow) and also a small amount along the stratum pyramidale of CA1 of the non-injected side (arrowheads). Scale bars = 1 mm. (B) Low power composite images of OX42 immunostaining and DAPI staining in the hippocampus (left and right panels, respectively) around the injection track of a saline-injected rat (Sal3). Scale bars = 0.5 mm.

interneurons has been demonstrated 8 weeks following tetanus toxin injection (Jefferys et al., 1992). One consistent structural change is axonal sprouting, both in the dentate gyrus (mossy fiber sprouting shown by Timm staining) and in the CA1 region. Based on the results obtained here, we can speculate that changes in $\alpha_2\delta-1$ distribution observed in the kainate model can be attributed to status epilepticus and/or associated neuronal cell loss.

$\alpha_2\delta-1$ up-regulation is a marker of peripheral nerve injury

The $\alpha_2\delta-1$ subunit is expressed in primary afferent DRG neurons, and is particularly strongly expressed in small DRG neurons, which include C-fiber nociceptors (Bauer et al., 2009). *Cacna2d1*, encoding $\alpha_2\delta-1$, is one of many genes whose expression is altered following

experimental peripheral nerve damage. Indeed, $\alpha_2\delta-1$ mRNA and protein shows up-regulation following several different types of damage to peripheral axons of DRG neurons, including ligation, chemotherapy and diabetes-induced neuropathy (Luo et al., 2001, 2002; Newton et al., 2001; Davis-Taber and Scott, 2006; Matsumoto et al., 2006; Xiao et al., 2007; Bauer et al., 2009). Furthermore we have found an alteration in $\alpha_2\delta-1$ splice variant expression in DRGs following peripheral nerve damage (Lana et al., 2013).

The mechanism(s) involved in alteration of gene expression following somatosensory nerve damage are thought to involve propagation of an axonal injury-induced Ca^{2+} wave to the DRG soma, resulting in histone deacetylase 5 export from the nucleus, and activation of gene transcription, a process which has been found to occur in damaged peripheral but not central neurons (Cho et al., 2013).




Fig. 6. $\alpha_2\delta-1$ up-regulation following SNL in rat DRGs and their projections, and in motor neurons. (A) Diagram of the SNL model used to induce mechanical hyperalgesia. Ipsilateral ligation (ligature, left) of the spinal nerve at the level of L5 (and L6, not shown) affects DRG neurons and their afferent projections innervating the dorsal horn of the spinal cord. Black circles indicate the location of DRG cell bodies, small black stars indicate motor neuron cell bodies. (B–E) Representative $\alpha_2\delta-1$ immunofluorescence images of L6 DRG section (B and C, ipsi- and contralateral respectively) and dorsal root section (D and E, ipsi- and contralateral respectively), 14 days after SNL. $\alpha_2\delta-1$ is upregulated in the ipsilateral DRG neurons of all sizes and also in their projections (green: $\alpha_2\delta-1$, blue: DAPI). Scale bars = 20 μm . (F) Montage of $\alpha_2\delta-1$ immunoperoxidase-stained microphotographs covering a complete L5 spinal cord section 14 days after SNL. $\alpha_2\delta-1$ immunoreactivity is increased ipsilaterally, in the dorsal horn (dh), the fasciculus gracilis (fg), and in the cell bodies of motor neurons of the ventral side compared to the contralateral side. White dashed boxes indicate areas shown in G and H. Diamond (\blacklozenge) marks a hole in the tissue introduced to identify the contralateral site. Scale bar = 200 μm . Fi: Insert shows the absence of staining when the primary antibody was omitted. (G, H) Higher power images of the ventral spinal cord, ipsilateral (G) and contralateral (H) to SNL, showing increased $\alpha_2\delta-1$ immunostaining in motor neurons on the ipsilateral side (e.g. white arrows in G). Scale bars = 50 μm . The data for the figure were obtained during the course of our previous study (Bauer et al., 2009), but the images were not included in that study.

Comparison of changes in $\alpha_2\delta-1$ observed in the models of epilepsy and neuropathic pain

A major difference between the peripheral somatosensory nerve damage that results in $\alpha_2\delta-1$ up-regulation and kainate-induced neuronal damage is that in the latter there is loss of CNS neurons, in some cases accompanied by sprouting of neighboring neurons, whereas in the former situation, the DRG somata are not killed by the peripheral nerve insult, and the damaged peripheral axons can regenerate. However, although our results do not demonstrate overt or widespread up-regulation of $\alpha_2\delta-1$ in hippocampal neuronal somata, it is possible that there has been redistribution of $\alpha_2\delta-1$ at synaptic terminals in areas affected by neuronal cell loss.

Mechanism of action of gabapentinoid drugs in the alleviation of neuropathic pain

The high affinity binding site for ^3H -gabapentin in the brain was purified and identified to be $\alpha_2\delta-1$ (Gee et al., 1996; Brown and Gee, 1998), and it was subsequently found to bind to $\alpha_2\delta-2$ with similar affinity (Marais et al., 2001;

Gong et al., 2001). Autoradiographic studies then identified $\alpha_2\delta-1$ as the major binding site for ^3H pregabalin in the rat cerebral cortex, hippocampus and other brain regions, although $\alpha_2\delta-2$ represented the main binding site in the cerebellum and interpeduncular nucleus (Bian et al., 2006).

We have recently shown that the up-regulation of $\alpha_2\delta-1$ is required for the rapid development of mechanical hypersensitivity following PSNL, since this is markedly delayed in $\alpha_2\delta-1$ knockout mice (Patel et al., 2013). It is widely assumed that the up-regulation of $\alpha_2\delta-1$ in damaged somatosensory neurons is related to the efficacy of the gabapentinoid drugs in alleviating neuropathic pain in humans as well as animal models, since the gabapentinoids are ineffective in mice lacking $\alpha_2\delta-1$ (Patel et al., 2013), and in knockin mice in which $\alpha_2\delta-1$ is mutated so that it does not bind gabapentinoid drugs (Field et al., 2006). Nevertheless the mechanism of action of these drugs at the molecular level still remains unclear, as they are generally found to produce little acute inhibition of calcium currents or synaptic transmission (Sutton et al., 2002; Brown and Randall, 2005; Hendrich et al., 2008). However, some studies

have found acute effects of these drugs on synaptic transmission (Uchitel et al., 2010).

We have found that the $\alpha_2\delta$ -subunits enhance plasma membrane expression of calcium channels; although the mechanism still remains unclear, it is thought to involve trafficking of the channels from their site of synthesis to the plasma membrane (Canti et al., 2005; Bauer et al., 2010). It is likely that in neurons $\alpha_2\delta$ subunits have multiple effects on calcium channel distribution, both associated with long-range calcium channel trafficking from their site of synthesis in the soma to their mainly presynaptic localization in nerve terminals (Bauer et al., 2009), and also local effects on calcium channel localization in membrane micro-domains such as the active zone and in lipid rafts (Davies et al., 2006; Hoppa et al., 2012), as well as influencing the recycling of calcium channels to the plasma membrane (Tran-Van-Minh and Dolphin, 2010).

Furthermore, we have found that the gabapentinoid drugs have an inhibitory effect on calcium currents when applied over longer time periods, in cultured cells and neurons (Hendrich et al., 2008; Tran-Van-Minh and Dolphin, 2010), which we infer is by inhibiting the trafficking of the $\alpha_2\delta$ subunits (Hendrich et al., 2008; Bauer et al., 2010; Tran-Van-Minh and Dolphin, 2010). We also observed *in vivo* that there was less up-regulation of $\alpha_2\delta$ -1 in nerve terminal zones, after the induction of somatosensory nerve injury when it was combined with chronic pregabalin treatment (Bauer et al., 2009), which might be an effect on long range axonal trafficking, or on lifetime of the protein and its local recycling at presynaptic terminals. Our evidence currently indicates that this interference by gabapentinoids with the function of $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2 results in a reduction of expression of the entire calcium channel complex at the plasma membrane (Tran-Van-Minh and Dolphin, 2010; Cassidy and Dolphin, 2014). In agreement with this, we have also observed inhibitory effects of chronically applied gabapentinoids on excitatory synaptic transmission from DRG neuron terminals (Hendrich et al., 2012).

Role of $\alpha_2\delta$ subunits and mechanism of action of gabapentinoids in epilepsy

The gabapentinoid drugs have therapeutic efficacy as antiepileptic drugs, although they are generally used in combination therapy (Marson et al., 2000; Glauser et al., 2006). In animal models of seizures, gabapentin has been shown to be effective (Cilio et al., 2001). However, the mechanism of action of the gabapentinoid drugs in epilepsy is poorly understood. Although both gabapentin and pregabalin were first developed to enhance GABA-ergic inhibition in the brain (Taylor et al., 1992, 2007; Silverman, 2008), it is now clear that they do not act by mechanisms involving inhibition of GABA breakdown, or activation of GABA-A or GABA-B receptors (Taylor et al., 2007; Li et al., 2011). Furthermore GABA itself does not bind to the $\alpha_2\delta$ subunits that are now known to be the target for gabapentinoid drugs (Li et al., 2011).

Although it has been found that $\alpha_2\delta$ -1 is the target for the gabapentinoid drugs in the alleviation of experimental neuropathic pain in rodents, (Field et al., 2006; Patel et al., 2013), this is not known for the efficacy of the gabapentinoids in animal models of epilepsy. An *in situ* hybridization study showed that $\alpha_2\delta$ -1 expression was often more associated with excitatory neurons, and $\alpha_2\delta$ -2 with inhibitory neurons (Cole et al., 2005). The $\alpha_2\delta$ -1 protein is strongly expressed in the hippocampus, therefore it is possible that a change in expression of $\alpha_2\delta$ -1 in epileptic foci might influence the effectiveness of the gabapentinoid drugs. In contrast $\alpha_2\delta$ -2 is expressed in a more restricted pattern, for example it is strongly expressed in cerebellar Purkinje neurons (Barclay et al., 2001; Brodbeck et al., 2002). The loss of expression of $\alpha_2\delta$ -2 in *cacna2d2* mutant mouse strains including “Ducky” results in cerebellar ataxia and spike-wave epilepsy, and is associated with severe Purkinje cell dysfunction (Barclay et al., 2001; Brodbeck et al., 2002; Brill et al., 2004; Donato et al., 2006). Furthermore *CACNA2D2* is disrupted in rare recessive human cases of epileptic encephalopathy (Edvardson et al., 2013; Pippucci et al., 2013). Therefore interference with $\alpha_2\delta$ -2 function might be intuitively less likely to be the therapeutic target of the gabapentinoids in epilepsy, compared to disruption of $\alpha_2\delta$ -1 function. However, this would not exclude the possibility that there is localized alteration of $\alpha_2\delta$ -2 expression, which could be a therapeutic target in focal (partial) epilepsy.

Thus it is possible that $\alpha_2\delta$ -2 levels or distribution might be affected in animal models of epilepsy. Unfortunately there are currently no available $\alpha_2\delta$ -2 antibodies that are effective in immunohistochemistry, so at present this cannot be easily tested. However, it would be extremely useful to examine whether gabapentin is effective in epilepsy models, using knockin mice in which either $\alpha_2\delta$ -1 (Field et al., 2006) or $\alpha_2\delta$ -2 is mutated so that it is gabapentin-insensitive (Lotarski et al., 2011).

Non-calcium channel functions of $\alpha_2\delta$ proteins

Recently, $\alpha_2\delta$ -1 has been found to interact with thrombospondins (Eroglu et al., 2009), and this interaction has been shown to be involved in synaptogenesis, a process which has been described as being independent of its function as a calcium channel subunit. Thrombospondins are a ubiquitous family of extracellular matrix proteins, which are secreted by many cell types, including microglia (Chamak et al., 1995; Eroglu et al., 2009). In a number of experimental and human epilepsies there is reactive gliosis (Seifert et al., 2010), microglial activation (Avignone et al., 2008), and axonal sprouting (Tauck and Nadler, 1985; McNamara and Routtenberg, 1995; Vreugdenhil et al., 2002; Sutula and Dudek, 2007). It has also been proposed that gabapentin inhibits the interaction between $\alpha_2\delta$ -1 and thrombospondins, and therefore inhibits synaptogenesis (Eroglu et al., 2009). Although this might be considered a plausible mechanism of action of gabapentinoid drugs in treatment of epilepsies, synaptic remodeling that occurs at epileptic foci (Lew

and Buckmaster, 2011) is likely to have already occurred before the onset of treatment with these drugs. Nevertheless, it is possible that the gabapentinoid drugs may also modify epileptogenesis and decrease the consequences of status epilepticus by reducing cellular damage and seizure frequency (Cilio et al., 2001; Li et al., 2012).

AUTHOR CONTRIBUTIONS

A.C.D. and J.G.R.J. conceived the study. P.J. performed *in vivo* procedures and monitoring. M.N.-R. performed and analyzed all hippocampal histology with the help of G.S. C.S.B. and M.N.-R. performed all DRG and spinal cord histology. A.C.D. and M.N.-R. wrote the paper, with input from all authors.

Acknowledgments—We thank Epilepsy Research UK (Grants P0803 to A.C.D. and J.J.; A0806 to P.J. and P1102 to J.J.), Wellcome Trust (077883/Z/05/Z and 098360/Z/12/Z to A.C.D.) and Czech Ministry of Health grant (Czech Republic, IGA NT 14489-3/2013; P.J.) for providing support for this study.

REFERENCES

- Akiyama H, Tooyama I, Kondo H, Ikeda K, Kimura H, McGeer EG, McGeer PL (1994) Early response of brain resident microglia to kainic acid-induced hippocampal lesions. *Brain Res* 635:257–268.
- Arroyo S, Anhut H, Kugler AR, Lee CM, Knapp LE, Garofalo EA, Messmer S (2004) Pregabalin add-on treatment: a randomized, double-blind, placebo-controlled, dose-response study in adults with partial seizures. *Epilepsia* 45:20–27.
- Avignone E, Ulmann L, Levavasseur F, Rassendren F, Audinat E (2008) Status epilepticus induces a particular microglial activation state characterized by enhanced purinergic signaling. *J Neurosci* 28:9133–9144.
- Barclay J, Balaguero N, Mione M, Ackerman SL, Letts VA, Brodbeck J, Canti C, Meir A, Page KM, Kusumi K, PerezReyes E, Lander ES, Frankel WN, Gardiner RM, Dolphin AC, Rees M (2001) Ducky mouse phenotype of epilepsy and ataxia is associated with mutations in the *Cacna2d2* gene and decreased calcium channel current in cerebellar Purkinje cells. *J Neurosci* 21:6095–6104.
- Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L, Kadurin I, Sri Ranjan Y, Fernandez-Alacid L, Millar NS, Dickenson AH, Lujan R, Dolphin AC (2009) The increased trafficking of the calcium channel subunit $\alpha 2\delta$ -1 to presynaptic terminals in neuropathic pain is inhibited by the $\alpha 2\delta$ ligand pregabalin. *J Neurosci* 29:4076–4088.
- Bauer CS, Tran-Van-Minh A, Kadurin I, Dolphin AC (2010) A new look at calcium channel $\alpha 2\delta$ subunits. *Curr Opin Neurobiol* 20:563–571.
- Bernard C, Anderson A, Becker A, Poolos NP, Beck H, Johnston D (2004) Acquired dendritic channelopathy in temporal lobe epilepsy. *Science* 305:532–535.
- Bian F, Li Z, Offord J, Davis MD, McCormick J, Taylor CP, Walker LC (2006) Calcium channel $\alpha 2\delta$ -1 subunit is the major binding protein for pregabalin in neocortex, hippocampus, amygdala, and spinal cord: an *ex vivo* autoradiographic study in $\alpha 2\delta$ -1 genetically modified mice. *Brain Res* 1075:68–80.
- Brill J, Klocke R, Paul D, Boison D, Gouder N, Klugbauer N, Hofmann F, Becker CM, Becker K (2004) Entla, a novel epileptic and ataxic *Cacna2d2* mutant of the mouse. *J Biol Chem* 279:7322–7330.
- Brodbeck J, Davies A, Courtney J-M, Meir A, Balaguero N, Canti C, Moss FJ, Page KM, Pratt WS, Hunt SP, Barclay J, Rees M, Dolphin AC (2002) The ducky mutation in *Cacna2d2* results in altered Purkinje cell morphology and is associated with the expression of a truncated *a2d-2* protein with abnormal function. *J Biol Chem* 277:7684–7693.
- Brown JP, Dissanayake VU, Briggs AR, Milic MR, Gee NS (1998) Isolation of the [3H]gabapentin-binding protein/ $\alpha 2\delta$ Ca²⁺ channel subunit from porcine brain: development of a radioligand binding assay for $\alpha 2\delta$ subunits using [3H]leucine. *Anal Biochem* 255:236–243.
- Brown JP, Gee NS (1998) Cloning and deletion mutagenesis of the $\alpha 2\delta$ calcium channel subunit from porcine cerebral cortex. *J Biol Chem* 273:25458–25465.
- Brown JT, Randall A (2005) Gabapentin fails to alter P/Q-type Ca²⁺ channel-mediated synaptic transmission in the hippocampus *in vitro*. *Synapse* 55:262–269.
- Buckmaster PS, Dudek FE (1997) Neuron loss, granule cell axon reorganization, and functional changes in the dentate gyrus of epileptic kainate-treated rats. *J Comp Neurol* 385:385–404.
- Cain SM, Snutch TP (2012) Voltage-gated calcium channels in epilepsy. In: Noebels J, Avoli M, Rogawski M, Olsen R, Delgado-Escueta A, editors. *Jasper's basic mechanisms of the epilepsies*. Oxford University Press. p. 66–84.
- Canti C, Nieto-Rostro M, Foucault I, Hebllich F, Wratten J, Richards MW, Hendrich J, Douglas L, Page KM, Davies A, Dolphin AC (2005) The metal-ion-dependent adhesion site in the Von Willebrand factor-A domain of $\alpha 2\delta$ subunits is key to trafficking voltage-gated Ca²⁺ channels. *Proc Natl Acad Sci U S A* 102:11230–11235.
- Cassidy J, Dolphin A (2014) Using exofacially tagged functional Cav2.2 to investigate the modulation of pore subunit trafficking by auxiliary calcium channel subunits. *US Biophysical Society Abstracts* 58:1672-Pos.
- Catterall WA (2011) Voltage-gated calcium channels. *Cold Spring Harb Perspect Biol* 3:a003947.
- Chamak B, Dobbertin A, Mallat M (1995) Immunohistochemical detection of thrombospondin in microglia in the developing rat brain. *Neuroscience* 69:177–187.
- Cheong E, Shin HS (2013) T-type Ca²⁺(+) channels in absence epilepsy. *Biochim Biophys Acta* 1828:1560–1571.
- Cho Y, Sloutsky R, Naegle KM, Cavalli V (2013) Injury-induced HDAC5 nuclear export is essential for axon regeneration. *Cell* 155:894–908.
- Cilio MR, Bolanos AR, Liu Z, Schmid R, Yang Y, Stafstrom CE, Mikati MA, Holmes GL (2001) Anticonvulsant action and long-term effects of gabapentin in the immature brain. *Neuropharmacology* 40:139–147.
- Cole RL, Lechner SM, Williams ME, Prodanovich P, Bleicher L, Varney MA, Gu G (2005) Differential distribution of voltage-gated calcium channel $\alpha 2\delta$ (alpha2delta) subunit mRNA-containing cells in the rat central nervous system and the dorsal root ganglia. *J Comp Neurol* 491:246–269.
- Davies A, Douglas L, Hendrich J, Wratten J, Tran-Van-Minh A, Foucault I, Koch D, Pratt WS, Saibil H, Dolphin AC (2006) The calcium channel $\alpha 2\delta$ -2 subunit partitions with Cav2.1 in lipid rafts in cerebellum: implications for localization and function. *J Neurosci* 26:8748–8757.
- Davis-Taber RA, Scott VE (2006) Transcriptional profiling of dorsal root ganglia in a neuropathic pain model using microarray and laser capture microdissection. *Drug Dev Res* 67:308–330.
- Dolphin AC (2012a) Calcium channel $\alpha 2\delta$ subunits in epilepsy and as targets for antiepileptic drugs. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV, editors. *Jasper's basic mechanisms of the epilepsies*. Oxford University Press. p. 959–973.
- Dolphin AC (2012b) Calcium channel auxiliary $\alpha 2\delta$ and β subunits: trafficking and one step beyond. *Nat Rev Neurosci* 13:542–555.
- Donato R, Page KM, Koch D, Nieto-Rostro M, Foucault I, Davies A, Wilkinson T, Rees M, Edwards FA, Dolphin AC (2006) The ducky2J mutation in *Cacna2d2* results in reduced spontaneous Purkinje cell activity and altered gene expression. *J Neurosci* 26:12576–12586.

- Dudek FE, Staley KJ (2012) The time course and circuit mechanisms of acquired epileptogenesis. In: Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV, editors. *Source Jasper's Basic Mechanisms of the Epilepsies* ([Internet]. 4th ed). Bethesda (MD): National Center for Biotechnology Information (US). p. 405–415.
- Edvardson S, Oz S, Abulhijaa FA, Taher FB, Shaag A, Zenvirt S, Dascal N, Elpeleg O (2013) Early infantile epileptic encephalopathy associated with a high voltage gated calcium channelopathy. *J Med Genet* 50:118–123.
- Eroglu C, Allen NJ, Susman MW, O'Rourke NA, Park CY, Ozkan E, Chakraborty C, Mulinyawe SB, Annis DS, Huberman AD, Green EM, Lawler J, Dolmetsch R, Garcia KC, Smith SJ, Luo ZD, Rosenthal A, Mosher DF, Barres BA (2009) Gabapentin receptor $\alpha 2\delta$ -1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell* 139:380–392.
- Escayg A, Jones JM, Kearney JA, Hitchcock PF, Meisler MH (1998) Calcium channel $\beta 4$ (*CACNB4*): human ortholog of the mouse epilepsy gene lethargic. *Genomics* 50:14–22.
- Fallon MT (2013) Neuropathic pain in cancer. *Br J Anaesth* 111:105–111.
- Field MJ, Cox PJ, Stott E, Melrose H, Offord J, Su TZ, Bramwell S, Corradini L, England S, Winks J, Kinloch RA, Hendrich J, Dolphin AC, Webb T, Williams D (2006) Identification of the $\alpha 2\delta$ -1 subunit of voltage-dependent calcium channels as a novel molecular target for pain mediating the analgesic actions of pregabalin. *Proc Natl Acad Sci U S A* 103:17537–17542.
- Finnerty GT, Jefferys JG (2002) Investigation of the neuronal aggregate generating seizures in the rat tetanus toxin model of epilepsy. *J Neurophysiol* 88:2919–2927.
- Gee NS, Brown JP, Dissanayake VUK, Offord J, Thurlow R, Woodruff GN (1996) The novel anticonvulsant drug, gabapentin (neurontin), binds to the $\alpha 2\delta$ subunit of a calcium channel. *J Biol Chem* 271:5768–5776.
- Glauser T, Ben Menachem E, Bourgeois B, Cnaan A, Chadwick D, Guerreiro C, Kalviainen R, Mattson R, Perucca E, Tomson T (2006) ILAE treatment guidelines: evidence-based analysis of antiepileptic drug efficacy and effectiveness as initial monotherapy for epileptic seizures and syndromes. *Epilepsia* 47:1094–1120.
- Gong HC, Hang J, Kohler W, Li L, Su TZ (2001) Tissue-specific expression and gabapentin-binding properties of calcium channel $\alpha 2\delta$ subunit subtypes. *J Membr Biol* 184:35–43.
- Hendrich J, Tran-Van-Minh A, Hebllich F, Nieto-Rostro M, Watschinger K, Striessnig J, Wratten J, Davies A, Dolphin AC (2008) Pharmacological disruption of calcium channel trafficking by the $\alpha 2\delta$ ligand gabapentin. *Proc Natl Acad Sci U S A* 105:3628–3633.
- Hendrich J, Bauer CS, Dolphin AC (2012) Chronic pregabalin inhibits synaptic transmission between rat dorsal root ganglion and dorsal horn neurons in culture. *Channels (Austin)* 6:124–132.
- Hoppa M, Lana B, Margas W, Dolphin AC, Ryan TA (2012) $\alpha 2\delta$ couples calcium channels to neurotransmitter release sites to control release probability. *Nature* 486:122–125.
- Imbrici P, Jaffe SL, Eunson LH, Davies NP, Herd C, Robertson R, Kullmann DM, Hanna MG (2004) Dysfunction of the brain calcium channel $\text{CaV}2.1$ in absence epilepsy and episodic ataxia. *Brain* 127:2682–2692.
- Jefferys JG (1992) Mechanism of tetanus toxin in neuronal cell death. *Trends Pharmacol Sci* 13:13–14.
- Jefferys JG, Borck C, Mellanby J (1995) Chronic focal epilepsy induced by intracerebral tetanus toxin. *Ital J Neurol Sci* 16: 27–32.
- Jefferys JG, Evans BJ, Hughes SA, Williams SF (1992) Neuropathology of the chronic epileptic syndrome induced by intrahippocampal tetanus toxin in rat: preservation of pyramidal cells and incidence of dark cells. *Neuropathol Appl Neurobiol* 18:53–70.
- Jiruska P, Finnerty GT, Powell AD, Lofti N, Cmejla R, Jefferys JG (2010) Epileptic high-frequency network activity in a model of non-lesional temporal lobe epilepsy. *Brain* 133:1380–1390.
- Jiruska P, Shtaya AB, Bodansky DM, Chang WC, Gray WP, Jefferys JG (2013) Dentate gyrus progenitor cell proliferation after the onset of spontaneous seizures in the tetanus toxin model of temporal lobe epilepsy. *Neurobiol Dis* 54:492–498.
- Klugbauer N, Marais E, Hofmann F (2003) Calcium channel $\alpha 2\delta$ subunits: differential expression, function, and drug binding. *J Bioenerg Biomembr* 35:639–647.
- Kurshan PT, Oztan A, Schwarz TL (2009) Presynaptic $\alpha(2)\delta$ -3 is required for synaptic morphogenesis independent of its $\text{Ca}(2+)$ -channel functions. *Nat Neurosci* 12:1415–1423.
- Lana B, Schlick B, Martin S, Pratt WS, Page KM, Goncalves L, Rahman W, Dickenson AH, Bauer CS, Dolphin AC (2013) Differential up-regulation in DRG neurons of an α delta-1 splice variant with a lower affinity for gabapentin after peripheral sensory nerve injury. *Pain* 155:522–533.
- Lew FH, Buckmaster PS (2011) Is there a critical period for mossy fiber sprouting in a mouse model of temporal lobe epilepsy? *Epilepsia* 52:2326–2332.
- Li CY, Song YH, Higuera ES, Luo ZD (2004) Spinal dorsal horn calcium channel $\alpha 2\delta$ -1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia. *J Neurosci* 24:8494–8499.
- Li H, Graber KD, Jin S, McDonald W, Barres BA, Prince DA (2012) Gabapentin decreases epileptiform discharges in a chronic model of neocortical trauma. *Neurobiol Dis* 48:429–438.
- Li Z, Taylor CP, Weber M, Piechan J, Prior F, Bian F, Cui M, Hoffman D, Donevan S (2011) Pregabalin is a potent and selective ligand for $\alpha(2)\delta$ -1 and $\alpha(2)\delta$ -2 calcium channel subunits. *Eur J Pharmacol* 667:80–90.
- Lotarski SM, Donevan S, El Kattan A, Osgood S, Poe J, Taylor CP, Offord J (2011) Anxiolytic-like activity of pregabalin in the Vogel conflict test in $\alpha 2\delta$ -1 (R217A) and $\alpha 2\delta$ -2 (R279A) mouse mutants. *J Pharmacol Exp Ther* 338:615–621.
- Luo ZD, Calcutt NA, Higuera ES, Valder CR, Song YH, Svensson CI, Myers RR (2002) Injury type-specific calcium channel $\alpha 2\delta$ -1 subunit up-regulation in rat neuropathic pain models correlates with allodynic effects of gabapentin. *J Pharmacol Exp Ther* 303:1199–1205.
- Luo ZD, Chaplan SR, Higuera ES, Sorkin LS, Stauderman KA, Williams ME, Yaksh TL (2001) Upregulation of dorsal root ganglion $\alpha 2\delta$ calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. *J Neurosci* 21:1868–1875.
- Luttjohann A, Fabene PF, van Luijtelaar G (2009) A revised Racine's scale for PTZ-induced seizures in rats. *Physiol Behav* 98:579–586.
- Marais E, Klugbauer N, Hofmann F (2001) Calcium channel $\alpha(2)\delta$ subunits – structure and gabapentin binding. *Mol Pharmacol* 59:1243–1248.
- Marson AG, Kadir ZA, Hutton JL, Chadwick DW (2000) Gabapentin add-on for drug-resistant partial epilepsy. *Cochrane Database Syst Rev*:CD001415.
- Matsumoto M, Inoue M, Hald A, Xie W, Ueda H (2006) Inhibition of paclitaxel-induced A-fiber hypersensitization by gabapentin. *J Pharmacol Exp Ther* 318:735–740.
- McNamara RK, Routtenberg A (1995) NMDA receptor blockade prevents kainate induction of protein F1/GAP-43 mRNA in hippocampal granule cells and subsequent mossy fiber sprouting in the rat. *Brain Res Mol Brain Res* 33:22–28.
- Moore RA, Straube S, Wiffen PJ, Derry S, McQuay HJ (2009) Pregabalin for acute and chronic pain in adults. *Cochrane Database Syst Rev*:CD007076.
- Moore RA, Wiffen PJ, Derry S, McQuay HJ (2011) Gabapentin for chronic neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev*:CD007938.
- Newton RA, Bingham S, Case PC, Sanger GJ, Lawson SN (2001) Dorsal root ganglion neurons show increased expression of the calcium channel $\alpha 2\delta$ -1 subunit following partial sciatic nerve injury. *Brain Res Mol Brain Res* 95:1–8.
- Patel R, Bauer CS, Nieto-Rostro M, Margas W, Ferron L, Chaggar K, Crews K, Ramirez JD, Bennett DL, Schwartz A, Dickenson AH, Dolphin AC (2013) $\alpha 2\delta$ -1 gene deletion affects somatosensory

- neuron function and delays mechanical hypersensitivity in response to peripheral nerve damage. *J Neurosci* 33: 16412–16426.
- Paxinos G, Watson C (2005) *The rat brain in stereotaxic coordinates*. Amsterdam: Elsevier.
- Pippucci T, Parmeggiani A, Palombo F, Maresca A, Angius A, Crisponi L, Cucca F, Liguori R, Valentino ML, Seri M, Carelli V (2013) A novel null homozygous mutation confirms CACNA2D2 as a gene mutated in epileptic encephalopathy. *PLoS ONE* 8:e82154.
- Poolos NP, Johnston D (2012) Dendritic ion channelopathy in acquired epilepsy. *Epilepsia* 53(Suppl 9):32–40.
- Powell KL, Cain SM, Snutch TP, O'Brien TJ (2013) Low threshold T-type calcium channels as targets for novel epilepsy treatments (in press). *Br J Clin Pharmacol*. <http://dx.doi.org/10.1111/bcp.12205>.
- Rajakulendran S, Kaski D, Hanna MG (2012) Neuronal P/Q-type calcium channel dysfunction in inherited disorders of the CNS. *Nat Rev Neurol* 8:86–96.
- Robinson AP, White TM, Mason DW (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. *Immunology* 57:239–247.
- Schlick B, Flucher BE, Obermair GJ (2010) Voltage-activated calcium channel expression profiles in mouse brain and cultured hippocampal neurons. *Neuroscience* 167:786–798.
- Schutz SG, Robinson-Papp J (2013). HIV-related neuropathy: current perspectives. *HIV AIDS (Auckl)* 5:243–251.
- Seifert G, Carmignoto G, Steinhauser C (2010) Astrocyte dysfunction in epilepsy. *Brain Res Rev* 63:212–221.
- Shah MM, Anderson AE, Leung V, Lin X, Johnston D (2004) Seizure-induced plasticity of h channels in entorhinal cortical layer III pyramidal neurons. *Neuron* 44:495–508.
- Shaw JA, Perry VH, Mellanby J (1990) Tetanus toxin-induced seizures cause microglial activation in rat hippocampus. *Neurosci Lett* 120:66–69.
- Silverman RB (2008) From basic science to blockbuster drug: the discovery of Lyrica. *Angew Chem Int Ed Engl* 47:3500–3504.
- Su H, Sochivko D, Becker A, Chen J, Jiang Y, Yaari Y, Beck H (2002) Upregulation of a T-type Ca²⁺ channel causes a long-lasting modification of neuronal firing mode after status epilepticus. *J Neurosci* 22:3645–3655.
- Sutton KG, Martin DJ, Pinnock RD, Lee K, Scott RH (2002) Gabapentin inhibits high-threshold calcium channel currents in cultured rat dorsal root ganglion neurones. *Br J Pharmacol* 135:257–265.
- Sutula TP, Dudek FE (2007) Unmasking recurrent excitation generated by mossy fiber sprouting in the epileptic dentate gyrus: an emergent property of a complex system. *Prog Brain Res* 163:541–563.
- Tauck DL, Nadler JV (1985) Evidence of functional mossy fiber sprouting in hippocampal formation of kainic acid-treated rats. *J Neurosci* 5:1016–1022.
- Taylor CP, Angelotti T, Fauman E (2007) Pharmacology and mechanism of action of pregabalin: the calcium channel alpha2-delta (alpha2-delta) subunit as a target for antiepileptic drug discovery. *Epilepsy Res* 73:137–150.
- Taylor CP, Garrido R (2008) Immunostaining of rat brain, spinal cord, sensory neurons and skeletal muscle for calcium channel alpha2-delta (alpha2-delta) type 1 protein. *Neuroscience* 155:510–521.
- Taylor CP, Vartanian MG, Andruszkiewicz R, Silverman RB (1992) 3-Alkyl GABA and 3-alkylglutamic acid analogues: two new classes of anticonvulsant agents. *Epilepsy Res* 11:103–110.
- Tedeschi A (2011) Tuning the orchestra: transcriptional pathways controlling axon regeneration. *Front Mol Neurosci* 4:60.
- Tran-Van-Minh A, Dolphin AC (2010) Gabapentin inhibits the Rab11-dependent recycling of the calcium channel subunit alpha2delta-2. *J Neurosci* 30:12856–12867.
- Tringham E, Powell KL, Cain SM, Kuplast K, Mezeyova J, Weerapura M, Eduljee C, Jiang X, Smith P, Morrison JL, Jones NC, Braine E, Rind G, Fee-Maki M, Parker D, Pajouhesh H, Parmar M, O'Brien TJ, Snutch TP (2012) T-type calcium channel blockers that attenuate thalamic burst firing and suppress absence seizures. *Sci Transl Med* 4:121ra19.
- Uchitel OD, Di Guilmi MN, Urbano FJ, Gonzalez-Inchausti C (2010) Acute modulation of calcium currents and synaptic transmission by gabapentinoids. *Channels (Austin)* 4:490–496.
- Vreugdenhil M, Hack SP, Draguhn A, Jefferys JG (2002) Tetanus toxin induces long-term changes in excitation and inhibition in the rat hippocampal CA1 area. *Neuroscience* 114:983–994.
- Xiao W, Boroujerdi A, Bennett GJ, Luo ZD (2007) Chemotherapy-evoked painful peripheral neuropathy: analgesic effects of gabapentin and effects on expression of the alpha-2-delta type-1 calcium channel subunit. *Neuroscience* 144:714–720.