

RESEARCH ARTICLE

By-Product Formation in Repetitive PCR Amplification of DNA Libraries during SELEX

Fabian Tolle¹, Julian Wilke¹, Jesper Wengel², Günter Mayer^{1*}

1. LIMES Institute, University of Bonn, Bonn, Germany, 2. Nucleic Acid Center, Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Odense M, Denmark

*gmayer@uni-bonn.de



CrossMark
click for updates

 OPEN ACCESS

Citation: Tolle F, Wilke J, Wengel J, Mayer G (2014) By-Product Formation in Repetitive PCR Amplification of DNA Libraries during SELEX. PLoS ONE 9(12): e114693. doi:10.1371/journal.pone.0114693

Editor: Mark Isalan, Imperial College London, United Kingdom

Received: August 30, 2014

Accepted: November 12, 2014

Published: December 9, 2014

Copyright: © 2014 Tolle et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper and its Supporting Information files.

Funding: Funded by Deutsche Forschungsgemeinschaft to G.M. (DFG Ma 3442/4.1). J.W. acknowledge support from the European Research Council under the European Union's Seventh Framework program (FP7/2007–2013)/ERC Grant agreement no. 268776. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

The selection of nucleic acid aptamers is an increasingly important approach to generate specific ligands binding to virtually any molecule of choice. However, selection-inherent amplification procedures are prone to artificial by-product formation that prohibits the enrichment of target-recognizing aptamers. Little is known about the formation of such by-products when employing nucleic acid libraries as templates. We report on the formation of two different forms of by-products, named ladder- and non-ladder-type observed during repetitive amplification in the course of *in vitro* selection experiments. Based on sequence information and the amplification behaviour of defined enriched nucleic acid molecules we suppose a molecular mechanism through which these amplification by-products are built. Better understanding of these mechanisms might help to find solutions minimizing by-product formation and improving the success rate of aptamer selection.

Introduction

Aptamers are short single stranded oligonucleotides with the ability to bind their target with high affinity and high specificity. They are selected against a broad range of targets including small molecules and proteins [1–3]. Therefore, aptamers can be conducive to a myriad of applications in various fields such as pharmaceutical sciences, e.g. as therapeutics, for drug discovery, as delivery and targeting units [4, 5], diagnostics, e.g. as sensors [6–9] or as tools in chemical biology [10–12].

Aptamers are usually generated by an iterative *in vitro* selection process called “systematic evolution of ligands by exponential enrichment” (SELEX) [1, 3, 13]. In

a typical SELEX experiment a library of up to 10^{15} different nucleic acid strands is exposed to a selection pressure where the best target binding sequences are partitioned from the rest. The best binding sequences are recovered, amplified by PCR and the enriched library is employed in the next selection cycle. After several rounds of selection and amplification the library is sequenced and monoclonal aptamer sequences are identified.

SELEX libraries typically comprise up to 100 nucleotides in length, containing a random region flanked by two constant primer-binding sites, which are used for PCR amplification of the library. The presence of up to 10^{15} different strands with undefined random regions can lead to several issues such as by-product formation in the PCR amplification. In conventional PCR approaches the major source for by-product formation is primer-primer hybridization leading to shorter double stranded products. In contrast to that, during amplification of complex libraries, single stranded sequences can bind to complementary bases in the random region serving as a primer, which is then extended by the polymerase, yielding longer ssDNA products [4, 14, 15]. This mechanism can lead to a full conversion of the product to longer by-products, even before the primers are exhausted.

Although being a common problem in the amplification of complex libraries [6, 8, 9, 16, 17], especially in the *in vitro* evolution community only a few studies about the molecular constitution and the underlying mechanisms of by-product formation have been published. In one example Marshall and Ellington reported molecular parasites enriched during a continuous isothermal amplification reaction that are longer than their parental molecules, yet replicate much more efficiently [10]. To our knowledge, albeit being detrimental to the enrichment process, there is no published report yet describing the mechanism for the formation of longer by-products during PCR amplification in SELEX experiments.

Herein we report two special kinds of PCR by-products observed during SELEX experiments conducted in our laboratory. Based on sequencing data we suggest a mechanistic model for the generation and amplification of this kind of by-products.

Material and Methods

Chemicals and Materials

All chemicals and the single-strand binding protein were purchased from Sigma-Aldrich (Munich, Germany). Human activated protein C was purchased from CellSystems (Troisdorf, Germany). Streptavidin coupled magnetic beads (Dynabeads M280) were purchased from Life Technologies (Darmstadt, Germany). Lambda exonuclease and T4 polynucleotide kinase were purchased from New England Biolabs (Frankfurt, Germany). Gamma ^{32}P ATP was purchased from PerkinElmer (Rodgau, Germany). NHS-PEG4-Biotin was purchased from Thermo Scientific (Rockford, USA). DNA purifications were

done with NucleoSpin Clean-Up/Plasmid kits from Macherey-Nagel (Düren, Germany) according to the manufacturers recommendation.

Oligonucleotides

Library: 5'-TGACCTTCGTGACTACAGNNNNNNNNNNNNNNNNNNNNNNNNNNNNNAC-TTAGTCAACT-GGCACAAT-3'; Forward primer: 5'-TGACCTTCGTGACTAC-AG-3'; Reverse primer: 5'-Phosphate-ATTGTGCCAGTTGACTAAGT-3'; Bold bases represent LNA building blocks. For subsequent lambda exonuclease digestion the reverse primer is labelled with a 5' phosphate group. Prior each selection cycle or binding assay the DNA was heat conditioned (refolded) by heating to 95°C for 3 min and slowly cooling down to 25°C in D-PBS.

PCR

PCR was done in a Mastercycler Personal/Gradient from Eppendorf (Hamburg, Germany). GoTaq DNA polymerase from Promega (Mannheim, Germany) and the supplied buffer containing 5 mM Mg²⁺ was used with the following cycling program (5 min 95°C; 1 min 95°C, 1 min 54°C, 1 min 72°C; hold 4°C). The optimal Mg²⁺ concentration and annealing temperature for the library were screened prior selection. The Mg²⁺ concentration was screened from 1.5 mM–5 mM, with increasing concentrations yielding more PCR-product and the highest product yield at 5 mM Mg²⁺ ([S1 Figure](#)). The annealing temperature was screened from 45°C to 65°C. 54°C was chosen as the optimal annealing temperature being the highest temperature to give optimal yields due to strongly decreasing yields from 57°C upwards ([S1 Figure](#)). 0.5 μM of both primers and 250 μM of dNTPs were used.

SELEX

The selections were carried out in selection buffer based on Dulbeccos' phosphate buffered saline (D-PBS) enriched with 1 mg/ml bovine serum albumin (BSA).

As target activated protein C (APC) immobilized on magnetic beads was used. For the selection, 100 μg APC were biotinylated with NHS-PEG4-Biotin and immobilized on 5 mg streptavidin coated magnetic beads. 500 pmol of the start library were incubated with 500 μg of empty streptavidin beads to remove the unwanted matrix binding sequences. The supernatant was incubated for 30 min at 37°C with 500 μg APC loaded beads in a total volume of 200 μl. The beads were thoroughly washed two times (30 sec and 5 min) with 200 μl selection buffer to remove unspecific binding DNA and the enriched library was eluted with 85 μl water from the beads by heat denaturation (80°C for 3 min). The eluted DNA was PCR amplified and purified via spin-columns. PCR product formation (18–22 PCR cycles) was carefully monitored by gel electrophoresis and stopped as soon as the desired product band could be detected on ethidium bromide stained gels. Single strand displacement of the double stranded PCR product was carried out

by lambda exonuclease digestion with 25 U lambda exonuclease [16, 17] for 1 h at 37°C and the resulting single strand DNA was purified via spin-columns. The purified single strand DNA (approx. 10 pmol) was applied in the next SELEX round. For each selection cycle the PCR product and the purified single strand DNA were analysed by gel electrophoresis on a 4% agarose gel. To increase the selection pressure the stringency of the washing steps was gradually increased by 2 to a maximum of 8 washing steps during the selection.

Cloning and Sequencing

Cloning was done with a TOPO TA Cloning kit from Life Technologies (Darmstadt, Germany) according to the manufacturers recommendation. 24 clones were sent to GATC-Biotech (Konstanz, Germany) for Sanger sequencing.

Binding Affinity Assay

Equilibrium dissociation constants were determined by nitrocellulose filter retention with radioactive labelled DNA. The single stranded DNA was 5' labelled with ^{32}P from gamma ^{32}P ATP using T4 polynucleotide kinase. The labelled DNA was purified from unreacted ATP and incubated in selection buffer with increasing concentrations of APC for 30 min at 37°C. The solution was filtered through nitrocellulose and the amount of bound aptamer was quantified by autoradiographie in a phosphorimager (FLA 5100 from Fuji, Japan). Raw binding data were corrected for unspecific background binding and equilibrium dissociation constants were calculated via nonlinear regression assuming a 1:1 binding model with Prism 5.0f (GraphPad Software, La Jolla, USA).

Results and Discussion

This study originated in the attempt to select DNA aptamers from constrained libraries with pre-defined stem regions. By pre-defining stem structures in the primer binding sites we endeavour to select aptamers that can easily be truncated in a post-SELEX maturation step, resulting in a fast and reliable method for the generation of short and versatile employable aptamers. Encouraged by the successful results previously obtained with RNA libraries [13] we set out to select compact DNA aptamers with a very short, only 6 bp long stem region, which is stabilized by three “Locked Nucleic Acids” (LNA) [14, 15] building blocks. As a model target the serine protease activated protein C (APC), which plays an important role in blood clotting, inflammation and cell death, was chosen.

During the course of a SELEX experiment we observed an emerging by-product pattern. From the 4th round on, two different kinds of dsDNA by-products after PCR amplification were observed (Fig. 1). Besides the correctly sized product band reflecting the length of the 58 base pair (bp) comprising dsDNA starting library, an intense second band migrating between the 75 bp and 100 bp marker lanes was observed. Additionally, a ladder-like product pattern was observed.




Fig. 1. Agarose gel analysis of the PCR products of the 8 SELEX cycles (lanes 1–8) showing the ladder-type and non-ladder-type by-products. Visualization by ethidium bromide staining; 4% agarose gel.

doi:10.1371/journal.pone.0114693.g001

Despite this by-product formation the SELEX was carried on for a total of eight rounds. After the 5th round a gel extraction of the correctly sized (58 bp) product band was performed in order to eliminate the by-products. However, despite this effort the same by-product pattern emerged after the next PCR amplification round. Subsequent filter retention binding assays with the 8th round library showed, as expected, no enrichment of target binders. This finding demonstrates that by-product formation is detrimental to a successful enrichment process.

In order to rule out that the nature of the library, containing LNAs caused this pattern formation, a control experiment with unmodified DNA primers was performed ([S2 Figure](#)). After consecutive amplification of the unmodified DNA library the same by-product pattern could be observed, showing that by-product formation is not due to the use of LNA-containing primer binding sites.

Similar by-product patterning has been previously observed in our laboratory using various nucleic acid libraries. In one example a selection against a small organic molecule with a different library containing a pre-defined stem (5'-AGCATAGAGACATCTGCTATTGGTAGCACA-N25-TGTGCAACCGTAGA-CTCCAGACTTCAGGTA-3') was rendered unsuccessfully due to the formation of similar ladder type by-products. Albeit vigorous gel purification attempts no enrichment could be observed. In our experience by-product formation is especially pronounced in constrained libraries, but not restricted to this kind of libraries only. The same by-product pattern was also observed with two other conventional unstructured libraries (5'-GCCTGTTGTGAGCCTCCTAAC-N49-CATGCTTATTCTTGTCTCCC-3' and 5'-CACGACGCAAGGGACCACAGG-N42-CAGCAGACACCGCAGAGGCA-3') ([S3 Figure](#)). This is especially noteworthy since both libraries were previously successfully used in other selection experiments [[2](#), [18](#)].

Having observed ladder type by-products in several selection attempts, we set out to investigate the nature of this kind of by-products. To understand why such elongated by-products are preferentially formed, the 8th round SELEX library was cloned and 24 clones were sent for sequencing. Representative sequences of the different classes of by-products identified are depicted in [Fig. 2](#) (for a complete list of sequencing data please refer to the [S4 Figure](#)).




Fig. 3. Proposed mechanism of "ladder type" by-product formation.

doi:10.1371/journal.pone.0114693.g003

primer binding sites. Since long stretches of such repeats allow for an efficient annealing of DNA, ever-longer by-products can be formed and will eventually dominate the library composition (Fig. 3). This mechanism leads to increasingly longer by-products but is only possible when a gap free full extension of the primer-binding site occurs. If the initial priming event happens in the middle of the random region (Fig. 4), the resulting by-product will have random gaps, preventing it to act as a template for further extensions.




Fig. 4. Proposed mechanism of “non-ladder type” by-product formation.

doi:10.1371/journal.pone.0114693.g004

In order to test our proposed by-product formation mechanism we conducted PCR amplification experiments with some of the representative monoclonal sequences (Fig. 5). In agreement with the proposed mechanism, the clones that reflect the length of the initial library (clones 2 and 5) yielded only one product of the expected size. However, amplification of clones with multiple reverse primer binding sites yielded ladder type by-products (clones 1, 8, 21, and 24). In turn, clones with a duplication of the 5' region (6 and 10), supposedly formed by the mechanisms described in Fig. 4, showed two products, but no formation of the ladder type by-products.

Having observed the ladder type by-products in several cases we believe that understanding the mechanism leading to the formation of this type of by-products is of general interest for the SELEX community and that the insights we could obtain might help to find solutions for this problem.

In a first attempt to overcome, or at least reduce, by-product formation several optimization attempts were executed. To disfavour the formation of long PCR products a special PCR program, completely skipping the elongation step, was investigated (5 min 95°C; 1 min 95°C, 30 sec 54°C; hold 4°C). Using this amplification strategy in consecutive PCR reactions the formation of ladder type by-products could be delayed to the 6. cycle.




Fig. 5. PCR amplification of representative mono-clones, the start library and a non-template control (NTC). Visualization by ethidium bromide staining; 4% agarose gel.

doi:10.1371/journal.pone.0114693.g005

Furthermore the use of single strand binding protein (SSB) was explored. In a concentration of 15 ng/ μ l the addition of SSB could delay the formation of ladder type by-products by one cycle, again alleviating, but not solving the problem. Additionally a study employing a shorter, only 13 nt long, reverse primer was employed. This time no improvement could be observed.

Since the addition of SSB and the optimized PCR program showed some improvement in delaying the formation of by-products in consecutive PCR cycles, a new SELEX attempt was started. Unfortunately despite the effort to reduce PCR by-products, also this attempt had to be aborted after 4 cycles of selection due to the formation of ladder type by-products.

To rule out a general problem with our SELEX methodology, a selection with a library containing different primer binding sites was carried out (5'-CACGACGCAAGGGACCACAGG-N42-CAGCACGACACCGCAGAGGCA-3'). In this case strong enrichment was observed (Figure 6A), thereby proving our SELEX methodology itself to be very effective for the selection of aptamers against APC. Cloning and sequencing the library of the eleventh selection cycle revealed three families of sequences. Interestingly, all sequences identified share the same consensus motive that was previously described for high affinity aptamers against APC [5, 19] (Figure 6B).

In conclusion, we demonstrate that amplification reactions during SELEX experiments can yield artificial by-product formation, which can be overcome by primer design. Our data reveal that these by-products build according to a special base-pairing mechanism, which results in distinct amplification patterns. Importantly, enrichment of specific target sequences is prohibited, most likely due to loss of potential binding sequences. So far all our attempts to reduce the by-product formation with this library failed, but the selection with a library containing different primer binding sites was successful. Noteworthy, in a different context, repetitive amplification of this library eventually also led to by-product formation (S3 Figure). This is an indication that besides primer design,




Fig. 6. Filter retention assay, A), with radioactive labelled DNA from the start library, 7th and 11th selection cycle. A strong increase in binding affinity is observable, demonstrating a successful enrichment during the SELEX. **B)** Sequencing data from the 11th selection cycle without primer binding sites. All mono-clones share the previously for high affinity APC aptamers described consensus motive (5'-TATCMCGNATGGGS-3') [11, 12, 19].

doi:10.1371/journal.pone.0114693.g006

the target also plays an important role. Since some desired library/target pairs are especially problematic, more solutions to generally increase the likelihood of a successful SELEX in these difficult cases are needed. Maybe the amplification related problems during SELEX might be overcome by special PCR methods and tweaked library design. Especially emulsion PCR has proven to be successful for the reduction of by-product formation in complex mixtures [7, 20-22] and should be further investigated in this context.

Supporting Information

S1 Figure. Optimisation of PCR conditions prior selection. A) The Mg²⁺ concentration was screened from 1.5 mM–5 mM B) The annealing temperature was screened from 45 °C to 65 °C. Visualization by ethidium bromide staining; 4% agarose gel.

[doi:10.1371/journal.pone.0114693.s001](https://doi.org/10.1371/journal.pone.0114693.s001) (EPS)

S2 Figure. Comparison of by-products using LNA and DNA primer in consecutive rounds of amplification. In both cases ladder-type by-products can be observed after 4 rounds of consecutive amplification. Visualization by ethidium bromide staining; 4% agarose gel.

[doi:10.1371/journal.pone.0114693.s002](https://doi.org/10.1371/journal.pone.0114693.s002) (EPS)

S3 Figure. By-product formation observed with another library (5'-CACGACGCAAGGGACCACAGG-N42-CAGCACGACACCGCAGAGGCA-3')

during a different selection experiment. Visualization by ethidium bromide staining; 4% agarose gel.

[doi:10.1371/journal.pone.0114693.s003](https://doi.org/10.1371/journal.pone.0114693.s003) (EPS)

S4 Figure. Complete list of sequencing data. Clones 4, 7, 13 and 20 did not yield utilisable sequencing results.

[doi:10.1371/journal.pone.0114693.s004](https://doi.org/10.1371/journal.pone.0114693.s004) (EPS)

Author Contributions

Conceived and designed the experiments: J. Wengel GM FT J. Wilke. Performed the experiments: FT J. Wilke. Analyzed the data: J. Wengel FT GM J. Wilke. Contributed reagents/materials/analysis tools: J. Wengel FT GM J. Wilke. Wrote the paper: J. Wengel FT GM J. Wilke.

References

1. Tuerk C, Gold L (1990) Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 249: 505–510.
2. Klug SJ, Famulok M (1994) All you wanted to know about SELEX. *Mol Biol Rep* 20: 97–107.
3. Ellington AD, Szostak JW (1990) In vitro selection of RNA molecules that bind specific ligands. *Nature* 346: 818–822. doi:10.1038/346818a0
4. Mushev MU, Krylov SN (2006) Selection of aptamers by systematic evolution of ligands by exponential enrichment: addressing the polymerase chain reaction issue. *Analytica Chimica Acta* 564: 91–96. doi:10.1016/j.aca.2005.09.069
5. Keefe AD, Pai S, Ellington A (2010) Aptamers as therapeutics. *Nature Reviews Drug Discovery* 9: 537–550. doi:10.1038/nrd3141
6. Kanagawa T (2003) Bias and artifacts in multitemplate polymerase chain reactions (PCR). *Journal of Bioscience and Bioengineering* 96: 317–323. doi:10.1016/S1389-1723(03)90130-7
7. Plaxco KW, Soh HT (2011) Switch-based biosensors: a new approach towards real-time, in vivo molecular detection. *Trends Biotechnol* 29: 1–5. doi:10.1016/j.tibtech.2010.10.005
8. Acinas SG, Sarma-Rupavtarm R, Klepac-Ceraj V, Polz MF (2005) PCR-induced sequence artifacts and bias: insights from comparison of two 16S rRNA clone libraries constructed from the same sample. *Appl Environ Microbiol* 71: 8966–8969. doi:10.1128/AEM.71.12.8966-8969.2005
9. Kalle E, Gulevich A, Rensing C (2013) External and semi-internal controls for PCR amplification of homologous sequences in mixed templates. *J Microbiol Methods* 95: 285–294. doi:10.1016/j.mimet.2013.09.014
10. Marshall KA, Ellington AD (1999) Molecular parasites that evolve longer genomes. *J Mol Evol* 49: 656–663.
11. Mayer G (2009) The chemical biology of aptamers. *Angew Chem Int Ed Engl* 48: 2672–2689. doi:10.1002/anie.200804643
12. Tolle F, Mayer G (2013) Dressed for success—applying chemistry to modulate aptamer functionality. *Chem Sci* 4: 60–67. doi:10.1039/c2sc21510a
13. Mayer G, Wulffen B, Huber C, Brockmann J, Flicke B, et al. (2008) An RNA molecule that specifically inhibits G-protein-coupled receptor kinase 2 in vitro. *RNA* 14: 524–534. doi:10.1261/rna.821908
14. Koshkin AA, Rajwanshi VK, Wengel J (1998) Novel convenient syntheses of LNA [2.2. 1] bicyclo nucleosides. *Tetrahedron Letters* 39: 4381–4384. doi:10.1016/S0040-4039(98)00706-0

15. **Koshkin AA, Singh SK, Nielsen P, Rajwanshi VK, Kumar R, et al.** (1998) LNA (Locked Nucleic Acids): Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclonucleoside monomers, oligomerisation, and unprecedented nucleic acid recognition. *Tetrahedron* 54: 3607–3630. doi:10.1016/S0040-4020(98)00094-5
16. **Avci-Adali M, Paul A, Wilhelm N, Ziemer G, Wendel HP** (2010) Upgrading SELEX technology by using lambda exonuclease digestion for single-stranded DNA generation. *Molecules* 15: 1–11. doi:10.3390/molecules15010001
17. **Svobodová M, Pinto A, Nadal P, O' Sullivan CK** (2012) Comparison of different methods for generation of single-stranded DNA for SELEX processes. *Anal Bioanal Chem* 404: 835–842. doi:10.1007/s00216-012-6183-4
18. **Raddatz M-SL, Dolf A, Endl E, Knolle P, Famulok M, et al.** (2008) Enrichment of Cell-Targeting and Population-Specific Aptamers by Fluorescence-Activated Cell Sorting. *Angewandte Chemie* 120: 5268–5271. doi:10.1002/ange.200800216
19. **Müller J, Isermann B, Dücker C, Salehi M, Meyer M, et al.** (2009) An exosite-specific ssDNA aptamer inhibits the anticoagulant functions of activated protein C and enhances inhibition by protein C inhibitor. *Chemistry & biology* 16: 442–451. doi:10.1016/j.chembiol.2009.03.007
20. **Williams R, Peisajovich SG, Miller OJ, Magdassi S, Tawfik DS, et al.** (2006) Amplification of complex gene libraries by emulsion PCR. *Nat Methods* 3: 545–550. doi:10.1038/nmeth896
21. **Schütze T, Rubelt F, Repkow J, Greiner N, Erdmann VA, et al.** (2011) A streamlined protocol for emulsion polymerase chain reaction and subsequent purification. *Analytical biochemistry* 410: 155–157. doi:10.1016/j.ab.2010.11.029
22. **Shao K, Ding W, Wang F, Li H, Ma D, et al.** (2011) Emulsion PCR: a high efficient way of PCR amplification of random DNA libraries in aptamer selection. *PLoS ONE* 6: e24910. doi:10.1371/journal.pone.0024910