

Investigation of *In Vitro* Activity of Five Antifungal Drugs against Dermatophytes Species Isolated from Clinical Samples Using the E-Test Method

Beş Antifungal İlacın Klinik Örneklerden İzole Edilen Dermatofit Türlerine Karşı *In Vitro* Aktivitesinin E-Test Yöntemi ile Araştırılması

Ayşe Esin Aktas¹, Nimet Yigit², Akin Aktas³, Sultan Gamze Gozubuyuk¹

¹Department of Microbiology and Clinical Microbiology, Faculty of Medicine, Ataturk University, Erzurum, Turkey

²Department of Medical Laboratory, Ataturk University Health Services Vocational School, Erzurum, Turkey

³Department of Dermatology, Faculty of Medicine, Ataturk University, Erzurum, Turkey

Abstract

Objective: Dermatophytosis is an infection with fungi related to the skin: glabrous skin, hair and/or nails. Oral treatment of fungal infections in dermatology has become a preferred modality for the management of these very common conditions. Although there are increasing numbers of antifungals available for treatment of dermatophytes, some cases and relapses have been unresponsive to treatment. The determination of fungus *in-vitro* antifungal susceptibility has been reported to be important for the ability to eradicate dermatophytes. It is necessary to perform antifungal susceptibility testing of dermatophytes. E-test (AB Biodisk, Sweden) is a rapid, easy-to-perform *in-vitro* antifungal susceptibility test. The aim of this study was to investigate the susceptibility of the different species of dermatophyte strains isolated clinical specimens to five antifungal agents using the E-test method.

Materials and Methods: A total of 66 specimens were collected from the nails, feet, inguinal region, trunk and hands. These strains tested MIC endpoints of E-test for amphotericin B, fluconazole, itraconazole, caspofungin, and ketoconazole were read after 72, and 96 hours incubation for each strain on RPMI 1640 agar.

Results: The dermatophytes tested included *Trichophyton rubrum* 43 (65.1%), *Trichophyton mentagrophytes* 7 (10.7%), *Microsporium canis* 5 (7.6%), *Trichophyton tonsurans* 5 (7.6%), *Epidermophyton floccosum* 4 (6.0%) and *Trichophyton violaceum* 2 (3.0%). The most active agent against all dermatophytes species was caspofungin with a minimal inhibitory concentration (MIC) range ($\mu\text{g}/\text{mL}^{-1}$) (0.02-3, 0.032-4, 0.125-0.50, 0.032-2, 0.25-0.50, 0.125-0.50) and itraconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.038-1.5, 0.094-1.5, 1-32, 0.016-0.50, 0.25-0.50, 0.125-0.50). The least active agent was fluconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0, 19-48, 2-256, 2-8, 256, 256, 8-24).

Conclusion: E-test seems to be an alternative method to MIC-determination of antifungal drugs for dermatophytes species, since it is a less-laborious methodology and results could be obtained faster.

Key Words: Dermatophytes, E-test, antifungal

Özet

Amaç: Dermatofitler mantarlar tarafından oluşturulan deri, saç ve tırnakları tutan enfeksiyonlardır. Dermatolojide mantar enfeksiyonlarının oral tedavisi yaygın olarak tercih edilen yöntem haline gelmiştir. Dermatofitlerin tedavisi için kullanılan antifungal ilaçların sayısının artmasına rağmen bazı durumlarda tedaviye yanıtızlık ve nüksler gelişmektedir. *In-vitro* antifungal duyarlılığın belirlenmesinin dermatofitlerin etkili olarak tedavi edilmesi için önemli olduğu bildirilmektedir. Antifungal duyarlılığın belirlenmesinde yöntemler önemlidir. E-test (AB Biodisk, İsveç) hızlı ve kolay uygulanabilir bir duyarlılık yöntemidir. Bu çalışmanın amacı, klinik örneklerden izole edilen farklı dermatofit türlerinin beş antifungal ilaca karşı duyarlılıklarını E-test yöntemi ile belirlemektir.

Gereç ve Yöntem: Tırnak, ayak, inguinal bölge, gövde ve el/bilek olmak üzere değişik anatomik bölgelerden alınan toplam 66 klinik örnek çalışmaya dâhil edildi. Bu örneklerden izole edilen dermatofit suşlarının duyarlılıkları, amfoterisin B, flukonazol, itraconazol, kaspo-fungin ve ketokonazol'e karşı RPMI 1640 agar besiyerinde 72 ve 96 saat süre ile E-test yöntemi kullanılarak çalışıldı.

Bulgular: Çalışmaya dâhil edilen dermatofit türleri *Trichophyton rubrum* 43 (%65,1), *Trichophyton mentagrophytes* 7 (%10,7), *Microsporium canis* 5 (%7,6), *Trichophyton tonsurans* 5 (%7,6), *Epidermophyton floccosum* 4 (%6,0) and *Trichophyton violaceum* 2 (%3,0) olarak belirlendi. Tüm dermatofit türlerine karşı en etkili antifungal ilaçlar (0,02-3, 0,032-4, 0,125-0,50, 0,032-2, 0,25-0,50, 0,125-0,50) MIC ($\mu\text{g}/\text{mL}^{-1}$) aralıkları ile kaspo-fungin ve (0,038-1,5, 0,094-1,5, 1-32, 0,016-0,50, 0,25-0,50, 0,125-0,50) MIC ($\mu\text{g}/\text{mL}^{-1}$) aralıkları ile itraconazol olarak bulundu. En az etkili antifungal ise (0, 19-48, 2-256, 2-8, 256, 256, 8-24) MIC ($\mu\text{g}/\text{mL}^{-1}$) aralıkları ile flukonazol olarak bulundu.

Sonuç: E-test dermatofit türlerinin antifungal duyarlılığının belirlenmesinde daha az zahmetli ve sonuçların daha hızlı elde edilebilir olması ile alternatif bir yöntem gibi görünmektedir.

Anahtar Kelimeler: Dermatofitler, E-test, antifungal



Received: May 21, 2013 / Accepted: August 2, 2013

Correspondence to: Ayşe Esin Aktas, Department of Microbiology and Clinical Microbiology, Faculty of Medicine, Ataturk University, Erzurum, Turkey
Phone: +90 442 344 73 45 e-mail: aeaktas@yahoo.co.uk

©Copyright 2014 by the Ataturk University School of Medicine - Available online at www.eajm.org
doi:10.5152/eajm.2014.05

Introduction

Dermatophytes are a specialized group of fungi, which effect keratinous tissue of humans and other vertebrates, causing superficial infections. The organisms belong to three genera, Trichophyton, Epidermophyton, and Microsporum. Infections caused by these fungi are among the most prevalent cutaneous infections globally and the recent increase in the number of patients with immunocompromised states, such as AIDS, diabetes mellitus, cancer and organ transplantation has given these infections more prominence [1-6].

The treatment of dermatophytosis is based on the use of topical and systemic antifungal agents. In recent years, a number of safe and highly effective antifungal agents have been introduced into clinical practice. Although an increasing number of antimycotics have become available for the treatment of dermatophytosis, there are reports suggesting recalcitrant to therapy or possibly resistance of dermatophytes to antimicrobial agents. In order to predict the ability of a given antimycotic agent to eradicate dermatophytes and help managing patients, determination of the *in vitro* antifungal susceptibility of dermatophytes would be helpful in understanding a failed or successful treatment. However, not all species have the same susceptibility pattern and it may be necessary to perform *in vitro* susceptibility testing for selection and monitoring of antifungal therapy. Although a reference method is not yet available, various techniques have been used to test dermatophytes, including broth macro- and micro dilution methods, agar dilution and disc diffusion. However, these methods are time-consuming and labour-intensive, and are not practical for the clinical laboratory. Therefore, simple alternative approaches are needed [1, 3, 4, 6-8].

The E-test is a simple, agar-based, quantitative minimal inhibitory concentration (MIC) method. The reagent consists of a thin, calibrated plastic strip with a predefined, exponential and continuous gradient of antifungal agent across 15 two-fold dilutions. The E-test has been satisfactorily used to test bacteria, yeasts and moulds. However, there is limited data available on the performance of the E-test for antifungal susceptibility of dermatophytes [4, 9-11].

The aim of this study was to investigate the susceptibility of the different species of dermatophyte strains isolated clinical specimens to five antifungal agents (amphotericin B, fluconazole, itraconazole, caspofungin, and ketoconazole) using the E-test method.

Materials and Methods

Strains and Specimens: Sixty-six strains were isolated from infected skin and nails in the Microbiology and Clinical

Microbiology Department of School of Medicine, Ataturk University. Isolates were collected over a one-year period in Mycology Laboratory. They included *T. rubrum*, *T. mentagrophytes*, *M. canis*, *T. tonsurans*, *E. floccosum* and *T. violaceum*. All strains were identified by standard methods, which included identification based on the macroscopic and microscopic characteristics of the culture strains. Additional tests included those for the ability to produce a red pigment when the strains were grown on Potato Dextrose Agar (PDA) and for the ability to produce urease, as well as the hair perforation test. Strains were stored -70°C until the time of use, and prior to testing were sub-cultured on PDA at 28°C for 15 days to ensure optimal growth characteristics [1, 3, 6]. All procedures in the experimental protocol were approved by The Ethics Committee of Medical Faculty.

E-Test Method

Medium: The test was performed in RPMI 1640 medium with L-glutamine, although without bicarbonate (Gibco, New York, USA), pH 7.0 supplemented with 2% glucose, buffered 0.165 M morpholinepropanesulfonic acid (MOPS) (Fisher Biotech, New Jersey, USA) and 1.8% agar (Difco, Sparks, USA). The 15-cm diameter petri plates contained RPMI 1640 at a depth of 4.0 mm [4].

Antifungal Agents: E-test strips were obtained from AB Biodisk (Solna, Sweden) and stored at -20°C until tests were performed. The concentrations assayed ranged from 0.002 to 32.000 µg/mL⁻¹ for amphotericin B, itraconazole, caspofungin, and ketoconazole and 0.016 to 256.000 µg/mL⁻¹ for fluconazole.

Procedure: All isolates were tested against five antifungal agents using the E-test according to the manufacturer's instructions. The inoculum suspensions were prepared and adjusted to 65-70% transmittance at a wavelength of 530 nm corresponding to a concentration of 10⁵-10⁶ cfu/mL⁻¹ verified by quantitative plate counts. The RPMI agar surface was inoculated by dipping a sterile swab into the inoculum suspension and streaking it evenly in three directions. After excess moisture was absorbed into the agar and the surface was completely dry, an E-test strip was applied to each plate. The plates were incubated at 28°C and the results were read at 72-96 hour [4].

Determination of MIC endpoints: In general, MIC was defined as the lowest drug concentration at which the border of the elliptical inhibition zone intercepted the MIC scale on the E-test strip. When a double halo of growth was observed, the MIC was read at the point where growth was completely inhibited. When different intersections were observed on either side of the strip, the highest MIC value was read [4].

Results

The isolated dermatophytes were obtained from the toe-nails 16 (24.2%), feet 33 (50.0%), inguinal region 7 (10.7%), trunk 5 (7.6%) and hands 5 (7.6%). The distribution of isolated species 66 dermatophytes were *T. rubrum* 43 (65.1%), *T. mentagrophytes* 7 (10.7%), *M. canis* 5 (7.6%), *T. tonsurans* 5 (7.6%), *E. floccosum* 4 (6.0%) and *T. violaceum* 2 (3.0%) (Table 1).

All strains tested grew well on RPMI glucose, supplement agar plated. They were read in the E-test method after 96 hours of incubation, except in the case of *T. mentagrophytes*, which required only 72 hours of incubation.

Table 2 summarizes the *in vitro* susceptibilities of 66 clinical isolates of dermatophytes to five antifungal agents as determined by E-test. The most active agent against all dermatophytes species was caspofungin with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.02-3, 0.032-4, 0.032-4, 0.125-0.50, 0.25-0.50, 0.125-0.50) and itraconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.038-1.5, 0.094-1.5, 1-32, 0.016-0.50, 0.25-0.50, 0.125-0.50). The least active agent was fluconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0,19-48, 2-256, 2-8, 256, 256, 8-24). Test results of the susceptibility to amphotericin B and ketoconazole were as follows; respectively, 0,012-8, 0,19-8, 0,50-3, 0,125-6, 32, 0,75 and 0,032-8, 0,064-8, 32, 32, 32, 32.

In general, the species of dermatophytes showed similar patterns of susceptibility to each antifungal agent tested. High MIC values were found for some isolates, two dermatophytes strains (1 *T. rubrum* and 1 *T. mentagrophytes*) had MICs of caspofungine of 32 $\mu\text{g}/\text{mL}$, 16 strains (11 *T. rubrum*, 4 *E. floccosum* and 1 *T. mentagrophytes*) had MICs of Amphotericin B of 32 $\mu\text{g}/\text{mL}$, 53 strains (36 *T. rubrum*, 5 *T. tonsurans*, 4 *E. floccosum*, 2 *M. canis*, and 6 *T. mentagrophytes*) had MICs of fluconazole of 256 $\mu\text{g}/\text{mL}$, 2 strains (2 *M. canis*) had MICs of itraconazole of 32 $\mu\text{g}/\text{mL}$, and 33 strains (18 *T. rubrum*, 1 *T. tonsurans*, 4 *E. floccosum*, 5 *M. canis*, 2 *T. violaceum*, and 3 *T. mentagrophytes*) had MICs of ketoconazole of 32 $\mu\text{g}/\text{mL}$. Table 2 summarizes the MIC ranges, concentrations inhibiting 50% (MIC 50) and 90% (MIC 90) of the

isolates of the five antifungal drugs against 66 strains of dermatophytes.

Discussion

Infections caused by dermatophytes occur worldwide and can be very severe and difficult to treat in patients whose immunological response is impaired. These infections represented an important public health problem as yet unresolved [4, 7].

Dermatophytes are responsible for the majority of fungal infections involving the skin, hair and nails. They comprise a phylogenetically closely related group of genera with numerous species. They attack the keratinized tissues and cause a wide spectrum of clinical manifestations that vary from mild to severe [6].

The distribution of the dermatophytes and their etiological agents has unequal frequencies, with variations of their prevalence according to the countries and even the regions of the same country. In this study, *T. rubrum* was the most frequently isolated organism 43 (65.1%), followed by *T. mentagrophytes* 7 (10.7%), *M. canis* 5 (7.6%), *T. tonsurans* 5 (7.6%), *E. floccosum* 4 (6.0%) and *T. violaceum* 2 (3.0%). These results are in agreement with many other local [3, 12-19] and international studies [1, 4, 7, 9, 20-25].

Most superficial infections caused by dermatophytes can be rapidly eradicated with topical and systemic antifungals. Oral antifungal therapy with newer agents, such as terbinafine, itraconazole and fluconazole, is the treatment of choice for dermatophytosis that does not respond to topical therapies. The activity spectrum to these drugs is variable, leading to treatment failure in 25-40% of treated patients, possibly due to poor patient compliance, lack of drug penetration into nail, medication bioavailability or drug interactions and resistance [26].

In vitro analysis of the antifungal activity of anti-fungal agents enables comparison between different antimycotics, which in turn may clarify the reasons for lack of clinical

Table 1. Isolated dermatophyte strains in relation to localization

Dermatophytes	n	%	Localization				
			Toe nail	Foot	Inguinal region	Trunk	Hands
<i>T. rubrum</i>	43	65.1%	12	24	4	3	-
<i>T. mentagrophytes</i>	7	10.7%	1	2	-	2	2
<i>M. canis</i>	5	7.6%	1	3	-	-	1
<i>T. tonsurans</i>	5	7.6%		2	3	-	-
<i>E. floccosum</i>	4	6.0%	2	2	-	-	-
<i>T. violaceum</i>	2	3.0%	-	-	-	-	2
Total	66	100.0%	16 (24.2%)	33 (50.0%)	7 (10.7%)	5 (7.6%)	5 (7.6%)

Table 2. Susceptibility data for dermatophytes species against five antifungal agents using the E-test method

Species (n)	Antifungal agent	MIC range*	MIC ₅₀	MIC ₉₀
<i>T. rubrum</i> (43)	Amphotericin B	0.012-8	0.50	1.5
	Fluconazole	0.19-48	-	-
	Itraconazole	0.038-1.5	0.50	0.19
	Caspofungine	0.02-3	1	0.064
	Ketoconazole	0.032-8	-	-
<i>T. mentagrophytes</i> (7)	Amphotericin B	0.19-8	0.70	4
	Fluconazole	2-256	-	-
	Itraconazole	0.094-1.5	0.25	1.5
	Caspofungine	0.032-4	0.25	2
	Ketoconazole	0.064-8	2	8
<i>M. canis</i> (5)	Amphotericin B	0.50-3	0.50	1
	Fluconazole	2-8	-	-
	Itraconazole	1-32	-	-
	Caspofungine	0.125-0.50	0.50	0.125
	Ketoconazole	32	-	-
<i>T. tonsurans</i> (5)	Amphotericin B	0.125-6	0.50	0.50
	Fluconazole	256	-	-
	Itraconazole	0.016-0.50	0.125	0.125
	Caspofungine	0.032-2	0.032	0.032
	Ketoconazole	32	-	-
<i>E. floccosum</i> (4)	Amphotericin B	32	-	-
	Fluconazole	256	-	-
	Itraconazole	0.25-0.50	0.25	0.50
	Caspofungine	0.25-0.50	0.25	-
	Ketoconazole	32	-	-
<i>T. violaceum</i> (2)	Amphotericin B	0.75	-	-
	Fluconazole	8-24	-	-
	Itraconazole	0.125-0.50	-	-
	Caspofungine	0.125-0.50	-	-
	Ketoconazole	32	-	-

MIC: Minimal inhibitory concentration
*MIC (µg/mL⁻¹)

response and assist clinicians in choosing an effective therapy for their patients. However, it is important that the methodologies used for *in vitro* testing be standardized to facilitate the establishment of quality control parameters and interpretive break points [27].

Currently, no reference method has been established to test drug susceptibilities of dermatophytes. The development of simple and reproducible techniques is required for clinical

testing of these important pathogens. The E-test is a new and promising method with broad applications in clinical laboratory practice, and is supported by the results of extensive testing of bacteria and yeasts. However, there are only a few reports describing the use of this method for dermatophytes [4, 9, 20-22].

In this study, we investigated MIC values of five antifungal agents (amphotericin B, fluconazole, itraconazole,

casposfungin, and ketoconazole) to the different species of dermatophyte strains isolated clinical specimens using the E-test method.

In our study, the most active agent against all dermatophytes species was casposfungin with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.02-3, 0.032-4, 0.125-0.50, 0.032-2, 0.25-0.50, 0.125-0.50) and itraconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.038-1.5, 0.094-1.5, 1-32, 0.016-0.50, 0.25-0.50, 0.125-0.50). The least active agent was fluconazole with an MIC range ($\mu\text{g}/\text{mL}^{-1}$) (0.19-48, 2-256, 2-8, 256, 256, 8-24). Test results of the susceptibility to amphotericin B and ketoconazole were as follows; respectively, 0.012-8, 0.19-8, 0.50-3, 0.125-6, 32, 0.75 and 0,032-8, 0,064-8, 32, 32, 32, 32.

With respect to itraconazole, all of *T. rubrum* isolates were inhibited in concentrations ranging from 0.038 to 1.5 $\mu\text{g}/\text{mL}^{-1}$. The other species, except *M. canis* showed similar sensitivity ranges. Two *M. canis* strains had MICs of itraconazole of 32 $\mu\text{g}/\text{mL}$. However, for fluconazole, we observed that high MIC values. Fifty-three strains (36 *T. rubrum*, 5 *T. tonsurans*, 4 *E. floccosum*, 2 *M. canis*, and 6 *T. mentagrophytes*) had MICs of fluconazole of 256 $\mu\text{g}/\text{mL}$. In general, our data are in agreement with studies of Don Santos et al. [22], Fernandez-Torres et al. [4], Silva-Barros et al. [9], Kang et al. [21] and Abdel-Aal et al. [20].

Casposfungin the other most active agents for all dermatophytes species in our study with an MIC range (0.02-3 for *T. rubrum*, 0.032-4 for *T. mentagrophytes*, 0.032-4 for *M. canis*, 0.032-2 for *T. tonsurans*, 0.25-0.50 for *E. floccosum*, 0.125-0.50 for *T. violaceum*).

In our study, 33 (50%) isolates of tested dermatophytes by E-test (18 *T. rubrum*, 1 *T. tonsurans*, 4 *E. floccosum*, 5 *M. canis*, 2 *T. violaceum*, and 3 *T. mentagrophytes*) were resistant with an MIC range 32 $\mu\text{g}/\text{mL}$ of ketoconazole. These results were obtained other researchers [4, 9, 20-22].

Amphotericin B, the other drug with an MIC range (0.012-8, 0.19-8, 0.50-3, 0.125-6, 32, 0.75) in the present study. 16 strains (11 *T. rubrum*, 4 *E. floccosum* and 1 *T. mentagrophytes*) had MICs of amphotericin B of 32 $\mu\text{g}/\text{mL}$. Kang et al.[21] observed that amphotericin B was 0.094~0.5 $\mu\text{g}/\text{mL}$ on *T. rubrum*, 0.032~1.0 $\mu\text{g}/\text{mL}$ on *T. mentagrophytes*, 0.19 $\mu\text{g}/\text{mL}$ on *M. canis*, and 0.032 $\mu\text{g}/\text{mL}$ on *M. gypseum*.

Antifungal susceptibility testing is a dynamic field of medical mycology. Development and standardization of antifungal susceptibility test have shown remarkable progress in the field of medical mycology [6], although, studies using the E-test method for dermatophytes susceptibilities is not yet sufficient. In a limited number of studies, showed that E-test seems to be an alternative method to MIC-determination of antifungal drugs for dermatophytes, since it is a less-laborious methodology and results could be obtained faster [4, 9, 21, 22].

In conclusion, this study showed that the E-test represented a simple and efficacious method for antifungal susceptibility testing of dermatophytes. Regarding its performance, the E-test was not labour demanding, was easy to interpret, and with the potential of being used as an alternative assay for azole antifungal susceptibility testing of dermatophytes.

Conflict of Interest: No conflict of interest was declared by the authors.

Peer-review: Externally peer-reviewed.

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Author Contributions: Concept - A.E.A., N.Y.; Design - A.E.A., N.Y.; Supervision - A.E.A., A.A.; Funding - A.E.A., ; Materials - A.A., A.E.A.; Data Collection and/ or Processing - S.G.G., A.E.A., A.A., N.Y.; Analysis and/or Interpretation - A.E.A., N.Y.; Literature Review - A.E.A., N.Y.; Writer - A.E.A., N.Y.; Critical Review - A.E.A., A.A., N.Y.

Financial Disclosure: The authors declared that this study has received no financial support.

References

1. Ebrahim HM, Asaad AM, Amer A. Antifungal susceptibility patterns of dermatophytes clinical isolates from dermatophytosis patients before and after therapy. *Egyptian J of Med Microbiol* 2010; 19: 41-46.
2. Fernandez- Torres B, Carrillo AJ, Martin E, et al. In vitro activities of 10 antifungal drugs against 508 dermatophytes strains. *Antimicrob Agents and Chemother* 2001; 45: 2524-8.
3. Karaca N, Koç AN. In vitro susceptibility testing of dermatophytes: comparison of disk diffusion and reference broth dilution method. *Diagn Microbiol Infect Dis* 2004; 48: 259-64.
4. Fernandez- Torres B, Carrillo-Munoz A, Ortoneda M, et al. Interlaboratory evaluation of the E-test for antifungal susceptibility testing of dermatophytes. *Med Mycol* 2003; 41: 125-130.
5. Fernandez-Torres B, Carrillo-Munoz A, Inza I, et al. Effect of culture medium on the disk diffusion method for determining antifungal susceptibilities of dermatophytes. *Antimicrob Agents and Chemother* 2006; 50: 2222-4.
6. Pakshir K, Bahaedinie L, Rezaei Z, et al. In vitro activity of six antifungal against clinically important dermatophytes. *Jundishapur Journal of Microbiology* 2009; 2: 158-163.
7. Araujo CR, Miranda KC, Fernandes OFL, et al. In-vitro susceptibility testing of dermatophytes isolated in Goiana, Brazil, against five antifungal agents by broth microdilution method. *Rev Inst Med Trop Sao Paulo* 2009; 51: 9-12.
8. Siqueira ER, Ferreira JC, Pedrosa RS, et al. Dermatophyte susceptibilities to antifungal azole agents tested in vitro by broth

- macro and microdilution methods. *Rev Inst Med Trop Sao Paulo* 2008; 50: 1-5.
9. da Silva Barros ME, de Assis Santos D, Soares Hamdan J. Antifungal susceptibility testing of *trichophyton rubrum* by E-test. *Arch Dermatol Res* 2007; 299: 107-9.
 10. Pfaller MA, Messer SA, Mills K, et al. In-vitro Susceptibility Testing of Filamentous Fungi: comparison of E-test and Reference Microdilution Methods for Determining Itraconazole MICs. *J Clin Microbiol* 2000; 38: 3359-61.
 11. Favel A, Michel-Nguyen A, Chastin C, et al. In-vitro susceptibility patterns of *candida lusitanae* and evaluation of the E-test method. *J Antimicrob Chemother* 1997; 39: 591-6.
 12. Sarifakioglu E, Seçkin D, Demirbilek M, et al. In vitro antifungal susceptibility patterns of dermatophyte strains causing tinea unguium. *Clin Exp Dermatol* 2007; 32: 675-9.
 13. Çetinkaya Z, Kiraz N, Karaca S, et al. Antifungal susceptibilities of dermatophytic agents isolated from clinical specimens. *Eur J Dermatol* 2005; 15: 258-61.
 14. Bilgili ME, Sabuncu İ, Saraçoğlu ZN, et al. Dermatophyte types isolated from patients presented with dermatophytosis in our clinic. *Türkiye Klinikleri J Dermatol* 2001; 11: 185-90.
 15. Dilek N, Yücel AY, Dilek AR, et al. Dermatophytosis agents in patients who attending to dermatology clinic of Firat University Hospital. *Turkish J of Dermatol* 2009; 3: 27-31.
 16. Gürcan Ş, Tikveşli M, Eskiocak M, et al. Investigation of the agents and risk factors of dermatophytosis: a hospital-based study. *Mikrobiyol Bul* 2008; 42: 95-102.
 17. İnci M, Özer B, Duran N, et al. Onikomikoz ön tanısıyla gönderilen örneklerden izole edilen dermatofitlerin değerlendirilmesi. *Türk Mikrobiyol Cem Derg* 2011; 41: 61-64.
 18. Özekinci T, Özbek E, Gedik M, et al. Dicle Üniversitesi Tıp Fakültesi mikrobiyoloji laboratuvarına başvuran hastalarda dermatofitoz etkenleri. *Dicle Tıp Dergisi* 2006; 33: 19-22.
 19. Ecemiş T, Değerli K, Ertmercan AT, et al. The study of retrospective onychomycosis in Manisa: 2003-2010. *Kocatepe Medical Journal* 2013; 14: 65-68.
 20. Abdel-Aal AM, Taha MM, El-Mashad N, et al. Antifungal susceptibility testing: new trends. *Egyptian Dermatology Online Journal* 2007; 3: 1-10.
 21. Kang GS, Suh MK, Ha GY. Antifungal susceptibility testing of dermatophytes using E-test. *Korean J Med Mycol* 2010; 15: 124-33.
 22. Dos Santos JI, Paulo CR, Viani FC, Gambale W. Susceptibility testing of *Trichophyton rubrum* and *Microsporum canis* to three azoles by E-test. *Journal de Mycologie Medicale* 2001; 1: 42-3.
 23. Carrillo-Munoz AJ, Cardenes CD, Carrillo-Orive B, et al. In vitro antifungal activity of voriconazole against dermatophytes and superficial isolates of *scopulariopsis brevicaulis*. *Rev Iberoam Micol* 2005; 22: 110-3.
 24. Carrillo-Munoz AJ, Giusiano G, Guarro J, et al. In vitro activity of voriconazole against dermatophytes, *scopulariopsis brevicaulis* and other opportunistic fungi as agents of onychomycosis. *Intl J Antimicrob Agents* 2007; 30: 157-61.
 25. Favre B, Hotbauer B, Hildering KS, et al. Comparison of in vitro activities of 17 antifungal drugs against a panel of 20 dermatophytes by using a microdilution assay. *J Clin Microbiol* 2003; 41: 4817-9.
 26. Bueno JG, Martinez C, Zapata B, et al. In vitro activity of fluconazole, itraconazole, voriconazole and terbinafine against fungi causing onychomycosis. *Clin Exp Dermatol* 2010; 35: 658-63.
 27. Gupta AK, Kohli Y. In vitro susceptibility testing of ciclopirox, terbinafine, ketoconazole and itraconazole against dermatophytes and nondermatophytes, and in vitro evaluation of combination antifungal activity. *British J of Dermatol* 2003; 149: 296-305.