

# Skeletal muscle ATP turnover by $^{31}\text{P}$ magnetic resonance spectroscopy during moderate and heavy bilateral knee extension

Daniel T. Cannon<sup>1,2</sup>, William E. Bimson<sup>3</sup>, Sophie A. Hampson<sup>4</sup>, T. Scott Bowen<sup>2,5</sup>, Scott R. Murgatroyd<sup>2</sup>, Simon Marwood<sup>4</sup>, Graham J. Kemp<sup>3,6</sup> and Harry B. Rossiter<sup>1,2</sup>

<sup>1</sup>Rehabilitation Clinical Trials Center, Division of Respiratory & Critical Care Physiology & Medicine, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA, USA

<sup>2</sup>School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, UK

<sup>3</sup>Magnetic Resonance & Image Analysis Research Centre, University of Liverpool, Liverpool, UK

<sup>4</sup>School of Health Sciences, Liverpool Hope University, Liverpool, UK

<sup>5</sup>Department of Internal Medicine and Cardiology, University of Leipzig – Heart Center, Leipzig, DE

<sup>6</sup>Department of Musculoskeletal Biology, University of Liverpool, Liverpool, UK

## Key points

- Heavy-intensity exercise causes a progressive increase in energy demand that contributes to exercise limitation.
- This inefficiency arises within the locomotor muscles and is thought to be due to an increase in the ATP cost of power production; however, the responsible mechanism is unresolved.
- We measured whole-body  $\text{O}_2$  uptake and skeletal muscle ATP turnover by combined pulmonary gas exchange and magnetic resonance spectroscopy during moderate and heavy exercise in humans.
- Muscle ATP synthesis rate increased throughout constant-power heavy exercise, but this increase was unrelated to the progression of whole-body inefficiency.
- Our data indicate that the increased ATP requirement is not the sole cause of inefficiency during heavy exercise, and other mechanisms, such as increased  $\text{O}_2$  cost of ATP resynthesis, may contribute.

**Abstract** During constant-power high-intensity exercise, the expected increase in oxygen uptake ( $\dot{V}_{\text{O}_2}$ ) is supplemented by a  $\dot{V}_{\text{O}_2}$  slow component ( $\dot{V}_{\text{O}_{2\text{sc}}}$ ), reflecting reduced work efficiency, predominantly within the locomotor muscles. The intracellular source of inefficiency is postulated to be an increase in the ATP cost of power production (an increase in P/W). To test this hypothesis, we measured intramuscular ATP turnover with  $^{31}\text{P}$  magnetic resonance spectroscopy (MRS) and whole-body  $\dot{V}_{\text{O}_2}$  during moderate (MOD) and heavy (HVY) bilateral knee-extension exercise in healthy participants ( $n = 14$ ). Unlocalized  $^{31}\text{P}$  spectra were collected from the quadriceps throughout using a dual-tuned ( $^1\text{H}$  and  $^{31}\text{P}$ ) surface coil with a simple pulse-and-acquire sequence. Total ATP turnover rate ( $\text{ATP}_{\text{tot}}$ ) was estimated at exercise cessation from direct measurements of the dynamics of phosphocreatine (PCr) and proton handling. Between 3 and 8 min during MOD, there was no discernable  $\dot{V}_{\text{O}_{2\text{sc}}}$  (mean  $\pm$  SD,  $0.06 \pm 0.12 \text{ l min}^{-1}$ ) or change in [PCr] ( $30 \pm 8$  vs.  $32 \pm 7 \text{ mM}$ ) or  $\text{ATP}_{\text{tot}}$  ( $24 \pm 14$  vs.  $17 \pm 14 \text{ mM min}^{-1}$ ; each  $P = \text{n.s.}$ ). During HVY, the  $\dot{V}_{\text{O}_{2\text{sc}}}$  was  $0.37 \pm 0.16 \text{ l min}^{-1}$  ( $22 \pm 8\%$ ), [PCr] decreased ( $19 \pm 7$  vs.  $18 \pm 7 \text{ mM}$ , or  $12 \pm 15\%$ ;  $P < 0.05$ ) and  $\text{ATP}_{\text{tot}}$  increased ( $38 \pm 16$  vs.  $44 \pm 14 \text{ mM min}^{-1}$ , or  $26 \pm 30\%$ ;  $P < 0.05$ ) between 3 and 8 min. However, the increase in  $\text{ATP}_{\text{tot}}$  ( $\Delta\text{ATP}_{\text{tot}}$ ) was not correlated with the  $\dot{V}_{\text{O}_{2\text{sc}}}$  during HVY ( $r^2 = 0.06$ ;  $P = \text{n.s.}$ ). This lack of relationship between  $\Delta\text{ATP}_{\text{tot}}$  and  $\dot{V}_{\text{O}_{2\text{sc}}}$  together

with a steepening of the  $[\text{PCr}] - \dot{V}_{\text{O}_2}$  relationship in HVY, suggests that reduced work efficiency during heavy exercise arises from both contractile (P/W) and mitochondrial sources (the  $\text{O}_2$  cost of ATP resynthesis; P/O).

(Received 11 June 2014; accepted after revision 22 September 2014; first published online 3 October 2014)

**Corresponding author** H. B. Rossiter: Division of Respiratory & Critical Care Physiology & Medicine, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, 1124 W Carson Street, CDCRC Building, Torrance, CA 90502, USA. Email: hrossiter@ucla.edu

**Abbreviations** A, amplitude;  $\text{ATP}_{\text{tot}}$ , total ATP turnover rate; D, ATP production from phosphocreatine breakdown; k, rate constant; L, ATP production from glycogenolysis; LT, lactate threshold; MRS, magnetic resonance spectroscopy; PCr, phosphocreatine;  $\text{PCr}_{\text{sc}}$ , phosphocreatine slow component; pH<sub>i</sub>, intramuscular pH; P<sub>i</sub>, inorganic phosphate; P/O, ATP yield per  $\text{O} \rightarrow \text{H}_2\text{O}$ ; P/W, ATP cost per unit power output; Q, ATP production from oxidative phosphorylation; RI, ramp incremental;  $S_{\text{pO}_2}$ , arterial oxygenation;  $\tau$ , time constant;  $V_{[\text{PCr}]}$ , initial rate of phosphocreatine resynthesis;  $\dot{V}_{\text{O}_2}$ , rate of whole-body  $\text{O}_2$  uptake;  $\dot{V}_{\text{O}_{2\text{peak}}}$ , peak rate of  $\text{O}_2$  uptake;  $\dot{V}_{\text{O}_{2\text{sc}}}$ , slow component of  $\text{O}_2$  uptake.

## Introduction

During constant-power exercise below the lactate threshold (LT; moderate intensity), the rate of pulmonary oxygen uptake ( $\dot{V}_{\text{O}_2}$ ) increases exponentially, reaching a steady state within 2–3 min. A steady-state  $\dot{V}_{\text{O}_2}$  indicates that the exercise-related energy transfer is accounted for by oxidative phosphorylation. However, above the LT (heavy intensity), the dynamics of  $\dot{V}_{\text{O}_2}$  become complicated by an additional, slow component ( $\dot{V}_{\text{O}_{2\text{sc}}}$ ; Poole *et al.* 1994). This becomes especially important at power outputs above critical power, where the  $\dot{V}_{\text{O}_{2\text{sc}}}$  will draw  $\dot{V}_{\text{O}_2}$  inexorably towards its physiological maximum. In this intensity domain, the limit of tolerance is reached rapidly, and the exercise cannot continue unless the power output is reduced below critical power (Coats *et al.* 2003). Although the  $\dot{V}_{\text{O}_{2\text{sc}}}$  is intimately related to exercise intolerance (Murgatroyd *et al.* 2011), the aetiology of the  $\dot{V}_{\text{O}_{2\text{sc}}}$  remains poorly understood.

The  $\dot{V}_{\text{O}_{2\text{sc}}}$  represents an impairment of exercise economy and is predominantly (~85%) due to increased  $\text{O}_2$  consumption in the muscles engaged in the locomotor work (Poole *et al.* 1991; Rossiter *et al.* 2002; Krusturp *et al.* 2009). However, the intracellular source of this inefficiency is uncertain. It has been postulated that the  $\dot{V}_{\text{O}_{2\text{sc}}}$  is related to an increased phosphate cost of force or power production; that is, an increase in the rate of ATP consumption per unit power output (or P/W) is met instantaneously by phosphocreatine (PCr; via the Lohmann reaction), the breakdown of which signals an increase in the rate of oxidative phosphorylation (Rossiter *et al.* 2002). However, distinguishing between this and the alternative hypothesis, that supra-LT exercise is associated with reductions in mitochondrial coupling (Krusturp *et al.* 2003), i.e. the ratio of the ATP resynthesized per oxygen converted to water (P/O), is technically challenging in humans.

To test these two hypotheses requires knowledge of dynamic changes in total ATP turnover rate ( $\text{ATP}_{\text{tot}}$ ) in

concert with power output and  $\dot{V}_{\text{O}_2}$ . Specifically, were the intramuscular source of the  $\dot{V}_{\text{O}_{2\text{sc}}}$  to be caused by an increase in P/W (in line with current views; Rossiter, 2011; Poole & Jones, 2012), then the magnitude of the  $\dot{V}_{\text{O}_{2\text{sc}}}$  during heavy exercise would be strongly related to the magnitude of the change in  $\text{ATP}_{\text{tot}}$ . Alternatively, if no proportionality between the  $\dot{V}_{\text{O}_{2\text{sc}}}$  and the change in  $\text{ATP}_{\text{tot}}$  were evident, then changes in P/W could not be the sole source of the  $\dot{V}_{\text{O}_{2\text{sc}}}$ .

The technical challenge thus becomes, how best to establish  $\text{ATP}_{\text{tot}}$  during heavy-intensity exercise that elicits a  $\dot{V}_{\text{O}_{2\text{sc}}}$ ? One approach uses  $^{31}\text{P}$  magnetic resonance spectroscopy (MRS; Kemp *et al.* 2001; Layec *et al.* 2009a) to partition ATP delivery from oxidative phosphorylation, PCr breakdown and glycogenolysis.  $^{31}\text{P}$  MRS provides direct measurement of  $[\text{PCr}]$  and allows the glycolytic rate (a relatively minor component of  $\text{ATP}_{\text{tot}}$  in exercise of this kind) to be estimated using reasonable assumptions about muscle  $\text{H}^+$  buffering (Kemp *et al.* 2001, 2014). Several methods have been proposed to calculate oxidative ATP yield using  $^{31}\text{P}$  MRS, but these show poor agreement (Layec *et al.* 2011). Previous studies to estimate  $\text{ATP}_{\text{tot}}$  during supra-LT exercise have assumed a linear  $\dot{V}_{\text{O}_2} - [\text{PCr}]$  relationship and a fixed time constant ( $\tau$ ) of PCr breakdown and resynthesis (Meyer, 1988; Walter *et al.* 1999; Lanza *et al.* 2005; Faraut *et al.* 2007) or first-order  $[\text{ADP}] - \dot{V}_{\text{O}_2}$  relationship in order to transform  $[\text{PCr}]$  into a rate of oxidative ATP turnover (Layec *et al.* 2009a). However, it is clear that the  $\dot{V}_{\text{O}_2} - [\text{PCr}]$  relationship is not linear through the intensity domains (Kemp, 2008; Wüst *et al.* 2011; Kemp *et al.* 2014), and accordingly,  $\tau\text{PCr}$  is not invariant across exercise intensities (Yoshida & Watari, 1993, 1994; Rossiter *et al.* 2002; Jones *et al.* 2008), making this an unreliable assumption on which to base estimation of  $\text{ATP}_{\text{tot}}$ . Assuming  $\tau\text{PCr}$  to be invariant is equivalent to assuming that any change in  $[\text{PCr}]$  is directly proportional to change in  $\text{ATP}_{\text{tot}}$ ; when  $\tau\text{PCr}$  changes across exercise intensity and/or duration, this proportionality is lost (Kemp *et al.* 2014). These new findings mean that the

close coherence between [PCr] and  $\dot{V}_{O_2}$  during the slow component phase (Rossiter *et al.* 2002; Layec *et al.* 2009a) is no longer sufficient evidence to imply that an increase in P/W alone is the responsible mechanism. Consequently, a direct measurement of oxidative ATP yield during supra-LT exercise that does not rely on these assumptions is required to distinguish whether change in P/W is the dominant mechanism for the  $\dot{V}_{O_{2sc}}$ .

Oxidative ATP turnover (the dominant proportion of  $ATP_{tot}$ ) at exercise cessation may be assessed directly from the initial rate of postexercise PCr resynthesis ( $V_{i[PCr]}$ ), easily measured by  $^{31}P$  MRS; the only assumptions required (the evidence for which is reviewed elsewhere; Kemp *et al.* 2014) are that PCr recovery is driven overwhelmingly by oxidative ATP synthesis and that any basal component of ATP turnover (i.e. ATP production not available for use by myosin ATPase, sarco(endo)plasmic reticulum  $Ca^{2+}$ -ATPase or  $Na^+$ - $K^+$ -ATPase during exercise or PCr resynthesis during recovery) is small and reasonably constant. Therefore, temporal characterization of oxidative energy yield during dynamic exercise can be made simply by halting the exercise and measuring  $V_{i[PCr]}$ . Although this method has inherently poor temporal resolution (it is valid only at the instant of exercise cessation), it provides the accuracy necessary to isolate the intracellular source of inefficiency during high-intensity exercise. The other, much smaller, components of  $ATP_{tot}$  can be estimated at the end of exercise by  $^{31}P$  MRS in ways that are relatively robust against uncertainty or changes in the underpinning assumptions.

The purpose of this study, therefore, was to characterize the rate of ATP turnover during sub- and supra-LT exercise in human quadriceps during bilateral, prone, knee-extension exercise using  $^{31}P$  MRS. The rate of pulmonary oxygen uptake was measured in the same conditions to quantify the  $\dot{V}_{O_{2sc}}$ . We hypothesized that the close association between the dynamics of the [PCr] and  $\dot{V}_{O_2}$  slow components during supra-LT exercise would be reflected in the dynamics of  $ATP_{tot}$  (measured independently), thereby confirming the hypothesis that increased P/W during heavy-intensity exercise is the predominant mechanism of the  $\dot{V}_{O_{2sc}}$ .

## Methods

### Ethical approval

The Biological Sciences Faculty Research Ethics Committee, University of Leeds, and the University of Liverpool Committee on Research Ethics approved this study, and all procedures complied with the latest revision of the *Declaration of Helsinki*. Written informed consent was obtained from all volunteers prior to their participation in the study.

### Participants

Fourteen healthy volunteers (one female, 13 male) agreed to participate in this study [mean  $\pm$  SD: age  $27 \pm 8$  years; height  $177 \pm 8$  cm; mass  $75 \pm 12$  kg; bilateral knee-extension peak  $\dot{V}_{O_2}$  ( $\dot{V}_{O_{2peak}}$ )  $2.0 \pm 0.5$  l  $min^{-1}$ ]. All participants were undertaking a regular exercise regimen, ranging from recreational fitness to amateur competitive sport. Volunteers were screened for cardiovascular disease risk with a resting ECG and a health history questionnaire.

### Exercise protocols

All exercise tests were undertaken on an MR-compatible computer-controlled electromagnetically braked knee-extension ergometer (MRI Ergometer Up/Down, Lode BV, Groningen, The Netherlands) customized for use at 3 T by the addition of extended carbon-fibre lever arms. The participants lay prone, with their feet strapped into moulded plastic stirrups, which were attached to carbon-fibre/aluminium arms, linking to the ergometer crank arms. The participants' hips were secured to the patient bed with nylon and Velcro straps in order to isolate power production to the quadriceps and minimize movement from hip flexion/extension. Knee movements were constrained by the scanner bore, allowing for  $\sim 35$  deg of bilateral knee extension (Whipp *et al.* 1999; Cannon *et al.* 2013). No resistance was applied during knee flexion, other than the constant work required to lift the mass of the lower leg.

The testing protocol began with a rigorous familiarization phase that took place in a temperature-controlled laboratory with pulmonary gas exchange measurements. Ramp incremental (RI) and constant-power protocols were completed until reproducible physiological measurements were obtained across two consecutive visits for each condition. The second phase of the study took place within the bore of an MR scanner for measurement of muscle phosphates. The same MRI ergometer was used for both phases of the protocol.

Initially, participants completed an RI exercise test to the limit of tolerance. For this, participants lay at rest for  $\sim 3$ – $4$  min, followed by a low-power exercise (5 W) for  $\sim 2$ – $4$  min. The power was then increased as a function of time at  $2$ – $5$  W  $min^{-1}$  (the rate of increase was dependent on the volunteer's size and strength) until the limit of tolerance was reached. Ramp rates were adjusted using 'trial and error' to determine a ramp rate that resulted in a ramp duration of  $\sim 10$ – $12$  min. The frequency of knee extension was constrained at  $90$   $min^{-1}$  with the use of a metronome. This cadence was chosen to allow synchronization with the MR scanner acquisitions (one pulse per two knee extensions) and also acted to ensure that the ergometer flywheel was maintained above its minimal operating speed. The RI was terminated upon

the participant being unable to maintain the required cadence, despite strong verbal encouragement. The results of the RI were used to determine the  $\dot{V}_{O_{2peak}}$  and to calculate power for subsequent tests. There is a substantial learning effect with the exercise model (large gains in peak power were achieved with consecutive tests), and therefore typically, more than three RI tests were completed by each participant until reproducible performances were achieved.

A series of constant-power exercise tests were then undertaken. These consisted of an 8 min moderate-intensity bout, followed by a 6 min rest and an 8 min heavy-intensity exercise bout. During moderate-intensity exercise, the target power was 80% of estimated LT (LT was  $\sim 60\text{--}70\%$   $\dot{V}_{O_{2peak}}$ ), and during heavy-intensity bouts the target power was halfway between estimated LT and  $\dot{V}_{O_{2peak}}$ . These intensity domains were confirmed *post hoc* from the profile of  $\dot{V}_{O_2}$  during constant-power bouts (Whipp, 1996). If necessary, power was adjusted in subsequent familiarization tests to ensure the absence (moderate) or presence (heavy) of the  $\dot{V}_{O_{2sc}}$ . Once familiarized, participants repeated this protocol three times on separate days to combine respired gas exchange data and improve the signal-to-noise ratio.

During the second phase of experiments, participants completed constant-power bouts within the bore of the superconducting magnet for  $^{31}\text{P}$  MRS. Two trials of constant-power tests were completed in a random order consisting of the following sequences: (i) 4 min of rest, followed by 3 min of moderate exercise, 6 min of rest and 3 min of heavy exercise; and (ii) 4 min of rest, followed by 8 min of moderate exercise, 6 min of rest and 8 min of heavy exercise. Each protocol was preceded by  $\sim 10$  min of magnet shimming to optimize the MRS signal, and separated by at least 30 min outside of the MR scanner. Therefore,  $\sim 60\text{--}90$  min elapsed between the two exercise trials.

### Pulmonary gas exchange

Participants breathed through a low-resistance ( $<0.1$  kPa  $\text{l}^{-1} \text{s}^{-1}$  up to  $15 \text{ l s}^{-1}$ ), low-dead-space (90 ml) mouthpiece for the measurement of respired gases. Flow rates and volumes were measured with an infrared turbine flow sensor (Interface Associates, Laguna Niguel, CA, USA), while a quadrupole mass spectrometer was used to measure respired gas concentrations after sampling air at  $0.5 \text{ ml s}^{-1}$  from the mouthpiece (MSX; nSpire Health Ltd, Hertford, UK). Gas concentration signals were time aligned with the flow sensor signal using proprietary software for the calculation of breath-by-breath gas exchange. These algorithms identified the end of each breath with the flow sensor and time aligned the changes in respired gases.

Prior to each experiment, the flow sensor and gas analysers were calibrated according to the manufacturers' instructions. The turbine volume transducers were calibrated with a 3 l syringe (Hans Rudolph Inc., Shawnee, KS, USA). The calibration was completed with flow rates ranging from  $0.2$  to  $6 \text{ l s}^{-1}$ , mimicking flow rates expected for humans at rest and during exercise. After the completion of the flow sensor calibration, the flow volumes were verified over 10 syringe strokes of varying flow rates and accepted when the means were within  $\pm 0.01 \text{ l}$ , with an SD and coefficient of variation of  $0.02 \text{ l}$  and  $1\%$ , respectively. Additionally, the mass spectrometer was calibrated with atmospheric air and precision-verified gases with concentrations of  $\text{O}_2$ ,  $\text{CO}_2$  and  $\text{N}_2$  spanning the physiological range. Following each experiment, mass spectrometer calibration factor drift was verified as negligible by sampling the calibration gases.

**Data analyses for pulmonary measures.** Breath-by-breath  $\dot{V}_{O_2}$  was filtered for errant breaths (i.e. values resulting after sighs, swallows, coughs etc., defined as residing outside of 99% prediction limits; Lamarra *et al.* 1987). Responses from like transitions were combined to improve the signal-to-noise ratio using an averaging technique that preserves the breath-by-breath density measured during the exercise transition. This method requires time aligning and sorting of all  $\dot{V}_{O_2}$  data from exercise transitions in the time domain. Time and  $\dot{V}_{O_2}$  are then averaged into bins of  $n$  breaths, where  $n$  is the number of exercise transitions completed (Murgatroyd *et al.* 2011). The magnitude of the  $\dot{V}_{O_{2sc}}$  was expressed as the difference in  $\dot{V}_{O_2}$  between 3 and 8 min of exercise.

Power output and flywheel speed from the ergometer were sampled continuously and digitized by a data-recording system and stored on a PC (PowerLab 8/30 with LabChart Pro; ADInstruments Pty Ltd, Bella Vista, NSW, Australia).

### $^{31}\text{P}$ Magnetic resonance spectroscopy

Muscle phosphorus-containing metabolites were measured with a 3 T superconducting magnet (Magnetom Trio; Siemens AG, Erlangen, Germany). A one-pulse MRS acquisition was employed using a dual-tuned ( $^1\text{H}$  and  $^{31}\text{P}$ ) 15- and 18-cm-diameter surface RF coil (RAPID Biomedical GmbH, Rimpfing, Germany), which was placed under the knee extensors, halfway between the hip and knee. The concave RF coil was stabilized with sandbags and was secured to the table once the participants' hips were strapped to the scanner table. A series of axial, sagittal and coronal gradient-recalled echo images of the thigh were acquired to confirm the placement of the RF coil relative to the knee-extensor muscles and to prescribe the volume over which shimming was achieved.

Subsequently, a standard  $^1\text{H}$  shimming protocol was used to optimize the homogeneity of the magnetic field ( $\beta_0$ ). A fully relaxed spectrum (repetition time of 10 s; number of scans = 4) was initially obtained to provide a high-resolution unsaturated resting spectrum along with a 32 scan spectrum with a repetition time of 2 s. Following this, free induction decays for  $^{31}\text{P}$  spectra were collected every 2 s with a spectral width of 3200 Hz and 1024 data points throughout the rest-to-exercise-to-rest transitions. The  $^{31}\text{P}$  data were averaged over four acquisitions, yielding a datum every 8 s.

### Kinetic analysis of $^{31}\text{P}$ MRS data

Signal intensities, frequencies and line widths of inorganic phosphate ( $\text{P}_i$ ), PCr,  $\gamma$ -ATP,  $\alpha$ -ATP and  $\beta$ -ATP were determined using Java-based Magnetic Resonance User Interface (jMRUI; Naressi *et al.* 2001) in order to transform the raw data into a time series for each of the phosphates of interest. Intramuscular pH ( $\text{pH}_i$ ) was estimated from the chemical shift of  $\text{P}_i$  (Moon & Richards, 1973), as follows:

$$\text{pH}_i = 6.75 + \log(\delta - 3.27/5.69 - \delta) \quad (1)$$

where  $\delta$  is the chemical shift of the  $\text{P}_i$  peak, relative to PCr.

Phosphocreatine kinetics were modelled using non-linear least-squares regression (OriginPro 7.5; OriginLab Corp., Northampton, MA, USA). The  $^{31}\text{P}$  MRS data were filtered for errant values resulting from artefacts (Rossiter *et al.* 2000) prior to characterization with the following function:

$$[\text{PCr}]_{(t)} = [\text{PCr}]_0 + A(1 - e^{-t/\tau}) \quad (2)$$

where  $\tau$  is a time constant and  $[\text{PCr}]_{(t)}$ ,  $[\text{PCr}]_0$ , and  $A$  are the time-variant form, baseline and fundamental amplitude, respectively. The fitting window was determined from an iterative process (Rossiter *et al.* 2001) to ensure the exclusion of phase III (steady state or slow component, depending on the intensity domain). The magnitude of the PCr slow component ( $[\text{PCr}]_{sc}$ ) was expressed as the difference in  $[\text{PCr}]$  between the third and eighth minute of exercise.

The  $\text{ATP}_{\text{tot}}$  was estimated from the contributions from oxidative phosphorylation ( $Q$ ), PCr breakdown ( $D$ ) and glycogenolysis ( $L$ ), which were determined from the  $^{31}\text{P}$  MRS data acquired during exercise and recovery, using methods described elsewhere (Kemp *et al.* 2001, 2007, 2014; Layec *et al.* 2011) and outlined below.

**Production of ATP from PCr breakdown ( $D$ ).** The rate of PCr breakdown by creatine kinase ( $D$ ) yields one component of ATP production (millimolar per minute) and was determined over 32 s (four spectra) immediately

prior to exercise cessation, according to the following equation:

$$D = d[\text{PCr}]/dt \quad (3)$$

In the present experiments, where  $[\text{PCr}]$  is either close to steady state or changing only slowly by the end of exercise,  $D$  is a very small component of end-exercise  $\text{ATP}_{\text{tot}}$ .

### Production of ATP from oxidative phosphorylation ( $Q$ ).

The rate of oxidative ATP yield ( $Q$ ) is reflected in the rate of  $[\text{PCr}]$  recovery at the instant of exercise cessation ( $V_{i[\text{PCr}]}$ ) and was calculated (millimolar per minute) as follows:

$$V_{i[\text{PCr}]} = kA \quad (4)$$

where  $A$  is the amplitude of  $[\text{PCr}]$  change (millimolar). The rate constant ( $k$ ) was estimated by fitting the PCr recovery kinetics with the following function:

$$[\text{PCr}]_{(t)} = [\text{PCr}]_{\text{end}} - A(1 - e^{-kt}) \quad (5)$$

where  $[\text{PCr}]_{(t)}$  is the time-dependent variant of  $[\text{PCr}]$ , and  $[\text{PCr}]_{\text{end}}$  is the concentration of PCr measured at the end of exercise. We make the well-evidenced assumption (Kemp *et al.* 2014) that the rate of suprabasal oxidative synthesis at the start of recovery [ $V_{i[\text{PCr}]}$  from eqn (4)] is a good estimate of the suprabasal rate of oxidative synthesis at the end of exercise ( $Q_{\text{end}}$ ).

### Production of ATP from anaerobic glycolysis ( $L$ ).

During exercise, glycogenolysis and the resulting lactate and  $\text{H}^+$  production cause disturbances in  $\text{pH}_i$ . These changes in  $\text{pH}_i$  are readily measured by  $^{31}\text{P}$  MRS data and can therefore be used to estimate ATP production from glycogenolysis; 1 mol of  $\text{H}^+$  resulting in 1.5 mol of ATP. This requires estimation of the flux rates as follows:  $\text{H}^+$  production accompanying changes in PCr concentration via the creatine kinase reaction ( $\text{H}_{\text{CK}}^+$ , which is positive, i.e.  $\text{H}^+$  'consumption', when  $[\text{PCr}]$  is falling in exercise, and negative, i.e.  $\text{H}^+$  generation, when  $[\text{PCr}]$  is rising in recovery); by the buffers of the muscle cytosol ( $\text{H}_{\beta}^+$ , which is positive, i.e.  $\text{H}^+$  'buffering' when  $\text{pH}_i$  is falling in exercise and negative, i.e.  $\text{H}^+$  'unbuffering' when  $\text{pH}_i$  is rising in recovery); and proton efflux from the cells ( $\text{H}_{\text{efflux}}^+$ ). Together, these sum to the total proton yield ( $P$ ) during exercise:

$$P = \text{H}_{\text{CK}}^+ + \text{H}_{\beta}^+ + \text{H}_{\text{efflux}}^+ \quad (6)$$

From which:

$$L = 1.5 \times P \quad (7)$$

The number of protons consumed at the creatine kinase reaction was calculated from the time-dependent changes

in [PCr] using the proton stoichiometric coefficient,  $\gamma$  (Kushmerick, 1997), as follows:

$$H_{CK}^+ = -\gamma D \quad (8)$$

Protons buffered ( $H_{\beta}^+$ , millimolar per minute) was calculated from the apparent buffering capacity,  $\beta_{total}$  (in millimoles of acid added per unit change in  $pH_i$ ) and from the (smoothed) rate of pH change during exercise, as follows:

$$H_{\beta}^+ = -\beta_{total} (dpH_i/dt) \quad (9)$$

where

$$\beta_{total} = \beta_{non-bicarbonate-non-P_i} + \beta_{P_i} \quad (10)$$

The intrinsic cytosolic buffering capacity ( $\beta_{non-bicarbonate-non-P_i}$ ) is calculated from initial-exercise data:

$$\beta_{non-bicarbonate-non-P_i} = \beta_a - \beta_{P_i} \quad (11)$$

where the apparent  $\beta$  ( $\beta_a$ ) is obtained from the initial rate of change in [PCr] ( $\Delta PCr_i$ ) and alkalinization of pH ( $\Delta pH_i$ ):

$$\beta_a = \gamma \Delta PCr_i / \Delta pH_i \quad (12)$$

The value of  $\beta_{P_i}$  was calculated as follows:

$$\beta_X = (2.303 \times H^+ \times K \times [P_i]) / (K + H^+)^2 \quad (13)$$

where  $K = 1.77 \times 10^{-7}$  (Conley *et al.* 1998). The  $\beta_{bicarbonate}$  was neglected, which assumes that muscle is a closed system during short-duration exercise *in vivo* (Kemp *et al.* 1993). Proton efflux ( $H_{efflux}^+$ , millimolar per minute) was estimated for each time point of exercise assuming a linear pH-dependence constant,  $\lambda$ , as follows:

$$H_{efflux}^+ = -\lambda \Delta pH_i \quad (14)$$

This proportionality constant,  $\lambda$  (millimolar per minute per pH unit) was estimated from initial recovery after exercise cessation, as follows:

$$\lambda = -V_{eff} / \Delta pH_i \quad (15)$$

At the cessation of exercise, the PCr resynthesized in the creatine kinase reaction is essentially a consequence solely of oxidative ATP production (Kemp *et al.* 2014). Therefore,  $H_{efflux}^+$  can be calculated from the rate of proton production from creatine kinase ( $H_{CK}^+$ ) and the rate of pH change on the other side, as follows:

$$V_{eff} = \beta_{total} (dpH_i/dt) = \gamma V_{i,PCr} \quad (16)$$

Where  $\Delta pH_i$  is very low, eqn (14) becomes unreliable, and the end-exercise rate of  $H_{efflux}^+$  is simply assumed to be equal to  $H_{efflux}^+$  calculated in from the initial recovery data by eqn (16).

In the present experiments, where  $pH_i$  is close to steady state or changing only slowly by the end of exercise,  $L$  is a very small component of  $ATP_{tot}$ .

### Statistical analyses

Relationships between variables were assessed with a Pearson correlation coefficient, where appropriate. The differences between  $^{31}P$  measures at discrete time points and across exercise intensities were compared with a two-factor (time  $\times$  intensity domain) repeated-measures ANOVA. Bonferroni-corrected Student's paired *t* tests were used *post hoc* to identify simple effects in the case of a significant main effect. For all tests,  $\alpha = 0.05$ . Analyses were completed using the Statistical Package for the Social Sciences (SPSS Inc., Chicago, IL, USA).


### Results

During RI exercise, participants attained a peak power output of  $47 \pm 11$  W and a  $\dot{V}_{O_{2peak}}$  of  $2.00 \pm 0.48$  l  $min^{-1}$ . Based on peak power output and estimated LT ( $\sim 60$ – $70\%$   $\dot{V}_{O_{2peak}}$ ), moderate (sub-LT;  $19 \pm 4$  W) and heavy (supra-LT;  $46 \pm 11$  W) constant-power exercise bouts were assigned. The dynamics of  $\dot{V}_{O_2}$  were examined *post hoc* to confirm the appropriate intensity assignment (Whipp, 1996; Rossiter, 2011).

During moderate exercise, there was no discernable pulmonary  $\dot{V}_{O_{2sc}}$  ( $0.06 \pm 0.12$  l  $min^{-1}$ ). However, during heavy exercise the  $\dot{V}_{O_{2sc}}$  was  $0.37 \pm 0.16$  l  $min^{-1}$  (Fig. 1A), or a  $22 \pm 8\%$  increase. The [PCr] did not change between 3 and 8 min of moderate-intensity exercise ( $30 \pm 8$  vs.  $32 \pm 7$  mM; n.s.). Conversely, during heavy exercise [PCr] fell from 3 to 8 min ( $19 \pm 7$  vs.  $18 \pm 7$  mM or  $12 \pm 15\%$  fall;  $P < 0.05$ ; Fig. 1B).

The ATP yield during moderate and heavy exercise from oxidative phosphorylation ( $Q$ ), PCr hydrolysis ( $D$ ), lactate production ( $L$ ) and, consequently,  $ATP_{tot}$  are presented in Table 1. The  $V_{i[PCr]}$ , calculated as described in eqn (4) is shown, along with the rate constant of PCr resynthesis ( $k$ ) and the amplitude of PCr recovery ( $A$ ), in Figs 2 (moderate) and 3 (heavy).

Comparisons of  $ATP_{tot}$  revealed a significant interaction (time  $\times$  intensity domain;  $F(1,13) = 17.2$ ;  $P < 0.01$ ;  $\eta^2 = 0.57$ ). The  $ATP_{tot}$  was not different between 3 and 8 min of moderate exercise (n.s.; Fig. 2 and Table 1), but  $ATP_{tot}$  increased ( $\Delta ATP_{tot}$ ) during heavy exercise from 3 to 8 min ( $CI_{95}$  of the difference;  $CI_{Difference}$  1.9, 12.6 mM  $min^{-1}$ ;  $P < 0.05$ ; Fig. 3 and Table 1), equating to a  $26 \pm 30\%$  increase in  $ATP_{tot}$  from 3 to 8 min (Fig. 4A). This percentage change in  $ATP_{tot}$  was not different to that measured in both the  $[PCr]_{sc}$  and the  $\dot{V}_{O_{2sc}}$  ( $F(2,26) = 2.4$ ; n.s.;  $\eta^2 = 0.16$ ; Fig. 4A). However, among participants the individual values of  $\Delta ATP_{tot}$  during heavy exercise were



**Figure 1.** Rate of whole-body  $\dot{V}_{O_2}$  (A), phosphocreatine concentration ([PCr]; B) and pH (C) plotted as a function of time for moderate-intensity (filled circles) and heavy-intensity prone bilateral knee-extension exercise (open circles)

Black bar denotes exercise bout from time 0 to 8 min. Data points are 8 s means, with error bars representing SD.

**Table 1.** Rates of ATP turnover from oxidative phosphorylation (Q), phosphocreatine hydrolysis (D), lactate production (L) and the sum (ATP<sub>tot</sub>) during moderate and heavy constant-power exercise at two time points

Parameter	Moderate exercise		Heavy exercise	
	3 min	8 min	3 min	8 min
Q (mmol min <sup>-1</sup> ) <sup>†</sup>	23 (14)	17 (13)	35 (17)	42 (13)*
D (mmol min <sup>-1</sup> )	0.6 (1.2)	0.2 (1.0)	1.1 (2.6)	0.7 (0.9)
L (mmol min <sup>-1</sup> )	1.0 (1.3)	0.3 (0.6)	1.5 (1.3)	1.3 (1.7)
ATP <sub>tot</sub> (mmol min <sup>-1</sup> ) <sup>‡</sup>	24 (14)	17 (14)	38 (16)	44 (14)*

Values are presented in millimolar per minute as means (SD).

\*Different from 3 min;  $P < 0.05$ . <sup>†</sup>Time  $\times$  intensity interaction;  $P < 0.01$ ;  $F(1,13) = 17.2$ ;  $r^2 = 0.57$ . <sup>‡</sup>Time  $\times$  intensity interaction;  $P < 0.01$ ;  $F(1,13) = 17.2$ ;  $r^2 = 0.57$

not significantly correlated with the magnitude of the  $\dot{V}_{O_{2sc}}$  (Fig. 4B).

To examine the relationship between  $\dot{V}_{O_2}$  and [PCr], a correction for the transit delay from muscle to lung was applied. The  $\dot{V}_{O_2}$  data were time corrected using 12 s difference with respect to <sup>31</sup>P measures (Rossiter *et al.* 1999; Krstrup *et al.* 2009). The relationship between  $\dot{V}_{O_2}$  and [PCr] was linear during moderate exercise and the first 3 min of heavy exercise ( $r^2 = 0.94$ ; Fig. 5). However, the slope of the [PCr]– $\dot{V}_{O_2}$  relationship was significantly steeper when data from 8 min of heavy exercise were included ( $-67 \pm 25$  vs.  $-61 \pm 25$  ml min mmol<sup>-1</sup>;  $P < 0.05$ ).

## Discussion

The [PCr] slow component ([PCr]<sub>sc</sub>), like the  $\dot{V}_{O_{2sc}}$ , is present only during exercise above the LT. The finding that the [PCr]<sub>sc</sub> and  $\dot{V}_{O_{2sc}}$  are of similar magnitude (Rossiter *et al.* 2002) led to the argument that the  $\dot{V}_{O_{2sc}}$  is caused by an increased phosphate cost of power production (P/W) during heavy-intensity exercise. However, this is at odds with the observed dissociation between the [PCr]<sub>sc</sub> and  $\dot{V}_{O_{2sc}}$  in endurance-trained individuals (Layec *et al.* 2009b, 2012), and both observations relied upon equivocal assumptions about the dynamic relationships between [ADP] and  $\dot{V}_{O_2}$  or  $\dot{V}_{O_2}$  and [PCr] (Yoshida & Watari, 1993, 1994; Rossiter *et al.* 2002; Jones *et al.* 2008; Kemp, 2008; Wüst *et al.* 2011). Our present data agree with previous reports that mean [PCr]<sub>sc</sub> and  $\dot{V}_{O_{2sc}}$  magnitudes were not statistically different. Crucially, however, the data add that, among individuals, the increase in the  $\dot{V}_{O_{2sc}}$  during heavy-intensity exercise (averaging  $\sim 22\%$ ) is not correlated with the increase in the phosphate cost of power production, ATP<sub>tot</sub> (average  $\sim 26\%$ ). Thus, while the exercising limb is likely to remain the major source of the  $\dot{V}_{O_{2sc}}$  (Poole *et al.* 1991; Rossiter *et al.* 2002; Bailey *et al.* 2010; Dimenna *et al.* 2010), the observed dissociation

between  $\dot{V}_{O_{2sc}}$  and  $\Delta ATP_{tot}$  (Fig. 4B) strongly suggests that the progressive increase in  $\dot{V}_{O_2}$  during heavy exercise is not solely due to contractile inefficiency (P/W). Thus, other explanations, such as a reduction in mitochondrial efficiency (P/O), should also be considered.


### ATP turnover during moderate and heavy constant-power exercise

The primary aim of this investigation was to estimate the ATP turnover rate for exercise below and above the LT and over time without assumptions about the  $[ADP]-\dot{V}_{O_2}$  or  $\dot{V}_{O_2}-[PCr]$  relationships. By using the most robust estimations of  $ATP_{tot}$  (Kemp *et al.* 1995; Walter *et al.* 1999; Lanza *et al.* 2005; Faraut *et al.* 2007), we provided  $^{31}P$  MRS-derived estimates of ATP yield from oxidative phosphorylation, lactate production and PCr hydrolysis at 3 and 8 min of exercise that were unencumbered by the recently challenged assumptions about the  $[ADP]-\dot{V}_{O_2}$  relationship (Kemp, 2008; Wüst *et al.* 2011; Glancy & Balaban, 2012; Kemp *et al.* 2014).

Unsurprisingly, there were no changes in  $ATP_{tot}$  during exercise below the lactate threshold, where negligible muscle fatigue is expected (Sargeant & Dolan, 1987; Yano

*et al.* 2001), reflecting steady-state conditions. Conversely, during heavy exercise in which the  $\dot{V}_{O_{2sc}}$  and  $[PCr]_{sc}$  were present,  $ATP_{tot}$  was increased between 3 and 8 min of exercise. This is consistent with the suggestions that the  $\dot{V}_{O_{2sc}}$  is consequent to increased P/W in the large locomotor muscles during supra-LT exercise (Rossiter *et al.* 2002), perhaps associated with muscle fatigue and a reduction in contractile efficiency. However, the lack of relationship between  $\Delta ATP_{tot}$  and  $\dot{V}_{O_{2sc}}$  is in contrast to this postulate and challenges the current understanding of the aetiology of  $\dot{V}_{O_{2sc}}$  (Rossiter, 2011; Poole & Jones, 2012).

Dissociation of the  $\dot{V}_{O_{2sc}}$  and changes in the phosphate cost of exercise may have a few explanations. It may indicate an increase in  $\dot{V}_{O_2}$  originating from regions within the knee extensors that are not interrogated by the surface coil. While we can only speculate on this, a similar finding has been reported where the  $\dot{V}_{O_2}$  and  $[PCr]$  slow components were dissociated in endurance-trained participants but not in sedentary control subjects, despite increasing EMG activity in both participant groups during the  $\dot{V}_{O_{2sc}}$  (Layec *et al.* 2009b, 2012). It was hypothesized that the exercise-trained volunteers may be better able to optimize motor unit recruitment patterns to maintain high-intensity exercise (e.g. compared with active but



**Figure 2.** Moderate-intensity exercise recovery rate constant ( $k$ ; A), amplitude of PCr resynthesis (termed 'A'; B), initial rate of PCr resynthesis ( $V_{i[PCr]}$ ; C) and total ATP turnover rate ( $ATP_{tot}$ ; D) at 8 min of exercise, plotted as a function of 3 min of exercise. Dashed line is  $y = x$ .



untrained subjects; Rossiter *et al.* 2002), thereby recruiting motor unit pools that reside outside of the muscle volume being interrogated by MRS. It should be noted, however, that our surface coil interrogated a large muscle volume (~300 g) compared with alternative techniques, e.g. biopsy (~200 mg). Additionally, controversy exists about whether progressive recruitment itself is even responsible for the slow component (Zoladz *et al.* 2008; Cannon *et al.* 2011; Vanhatalo *et al.* 2011), in which case recruitment of muscle outside the surface coil view would seem to be an unlikely explanation if the recruitment pattern is stable.


The source of the  $\dot{V}_{O_{2sc}}$  may even reside outside of the locomotor muscles. Progressive increases in respiratory (Wasserman *et al.* 1995; Żołądź & Korzeniewski, 2001) or cardiac work, or even work from non-power-producing musculature, such as stabilizing effort during cycling (Billat *et al.* 1998), may contribute to a reduction in exercise efficiency during the slow component. It is unlikely that the stabilizing effort would contribute to prone knee extension, where the work of stabilizing the torso is minimized by the body position, the ergometer and the heavy strapping used to isolate quadriceps activity. Nonetheless, the work of ventilation during prone knee extension may still contribute a meaningful proportion, particularly as the locomotor muscle mass in

our study is relatively small in comparison to cycling or running.

Finally, dissociation of the  $\dot{V}_{O_{2sc}}$  and  $\Delta ATP_{tot}$  could result from mitochondrial uncoupling (reduced P/O; Fig. 5). In this scenario, an increased  $O_2$  cost of ATP resynthesis may contribute to driving the increase in  $\dot{V}_{O_2}$  during heavy exercise, rather than it coming exclusively from an increased ATP cost of muscle power generation.

### The $\dot{V}_{O_2}$ -[PCr] relationship and mitochondrial coupling during heavy-intensity exercise


Without an invasive measure of  $\dot{V}_{O_2}$  across the volume of tissue interrogated by MRS, the relationship between whole-body  $\dot{V}_{O_2}$  and localized [PCr] is the next best estimate for coupling of  $O_2$  uptake and ATP turnover. Our data show that the mean  $\dot{V}_{O_2}$ -[PCr] relationship was linear over the moderate intensity and during the first minutes of heavy exercise ( $r^2 = 0.94$ ; Fig. 5). Importantly, this relationship became steeper ( $P < 0.05$ ) with the inclusion of data from the final minutes of heavy exercise. With some important assumptions, these data suggest a reduced P/O between 3 and 8 min of heavy exercise, implicating mitochondrial uncoupling as an additional mechanism of the  $\dot{V}_{O_{2sc}}$ .



**Figure 3.** Heavy-intensity exercise recovery rate constant ( $k$ ; A), amplitude of PCr resynthesis (termed 'A'; B),  $V_{i[PCr]}$  (C) and  $ATP_{tot}$  (D) at 8 min of exercise, plotted as a function of 3 min of exercise. Dashed line is  $y = x$ .

It is important to recognize that the slope of the  $\dot{V}_{O_2}$ -[PCr] relationship reflects the combined influence of mitochondrial density, the rate constant ( $k$ ) of [PCr] breakdown relative to  $k$  of  $\dot{V}_{O_2}$ , the total [creatine] and the P/O (Meyer, 1988; Kemp *et al.* 2014). Mitochondrial density and total [creatine] are constant during acute exercise, and therefore any divergence in  $\dot{V}_{O_2}$ -[PCr] slope would result from changes in  $k$ [PCr] and/or P/O over the exercise intensities. While the  $k$ [PCr] was not different between 3 and 8 min of heavy-intensity exercise ( $p = \text{n.s.}$ ), there was variance among individuals (Fig. 3A). Therefore, while we base our interpretation on the group mean, we cannot rule out the influence of variance in the individual changes in  $k$ [PCr] in interpreting the  $\dot{V}_{O_2}$ -[PCr] slope. In addition, we used a fixed transit delay to phase align the  $\dot{V}_{O_2}$  and [PCr] measurements in the time domain. This correction provided the best fit to the kinetics that we could make, but it is a limitation for interpreting the  $\dot{V}_{O_2}$ -[PCr] relationship. Specifically, small errors in transit


delay adjustment result in non-linear distortion when plotting single participant data, although this influence is greater during the early kinetics (first 2 min) than between 3 and 8 min of exercise, when the kinetics are slower. Finally, the progressive intramuscular acidification during exercise would be expected to dissociate the dynamics of  $\dot{V}_{O_2}$  and [PCr], speeding the former and slowing the latter (Iotti *et al.* 1993; Gerbino *et al.* 1996; Layec *et al.* 2013). Therefore, while substantial assumptions necessarily underlie the interpretation of the  $\dot{V}_{O_2}$ -[PCr] relationship, it is currently the only way to examine change in P/O as a potential mechanism explaining the lack of relationship between the magnitude of the  $\dot{V}_{O_2\text{sc}}$  and  $\Delta\text{ATP}_{\text{tot}}$ . These data suggest that P/O is stable during moderate-intensity exercise and the first 3 min of heavy-intensity exercise, in agreement with the other  $^{31}\text{P}$  MRS studies (e.g. where the  $\dot{V}_{O_2}$ -[PCr] relationship is strikingly linear throughout the metabolic rate range; Bailey *et al.* 2010), but that sustained heavy-intensity exercise beyond 3 min may be accompanied by a reduction in P/O. Consequently, contrary to the prevailing hypothesis (Rossiter *et al.* 2002), the  $\dot{V}_{O_2\text{sc}}$  may be, at least in part, a result of mitochondrial uncoupling in the active muscle during acidifying exercise.



**Figure 4. Magnitudes and relationship between  $\dot{V}_{O_2}$  slow component and muscle [PCr] and  $\text{ATP}_{\text{tot}}$  slow components**  $\dot{V}_{O_2\text{sc}}$  and  $[\text{PCr}]_{\text{sc}}$  and  $\Delta\text{ATP}_{\text{tot}}$  from minute 3 to 8 of heavy exercise expressed as a percentage change (A). B shows  $\Delta\text{ATP}_{\text{tot}}$  during heavy exercise plotted as a function of the  $\dot{V}_{O_2\text{sc}}$ .

#### Potential mechanisms of mitochondrial uncoupling

There are various mechanisms that might cause the mitochondrial transmembrane proton gradient to dissipate during exercise. This proton 'leak' is regulated by uncoupling proteins and contributes to setting the resting P/O. If this process is augmented during exercise,



**Figure 5. Relationship between pulmonary  $\dot{V}_{O_2}$  and [PCr] during moderate (filled circles) and heavy exercise (open circles)**

The regression shown (continuous line) was fitted to data from moderate exercise and from the first 3 min of heavy exercise and extrapolated (dashed line) to 8 min of heavy exercise. Error bars represent SD. The  $\dot{V}_{O_2}$  data were phase aligned with respect to [PCr] measurements.

the ATP yield per atomic oxygen consumed would fall. Others have shown upregulation of uncoupling proteins 2 and 3 (both expressed in skeletal muscle) with an acute bout of exercise, and these can induce mitochondrial uncoupling, which is likely to minimize production of, and damage from, reactive oxygen species (Brand *et al.* 2004; Bo *et al.* 2008; Jiang *et al.* 2009). This effect may be akin to the chronic uncoupling reported with ageing, posited as a protective mechanism against damage from reactive oxygen species (Brand *et al.* 2004; Amara *et al.* 2007), particularly as leak respiration comprises a large proportion of resting  $\dot{V}_{O_2}$ . However, the kinetics of upregulation of uncoupling proteins are relatively slow in comparison with the duration of exercise in our study; upregulation of uncoupling proteins is typically present ~45–90 min after acute exercise. Additionally, investigations into mitochondrial uncoupling have relied on relatively long bouts of exercise (>30 min), and evidence from human muscle suggests that acute exercise may not be sufficient to elicit the same effect size for upregulation as seen in the rat (Fernström *et al.* 2004). Therefore, upregulation of uncoupling proteins seems less likely to explain fully the lack of relationship between  $\dot{V}_{O_{2sc}}$  and  $\Delta ATP_{tot}$  during heavy exercise.

Alternatively, dissociation of the  $\dot{V}_{O_{2sc}}$  and  $\Delta ATP_{tot}$  may result from high  $[H^+]$  or  $[P_i]$  during exercise (Walsh *et al.* 2002). Low pH can reduce  $[ADP]$  from a shift in the creatine kinase equilibrium (Conley *et al.* 2001) and also serve to dissociate creatine kinase from the mitochondrial membrane, leading to a disruption in oxidative phosphorylation (Walsh *et al.* 2002). While evidence for a direct effect of acidosis is certainly not conclusive (Suleymanlar *et al.* 1992; Kemp *et al.* 2014), numerous studies show disturbances to oxidative phosphorylation through the inhibition of respiratory enzymes or reductions in the proton motive force (Hillered *et al.* 1984; Harkema & Meyer, 1997; Jubrias *et al.* 2003), but fail to result in change to P/O alone (Tonkonogi & Sahlin, 1999). Nevertheless, the variable relationships between the magnitude of the  $\dot{V}_{O_{2sc}}$  and  $\Delta ATP_{tot}$ , together with a steepened  $\dot{V}_{O_2}$ -[PCr] relationship, suggest P/O change as a possible scenario during heavy exercise.

### Technical considerations and study limitations

While limitations accompany the estimations, our study design provides an advantage over previous reports of ATP turnover rate in the literature. Prior estimations have relied on extrapolation of  $V_{i[PCr]}$ , which is assumed to be affected only by the [PCr] recovery amplitude. This model constrains P/O with a linear  $\dot{V}_{O_2}$ -[PCr] relationship, by definition (Layec *et al.* 2009a, which is in contrast with recent findings (Kemp, 2008; Wüst *et al.* 2011; Glancy & Balaban, 2012) and the observations in this study (Fig. 5). Conversely, [PCr] recovery dynamics

may be plastic during supra-LT exercise where intracellular acidification (Yoshida & Watari, 1993, 1994), fatigue-related metabolite accumulation (Jones *et al.* 2008) and muscle fatigue (Yano *et al.* 2001; Cannon *et al.* 2011) have been reported. While the group mean for  $k[PCr]$  resynthesis (or time constant,  $\tau = 1/k$ ) is not different following sub- and supra-LT exercise in this study and others (Rossiter *et al.* 2002), our data suggest that  $k[PCr]$  is not constant within an individual. Therefore, in our study,  $V_{i[PCr]}$  (and, thus,  $Q$  and  $ATP_{tot}$ ) were not constrained to increase in response only to changes in [PCr]. In other words, the augmented amplitude of [PCr] during the slow component did not result in an obligatorily faster initial rate of change following the cessation of exercise; our measurement was dependent on the recovery dynamics characterized and specific to that moment in time. Consequently, the estimations provided for oxidative ATP yield in our study are devoid of the assumptions about the  $\dot{V}_{O_2}$ -[ADP] and  $\dot{V}_{O_2}$ -[PCr] relationships.

$ATP_{tot}$  is most heavily weighted on changes in  $V_{i[PCr]}$ , a measure that is sensitive to noise in the MRS signal (e.g. Fig. 7 of Rossiter *et al.* 2000); this initial rate is derived from characterization of the kinetics of [PCr] recovery. The influence of noise in [PCr] recovery kinetics, particularly in the early transient, is likely to be the largest source of variability to resolve  $ATP_{tot}$ . Conversely, the confidence in characterizing [PCr] off-kinetics is substantially greater than for pulmonary  $\dot{V}_{O_2}$  or even [PCr] during the on-transient. Any improvement in the characterization of  $^{31}P$  dynamics will take a considerable leap in signal-to-noise ratio and more rapid acquisition of spectra.

The heterogeneous nature of skeletal muscle metabolism (Koga *et al.* 2007; Damon *et al.* 2008; Saitoh *et al.* 2009; Cannon *et al.* 2013) may have obscured the characterization of [PCr] dynamics, and therefore  $ATP_{tot}$ . Using  $^{31}P$  MRS, we measured a volume of tissue (~300 g) that may not be representative of the entire knee extensor group responsible for the power output or the diversity of metabolic strain within this group. Finally, the unmeasured work of knee flexion is not accounted for with our ergometer. Therefore, the work of knee flexion (to lift the leg) is assumed to be constant in our experiments, but does contribute to the pulmonary  $\dot{V}_{O_2}$  signal.

### Conclusions

Similar to previous studies, the mean magnitudes of the  $\dot{V}_{O_2}$  and [PCr] slow components were not different during heavy exercise, consistent with the prevailing hypothesis for the intramuscular source of the  $\dot{V}_{O_{2sc}}$ , i.e. an increase in the phosphate cost of force production. Although the magnitude of the  $\dot{V}_{O_{2sc}}$  (~22%) was similar to the increase in  $ATP_{tot}$  (~26%) from 3 to 8 min during heavy exercise,

there was no relationship detected between these measures among individuals. Therefore, our data suggest that the pulmonary  $\dot{V}_{O_{2sc}}$  does not originate solely from increases in the phosphate cost of power production (increased P/W). Other mechanisms, such as an increased  $O_2$  cost of ATP resynthesis (reduced P/O) during acidifying exercise, may also contribute to generating the  $\dot{V}_{O_{2sc}}$ .

## References

- Amara CE, Shankland EG, Jubrias SA, Marcinek DJ, Kushmerick MJ & Conley KE (2007). Mild mitochondrial uncoupling impacts cellular aging in human muscles *in vivo*. *Proc Natl Acad Sci U S A* **104**, 1057–1062.
- Bailey SJ, Fulford J, Vanhatalo A, Winyard PG, Blackwell JR, DiMenna FJ, Wilkerson DP, Benjamin N & Jones AM (2010). Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *J Appl Physiol* **109**, 135–148.
- Billat VL, Richard R, Binsse VM, Koralsztein JP & Haouzi P (1998). The  $\dot{V}_{O_2}$  slow component for severe exercise depends on type of exercise and is not correlated with time to fatigue. *J Appl Physiol* **85**, 2118–2124.
- Bo H, Jiang N, Ma G, Qu J, Zhang G, Cao D, Wen L, Liu S, Ji LL & Zhang Y (2008). Regulation of mitochondrial uncoupling respiration during exercise in rat heart: role of reactive oxygen species (ROS) and uncoupling protein 2. *Free Radic Biol Med* **44**, 1373–1381.
- Brand MD, Buckingham JA, Esteves TC, Green K, Lambert AJ, Miwa S, Murphy MP, Pakay JL, Talbot DA & Echtay KS (2004). Mitochondrial superoxide and aging: uncoupling-protein activity and superoxide production. *Biochem Soc Symp*, 203–213.
- Cannon DT, Howe FA, Whipp BJ, Ward SA, McIntyre DJ, Ladroue C, Griffiths JR, Kemp GJ & Rossiter HB (2013). Muscle metabolism and activation heterogeneity by combined  $^{31}P$  chemical shift and T2 imaging, and pulmonary  $O_2$  uptake during incremental knee-extensor exercise. *J Appl Physiol* **115**, 839–849.
- Cannon DT, White AC, Andriano MF, Kolkhorst FW & Rossiter HB (2011). Skeletal muscle fatigue precedes the slow component of oxygen uptake kinetics during exercise in humans. *J Physiol* **589**, 727–739.
- Coats EM, Rossiter HB, Day JR, Miura A, Fukuba Y & Whipp BJ (2003). Intensity-dependent tolerance to exercise after attaining  $\dot{V}_{O_{2\max}}$  in humans. *J Appl Physiol* **95**, 483–490.
- Conley KE, Kemper WF & Crowther GJ (2001). Limits to sustainable muscle performance: interaction between glycolysis and oxidative phosphorylation. *J Exp Biol* **204**, 3189–3194.
- Conley KE, Kushmerick MJ & Jubrias SA (1998). Glycolysis is independent of oxygenation state in stimulated human skeletal muscle *in vivo*. *J Physiol* **511**, 935–945.
- Damon BM, Wadington MC, Lansdown DA & Hornberger JL (2008). Spatial heterogeneity in the muscle functional MRI signal intensity time course: effect of exercise intensity. *Magn Reson Imaging* **26**, 1114–1121.
- DiMenna FJ, Fulford J, Bailey SJ, Vanhatalo A, Wilkerson DP & Jones AM (2010). Influence of priming exercise on muscle [PCr] and pulmonary  $O_2$  uptake dynamics during ‘work-to-work’ knee-extension exercise. *Respir Physiol Neurobiol* **172**, 15–23.
- Faraut B, Giannesini B, Matarazzo V, Marqueste T, Dalmasso C, Rougon G, Cozzone PJ & Bendahan D (2007). Downregulation of uncoupling protein-3 *in vivo* is linked to changes in muscle mitochondrial energy metabolism as a result of capsiate administration. *Am J Physiol Endocrinol Metab* **292**, E1474–E1482.
- Fernström M, Tonkonogi M & Sahlin K (2004). Effects of acute and chronic endurance exercise on mitochondrial uncoupling in human skeletal muscle. *J Physiol* **554**, 755–763.
- Gerbino A, Ward SA & Whipp BJ (1996). Effects of prior exercise on pulmonary gas-exchange kinetics during high-intensity exercise in humans. *J Appl Physiol* **80**, 99–107.
- Glancy B & Balaban RS (2012). Role of mitochondrial  $Ca^{2+}$  in the regulation of cellular energetics. *Biochemistry* **51**, 2959–2973.
- Harkema SJ & Meyer RA (1997). Effect of acidosis on control of respiration in skeletal muscle. *Am J Physiol Cell Physiol* **272**, C491–C500.
- Hillered L, Ernster L & Siesjö BK (1984). Influence of *in vitro* lactic acidosis and hypercapnia on respiratory activity of isolated rat brain mitochondria. *J Cereb Blood Flow Metab* **4**, 430–437.
- Iotti S, Lodi R, Frassinetti C, Zaniol P & Barbiroli B (1993). *In vivo* assessment of mitochondrial functionality in human gastrocnemius muscle by  $^{31}P$  MRS. The role of pH in the evaluation of phosphocreatine and inorganic phosphate recoveries from exercise. *NMR Biomed* **6**, 248–253.
- Jiang N, Zhang G, Bo H, Qu J, Ma G, Cao D, Wen L, Liu S, Ji LL & Zhang Y (2009). Upregulation of uncoupling protein-3 in skeletal muscle during exercise: a potential antioxidant function. *Free Radic Biol Med* **46**, 138–145.
- Jones AM, Wilkerson DP, DiMenna F, Fulford J & Poole DC (2008). Muscle metabolic responses to exercise above and below the ‘critical power’ assessed using  $^{31}P$ -MRS. *Am J Physiol Regul Integr Comp Physiol* **294**, R585–R593.
- Jubrias SA, Crowther GJ, Shankland EG, Gronka RK & Conley KE (2003). Acidosis inhibits oxidative phosphorylation in contracting human skeletal muscle *in vivo*. *J Physiol* **553**, 589–599.
- Kemp G (2008). Physiological implications of linear kinetics of mitochondrial respiration *in vitro*. *Am J Physiol Cell Physiol* **295**, C844–C846; author reply C847–C848.
- Kemp GJ, Ahmad RE, Nicolay K & Prompers JJ (2014). Quantification of skeletal muscle mitochondrial function by  $^{31}P$  magnetic resonance spectroscopy techniques: a quantitative review. *Acta Physiol (Oxf)* in press, doi: 10.1111/apha.12307.
- Kemp GJ, Meyerspeer M & Moser E (2007). Absolute quantification of phosphorus metabolite concentrations in human muscle *in vivo* by  $^{31}P$  MRS: a quantitative review. *NMR Biomed* **20**, 555–565.

- Kemp GJ, Roussel M, Bendahan D, Le Fur Y & Cozzone PJ (2001). Interrelations of ATP synthesis and proton handling in ischaemically exercising human forearm muscle studied by  $^{31}\text{P}$  magnetic resonance spectroscopy. *J Physiol* **535**, 901–928.
- Kemp GJ, Taylor DJ, Styles P & Radda GK (1993). The production, buffering and efflux of protons in human skeletal muscle during exercise and recovery. *NMR Biomed* **6**, 73–83.
- Kemp GJ, Thompson CH, Taylor DJ & Radda GK (1995). ATP production and mechanical work in exercising skeletal muscle: a theoretical analysis applied to  $^{31}\text{P}$  magnetic resonance spectroscopic studies of dialyzed uremic patients. *Magn Reson Med* **33**, 601–609.
- Koga S, Poole DC, Ferreira LF, Whipp BJ, Kondo N, Saitoh T, Ohmae E & Barstow TJ (2007). Spatial heterogeneity of quadriceps muscle deoxygenation kinetics during cycle exercise. *J Appl Physiol* **103**, 2049–2056.
- Krustrup P, Ferguson RA, Kjær M & Bangsbo J (2003). ATP and heat production in human skeletal muscle during dynamic exercise: higher efficiency of anaerobic than aerobic ATP resynthesis. *J Physiol* **549**, 255–269.
- Krustrup P, Jones AM, Wilkerson DP, Calbet JA & Bangsbo J (2009). Muscular and pulmonary  $\text{O}_2$  uptake kinetics during moderate- and high-intensity sub-maximal knee-extensor exercise in humans. *J Physiol* **587**, 1843–1856.
- Kushmerick MJ (1997). Multiple equilibria of cations with metabolites in muscle bioenergetics. *Am J Physiol Cell Physiol* **272**, C1739–C1747.
- Lamarra N, Whipp BJ, Ward SA & Wasserman K (1987). Effect of interbreath fluctuations on characterizing exercise gas exchange kinetics. *J Appl Physiol* **62**, 2003–2012.
- Lanza IR, Befroy DE & Kent-Braun JA (2005). Age-related changes in ATP-producing pathways in human skeletal muscle in vivo. *J Appl Physiol* **99**, 1736–1744.
- Layec G, Bringard A, Le Fur Y, Vilmen C, Micallef JP, Perrey S, Cozzone PJ & Bendahan D (2009a). Effects of a prior high-intensity knee-extension exercise on muscle recruitment and energy cost: a combined local and global investigation in humans. *Exp Physiol* **94**, 704–719.
- Layec G, Bringard A, Le Fur Y, Vilmen C, Micallef JP, Perrey S, Cozzone PJ & Bendahan D (2011). Comparative determination of energy production rates and mitochondrial function using different  $^{31}\text{P}$  MRS quantitative methods in sedentary and trained subjects. *NMR Biomed* **24**, 425–438.
- Layec G, Bringard A, Vilmen C, Micallef JP, Le Fur Y, Perrey S, Cozzone PJ & Bendahan D (2009b). Does oxidative capacity affect energy cost? An in vivo MR investigation of skeletal muscle energetics. *Eur J Appl Physiol* **106**, 229–242.
- Layec G, Bringard A, Yashiro K, Le Fur Y, Vilmen C, Micallef JP, Perrey S, Cozzone PJ & Bendahan D (2012). The slow components of phosphocreatine and pulmonary oxygen uptake can be dissociated during heavy exercise according to training status. *Exp Physiol* **97**, 955–969.
- Layec G, Malucelli E, Le Fur Y, Manners D, Yashiro K, Testa C, Cozzone PJ, Iotti S & Bendahan D (2013). Effects of exercise-induced intracellular acidosis on the phosphocreatine recovery kinetics: a  $^{31}\text{P}$  MRS study in three muscle groups in humans. *NMR Biomed* **26**, 1403–1411.
- Meyer RA (1988). A linear model of muscle respiration explains monoexponential phosphocreatine changes. *Am J Physiol Cell Physiol* **254**, C548–C553.
- Moon RB & Richards JH (1973). Determination of intracellular pH by  $^{31}\text{P}$  magnetic resonance. *J Biol Chem* **248**, 7276–7278.
- Murgatroyd SR, Ferguson C, Ward SA, Whipp BJ & Rossiter HB (2011). Pulmonary  $\text{O}_2$  uptake kinetics as a determinant of high-intensity exercise tolerance in humans. *J Appl Physiol* **110**, 1598–1606.
- Naressi A, Couturier C, Devos JM, Janssen M, Mangeat C, de Beer R & Graveron-Demilly D (2001). Java-based graphical user interface for the MRUI quantitation package. *MAGMA* **12**, 141–152.
- Poole DC, Barstow TJ, Gaesser GA, Willis WT & Whipp BJ (1994).  $\text{VO}_2$  slow component: physiological and functional significance. *Med Sci Sports Exerc* **26**, 1354–1358.
- Poole DC & Jones AM (2012). Oxygen uptake kinetics. *Compr Physiol* **2**, 933–996.
- Poole DC, Schaffartzik W, Knight DR, Derion T, Kennedy B, Guy HJ, Prediletto R & Wagner PD (1991). Contribution of exercising legs to the slow component of oxygen uptake kinetics in humans. *J Appl Physiol* **71**, 1245–1260.
- Rossiter HB (2011). Exercise: kinetic considerations for gas exchange. *Compr Physiol* **1**, 203–244.
- Rossiter HB, Howe FA, Ward SA, Kowalchuk JM, Griffiths JR & Whipp BJ (2000). Intersample fluctuations in phosphocreatine concentration determined by  $^{31}\text{P}$ -magnetic resonance spectroscopy and parameter estimation of metabolic responses to exercise in humans. *J Physiol* **528**, 359–369.
- Rossiter HB, Ward SA, Doyle VL, Howe FA, Griffiths JR & Whipp BJ (1999). Inferences from pulmonary  $\text{O}_2$  uptake with respect to intramuscular [phosphocreatine] kinetics during moderate exercise in humans. *J Physiol* **518**, 921–932.
- Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR & Whipp BJ (2001). Effects of prior exercise on oxygen uptake and phosphocreatine kinetics during high-intensity knee-extension exercise in humans. *J Physiol* **537**, 291–303.
- Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR & Whipp BJ (2002). Dynamic asymmetry of phosphocreatine concentration and  $\text{O}_2$  uptake between the on- and off-transients of moderate- and high-intensity exercise in humans. *J Physiol* **541**, 991–1002.
- Saitoh T, Ferreira LF, Barstow TJ, Poole DC, Ooue A, Kondo N & Koga S (2009). Effects of prior heavy exercise on heterogeneity of muscle deoxygenation kinetics during subsequent heavy exercise. *Am J Physiol Regul Integr Comp Physiol* **297**, R615–R621.
- Sargeant AJ & Dolan P (1987). Effect of prior exercise on maximal short-term power output in humans. *J Appl Physiol* **63**, 1475–1480.
- Suleymanlar G, Zhou HZ, McCormack M, Elkins N, Kucera R, Reiss OK & Shapiro JI (1992). Mechanism of impaired energy metabolism during acidosis: role of oxidative metabolism. *Am J Physiol Heart Circ Physiol* **262**, H1818–H1822.

- Tonkonogi M & Sahlin K (1999). Actively phosphorylating mitochondria are more resistant to lactic acidosis than inactive mitochondria. *Am J Physiol Cell Physiol* **277**, C288–C293.
- Vanhatalo A, Poole DC, DiMenna FJ, Bailey SJ & Jones AM (2011). Muscle fiber recruitment and the slow component of O<sub>2</sub> uptake: constant work rate vs. all-out sprint exercise. *Am J Physiol Regul Integr Comp Physiol* **300**, R700–R707.
- Walsh B, Tiiveli T, Tonkonogi M & Sahlin K (2002). Increased concentrations of P<sub>i</sub> and lactic acid reduce creatine-stimulated respiration in muscle fibers. *J Appl Physiol* **92**, 2273–2276.
- Walter G, Vandenborne K, Elliott M & Leigh JS (1999). *In vivo* ATP synthesis rates in single human muscles during high intensity exercise. *J Physiol* **519**, 901–910.
- Wasserman K, Stringer WW & Casaburi R. (1995). Is the slow component of exercise VO<sub>2</sub> a respiratory adaptation to anaerobiosis? *Adv Exp Med Biol* **393**, 187–194.
- Whipp BJ (1996). Domains of aerobic function and their limiting parameters. In *The Physiology and Pathophysiology of Exercise Tolerance*, ed. Steinacker JM & Ward SA, pp. 83–89. Plenum, New York.
- Whipp BJ, Rossiter HB, Ward SA, Avery D, Doyle VL, Howe FA & Griffiths JR (1999). Simultaneous determination of muscle <sup>31</sup>P and O<sub>2</sub> uptake kinetics during whole body NMR spectroscopy. *J Appl Physiol* **86**, 742–747.
- Wüst RC, Grassi B, Hogan MC, Howlett RA, Gladden LB & Rossiter HB. (2011). Kinetic control of oxygen consumption during contractions in self-perfused skeletal muscle. *J Physiol* **589**, 3995–4009.
- Yano T, Yunoki T & Ogata H (2001). Relationship between the slow component of oxygen uptake and the potential reduction in maximal power output during constant-load exercise. *J Sports Med Phys Fitness* **41**, 165–169.
- Yoshida T & Watari H (1993). Changes in intracellular pH during repeated exercise. *Eur J Appl Physiol Occup Physiol* **67**, 274–278.
- Yoshida T & Watari H (1994). Exercise-induced splitting of the inorganic phosphate peak: investigation by time-resolved <sup>31</sup>P-nuclear magnetic resonance spectroscopy. *Eur J Appl Physiol Occup Physiol* **69**, 465–473.
- Zoladz JA, Gladden LB, Hogan MC, Nieckarz Z & Grassi B (2008). Progressive recruitment of muscle fibers is not necessary for the slow component of  $\dot{V}O_2$  kinetics. *J Appl Physiol* **105**, 575–580.
- Zołądz JA & Korzeniewski B (2001). Physiological background of the change point in  $\dot{V}O_2$  and the slow component of oxygen uptake kinetics. *J Physiol Pharmacol* **52**, 167–184.

## Additional information

### Competing interests

None declared.

### Author contributions

D.T.C., G.J.K. and H.B.R. conceived and designed experiments and analysed data. All authors performed experiments and interpreted data. D.T.C. prepared the figures. D.T.C. and H.B.R. wrote the manuscript. All authors critically reviewed and approved the final version of the manuscript.

### Funding

This research was supported by Biotechnology and Biological Sciences Research Council (BBSRC) UK BB/I001174/1 and BB/I00162X/1.

### Acknowledgements

We are very grateful to Dr Peter Brooks and Emma Routeledge from the Department of Mechanical Engineering at the University of Leeds, for the design and construction of the customized carbon-fibre lever arms of the MR ergometer. We also wish to thank all volunteers for their time and dedication.