

Published in final edited form as:

Environ Res. 2015 January ; 136: 47–56. doi:10.1016/j.envres.2014.10.011.

Maternal blood and hair manganese concentrations, fetal growth, and length of gestation in the ISA cohort in Costa Rica

Ana M. Mora^{1,2,*}, Berna van Wendel de Joode¹, Donna Mergler³, Leonel Córdoba¹, Camilo Cano¹, Rosario Quesada¹, Donald R. Smith⁴, José A. Menezes-Filho⁵, and Brenda Eskenazi²

¹ Central American Institute for Studies on Toxic Substances (IRET), Universidad Nacional, P.O. Box 86-3000 Heredia, Costa Rica.

² Center for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, 1995 University Ave, Suite 265, Berkeley, CA 94704, USA.

³ Centre de Recherche Interdisciplinaire sur la Biologie, la Santé, la Société et l'Environnement (CINBIOSE), Université du Québec à Montréal, Pavillon des sciences, 141, Avenue du Président Kennedy, H2X 1Y4 Montréal, Québec, Canada.

⁴ Microbiology and Environmental Toxicology, University of California, Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, USA.

⁵ Laboratory of Toxicology, College of Pharmacy, Federal University of Bahia, Av. Barão de Jeremoabo s/n Campus Universitário de Ondina, 40170-115 Salvador, Bahia, Brazil.

Abstract

Background—Animal studies have shown that both deficiency and excess manganese (Mn) may result in decreased fetal size and weight, but human studies have reported inconsistent results.

Methods—We examined the association of blood and hair Mn concentrations measured at different times during pregnancy with fetal growth among term births and length of gestation in a cohort of 380 mother-infant pairs living near banana plantations aerially sprayed with Mn-

© 2014 Elsevier Inc. All rights reserved.

* **Corresponding author:** Ana María Mora, MD, PhD Central American Institute for Studies on Toxic Substances (IRET), Universidad Nacional. P.O. Box 86-3000 Heredia, Costa Rica. animora@yahoo.com, Phone: +506 2263-6375 ext. 114, Fax: +506 2263-6375.

Berna van Wendel de Joode: berendina.vanwendel.dejoode@una.cr

Donna Mergler: mergler.donna@uqam.ca

Leonel Córdoba: leonel.cordoba.gamboa@una.cr

Juan Camilo Cano: jucanoc@gmail.com

Rosario Quesada: maria.quesada.varela@una.cr

Donald R. Smith: drsmith@ucsc.edu

José A. Menezes-Filho: antomen@ufba.br

Brenda Eskenazi: eskenazi@berkeley.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

containing fungicides in Costa Rica. We used linear regression and generalized additive models to test for linear and nonlinear associations.

Results—Mean (\pm SD) blood Mn concentration was 24.4 ± 6.6 $\mu\text{g/L}$ and geometric mean (geometric SD) hair Mn concentration was 1.8 (3.2) $\mu\text{g/g}$. Hair Mn concentrations during the second and third trimesters of gestation were positively related to infant chest circumference (β for 10-fold increase = 0.62 cm; 95% CI: $0.16, 1.08$; and $\beta = 0.55$ cm; 95% CI: $-0.16, 1.26$, respectively). Similarly, average maternal hair Mn concentrations during pregnancy were associated with increased chest circumference (β for 10-fold increase = 1.19 cm; 95% CI: $0.43, 1.95$) in infants whose mothers did not have gestational anemia, but not in infants of mothers who had gestational anemia ($\beta = 0.39$ cm; 95% CI: $-0.32, 1.10$; $p_{\text{INT}} = 0.14$). All these associations were linear. Blood Mn concentrations did not show consistent linear nor nonlinear relationships with any of the birth outcomes.

Conclusions—Mn plays an important role in fetal development, but the extent to which environmental exposures may cause adverse health effects to the developing fetus is not well understood. Among women living near banana plantations in Costa Rica, we did not observe linear or nonlinear associations of Mn concentrations with lowered birth weight or head circumference, as reported in previous studies. However, we did find positive linear associations between maternal hair Mn concentrations during pregnancy and infant chest circumference.

Keywords

Birth outcomes; birth weight; manganese; pesticides; pregnancy; Costa Rica

Introduction

Manganese (Mn) is an important cofactor in enzymatic reactions involved in bone formation and metabolic regulation in animals and humans (ATDSR 2012). Both Mn deficiency and overexposure can disrupt the growth hormone/insulin-like growth factor axis and insulin metabolism (Clegg et al. 1998; Hiney et al. 2011; Keen et al. 1999). Mn can also induce cellular free radical damage and mitochondrial dysfunction at high concentrations (HaMai and Bondy 2004; Keen et al. 2000). Dietary intake is the main source of Mn for the general population, but elevated exposure to Mn can occur through the environment (ATDSR 2012). Environmental sources of Mn exposure include groundwater with naturally high concentrations of natural Mn or contaminated by industrial waste (Bouchard et al. 2007; Bouchard et al. 2011; He et al. 1994; Kondakis et al. 1989; Wasserman et al. 2006), combustion of anti-knock additives in gasoline (Zayed et al. 1999), dust emissions from industrial sources (Menezes-Filho et al. 2011) and Mn mines (Riojas-Rodriguez et al. 2010), and spraying of Mn-containing ethylene bisdithiocarbamate fungicides (Gunier et al. 2013; Mora et al. 2014).

Mn is an essential nutrient, but it is also a neurotoxicant at high doses (ATDSR 2012). Mn is closely regulated by homeostatic mechanisms, but blood Mn concentrations increase throughout pregnancy (Gunier et al. submitted; Hambridge and Droegemueller 1974; Mora et al. 2014; Spencer 1999; Takser et al. 2004; Tholin et al. 1995) and Mn crosses the placenta via active transport mechanisms (Krachler et al. 1999; Leazer and Klaassen 2003),

suggesting that Mn plays an important role in fetal development that requires elevated Mn concentrations. Notably, the extent to which environmental exposures may alter Mn concentrations during pregnancy and cause adverse health effects to the developing fetus is not well understood.

Several animal studies have found that both Mn deficiency and overexposure (via oral, subcutaneous, and intravenous routes) during gestation are associated with decreased fetal size and weight (Bolze et al. 1985; Colomina et al. 1996; Sanchez et al. 1993; Treinen et al. 1995). However, a small number of studies in pregnant rodents exposed to elevated doses of Mn orally or subcutaneously have not observed associations between dose of Mn and fetal size or weight (Gray and Laskey 1980; Molina et al. 2011; Pappas et al. 1997; Torrente et al. 2002). In fact, a recent study found that pups prenatally exposed to Mn in drinking water were born with significantly higher body weights compared to controls (Betharia and Maher 2012).

A number of studies have examined the effects of prenatal Mn exposure on birth outcomes in humans (Table S1) (Chen et al. 2014; Eum et al. 2014; Guan et al. 2013; Osada et al. 2002; Takser et al. 2004; Vigeh et al. 2008; Xu et al. 2011; Yu and Cao 2013; Zota et al. 2009). Cross-sectional studies in Oklahoma ($n = 470$ term newborns), China ($n = 172$ preterm and term infants), and Korea ($n = 331$ term newborns) have reported nonlinear associations between maternal blood Mn concentrations at delivery (medians = 22, 53.8, and 21.5 $\mu\text{g/L}$, respectively) and birth weight (Chen et al. 2014; Eum et al. 2014; Zota et al. 2009). Infant birth weight increased linearly with Mn concentrations up to 31 $\mu\text{g/L}$ in the Oklahoma study, 41.8 $\mu\text{g/L}$ in the Chinese study, and 30-35 $\mu\text{g/L}$ in the Korean study. At higher Mn concentrations, non-significant inverse relationships were observed between maternal Mn and birth weight in all three studies. A second study in China ($n = 125$ mother-child pairs) did not observe an association between maternal blood Mn concentrations at delivery (median = 50.6 $\mu\text{g/L}$) and birth weight, but found significant inverted U-shaped relationships between Mn concentrations, head circumference, and chest circumference (Guan et al. 2013). Additionally, a case-control study of 271 Iranian mother-child pairs found that mothers of newborns with intrauterine growth retardation had significantly lower blood Mn concentrations shortly after delivery compared to mothers of newborns with sizes appropriate for gestational age (means = 16.7 vs. 19.1 $\mu\text{g/L}$, respectively) (Vigeh et al. 2008). To date, only one epidemiological study has been published on the relationship between blood Mn concentrations measured at multiple time points during pregnancy (means in first, second, and third trimesters of gestation = 9.0, 9.9, and 16.3 $\mu\text{g/L}$, respectively) and birth outcomes (Takser et al. 2004). This study of 149 Canadian mothers and their children did not find any significant associations between Mn concentrations at any point in pregnancy and newborn growth parameters.

Previous studies have exclusively examined the association between birth outcomes and Mn concentrations measured in blood. In the present study, we measured Mn in maternal hair as well as blood samples collected multiple times over pregnancy and assessed its association with fetal growth and length of gestation in a mother-infant cohort living near banana plantations in Costa Rica aerially sprayed with the Mn-containing fungicide mancozeb.

2 Materials and methods

2.1 Study population

The Infants' Environmental Health Study ('*Infantes y Salud Ambiental*', ISA) is a prospective birth cohort study of fetal and early childhood exposure to pesticides and their impact on growth and neurodevelopment in children living near banana plantations in Matina County, Costa Rica. Subject recruitment and procedures for the ISA study have been described elsewhere (Mora et al. 2014; van Wendel de Joode et al. 2014). Briefly, pregnant women were recruited through meetings in local schools, communal groups, advertisements, and friends' referrals between March 2010 and June 2011. Women were eligible to participate in the study if they were < 33 weeks of gestation, > 15 years of age, and living in one of the 40 villages of Matina County located < 5 km of a banana plantation. A total of 451 women were enrolled. After losses due to miscarriage and stillbirths ($n = 21$), loss to follow-up ($n = 39$), and the exclusion of twins ($n = 2$) and women who did not have delivery medical records available at the time of the postpartum interview ($n = 9$), information on birth weight and length of gestation was available for 380 singleton liveborn infants. Participants included in this analysis did not differ significantly from the original full cohort on most socio-demographic factors, including maternal education, marital status, parity, family income, and blood and hair Mn concentrations during pregnancy. Written informed consent was obtained from all women and additional informed consent was obtained from the parents or legal guardians of participants under the age of 18 years. All study activities were approved by the Ethical Committee of the Universidad Nacional in Costa Rica.

2.2 Data collection

Women were interviewed one to three times during pregnancy (depending on their gestational age at enrollment) and following delivery. Interviews were conducted using structured questionnaires and occurred at enrollment (median = 19 weeks gestation), at the beginning and in the middle of the third trimester of pregnancy (medians = 30 and 33 weeks gestation, respectively), and after delivery (median = 7 weeks postpartum). Socio-demographic information, including maternal age, education, marital status, parity, and family income, was collected at the baseline interview. Information on smoking, alcohol consumption, drug use, caffeine consumption, occupational status, medical conditions, medications, pregnancy complications, and obstetric ultrasounds was gathered at each interview. Data completed by hospital/clinic personnel were abstracted from prenatal medical records provided to the study participants. The abstracted data included maternal pre-pregnancy weight, timing of prenatal care initiation, maternal weight and blood pressure levels at each prenatal care visit, and medical conditions documented in the medical records by the treating physician. Data on hemoglobin concentrations was abstracted for a subset of women ($n = 52$), and when grouped by maternal report of gestational anemia diagnosed at any point during pregnancy, those who reported anemia ($n = 20$) had significantly lower hemoglobin concentrations (p -Wilcoxon-Mann-Whitney test = 0.03). Pre-pregnancy body mass index (BMI) was calculated as (weight in kilograms)/(height in meters)², using maternal pre-pregnancy weight (if available) or weight at the first prenatal care visit (if < 14 weeks gestation) abstracted from medical records, and height measured by the study interviewers. Gestational hypertension was determined using the blood pressure levels

abstracted from prenatal medical records (blood pressure at least 140 mm Hg systolic and/or at least 90 mm Hg diastolic in at least two visits at > 20 weeks of pregnancy in women with previously normal blood pressure), diagnosis abstracted from medical records, and maternal report of hypertensive drug use during pregnancy. Women with gestational diabetes were identified using maternal report and diagnosis abstracted from medical records.

2.3 Fetal growth and length of gestation

Data on infant birth weight, body length, head circumference, and chest circumference were abstracted from delivery medical records provided to the study participants and used as recorded in grams or centimeters. Infant ponderal index, a measure of proportionality of growth, was calculated as (birth weight in grams \times 100)/(body length in centimeters)³. Chest circumference was measured by the hospital staff according to standard procedures (without clothes at the level of the nipples and below the inferior angle of the scapulae), but was only available for a subset of newborns ($n = 165$). Those with data on chest circumference and those missing these data were similar on most socio-demographic characteristics, such as maternal age, maternal education, parity, and family income. However, mothers of infants with chest circumference data had lower hair Mn concentrations during the second trimester, third trimester, and averaged concentrations during pregnancy compared to mothers of infants with no chest circumference data.

Length of gestation was determined using the date of the last menstrual period (LMP) reported by women at enrollment, early obstetric ultrasounds (< 14 weeks gestation), and/or estimates abstracted from medical records. If the gestational age estimated using the LMP date and the gestational age estimated using an early ultrasound differed by < 7 days, the LMP estimate was used as the length of gestation; otherwise, the ultrasound estimate was used. If the woman did not have an early ultrasound, was sure about her LMP date, and the LMP and medical record estimates differed by < 14 days, the LMP estimate was used; otherwise, the medical record estimate was used. Women with a gestational age at birth > 44 weeks and 6 days were excluded from all the analyses ($n = 4$). Gestational age at the time of biological sample collection was calculated using data on length of gestation and date of sample collection.

2.4 Blood Mn measurements

Blood samples were collected by venipuncture at enrollment or shortly after (median = 19, interquartile range = 14-25 weeks gestation) and in the latter half of pregnancy (median = 30, interquartile range = 26-33 weeks gestation). Whole blood samples were stored at -20°C until its shipment to the University of California, Santa Cruz, where they were analyzed for Mn using high-resolution inductively coupled plasma mass spectrometry (Finnigan XR ICP-MS) (Smith et al. 2007). Details of blood collection, analysis, and quality control procedures are described elsewhere (Mora et al. 2014). Whole blood Mn was measured in 613 samples collected from 375 women. A total of 238 women provided two samples during pregnancy and 137 provided only one (at enrollment or shortly after). Blood Mn concentrations measured in this study were all above the analytical limit of detection (LOD = 0.003 $\mu\text{g/L}$).

2.5 Hair Mn measurements

Hair samples (~ 20-30 strands) were collected from the occipital region, within 2 mm from the scalp, at enrollment (median = 19, interquartile range = 14-24 weeks gestation) and in the latter half of pregnancy (median = 31, interquartile range = 26-34 weeks gestation). Samples were stored in plastic bags at room temperature and shipped in batches to the Federal University of Bahia, Brazil. The one-centimeter closest to the scalp from each hair sample was cleaned as described elsewhere (Menezes-Filho et al. 2009) and analyzed for Mn using electrothermal atomic spectroscopy with Zeeman background correction (GTA-120, Varian Inc.). Mn was measured in 708 hair samples collected from 380 study participants; 328 women provided two hair samples during pregnancy and 52 provided only one (at enrollment). Only three hair samples had Mn concentrations below the limit of detection (LOD = 0.01 µg/g) and their values were set at LOD/2.

2.6 Statistical analysis

To evaluate the within- and between-woman variability of Mn concentrations in blood and hair samples during pregnancy, we calculated the intraclass correlation coefficient (ICC) using mixed effect models (Rabe-Hesketh and Skrondal 2012; Rosner 2006). To account for the well-known increase in blood Mn concentrations over pregnancy (Hambridge and Droegemueller 1974; Mora et al. 2014; Spencer 1999; Tholin et al. 1995) and the decrease in hair Mn observed in previous analyses (Mora et al. 2014), we grouped blood and hair Mn concentrations by trimester of gestation at the time of sample collection and then averaged the concentrations for each trimester. We also averaged the Mn concentrations across the repeated samples collected for each woman throughout pregnancy. Blood Mn concentrations were normally distributed, whereas hair Mn concentrations were skewed to the left. Thus, we transformed hair Mn concentrations (for each trimester of gestation and average during pregnancy) to the log₁₀ scale to normalize the residuals.

Several factors were identified *a priori* as potential confounders of the association of Mn with fetal growth or length of gestation on the basis of directed acyclic graphs (Greenland et al. 1999). These confounders included maternal age, pre-pregnancy BMI, timing of prenatal care initiation, caffeine consumption during pregnancy, and interpregnancy interval as continuous variables, and maternal marital status, maternal country of birth, maternal education, parity, maternal occupation during pregnancy, family income, alcohol consumption (< 1 vs. 1 drink per week), smoking, and drug use during pregnancy (any vs. none), caffeinated tea intake during pregnancy, iron intake during pregnancy, history of miscarriage, vaginal bleeding, gestational diabetes, previous low birth weight delivery (yes vs. no), and infant sex as categorical variables (categorized as shown in Table 1 unless specified above). Covariates were included in the regression models if they were associated with blood or hair Mn and any of the birth outcomes in bivariate analyses at a *p*-value < 0.10. Missing covariate values for parity (*n* = 12), history of abortion (*n* = 12), maternal height (*n* = 14), and caffeinated tea consumption during pregnancy (*n* = 2) were imputed at random based on observed probability distributions (< 5% missing) (Lubin et al. 2004). For women missing weight (*n* = 32), this variable was predicted from a regression model that included maternal age, history of abortion, parity, maternal education, maternal height, and timing of prenatal care initiation as predictor variables. Because birth weight in preterm

newborns may reflect growth restriction and/or prematurity (Hernandez-Diaz et al. 2008; Wilcox 2006), we restricted our analyses on exposure to Mn and birth weight, birth length, ponderal index, head circumference, and chest circumference to term newborns (< 37 weeks).

We ran statistical analyses for blood and hair Mn separately. First, linear regression models were used to examine linear associations of blood and hair Mn with birth weight, length, ponderal index, head circumference, chest circumference, and length of gestation (as continuous variables). We then assessed potential nonlinear associations between blood and hair Mn and the birth outcomes of interest by examining penalized splines for Mn using generalized additive models (Hastie and Tibshirani 1990). If a potentially nonlinear association between blood or hair Mn concentrations and any of the birth outcomes was identified, we created indicator variables for tertiles of each Mn biomarker distribution and included them in the adjusted regression models for blood and hair Mn.

We examined differences in the associations of mean blood and mean hair Mn concentrations with birth outcomes for boys and girls using cross-product terms. We also examined effect measure modification by gestational anemia because evidence suggests that iron deficiency may result in increased absorption and retention of Mn in various organs (Chandra and Shukla 1976; Finley 1999; Garcia et al. 2007; Mena et al. 1969; Thompson et al. 2006; Yokoi et al. 1991). We tested for effect modification using mean blood and hair Mn concentrations and not Mn concentrations by trimesters of pregnancy because of sample size limitations, and used a p -value < 0.20 as the cut-off point for significance.

We conducted several sensitivity analyses to assess the robustness of our results. First, we reran models after excluding outliers with studentized residuals (residuals divided by the model standard error) greater than three standard units. Second, we reran the analyses restricted to the subset of women with data available on all covariates ($n = 329$ and 310 mother-child pairs for blood and hair Mn, respectively) to evaluate the appropriateness of using imputation for missing data. Third, we included gestational hypertension as a covariate in the adjusted regression models to examine whether it might be a confounder or mediator of the association between Mn exposure and birth outcomes (Lee and Kim 2011; Taneja and Mandal 2007; Vigeh et al. 2013). Fourth, we reran the analyses including both blood and hair Mn in the same adjusted regression models. Fifth, we fitted the regression models using blood and hair Mn concentrations by halves of pregnancy (< 20 and ≥ 20 weeks gestation) to assess whether the associations with the birth outcomes of interest would change by increasing power. Lastly, we examined the associations of maternal blood and hair Mn concentrations with head circumference/birth weight, head circumference/body length, and head circumference/chest circumference ratios.

Statistical analyses were conducted using Stata version 11.2 (StataCorp, College Station, TX).

3 Results

Most of the women in the study were young (median age = 22 years), born in Costa Rica (81%), married or living as married (74%), and multiparous (64%) (Table 1). About half of them completed primary school (49%) and were overweight or obese before pregnancy (48%). More than half of the women were living below the poverty line (59%). Few women reported smoking (4%), consuming alcohol (3%, data not shown), or using illegal drugs (2%, data not shown) during pregnancy; and more than two-thirds began prenatal care in the first trimester (80%, data not shown). Approximately 3% of women had gestational hypertension, 4% had gestational diabetes, and 40% reported having been diagnosed with gestational anemia at some point during pregnancy. Mean (\pm SD) birth weight was $3,235 \pm 410$ g; mean body (crown-heel) length was 49.9 ± 2.2 cm; mean ponderal index was 2.6 ± 0.3 g/cm³; mean head circumference was 33.8 ± 1.4 cm; mean chest circumference was 33.0 ± 1.7 cm; and mean length of gestation was 39.5 ± 1.8 weeks. A total of 12 infants (3%) weighed $< 2,500$ g at birth and 24 infants (6%) were preterm. Lower birth weight infants were born to women who were born outside of Costa Rica, primiparous, lived below the poverty line, reported taking iron during pregnancy, did not have gestational diabetes, and had vaginal bleeding in the first trimester of pregnancy. Boys and girls had similar birth weights and body lengths, but girls had slightly smaller head circumferences and longer gestational durations (data not shown).

The distribution of maternal blood and hair Mn concentrations by trimester and averaged during pregnancy are shown in Table 2. Mean (\pm SD) blood Mn concentration during pregnancy was 24.5 ± 6.1 μ g/L and geometric mean (geometric SD) hair Mn concentration was 1.9 (2.8) μ g/g. Blood Mn concentrations increased significantly between the second and third trimesters of pregnancy (means = 23.8 and 26.5 μ g/g, respectively; *p*-paired t-test < 0.01), whereas hair Mn concentrations showed a slight decrease between the second and third trimesters (geometric means (GM) = 1.8 and 1.7 μ g/g, respectively; *p*-Wilcoxon signed-rank test = 0.04). Blood Mn concentrations by trimester of pregnancy were moderately correlated among women with repeated measurements during pregnancy ($r_s = 0.41$ - 0.44), with the exception of blood Mn concentrations during the first and third trimester. Hair Mn concentrations by trimester were also positively correlated among participants with repeated measurements ($r_s = 0.49$ - 0.64). Intraclass correlation coefficient (ICC) values were 0.42 and 0.58 for blood and hair Mn concentrations, respectively (data not shown), indicating that 58 and 42% of the variability in blood and hair Mn concentrations was due to within-woman variability. Maternal blood and hair Mn concentrations were not correlated ($r_s = -0.06$, *p* = 0.22, *n* = 375).

Table 3 presents the adjusted multivariable models for the associations of maternal blood and hair Mn concentrations with measures of fetal growth and length of gestation (see Table S2 for unadjusted models). We did not observe significant associations between average maternal blood Mn concentrations during pregnancy and any of the outcomes of interest in the combined analyses, nor did we find significant associations between Mn concentrations during the first trimester and any measure of fetal growth or length of gestation.

For maternal blood Mn concentrations during the second trimester, we observed nonlinear associations with birth weight, body length, and length of gestation in the adjusted generalized additive models ($p_{\text{GAM}} = 0.04, 0.02, \text{ and } 0.04$, respectively; Figure S1). However, the adjusted linear regression models fitted with indicator variables for each tertile of blood Mn concentrations did not show statistically significant associations for any of these endpoints (data not shown).

For maternal blood Mn concentrations during the third trimester, we observed a small but significant and positive association with birth weight (β for every $\mu\text{g/L}$ increase in Mn concentrations = 9.0 g; 95% CI: 0.5, 17.5; Table 3), after adjusting for covariates. Nevertheless, when we removed a single outlier, this result was no longer significant ($\beta = 6.3$ g; 95% CI: -2.0, 14.6). There were no linear or nonlinear associations between third trimester blood Mn concentrations and any of the other birth outcomes.

Maternal hair Mn concentrations averaged over the entire pregnancy and stratified by trimester were not related to birth weight, body length, ponderal index, or length of gestation (Table 3). However, maternal hair Mn concentrations during the second and third trimesters of pregnancy were positively associated with chest circumference (β for a ten-fold increase in Mn concentrations = 0.62 cm; 95% CI: 0.16, 1.08; and $\beta = 0.55$ cm; 95% CI: -0.16, 1.26, respectively).

We did not observe significant differences between boys and girls in the associations of mean maternal blood or hair Mn concentrations during pregnancy with the endpoints of interest (Table S3), but we did find evidence of effect modification by gestational anemia for the association between average maternal hair Mn concentrations during pregnancy and chest circumference (Table 4). Each ten-fold increase in mean maternal hair Mn concentrations during pregnancy was associated with a 1.19 cm (95% CI: 0.43, 1.95) increase in chest circumference among infants whose mothers did not have gestational anemia, but only a 0.39 cm (95% CI: -0.32, 1.10; $p_{\text{INT}} = 0.14$) increase among infants of mothers who had gestational anemia. We did not observe effect modification by gestational anemia for any other birth outcome.

Sensitivity analyses were conducted excluding outliers and excluding women with missing covariates. Findings were consistent with those from the main multivariate linear regression and generalized additive models, except for the linear association between maternal blood Mn concentrations during the third trimester of pregnancy and birth weight that became not statistically significant. When we reran the analyses including gestational hypertension as a covariate in the models and using blood and hair Mn concentrations by halves of pregnancy, results were not appreciably altered. When both blood and hair Mn were included as predictors of the birth outcomes of interest in the adjusted models, results were similar to those observed in the main multivariate linear regression and generalized additive models for both hair or blood concentrations, except for a positive association between hair Mn concentrations during the second trimester of pregnancy and birth weight (β for a ten-fold increase in Mn concentrations = 89.2 g; 95% CI: 0.4, 178.8) that became statistically significant. Finally, when we assessed the associations of maternal blood and hair Mn concentrations with head circumference/birth weight, head circumference/body length, and

head circumference/chest circumference ratios, results were consistent with those observed with the individual measures of fetal growth (data not shown).

4 Discussion

In this prospective cohort study of Costa Rican women living near banana plantations aerially sprayed with Mn-containing fungicides, we did not observe significant linear or nonlinear adverse associations of maternal Mn concentrations with birth weight and head circumference, as reported in previous studies (Chen et al. 2014; Eum et al. 2014; Guan et al. 2013; Zota et al. 2009). In fact, we observed that higher maternal hair Mn concentrations during the second and third trimesters of pregnancy were associated with larger chest circumferences. We also found that higher maternal hair Mn concentrations averaged during pregnancy were associated with increased chest circumference in infants born to mothers with no gestational anemia.

This study adds to the limited and somewhat conflicting data on the effects of prenatal Mn exposure on fetal growth and length of gestation (Table S1) (Chen et al. 2014; Eum et al. 2014; Guan et al. 2013; Osada et al. 2002; Takser et al. 2004; Vigeh et al. 2008; Xu et al. 2011; Yu and Cao 2013; Zota et al. 2009). Numerous epidemiological studies have examined the association between maternal blood Mn concentrations during pregnancy and birth weight, but only half of them have found significant associations (Chen et al. 2014; Eum et al. 2014; Vigeh et al. 2008; Zota et al. 2009). Cross-sectional studies in Oklahoma, China, and Korea, with different concentrations of Mn in maternal whole blood at delivery (medians = 22, 53.8, and 21.5 $\mu\text{g/L}$, respectively), reported inverted U-shaped relationships between Mn concentrations and birth weight (Chen et al. 2014; Eum et al. 2014; Zota et al. 2009). Maternal blood Mn concentrations in the third trimester and averaged over pregnancy in our study were similar to those observed at delivery in the Oklahoma and Korean study (Eum et al. 2014; Zota et al. 2009) and lower than those found in the Chinese study (Chen et al. 2014), but we did not observe linear or nonlinear associations of blood Mn concentrations with birth weight that would remain statistically significant after excluding outliers. Similarly, the only other prospective cohort study that has examined the relationship of maternal Mn concentrations at different time points during pregnancy (median for third trimester = 15.5 $\mu\text{g/L}$) with fetal growth, besides ours, did not observe any linear or nonlinear associations between blood Mn concentrations and birth weight (Takser et al. 2004). Inconsistencies between the cross-sectional studies that reported inverted U-shaped associations between maternal Mn concentrations and birth weight and our study may arise from differences in the time of biological sampling (at delivery versus third trimester or averaged over pregnancy), exposure pathways, socio-demographic context, and statistical analyses conducted by the researchers (e.g., how confounders were selected and modeled).

Consistent with previous studies (Guan et al. 2013; Takser et al. 2004; Yu and Cao 2013), we did not find associations of maternal blood Mn concentrations with body length and ponderal index. We did not observe an association between blood Mn concentrations during pregnancy and head circumference either. However, a study of 125 Chinese mother-child pairs found an inverted U-shaped relationship of maternal Mn concentrations with head

circumference in their unadjusted analyses (Guan et al. 2013). This Chinese study reported maternal blood Mn concentrations at delivery that were much higher than the Mn concentrations observed in our study during the third trimester (medians = 50.6 versus 26.4 $\mu\text{g/L}$, respectively). This cross-sectional study determined that the main sources of Mn exposure of their study population were living near major transportation routes and exposure to harmful occupational factors (i.e., industrial dust, decorating materials, chemical reagents, and gasoline). Disparities between the Chinese study and ours may be due to differences in the socio-demographic characteristics of the study populations, their exposure levels, and sources of Mn exposure, mainly because it remains unclear how different sources and pathways of exposure to Mn translate into Mn concentrations in various biological media (e.g., blood, hair, teeth) (Smith et al. 2007). Inconsistencies between the two studies may also be due to differences in the statistical methods used by the researchers (i.e., unadjusted vs. adjusted analyses).

To our knowledge, this is the first epidemiological study to assess the relation of hair Mn concentrations during pregnancy with fetal growth and length of gestation. We did not observe associations between maternal hair Mn concentrations and birth weight, body length, ponderal index, or head circumference. Nevertheless, we found positive linear associations of hair Mn concentrations during the second and third trimesters of pregnancy with chest circumference. The only other study that has examined chest circumference, the study of 125 Chinese mother-child pairs mentioned above, reported that chest circumference increased with maternal blood Mn concentrations at delivery up to 55.7 $\mu\text{g/L}$ and then decreased slightly after Mn concentrations exceeded this point (Guan et al. 2013); we had few women with blood Mn concentrations as high as this value and this may have affected our ability to observe a decrease in chest circumference. More importantly, the Chinese study did not measure Mn concentrations in maternal hair samples, and given that blood measurements reflect body Mn dynamics in adults (not specific to pregnant women) on the order of days or weeks whereas hair is thought to integrate circulating Mn concentrations over longer periods of time (Smith et al. 2007), it is not appropriate to compare their findings with ours. It is also important to mention that, in our study, data on chest circumference was only available for a subset of infants and that mothers of those infants had significantly lower hair Mn concentrations compared to mothers of infants for who did not have chest circumference data. We compared the subset of mother-child pairs with chest circumference data with the rest of the study population and did not find any significant differences in their socio-demographic characteristics, which suggests that the possibility of selection bias may be small. However, it is possible that the relationship between hair Mn concentrations and chest circumference may have been different if the entire population had been included in the analyses.

Evidence suggests that iron deficiency results in increased uptake and distribution of Mn in animals and humans (Finley 1999; Heilig et al. 2005; Mena et al. 1969; Smith et al. 2007; Thompson et al. 2007). We did not observe differences in blood or hair Mn concentrations in women who reported having gestational anemia at any time point during pregnancy and women who reported not having it. Nevertheless, we did find a positive association between mean maternal hair Mn concentrations during pregnancy and chest circumference in women without gestational anemia, but not in women with anemia. We hypothesize that the hair Mn

concentrations observed in pregnant women in our study (with and without anemia) may be within the range of Mn concentrations required for normal fetal growth and this is why we observed a positive association with chest circumference. Half of the published studies have adjusted for iron status or hemoglobin concentrations during pregnancy in their analyses of Mn exposure and fetal growth, but none has examined effect modification by anemia. Although some researchers have suggested that chest circumference may be especially sensitive to the environmental factors during late pregnancy (Silventoinen et al. 2012), few studies have been published on the importance of infant chest circumference as an indicator of health outcomes (e.g., it has been referred to as indicator of nutritional status and a screening tool for childhood obesity) (Akaboshi et al. 2012; Nichols 1996; Sundaram et al. 1995). In the absence of associations of blood and hair Mn concentrations with other measures of fetal growth in our study, it remains unclear what the clinical significance of an increased chest circumference may be.

This study has several limitations. To date, there is no consensus on which is the best biomarker to assess human exposure to Mn (Eastman et al. 2013; Smith et al. 2007) and how reliable the available biomarkers are for assessing prenatal exposure to the fetus. Blood and hair Mn concentrations measured in this study may not be the best surrogates for maternal Mn load or fetal exposure, but, unfortunately, cord blood Mn concentrations were not measured. In this study, although we adjusted Mn concentrations by gestational age at the time of sample collection, hair Mn concentrations measured in the one-centimeter hair sample closer to the scalp may not only reflect the exposure at the exact time of collection but over the prior ~ 30 days of hair growth (e.g., if the hair sample was collected in the middle of the first and second trimester the exposure period captured would be the early first and second trimester). In addition, the small sample sizes after stratifying by trimester of pregnancy and limited variability in blood Mn concentrations may have limited our ability to detect potential associations with the birth outcomes of interest. Lastly, we examined the association between Mn and birth growth parameters, but intrauterine growth is a multifactorial process that can be affected by multiple determinants. Unmeasured factors related to the study population, such as nutritional genetic, and lifestyle characteristics, infections, and exposure to other toxic substances during pregnancy, may have biased our results. Despite its limitations, this study has the benefits of a strong prospective design, with blood and hair Mn concentrations measured at different time points during pregnancy, and data on birth outcomes collected by hospital/clinic staff blinded to the Mn exposure status.

In summary, we did not observe the nonlinear associations between maternal blood Mn concentrations and measures of fetal growth that previous studies have reported. However, we did find significant linear associations of maternal hair Mn concentrations with increased chest circumference, but the clinical significance of this finding, in the absence of associations with other birth outcomes, is unknown. Given the complexity of Mn, an essential element that is also an environmental pollutant, the multifactorial nature of the intrauterine growth, and the inconsistencies between previous epidemiological studies, further studies are needed before a clear profile can be determined.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the ISA staff, students, community partners, and, especially, the ISA study participants and their families. We would also like to thank L. Pardo for her assistance with the literature review, C. Hernández and D. Viquez for conducting the postpartum study visits, T. Jursa for his analytical assistance, and R. Gunier for his statistical advice.

Funding sources

This work was funded by research grant numbers 105296-001 from Canada's International Development Research Center; 6807-05-2011/7300127 from Health Canada Global Health Program; D43 ES018745 and R01 ES015572 from the U.S. National Institute for Environmental Health Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the funders.

Abbreviations

BMI	Body Mass Index
CI	Confidence Interval
EBDC	Ethylene Bisdithiocarbamates
ETU	Ethylenethiourea
GAM	Generalized Additive Models
GM	Geometric Mean
GSD	Geometric Standard Deviation
ICC	Intraclass Correlation Coefficient
ICP-MS	Inductively Coupled Plasma Mass Spectrometry
ISA	Infants' Environmental Health study (<i>'Infantes y Salud Ambiental'</i> in Spanish)
LMP	Last Menstrual Period
LOD	Limit of Detection
Mn	Manganese
NS	Not Statistically Significant
SD	Standard Deviation

References

- Akaboshi I, Kitano A, Kan H, Haraguchi Y, Mizumoto Y. Chest circumference in infancy predicts obesity in 3-year-old children. *Asia Pac J Clin Nutr.* 2012; 21(4):495–501. [PubMed: 23017307]
- ATDSR. Toxicological profile for manganese. Agency for Toxic Substances and Disease Registry; Atlanta, GA: 2012.
- Betharia S, Maher TJ. Neurobehavioral effects of lead and manganese individually and in combination in developmentally exposed rats. *Neurotoxicology.* 2012; 33:1117–1127. [PubMed: 22732189]
- Bolze MS, Reeves RD, Lindbeck FE, Kemp SF, Elders MJ. Influence of manganese on growth, somatomedin and glycosaminoglycan metabolism. *J Nutr.* 1985; 115:352–358. [PubMed: 3882911]

- Bouchard M, Laforest F, Vandelac L, Bellinger D, Mergler D. Hair manganese and hyperactive behaviors: Pilot study of school-age children exposed through tap water. *Environ Health Perspect.* 2007; 115:122–127. [PubMed: 17366831]
- Bouchard MF, Sauve S, Barbeau B, Legrand M, Brodeur ME, Bouffard T, et al. Intellectual impairment in school-age children exposed to manganese from drinking water. *Environ Health Perspect.* 2011; 119:138–143. [PubMed: 20855239]
- Chandra SV, Shukla GS. Role of iron deficiency in inducing susceptibility to manganese toxicity. *Arch Toxicol.* 1976; 35:319–323. [PubMed: 989299]
- Chen L, Ding G, Gao Y, Wang P, Shi R, Huang H, et al. Manganese concentrations in maternal-infant blood and birth weight. *Environ Sci Pollut Res Int.* 2014; 21(9):6170–6175. [PubMed: 24477335]
- Clegg MS, Donovan SM, Monaco MH, Baly DL, Ensuna JL, Keen CL. The influence of manganese deficiency on serum IGF-1 and IGF binding proteins in the male rat. *Proc Soc Exp Biol Med.* 1998; 219:41–47. [PubMed: 9751221]
- Colomina MT, Domingo JL, Llobet JM, Corbella J. Effect of day of exposure on the developmental toxicity of manganese in mice. *Vet Hum Toxicol.* 1996; 38(1):7–9. [PubMed: 8825740]
- Eastman RR, Jursa TP, Benedetti C, Lucchini RG, Smith DR. Hair as a biomarker of environmental manganese exposure. *Environ Sci Technol.* 2013; 47:1629–1637. [PubMed: 23259818]
- Eum JH, Cheong HK, Ha EH, Ha M, Kim Y, Hong YC, et al. Maternal blood manganese level and birth weight: A MOCEH birth cohort study. *Environ Health.* 2014; 13(1):31. [PubMed: 24775401]
- Finley JW. Manganese absorption and retention by young women is associated with serum ferritin concentration. *Am J Clin Nutr.* 1999; 70:37–43. [PubMed: 10393136]
- Garcia SJ, Gellein K, Syversen T, Aschner M. Iron deficient and manganese supplemented diets alter metals and transporters in the developing rat brain. *Toxicol Sci.* 2007; 95:205–214. [PubMed: 17060373]
- Gray L Jr, Laskey JW. Multivariate analysis of the effects of manganese on the reproductive physiology and behavior of the male house mouse. *J Toxicol Environ Health.* 1980; 6:861–867. [PubMed: 7420485]
- Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology.* 1999; 10:37–48. [PubMed: 9888278]
- Guan H, Wang M, Li X, Piao F, Li Q, Xu L, et al. Manganese concentrations in maternal and umbilical cord blood: Related to birth size and environmental factors. *Eur J Public Health.* 2013; 24(1):150–157. [PubMed: 23543679]
- Gunier RB, Bradman A, Jerrett M, Smith DR, Harley KG, Austin C, et al. Determinants of manganese in prenatal dentin of shed teeth from CHAMACOS children living in an agricultural community. *Environ Sci Technol.* 2013; 47:11249–11257. [PubMed: 24053404]
- Gunier, RB.; Mora, AM.; Smith, DR.; Arora, M.; Austin, C.; Eskenazi, B., et al. Biomarkers of manganese exposure in pregnant women and children living in an agricultural community in California. submitted
- HaMai D, Bondy SC. Oxidative basis of manganese neurotoxicity. *Ann N Y Acad Sci.* 2004; 1012:129–141. [PubMed: 15105260]
- Hambridge KM, Droegemueller W. Changes in plasma and hair concentrations of zinc, copper, chromium, and manganese during pregnancy. *Obstet Gynecol.* 1974; 44(5):666–672. [PubMed: 4420576]
- Hastie, TJ.; Tibshirani, RJ. Generalized additive models. Chapman and Hall; New York, NY: 1990.
- He P, Liu DH, Zhang GQ. [Effects of high-level-manganese sewage irrigation on children's neurobehavior]. *Zhonghua Yu Fang Yi Xue Za Zhi.* 1994; 28:216–218. [PubMed: 7842882]
- Heilig E, Molina R, Donaghey T, Brain JD, Wessling-Resnick M. Pharmacokinetics of pulmonary manganese absorption: Evidence for increased susceptibility to manganese loading in iron-deficient rats. *Am J Physiol Lung Cell Mol Physiol.* 2005; 288(5):L887–893. [PubMed: 15618452]
- Hernandez-Diaz S, Wilcox AJ, Schisterman EF, Hernan MA. From causal diagrams to birth weight-specific curves of infant mortality. *Eur J Epidemiol.* 2008; 23(3):163–166. [PubMed: 18224448]

- Hiney JK, Srivastava VK, Dees WL. Manganese induces IGF-1 and cyclooxygenase-2 gene expressions in the basal hypothalamus during prepubertal female development. *Toxicol Sci.* 2011; 121:389–396. [PubMed: 21402727]
- Keen CL, Ensunsa JL, Watson MH, Baly DL, Donovan SM, Monaco MH, et al. Nutritional aspects of manganese from experimental studies. *Neurotoxicology.* 1999; 20:213–223. [PubMed: 10385885]
- Keen CL, Ensunsa JL, Clegg MS. Manganese metabolism in animals and humans including the toxicity of manganese. *Met Ions Biol Syst.* 2000; 37:89–121. [PubMed: 10693132]
- Kondakis XG, Makris N, Leotsinidis M, Prinou M, Papapetropoulos T. Possible health effects of high manganese concentration in drinking water. *Arch Environ Health.* 1989; 44:175–178. [PubMed: 2751354]
- Krachler M, Rossipal E, Micetic-Turk D. Trace element transfer from the mother to the newborn-- investigations on triplets of colostrum, maternal and umbilical cord sera. *Eur J Clin Nutr.* 1999; 53(6):486–494. [PubMed: 10403586]
- Leazer TM, Klaassen CD. The presence of xenobiotic transporters in rat placenta. *Drug Metab Dispos.* 2003; 31:153–167. [PubMed: 12527696]
- Lee BK, Kim Y. Relationship between blood manganese and blood pressure in the korean general population according to knhanes 2008. *Environ Res.* 2011; 111:797–803. [PubMed: 21601843]
- Lubin JH, Colt JS, Camann D, Davis S, Cerhan JR, Severson RK, et al. Epidemiologic evaluation of measurement data in the presence of detection limits. *Environ Health Perspect.* 2004; 112:1691–1696. [PubMed: 15579415]
- Mena I, Horiuchi K, Burke K, Cotzias GC. Chronic manganese poisoning. Individual susceptibility and absorption of iron. *Neurology.* 1969; 19:1000–1006. [PubMed: 5387706]
- Menezes-Filho JA, Paes CR, Pontes AM, Moreira JC, Sarcinelli PN, Mergler D. High levels of hair manganese in children living in the vicinity of a ferro-manganese alloy production plant. *Neurotoxicology.* 2009; 30:1207–1213. [PubMed: 19393689]
- Menezes-Filho JA, Novaes Cde O, Moreira JC, Sarcinelli PN, Mergler D. Elevated manganese and cognitive performance in school-aged children and their mothers. *Environ Res.* 2011; 111:156–163. [PubMed: 20943219]
- Molina RM, Phattananarudee S, Kim J, Thompson K, Wessling-Resnick M, Maher TJ, et al. Ingestion of mn and pb by rats during and after pregnancy alters iron metabolism and behavior in offspring. *Neurotoxicology.* 2011; 32:413–422. [PubMed: 21458486]
- Mora AM, van Wendel de Joode B, Mergler D, Cordoba L, Cano C, Quesada R, et al. Blood and hair manganese concentrations in pregnant women from the Infantes' Environmental Health study (ISA) in Costa Rica. *Environ Sci Technol.* 2014; 48:3467–3476. [PubMed: 24601641]
- Nichols MM. Routine chest measurement in the newborn. *Arch Pediatr Adolesc Med.* 1996; 150:771. [PubMed: 8673214]
- Osada H, Watanabe Y, Nishimura Y, Yukawa M, Seki K, Sekiya S. Profile of trace element concentrations in the feto-placental unit in relation to fetal growth. *Acta Obstet Gynecol Scand.* 2002; 81(10):931–937. [PubMed: 12366483]
- Pappas BA, Zhang D, Davidson CM, Crowder T, Park GA, Fortin T. Perinatal manganese exposure: Behavioral, neurochemical, and histopathological effects in the rat. *Neurotoxicol Teratol.* 1997; 19:17–25. [PubMed: 9088007]
- Rabe-Hesketh, S.; Skrondal, A. *Multilevel and longitudinal modeling using Stata.* 3rd ed.. Taylor & Francis; New York, NY: 2012.
- Riojas-Rodriguez H, Solis-Vivanco R, Schilman A, Montes S, Rodriguez S, Rios C, et al. Intellectual function in Mexican children living in a mining area and environmentally exposed to manganese. *Environ Health Perspect.* 2010; 118:1465–1470. [PubMed: 20936744]
- Rosner, B. *Fundamentals of biostatistics.* 6th ed.. Duxbury Press; Pacific Grove, CA: 2006.
- Sanchez DJ, Domingo JL, Llobet JM, Keen CL. Maternal and developmental toxicity of manganese in the mouse. *Toxicol Lett.* 1993; 69:45–52. [PubMed: 8356567]
- Silventoinen K, Kaprio J, Dunkel L, Yokoyama Y. Genetic and environmental influences on chest circumference during infancy: A longitudinal study of japanese twins. *Paediatric and perinatal epidemiology.* 2012; 26:553–560. [PubMed: 23061691]

- Smith D, Gwiazda R, Bowler R, Roels H, Park R, Taicher C, et al. Biomarkers of mn exposure in humans. *Am J Ind Med.* 2007; 50:801–811. [PubMed: 17924418]
- Spencer A. Whole blood manganese levels in pregnancy and the neonate. *Nutrition.* 1999; 15:731–734. [PubMed: 10501283]
- Sundaram KR, Seth V, Jena TK, Shukla DK. Age at which chest circumference overtakes head circumference in children. *Indian J Pediatr.* 1995; 62(1):89–94. [PubMed: 10829849]
- Takser L, Lafond J, Bouchard M, St-Amour G, Mergler D. Manganese levels during pregnancy and at birth: Relation to environmental factors and smoking in a southwest Quebec population. *Environ Res.* 2004; 95:119–125. [PubMed: 15147916]
- Taneja SK, Mandal R. Mineral factors controlling essential hypertension--a study in the Chandigarh, India population. *Biol Trace Elem Res.* 2007; 120:61–73. [PubMed: 17916956]
- Tholin K, Sandstrom B, Palm R, Hallmans G. Changes in blood manganese levels during pregnancy in iron supplemented and non supplemented women. *J Trace Elem Med Biol.* 1995; 9:13–17. [PubMed: 8846152]
- Thompson K, Molina R, Donaghey T, Brain JD, Wessling-Resnick M. The influence of high iron diet on rat lung manganese absorption. *Toxicol Appl Pharmacol.* 2006; 210:17–23. [PubMed: 15993455]
- Thompson K, Molina RM, Donaghey T, Schwob JE, Brain JD, Wessling-Resnick M. Olfactory uptake of manganese requires dmt1 and is enhanced by anemia. *Toxicol Appl Pharmacol.* 2007; 210(1-2): 17–23. [PubMed: 15993455]
- Torrente M, Colomina MT, Domingo JL. Effects of prenatal exposure to manganese on postnatal development and behavior in mice: Influence of maternal restraint. *Neurotoxicol Teratol.* 2002; 24:219–225. [PubMed: 11943509]
- Treinen KA, Gray TJ, Blazak WF. Developmental toxicity of mangafodipir trisodium and manganese chloride in sprague-dawley rats. *Teratology.* 1995; 52:109–115. [PubMed: 8588182]
- van Wendel de Joode B, Mora AM, Cordoba L, Cano C, Quesada R, Skytt A, et al. Mancozeb applications with light aircrafts are associated with increased ethylenethiourea (ETU) concentrations in pregnant women: Results from the Infants' Environmental Health study (ISA). *Environ Health Perspect.* 2014 doi:10.1289/ehp.1307679.
- Vigeh M, Yokoyama K, Ramezanzadeh F, Dahaghin M, Fakhriazad E, Seyedaghamiri Z, et al. Blood manganese concentrations and intrauterine growth restriction. *Reprod Toxicol.* 2008; 25:219–223. [PubMed: 18242051]
- Vigeh M, Yokoyama K, Ohtani K, Shahbazi F, Matsukawa T. Increase in blood manganese induces gestational hypertension during pregnancy. *Hypertens Pregnancy.* 2013; 32(3):214–224. [PubMed: 23905606]
- Wasserman GA, Liu X, Parvez F, Ahsan H, Levy D, Factor-Litvak P, et al. Water manganese exposure and children's intellectual function in Araihazar, Bangladesh. *Environ Health Perspect.* 2006; 114:124–129. [PubMed: 16393669]
- Wilcox AJ. Invited commentary: The perils of birth weight--a lesson from directed acyclic graphs. *Am J Epidemiol.* 2006; 164:1121–1123. discussion 1124–1125. [PubMed: 16931545]
- Xu L, Yokoyama K, Tian Y, Piao FY, Kitamura F, Kida H, et al. Decrease in birth weight and gestational age by arsenic among the newborn in Shanghai, China. *Nihon koshu eisei zasshi.* 2011; 58:89–95. [PubMed: 21473424]
- Yokoi K, Kimura M, Itokawa Y. Effect of dietary iron deficiency on mineral levels in tissues of rats. *Biol Trace Elem Res.* 1991; 29:257–265. [PubMed: 1726408]
- Yu X, Cao L. Elevated cord serum manganese level is associated with a neonatal high ponderal index. *Environ Res.* 2013; 121:79–83. [PubMed: 23164521]
- Zayed J, Vyskocil A, Kennedy G. Environmental contamination and human exposure to manganese--contribution of methylenecyclopentadienyl manganese tricarbonyl in unleaded gasoline. *Int Arch Occup Environ Health.* 1999; 72:7–13. [PubMed: 10029224]
- Zota AR, Eitinger AS, Bouchard M, Amarasiriwardena CJ, Schwartz J, Hu H, et al. Maternal blood manganese levels and infant birth weight. *Epidemiology.* 2009; 20:367–373. [PubMed: 19289966]

Research highlights

- We examined the relation of blood and hair Mn during pregnancy with birth outcomes
- Positive linear association between hair Mn and infant chest circumference
- No association of hair Mn with birth weight and head circumference
- No consistent linear nor nonlinear associations between blood Mn and birth outcomes

Table 1Characteristics of study population, ISA study, Costa Rica, 2010-2011 ($n = 380$).

	n (%)
<i>Maternal characteristics</i>	
Age (years)	
< 18	68 (17.9)
18-24	177 (46.6)
25-29	65 (17.1)
30-34	37 (9.7)
35	33 (8.7)
Education	
6th grade	194 (51.0)
> 6th grade	186 (49.0)
Marital status	
Married or living as married	283 (74.5)
Unmarried	97 (25.5)
Country of birth	
Costa Rica	309 (81.3)
Other Central American countries	71 (18.7)
Family income ^a	
Above poverty line	145 (41.4)
Below poverty line	205 (58.6)
Parity	
0	136 (35.8)
1	244 (64.2)
History of miscarriage	
No	314 (82.6)
Yes	66 (17.4)
Pre-pregnancy BMI (kg/m ²)	
Underweight (< 18.5)	12 (3.2)
Normal (18.5-24.9)	186 (49.0)
Overweight (25.0-29.9)	110 (28.9)
Obese (≥ 30)	72 (18.9)
Smoking during pregnancy	
No	363 (95.5)
Yes	17 (4.5)
Caffeinated tea consumption during pregnancy	
No	301 (79.2)
Yes	79 (20.8)
Iron intake during pregnancy	
No	30 (7.9)
Yes	350 (92.1)

	n (%)
Gestational hypertension ^a	
No	365 (96.3)
Yes	14 (3.4)
Gestational diabetes	
No	365 (96.0)
Yes	15 (4.0)
Gestational anemia	
No	227 (59.7)
Yes	153 (40.3)
Vaginal bleeding during the 1st trimester ^a	
No	338 (96.0)
Yes	14 (4.0)
Work status during pregnancy	
Did not work or worked at home	282 (74.2)
Agricultural work	33 (8.7)
Other work (non-agricultural)	65 (17.1)
Infant characteristics	
Sex	
Male	193 (50.8)
Female	187 (49.2)
Low birth weight	
No	368 (96.8)
Yes	12 (3.2)
Preterm birth	
No	356 (93.7)
Yes	24 (6.3)

Abbreviations: n, number of mother-child pairs; BMI, body mass index.

^aMissing values: family income ($m = 30$, 7.9%), gestational hypertension ($m = 1$, 0.3%), and vaginal bleeding during the 1st trimester ($m = 28$, 7.4%).

Table 2

Distribution of maternal blood and hair Mn concentrations during pregnancy, ISA study, Costa Rica, 2010-2011 ($n = 380$).

Biomarkers	<i>n</i>	Mean (SD)	GM (GSD)	Min	Percentile			Max
					25th	50th	75th	
Blood Mn ($\mu\text{g/L}$)								
By trimesters of pregnancy								
1st trimester	94	22.7 (6.4)	21.8 (1.3)	10.8	18.0	22.5	26.9	36.9
2nd trimester	276	23.8 (6.1)	23.0 (1.3)	9.4	19.8	24.0	27.7	56.3
3rd trimester	203	26.5 (6.6)	25.7 (1.3)	13.2	21.4	26.4	30.3	50.6
Mean during pregnancy ^a	375	24.5 (6.1)	23.7 (1.3)	10.7	20.4	24.0	28.0	50.6
Hair Mn ($\mu\text{g/g}$)								
By trimesters of pregnancy								
1st trimester	107	3.6 (6.4)	1.8 (3.0)	0.1	0.9	1.6	3.8	53.3
2nd trimester	309	3.4 (4.8)	1.8 (3.1)	0.05	0.9	1.7	3.8	36.0
3rd trimester	229	3.6 (6.2)	1.7 (3.2)	0.2	0.7	1.4	3.3	52.6
Mean during pregnancy ^a	380	3.5 (5.3)	1.9 (2.8)	0.1	0.9	1.8	4.0	53.3

Abbreviations: SD, standard deviation; GM, geometric mean; GSD, geometric standard deviation.

^aIn the women for whom only one Mn measurement was available, the single measurement was used in lieu of the average.

Table 3

Adjusted associations of maternal blood and hair (\log_{10}) Mn concentrations during pregnancy with fetal growth (among term births)^d and length of gestation, ISA study, Costa Rica, 2010-2011 ($n = 380$).

Exposure	Birth weight (g)		Body length (cm)		Ponderal index (g/cm ³)		Head circumference (cm)		Chest circumference (cm)		Length of gestation (days)	
	<i>n</i>	β (95% CI)	<i>n</i>	β (95% CI)	<i>n</i>	β (95% CI)	<i>n</i>	β (95% CI)	<i>n</i>	β (95% CI)	<i>n</i>	β (95% CI)
Blood Mn (ng/L),^{b, c}												
By trimesters												
Trimester 1	88	1.5 (-9.7,12.7)	87	0.03 (-0.03,0.09)	87	0.00 (-0.01,0.01)	86	0.02 (-0.03,0.06)	56	0.04 (-0.01,0.09)	94	-0.23 (-0.68,0.21)
Trimester 2	258	2.4 (-5.2,10.1) #	255	-0.02 (-0.06,0.03) #	255	0.01 (0.00,0.01) *	253	-0.01 (-0.04,0.02)	128	-0.02 (-0.06,0.03)	276	-0.06 (-0.33,0.20) #
Trimester 3	194	9.0 (0.5,17.5) **	190	0.03 (-0.02,0.08)	190	0.00 (0.00,0.01)	187	0.02 (-0.01,0.05)	79	0.01 (-0.04,0.07)	203	-0.02 (-0.24,0.21)
Mean during pregnancy	351	3.1 (-3.6,9.8)	347	0.00 (-0.04,0.04)	347	0.00 (0.00,0.01)	343	-0.01 (-0.03,0.02)	153	-0.02 (-0.06,0.02)	375	-0.10 (-0.31,0.12)
Hair Mn ($\mu\text{g/g}$),^{d, e}												
By trimesters												
Trimester 1	95	28.9 (-113.9,171.7)	94	-0.18 (-0.92,0.57)	94	0.07 (-0.03,0.18)	93	0.19 (-0.37,0.76)	60	0.26 (-0.48,0.99)	101	2.61 (-2.75,7.98)
Trimester 2	274	68.8 (-18.2,155.8)	270	0.13 (-0.37,0.63)	270	0.04 (-0.04,0.11)	268	-0.02 (-0.33,0.29)	133	0.62 (0.16,1.08) **	290	-0.04 (-2.82,2.73)
Trimester 3	202	32.5 (-75.6,140.5)	198	-0.01 (-0.66,0.65)	198	0.04 (-0.05,0.13)	196	-0.12 (-0.51,0.28)	82	0.55 (-0.16,1.26)	215	-0.72 (-3.88,2.43)
Mean during pregnancy	331	58.9 (-28.9,146.8)	326	0.01 (-0.48,0.50)	326	0.05 (-0.02,0.12)	323	-0.03 (-0.34,0.29)	154	0.66 (0.19,1.13) **	352	0.68 (-2.09,3.44)

Abbreviations: CI, confidence interval.

^aModels for birth weight, body length, ponderal index, head circumference, and chest circumference include only term births.

^bResults are for one-unit increase in blood Mn concentrations.

^cAdjusted for maternal education, parity, smoking during pregnancy, pre-pregnancy body mass index, infant sex, gestational diabetes, caffeinated tea intake during pregnancy, and timing of prenatal care initiation. Models for birth weight, body length, ponderal index, head circumference, and chest circumference also adjusted for length of gestation and length of gestation squared.

^dResults are for 10-fold increase in hair Mn concentrations.

^eAdjusted for maternal education, parity, smoking during pregnancy, pre-pregnancy body mass index, infant sex, maternal country of birth, vaginal bleeding in the first trimester of gestation ($m = 28$, 7.4%), caffeinated tea intake during pregnancy, iron intake during pregnancy, and maternal age. Models for birth weight, ponderal index, body length, ponderal index, head circumference, and chest circumference also adjusted for length of gestation and length of gestation squared.

* PLinear < 0.10

** pLinear < 0.05

pGAM < 0.05

Table 4

Adjusted associations of maternal blood and hair (\log_{10}) Mn concentrations during pregnancy with fetal growth (among term births)^a and length of gestation stratified by gestational anemia, ISA study, Costa Rica, 2010-2011 ($n = 380$).

Outcome	Mean blood Mn ($\mu\text{g/L}$) ^{b, c}			Mean hair Mn ($\mu\text{g/g}$) ^{d, e}		
	<i>n</i>	β (95% CI)	<i>p</i> _{INT}	<i>n</i>	β (95% CI)	<i>p</i> _{INT}
Birth weight (g)						
No gestational anemia	205	6.2 (-2.2,14.5)	0.32	195	42.8 (-76.5,162.2)	0.99
Gestational anemia	146	-3.1 (-14.4,8.1)		136	27.2 (-106.8,161.2)	
Body length (cm)						
No gestational anemia	204	0.00 (-0.05,0.05)	0.95	193	-0.22 (-0.90,0.47)	0.92
Gestational anemia	143	-0.01 (-0.07,0.05) [#]		133	-0.08 (-0.83,0.67)	
Ponderal index (g/cm^3)						
No gestational anemia	204	0.01 (0.00,0.01)	0.31	193	0.09 (-0.01,0.19) [*]	0.66
Gestational anemia	143	0.00 (-0.01,0.01)		133	0.02 (-0.07,0.11) [#]	
Head circumference (cm)						
No gestational anemia	202	0.00 (-0.03,0.03)	0.24	191	0.11 (-0.31,0.52)	0.52
Gestational anemia	141	-0.03 (-0.07,0.02)		132	-0.24 (-0.75,0.26)	
Chest circumference (cm)						
No gestational anemia	82	-0.02 (-0.07,0.04)	0.75	83	1.19 (0.43,1.95) ^{**}	0.14
Gestational anemia	71	-0.04 (-0.10,0.03)		71	0.39 (-0.32,1.10)	
Length of gestation (days)						
No gestational anemia	223	-0.06 (-0.33,0.21)	0.49	213	0.38 (-3.50,4.25)	0.92
Gestational anemia	152	-0.17 (-0.54,0.21)		139	1.05 (-2.87,4.97)	

Abbreviations: CI, confidence interval.

^aModels for birth weight, body length, ponderal index, head circumference, and chest circumference include only term births.

^bResults are for each one-unit increase in blood Mn concentrations.

^cAdjusted for maternal education, parity, smoking during pregnancy, pre-pregnancy body mass index, infant sex, gestational diabetes, caffeinated tea intake during pregnancy, and timing of prenatal care initiation. Models for birth weight, body length, ponderal index, head circumference, and chest circumference also adjusted for length of gestation and length of gestation squared.

^dResults are for each 10-fold increase in hair Mn concentrations.

^eAdjusted for maternal education, parity, smoking during pregnancy, pre-pregnancy body mass index, infant sex, maternal country of birth, vaginal bleeding in the first trimester of gestation ($m = 28$, 7.4%), caffeinated tea intake during pregnancy, iron intake during pregnancy, and maternal age. Models for birth weight, body length, ponderal index, head circumference, and chest circumference also adjusted for length of gestation and length of gestation squared.

* PLinear < 0.10

** PLinear < 0.05

PGAM < 0.05