

Published in final edited form as:

J Cell Sci. 2007 September 1; 120(0 17): 3138–3146. doi:10.1242/jcs.005298.

The distinct roles of Ras and Rac in PI 3-kinase-dependent protrusion during EGF-stimulated cell migration

Shu-Chin Yip^{1,2,*}, Mirvat El-Sibai^{1,*}, Salvatore J. Coniglio³, Ghassan Mouneimne², Robert J. Eddy², Beth E. Drees³, Paul O. Neilsen³, Sumanta Goswami⁴, Marc Symons⁵, John S. Condeelis², and Jonathan M. Backer^{1,‡}

¹Department of Molecular Pharmacology, Albert Einstein College of Medicine, Bronx, NY 10461, USA

²Department of Anatomy and Structural Biology, Albert Einstein College of Medicine, Bronx, NY 10461, USA

³Echelon Biosciences, Inc., Salt Lake City, UT, USA

⁴Department of Biology, Yeshiva University, NY, USA

⁵Center for Oncology and Cell Biology, Institute for Medical Research at North Shore-LIJ, Manhasset, NY, USA

Summary

Cell migration involves the localized extension of actin-rich protrusions, a process that requires Class I phosphoinositide 3-kinases (PI 3-kinases). Both Rac and Ras have been shown to regulate actin polymerization and activate PI 3-kinase. However, the coordination of Rac, Ras and PI 3-kinase activation during epidermal growth factor (EGF)-stimulated protrusion has not been analyzed. We examined PI 3-kinase-dependent protrusion in MTLn3 rat adenocarcinoma cells. EGF-stimulated phosphatidyl-inositol (3,4,5)-trisphosphate [PtdIns(3,4,5)P₃] levels showed a rapid and persistent response, as PI 3-kinase activity remained elevated up to 3 minutes. The activation kinetics of Ras, but not Rac, coincided with those of leading-edge PtdIns(3,4,5)P₃ production. Small interfering RNA (siRNA) knockdown of K-Ras but not Rac1 abolished PtdIns(3,4,5)P₃ production at the leading edge and inhibited EGF-stimulated protrusion. However, Rac1 knockdown did inhibit cell migration, because of the inhibition of focal adhesion formation in Rac1 siRNA-treated cells. Our data show that in EGF-stimulated MTLn3 carcinoma cells, Ras is required for both PtdIns(3,4,5)P₃ production and lamellipod extension, whereas Rac1 is required for formation of adhesive structures. These data suggest an unappreciated role for Ras during protrusion, and a crucial role for Rac in the stabilization of protrusions required for cell motility.

Keywords

Ras; Rac; Cdc42; Rho; Lamellipodia

[‡]Author for correspondence (backer@aecom.yu.edu).

^{*}These authors contributed equally to this work

Introduction

Cell migration is an essential property of cells engaged in embryogenesis, inflammation, wound healing and tumor metastasis. Motility toward a source of chemoattractant depends on localized extension of actin-rich protrusions, such as lamellipodia. Studies with metastatic rat carcinoma line MTLn3 have shown that actin polymerization at the leading edge of the lamellipod relies upon the synergistic interaction of Arp2/3 complex with actin-severing protein cofilin, which generates new actin barbed ends at the leading edge (Bailey et al., 2001; Chan et al., 2000; DesMarais et al., 2004; Ichetovkin et al., 2002). Arp2/3-mediated actin polymerization in MTLn3 is likely to be activated by Wiskott-Aldrich syndrome protein (WASP) family proteins, which are in turn regulated by upstream Rho GTPases (Bompard and Caron, 2004; Machesky et al., 1999; Miki and Takenawa, 2003; Weaver et al., 2003).

The kinetics of actin polymerization in globally epidermal growth factor (EGF)-stimulated MTLn3 cells, measured either by lamellipod extension or free barbed end formation, is highly reproducible. Lamellipod extension begins after 30-60 seconds of EGF stimulation, and is maximal within 180-300 seconds (Chan et al., 1998; Segall et al., 1996). Barbed end formation, visualized by the incorporation of fluorescently labeled biotin-actin, exhibits early and late transients, at 1 and 3 minutes (Chan et al., 1998; Mouneimne et al., 2004). Early barbed end formation at 1 minute is linked to directional sensing, whereas late barbed end formation at 3 minutes seems to be involved in the production of mechanical protrusive force (Ghosh et al., 2004; Mouneimne et al., 2004). Whereas phosphoinositide 3-kinase (PI 3-kinase) is required for protrusion in MTLn3 cells (Hill et al., 2000), inhibition of PI 3-kinase primarily suppresses the late, more than the early, barbed end transient (Mouneimne et al., 2004). A similar diminution of late actin polymerization was observed in *Dictyostelium pi3k1/2*-null cells (Chen et al., 2003; Funamoto et al., 2002). These data are consistent with recent data from a variety of systems, suggesting that PI 3-kinase activation at the leading edge of moving cells is required for maximal speed of migration, but is not involved in directional sensing (Andrew and Insall, 2007; Ferguson et al., 2007; Hoeller and Kay, 2007; Nishio et al., 2007).

How is PI 3-kinase activity regulated at the leading edge of the lamellipod? Class I PI 3-kinases can be directly activated by interactions with receptor tyrosine kinases and their substrates, with small GTPases, and with $\beta\gamma$ -subunits from activated trimeric G-proteins (Vanhaesebroeck et al., 1997). In polarized neutrophils, the asymmetrical accumulation of phosphatidylinositol (3,4,5)-trisphosphate [PtdIns(3,4,5) P_3] at the leading edge has been proposed to involve a PI 3-kinase-dependent positive feedback loop between PI 3-kinase and Rac (Bourne and Weiner, 2002; Srinivasan et al., 2003; Weiner et al., 2002), consistent with the fact that Rac activates PI 3-kinase by binding to the breakpoint cluster region (BCR) of p85 regulatory subunits (Bokoch et al., 1996; Zheng et al., 1994) and PtdIns(3,4,5) P_3 production enhances the activity of guanine nucleotide exchange factors for Rac activation (Hawkins et al., 1995; Welch et al., 2003). However, membrane localization of Akt-PH-green fluorescent protein (GFP), a probe for phosphatidylinositol (4,5)-bisphosphate [PtdIns(3,4) P_2] and PtdIns(3,4,5) P_3 , is blocked by the Rho GTPase inhibitor

toxin B in formyl-methionyl-leucyl-proline (fMLP)- but not in insulin-stimulated cells (Servant et al., 2000; Srinivasan et al., 2003; Weiner et al., 2002), suggesting that PI 3-kinase localization is differentially regulated by G-protein-coupled receptors as opposed to receptor tyrosine kinases. Although RacB and PI 3-kinase have similarly been proposed to constitute a positive feedback loop in *Dictyostelium* (Park et al., 2004; Sasaki et al., 2004), the mechanism is less clear because there is no homolog of the PI 3-kinase p85 regulatory subunit and therefore no obvious way for activated Rac to regulate PI 3-kinase activity.

Ras is also an upstream activator of PI 3-kinase, and has been implicated in growth factor-induced membrane ruffling (Bar-Sagi et al., 1987; Kundra et al., 1995; Price et al., 1999; Rodriguez-Viciana et al., 1994; Rodriguez-Viciana et al., 1997). In mammalian cells, EGF has been shown to activate Ras at the cell periphery in a study using a fluorescence resonance energy transfer (FRET)-based biosensor (Mochizuki et al., 2001). Overexpression of constitutively active Ras^{V12} induced strong membrane ruffling, whereas inhibition of endogenous Ras prevented scatter factor/hepatocyte growth factor (SF/HGF)-stimulated actin reorganization and spreading (Bar-Sagi and Feramisco, 1986; Ridley, 1995; Ridley et al., 1992; Rodriguez-Viciana et al., 1997). Similarly, cAMP-induced motility was defective in *Dictyostelium* cells expressing either Ras-null or RasG^{N17} mutants (Insall et al., 1996; Kae et al., 2004; Sasaki et al., 2004; Tuxworth et al., 1997), and platelet-derived growth factor (PDGF)-stimulated motility was dependant on Ras activity in 3T3 cells (Kundra et al., 1995).

p85/p110 α PI 3-kinase activity is required for EGF-stimulated lamellipod extension in MTLn3 cells (Condeelis, 2001; Hill et al., 2000; Mouneimne et al., 2004). However, the relative contribution of Ras and Rac to PI 3-kinase-mediated protrusion has not been examined, and the hypothesis that Rac and PI 3-kinase form a positive feedback loop at the leading edge has not been tested in EGF-responsive cells. In this study we have examined the mechanism of PI 3-kinase activation and PI 3-kinase-dependent lamellipod extension in EGF-stimulated cells. We find a crucial role of Ras but not Rac in EGF-stimulated PI 3-kinase activation at the leading edge of cells and lamellipod extension. Rac is required for cell motility, but this is because of its affects on formation of adhesions behind the edge of cell protrusions. Thus, Ras and Rac play distinct, but coordinated, roles during EGF-stimulated protrusion in carcinoma cells.

Results

EGF-induced PtdIns(3,4,5) P_3 production at the leading edge of carcinoma cells

We measured the production of PtdIns(3,4,5) P_3 at the leading edge of EGF-stimulated MTLn3 cells using a monoclonal anti-PtdIns(3,4,5) P_3 antibody (Echelon Bioscience, Salt Lake City, UT). In quiescent MTLn3 cells, minimal membrane-associated PtdIns(3,4,5) P_3 was observed (Fig. 1A, 0 minutes). Anti-PtdIns(3,4,5) P_3 staining increased rapidly after EGF stimulation, appearing as punctate structures in a narrow band at the cell edge. PtdIns(3,4,5) P_3 staining was sixfold above basal at 1 minute (Fig. 1B). Staining of EGF-stimulated cells was specifically blocked by pre-absorption of the antibody with PtdIns(3,4,5) P_3 , but not by other phosphoinositides (S.-C.Y. et al., unpublished). After the peak at 1 minute, leading-edge PtdIns(3,4,5) P_3 remained 2.5-fold above basal until 3

minutes, when the protrusion of the lamellipod in MTLn3 cells is maximal (Chan et al., 1998; Segall et al., 1996). PtdIns(3,4,5) P_3 staining returned to basal levels by 4-5 minutes of stimulation. These data show that EGF-induced PtdIns(3,4,5) P_3 production is rapid, persists throughout the first 3 minutes of EGF stimulation, and is evident at the leading edge of the lamellipod. Given that the half-life of PtdIns(3,4,5) P_3 in MTL3 cells is under 10 seconds (S.-C.Y. et al., unpublished), the sustained elevation of PtdIns(3,4,5) P_3 at the leading edge of the lamellipod requires the presence of sustained local activation of PI 3-kinase.

We next examined the kinetics of phosphotyrosine-associated PI 3-kinase activity. Phosphotyrosine-associated PI 3-kinase activity peaked sharply at \sim 4.3-fold above basal at 40 seconds, followed by a slower decline that returned to basal activity at 5 minutes (Fig. 1C). The correlation between phosphotyrosine-associated PI 3-kinase activity and the kinetics of leading-edge PtdIns(3,4,5) P_3 suggests that PI 3-kinase activation at the leading edge of the lamellipod is regulated at least in part by direct binding to tyrosine-phosphorylated proteins.

Kinetics of Ras and Rac activation in EGF-stimulated carcinoma cells

We compared the time-course of PtdIns(3,4,5) P_3 production to that of Rac and Ras activation, because both GTPases have been implicated in PI 3-kinase activation (Bokoch et al., 1996; Pacold et al., 2000; Rodriguez-Viciano et al., 1994). Rac and Ras activation in MTLn3 cells was measured in GST-pulldown assays using the CRIB domain from Pak1 and the Ras-binding domain (RBD) from Raf, respectively (Benard et al., 1999; de Rooij and Bos, 1997). Total levels of Ras and Rac in cells lysates did not change during 5 minutes of EGF stimulation (data not shown). In resting MTLn3 cells, the levels of activated Rac and Ras were low (Fig. 2A). Both GTPases were activated rapidly after EGF stimulation. Rac activity was maximal at 50-60 seconds, but the activation was transient and returned almost to basal levels by 3 minutes (Fig. 2A,B, open circles). Consistent with previous findings (Benard et al., 1999), Rac GTPase activity was inhibited by 100 nM wortmannin (Fig. 2A, top panel). Although EGF-induced Ras activation was also maximal at 40-60 seconds, Ras activation persisted throughout 5 minutes of stimulation (Fig. 2A,B, closed diamonds). Interestingly, Ras activation is also inhibited by wortmannin treatment (Fig. 2A), suggesting that Ras activity is regulated in part by PI 3-kinase. These data show that both Ras activation and leading-edge PtdIns(3,4,5) P_3 production exhibit a persistent activation, whereas Rac activation is largely gone by 3 minutes.

We also measured the localization of Rac and Ras. Staining of EGF-stimulated cells with anti-Rac antibody after various periods of EGF stimulation showed that Rac was rapidly but transiently recruited to the cell edge in EGF-stimulated cells (Fig. 2C). Anti-Rac staining was specific, as it was abolished by pre-absorption with GST-Rac or by small interfering RNA (siRNA) knockdown of Rac expression (supplementary material Fig. S1A,B). We also measured the intracellular distribution of activated Ras in EGF-stimulated cells by staining with GST-RBD followed by anti-GST antibodies. In resting cells, there was a minimal level of activated Ras detected in the plasma membrane. After EGF stimulation, an increase in fluorescent intensity was observed at both plasma membrane and perinuclear regions as previously reported (Chiu et al., 2002; Hancock, 2003; Mochizuki et al., 2001) (Fig. 2D).

GST-RBD immunostaining at the leading edge persisted for at least 3 minutes, and did not return to basal levels until 4-5 minutes of EGF stimulation (Fig. 2D). The Ras staining was specific, as it was abolished by pretreatment of cells with the farnesylation inhibitor manumycin A (data not shown), by incubating with recombinant GST prior to incubation with anti-GST antibody, and by combined knockdown of N-Ras and K-Ras (supplementary material Fig. S1C,D). Taken together, data in Figs 1 and 2 show that the kinetics of Ras activation and membrane localization is similar to the sustained kinetics of PI(3,4,5)P₃ production in EGF-stimulated cells, whereas Rac activation and membrane localization is transient.

Inhibition of Ras but not Rac reduces leading-edge PtdIns(3,4,5)P₃ production

To examine the requirement for Ras versus Rac in the regulation of PtdIns(3,4,5)P₃ production at the leading edge, we knocked down each GTPase using siRNA. Rac1 is the predominant Rac isoform in MTLn3 cells, as levels of Rac2 and Rac3 were barely detectable by quantitative RT-PCR (Q-RT-PCR) (M.S., unpublished). To inhibit Rac1, we transiently transfected MTLn3 cells with siRNA duplexes directed against luciferase (negative control) and Rac1. Optimal suppression of Rac1 occurred at 48 hours post-transfection, when Rac1 knockdown efficiency was approximately 80-85% as detected by anti-Rac immunoblots (see supplementary material Fig. S2A, upper panel). We measured the effect of siRNA-mediated Rac1 knockdown on EGF-stimulated production of PtdIns(3,4,5)P₃ at the leading edge. Control or Rac1 siRNA-treated cells were stimulated with EGF for 1 or 3 minutes, and then stained with anti-PtdIns(3,4,5)P₃ antibodies. Knockdown of Rac1 expression had no effect on PtdIns(3,4,5)P₃ production at the leading edge (Fig. 3A): in both control and Rac1 siRNA-treated cells, EGF-stimulated production of PtdIns(3,4,5)P₃ at the leading edge of both was maximal at 1 minute, and remained 3-4-fold above basal levels at 3 minutes (Fig. 3B). These results show that inhibition of Rac expression does not affect PtdIns(3,4,5)P₃ production at the leading edge of the lamellipod.

Q-RT-PCR analysis of MTLn3 cells showed that N-Ras and K-Ras were expressed at similar levels, and that we were able to achieve 70-80% knockdown of N-Ras and K-Ras using a pool of four distinct siRNA oligonucleotides for each protein (see supplementary material Fig. S2B). Although knockdown of H-Ras was only 40% efficient, control MTLn3 cells express 20-fold less H-Ras than N-Ras or K-Ras (supplementary material Fig. S2B). Anti-PtdIns(3,4,5)P₃ staining of EGF-stimulated cells showed that knockdown of K-Ras had little effect on PtdIns(3,4,5)P₃ levels at 1 minute, but markedly inhibited production at 3 minutes (Fig. 3C,D). By contrast, transfection of cells with control siRNA (luciferase) or H-Ras siRNA had no effect on PtdIns(3,4,5)P₃ production at either time point (Fig. 3D). siRNA knockdown of N-Ras has no effect at 1 minute and a modest effect at 3 minutes (Fig. 3D). Similarly, treatment of cells with the farnesyltransferase inhibitor manumycin A also reduced PI(3,4,5)P₃ production at 3 minutes (data not shown). Thus, EGF-stimulated PtdIns(3,4,5)P₃ production is inhibited by knockdown of K-Ras and, to a lesser extent, N-Ras, but not by knockdown of Rac1.

Specific inhibition of Ras but not Rac blocks EGF-stimulated protrusion

Both Rac and Ras have been implicated in membrane protrusion and PI 3-kinase activation (Bokoch et al., 1996; Rodriguez-Viciana et al., 1997; Welch et al., 2003). Moreover, overexpression of constitutively active Rac in MTLn3 cells is sufficient to induce lamellipod extension (El-Sibai et al., 2007). To determine whether Rac is necessary for EGF-stimulated protrusion in MTLn3 cells, we used time-lapse video microscopy to measure the surface area of control or Rac1 siRNA-treated cells during EGF stimulation. Although Rac1 siRNA-treated cells showed a mixed morphology, with both rounded and normally spread cells, all of these cells were capable of extending lamellipodia in response to EGF (supplementary material Movie 1). Although the mean area of quiescent Rac1 siRNA-treated cells was less than that of control cells ($P<0.01$), Rac1 siRNA-treated cells achieved a final area that was not significantly different than that of the control cells after 3 minutes of EGF stimulation (Fig. 4A). Thus, protrusion relative to initial area was greater in the Rac siRNA-treated cells than in control cells ($P<0.001$ at 1 and $P<0.01$ at 3 minutes; Fig. 4B), but this reflected the smaller initial size rather than a true hyperprotrusion. Similar results were obtained after knockdown of Rac1 with a pool of four distinct siRNA oligonucleotides (supplementary material Fig. S2A, upper panel; supplementary material Fig. S3) and in cells treated with the Rac-specific inhibitor NSC23766, whose efficacy against Rac was confirmed in a CRIB domain pulldown assay (see supplementary material Fig. S4A,B). These results show that Rac1 is not required for the protrusion step in EGF-stimulated cell motility in carcinoma cells.

To determine the effects of specific inhibition of Ras on EGF-stimulated lamellipod extension, we measured EGF-stimulated protrusion in cells transfected with pools of four siRNA oligonucleotides for each Ras isoform. Protrusion was measured in control or Ras siRNA-treated cells stimulated with EGF; data from a single experiment (Fig. 4B) and pooled data from two experiments (Fig. 4C) are shown. Whereas a slight decrease in cell protrusion is seen with H-Ras siRNA, protrusion was inhibited by almost 45% in cells treated with N-Ras siRNA, and nearly 60% in cells treated with K-Ras siRNA (Fig. 4C). Whereas inhibition by N-Ras siRNA did not reach statistical significance, inhibition by K-RAS siRNA was significant ($P<0.05$). Knockdown of both N-Ras and K-Ras did not exacerbate the phenotype seen with K-Ras siRNA alone. We also saw an inhibition of EGF-stimulated protrusion in MTLn3 cells microinjected with the Y13-259 inhibitory anti-Ras antibody (data not shown). These data clearly demonstrate a requirement for K-Ras, and to a lesser extent N-Ras, in EGF-stimulated protrusion.

To further explore the formation of lamellipodia in Rac and Ras knockdown cells, we measured the EGF-stimulated recruitment of Arp2/3 to the leading edge, which corresponds to the zone of new barbed end formation (Bailly et al., 1999). We found that Arp2/3 recruitment in Rac1 siRNA-treated cells was indistinguishable from that in control cells (Fig. 5). By contrast, in K-Ras knockdown cells, Arp2/3 recruitment to the cell edge was normal at 1 minute but was markedly suppressed at 3 minutes of EGF stimulation.

Rac is required for cell locomotion

As our data show that Ras, but not Rac, is required for EGF-stimulated PtdIns(3,4,5) P_3 production and protrusion, we tested whether Rac might be required for other steps in the motility cycle. We performed a motion analysis of time-lapse movies of control and Rac1 siRNA-treated cells, recorded over 1 hour in the presence of 5% serum (supplementary material Movie 2). The Rac1 siRNA-treated cells showed an approximately fourfold reduction in net path; data are provided for a single siRNA oligonucleotide and a pool of four distinct siRNA oligonucleotides (Table 1). Similar results were obtained using the Rac-specific inhibitor NSC23766 (Table 1), which also caused a fourfold decrease in net path.

We note that unlike control MTLn3 cells, in which EGF stimulation suppresses ruffling (Segall et al., 1996), we see significant membrane ruffling in Rac1 siRNA-treated cells (supplementary material Movie 1). Ruffling can result from the dissipation of a lamellipod, when membrane protrusions fail to adhere at the leading edge (BurrIDGE and Wennerberg, 2004). In addition, the surface area of unstimulated Rac knockdown cells is approximately 30% less than in control cells (Fig. 4A), which could result from a defect in cell adhesion. To test whether Rac1 knockdown affects formation of adhesive structures, we stained control and Rac siRNA-treated cells with anti-paxillin antibodies. In control cells, EGF stimulates an increase in cell area, with a proliferation of focal adhesions behind the edge of the protruding cell (Fig. 6, upper panels). However, knockdown of Rac1 leads to a significant reduction in focal adhesion in both unstimulated and EGF-stimulated cells (Fig. 6, lower panels). These data suggest that the reduced motility of Rac1 knockdown cells is because of a reduced capacity to form adhesions after protrusion.

Discussion

PtdIns(3,4,5) P_3 production at the leading edge of acutely EGF-stimulated carcinoma cells is rapid and persistent, as PtdIns(3,4,5) P_3 levels remain elevated for up to 3 minutes. Given that the half-life of PtdIns(3,4,5) P_3 in MTLn3 cells is less than 10 seconds (S.-C.Y. et al., unpublished), the kinetics of PtdIns(3,4,5) P_3 accumulation reflect the kinetics of PI 3-kinase activation. Although the mechanism of PI 3-kinase activation at the cell edge at 1 minute is not yet clear, phosphotyrosine-associated PI 3-kinase activity is also maximal at 1 minute, and PtdIns(3,4,5) P_3 accumulation at 1 minute is not significantly inhibited by suppression of either Ras or Rac. Thus, leading-edge PtdIns(3,4,5) P_3 at 1 minute is probably due in part to direct binding of Class IA PI 3-kinase to tyrosine phosphorylated EGF receptor substrates such as Gab1 (Holgado-Madruga et al., 1996). Recent data from our laboratory suggest that Cdc42 may also be involved in regulation of the early PtdIns(3,4,5) P_3 peak (El-Sibai et al., 2007).

By contrast, siRNA knockdown of Ras, but not Rac, inhibits PtdIns(3,4,5) P_3 production at the leading edge at 3 minutes. Consistent with this, EGF-stimulated PtdIns(3,4,5) P_3 production is kinetically correlated with the activation of Ras at the leading edge, whereas Rac is only transiently activated and its activity is markedly reduced by 3 minutes. The K-Ras isoform plays the major role in this process, although a modest reduction in PtdIns(3,4,5) P_3 levels was seen after N-Ras knockdown.

Unlike fMLP-stimulated neutrophils, in which a positive feedback loop between Rac and PI 3-kinase has been postulated (Servant et al., 2000; Srinivasan et al., 2003; Weiner et al., 2002), EGF-stimulated PtdIns(3,4,5) P_3 production in MTLn3 cells is Rac independent, whereas wortmannin inhibits Rac activation. Thus, in carcinoma cells, EGF-stimulated Rac activation requires PI 3-kinase, but PI 3-kinase activation at the leading edge does not require Rac. However, our data do support the possibility of a positive feedback loop between Ras and PI 3-kinase, which may contribute to PI 3-kinase-dependent lamellipod extension. Treatment of cells with Ras siRNA reduces PtdIns(3,4,5) P_3 levels at the leading edge at 3 minutes, and EGF-stimulated Ras activation is reduced in wortmannin-treated cells. The possibility of a local positive feedback loop between Ras and PI 3-kinase is supported by previous studies showing that Ras-GTP activates PI 3-kinase by direct interaction with the p110 catalytic subunit (Kodaki et al., 1994; Rodriguez-Viciana et al., 1994), and that PI 3-kinase contributes to Ras activation through the PtdIns(3,4,5) P_3 -mediated enhancement of Gab1 phosphorylation and recruitment of Grb2/Sos (Rodrigues et al., 2000; Yart et al., 2001).

We previously showed that PI 3-kinase is required for barbed end formation at 3 minutes, and is involved in the generation of protrusive force (Mouneimne et al., 2004). siRNA knockdown of K-Ras, and to a lesser extent N-Ras, inhibits EGF-stimulated PtdIns(3,4,5) P_3 production at 3 minutes but not 1 minute, and the effect of Ras knockdown on protrusion is most pronounced at later time points (3-5 minutes), whereas protrusion at early time points is relatively unaffected. These data support an important role for sustained Ras-mediated PtdIns(3,4,5) P_3 production in EGF-stimulated lamellipod extension. Ras has been previously implicated in growth factor-induced chemotaxis in both mammalian systems (Kundra et al., 1995; Price et al., 1999; Rodriguez-Viciana et al., 1997) and *Dictyostelium* (Insall et al., 1996; Kae et al., 2004; Sasaki et al., 2004; Tuxworth et al., 1997). A recent study in fibroblasts suggested that the K(B)-Ras isoform in particular was required for PDGF-stimulated migration (Liao et al., 2006).

The failure of Rac inhibition or siRNA knockdown to inhibit EGF-stimulated lamellipod formation is surprising, given that the formation of a Rac/IRSp53/WASP family verprolin-homologous protein 2 (WAVE2) complex and other Rac-dependent WAVE-activation complexes have been implicated in membrane ruffling and lamellipod extension (Gautreau et al., 2004; Miki et al., 1998; Miki et al., 2000; Weiner et al., 2006). It is unlikely that the residual 20% Rac activity after siRNA treatment accounts for the lack of effects on lamellipod extension at 3 minutes, given that we did see effects of Rac1 knockdown on motility. Interestingly, expression of constitutively active Rac does produce lamellipodia in MTLn3 cells (El-Sibai et al., 2007), suggesting that Rac can drive protrusion in these cells. However, in control MTLn3 cells, the kinetics of Rac activation after acute EGF stimulation are extremely transient, and other PI 3-kinase-dependent mechanisms appear to regulate protrusion. Whereas Rac is required for protrusion in many cell types, protrusion and chemotaxis in the absence of Rac activity have been previously described in colon carcinoma cells, macrophages and dendritic cells (O'Connor et al., 2000; West et al., 2000; Wheeler et al., 2006). Ruffling and/or protrusion induced by expression of activated Rab5 or the Cdc42 effector MSE55, or by infection with *Salmonella typhimurium*, are also Rac1-

independent (Burbelo et al., 1999; Jones et al., 1993; Spaargaren and Bos, 1999). We have not elucidated the mechanism of Rac-independent protrusion, but it could involve the recruitment of WAVE2 by PtdIns(3,4,5) P_3 production at the cell edge (Oikawa et al., 2004). Recent data from our laboratory have also implicated the PI 3-kinase-dependent activation of Cdc42 as playing an important role in protrusion in MTLn3 cells (El-Sibai et al., 2007).

If Rac siRNA-treated cells undergo EGF-stimulated protrusion, how can we explain their defect in motility? Time-lapse studies (supplementary material Movie 2) suggest that although Rac1 knockdown cells rapidly extend membrane protrusions, these structures are not stable and do not persist in any direction. Activated Rac has been reported to promote integrin-mediated adhesions in migrating cells, where new lamellipodia are stabilized by formation of adhesions (Kiosses et al., 2001), and we find that focal adhesions are greatly reduced in Rac1 knockdown cells. Thus, a likely explanation for the reduced motility of Rac1 siRNA-treated cells is the failure to establish stable adhesions after lamellipod extension. This is consistent with our previous report that overexpression of dominant-negative Rac greatly reduces both cell spreading and the number of focal adhesions during protrusion (Bouzahzah et al., 2001). Rac and PI 3-kinase activities have been shown to be required for recruitment of Arp2/3 complex to vinculin, an integrin-associated protein, at new adhesion sites at the leading edge of lamellipod in EGF-stimulated A431 cells (DeMali et al., 2002; DeMali and Burridge, 2003). This suggests that a coordination of membrane protrusion and Rac-mediated recruitment of adhesion complexes to the leading edge may be necessary for MTLn3 cells to migrate efficiently.

Although our data suggests that Ras-mediated activation of PI 3-kinase at 3 minutes is required for the generation of protrusive force, we do not yet understand the mechanistic role of EGF-stimulated leading-edge PtdIns(3,4,5) P_3 production at 1 minute. This event temporally coincides with the early transient of actin polymerization, which has been implicated in directional sensing in EGF-stimulated tumor cells (Ghosh et al., 2004; Mouneimne et al., 2004). EGF-induced PI 3-kinase activation at the leading edge at 1 minute may contribute to early barbed end formation, as residual barbed ends were seen after inhibition of cofilin with siRNA or blocking antibodies (approximately 20% of maximum) (Mouneimne et al., 2004). Alternatively, PI 3-kinase activation at 1 minute may initiate signaling events that contribute to the formation or stabilization of focal contacts at the leading edge at later times. The regulation and function of PI 3-kinase and Rac during the initial responses to EGF stimulation will be an important issue for future study.

Materials and Methods

Materials

Anti-PtdIns(3,4,5) P_3 antibody (IgG) was a generous gift from Echelon Biosciences (Salt Lake City, UT). Anti-phosphotyrosine antibody (PY20) was purchased from Signal Transduction Laboratories. Anti-EGF receptor antibody and anti-phospho-Erk antibody was from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-Rac antibody (clone 23A8) and anti-phospho-Ser473 AKT antibodies were purchased from Upstate Biotechnology (Charlottesville, VA). Anti-Ras antibody (Y13-259) was provided by T. Michaeli (Albert Einstein College of Medicine, Bronx, NY). Manumycin A, *Clostridium Dificile* toxin B and

wortmannin were obtained from CalBiochem (San Diego, CA). NSC23766 was a gift from Yi Zheng (Children's Hospital Research Foundation, Cincinnati, OH). siRNA duplexes were purchased from Dharmacon.

Cell culture and EGF stimulation

The rat mammary adenocarcinoma breast cancer cell line, MTLn3, has been previously described (Segall et al., 1996). In each experiment, MTLn3 cells were starved in L15 media (Gibco BRL) supplemented with 0.35% bovine serum albumin (BSA) in a 37°C, non-CO₂ incubator for 3-4 hours. Cells were then stimulated with 5 nM murine EGF (Upstate Biotechnology) for various times.

Microscopy and immunofluorescence analysis

For immunofluorescence staining procedures, cells were plated on collagen-coated coverslips 24 hours prior to the experiment. After incubation without or with EGF for various times, cells were rapidly fixed and permeabilized by immersion in 3.7% paraformaldehyde, 0.1% glutaraldehyde and 0.15 mg/ml saponin in fix buffer (5 mM KCl, 137 mM NaCl, 4 mM NaHCO₃, 0.4 mM KH₂PO₄, 1.1 mM Na₂HPO₄, 2 mM MgCl₂, 5 mM PIPES, pH 7.2, 2 mM EGTA and 5.5 mM glucose) for 1 hour at 37°C. The fixed cells were incubated with primary antibodies for 1 hour, followed by secondary antibodies for 45 minutes, and mounted in 6 mg/ml of N-propyl gallate prepared in 50% glycerol-phosphate-buffered saline (PBS) (v/v).

All fluorescent images were obtained using a 60 × 1.4 NA Olympus objective optic and a cooled CCD camera. For fluorescence quantification, all digital images were imported in NIH image software and analyzed using a previously described macro (DesMarais et al., 2002). This macro collects pixel intensities from the perimeter of the cell in a 0.22 μm stepwise manner. The pixel intensities in the leading-edge compartment, defined as 0.66 μm from the perimeter of the cells, were averaged and normalized to the edge intensity of non-stimulated cells.

Immunoprecipitation and western blots

For immunoprecipitations, MTLn3 cells were plated on 100 mm tissue culture dishes. EGF-stimulated cells were rinsed with cold PBS containing 1 mM sodium vanadate and lysed in 20 mM Tris, 137 mM NaCl, 1 mM MgCl₂, 1 mM CaCl₂, 1% NP-40, 10% glycerol, pH 7.5, supplemented with protease inhibitors [1 mM phenylmethylsulphonyl fluoride (PMSF), 1 μg/ml leupeptin, 1 μg/ml aprotinin]. Immunoprecipitated proteins were separated by SDS-PAGE and transferred to PVDF membranes. After blocking, membranes were incubated with primary antibodies at 4°C overnight followed by horseradish peroxidase (HRP)-conjugated secondary antibodies, and visualized using enhanced chemiluminescence (ECL) (Amersham Biosciences). For measurement of phosphotyrosine-associated PI 3-kinase activity, anti-phosphotyrosine (PY20; Signal Transduction Laboratory) immunoprecipitates were assayed for lipid kinase activity towards phosphatidylinositol, as described (Yu et al., 1998).

GST-CRIB and GST-RBD protein purification and pull-down assays

GST fusion proteins containing the PAK1 GTPase-binding domain [hPAK(67-150), CRIB domain] or the Raf1 RBD (amino acids 51-131, RBD domain) were generous gifts from Klaus M. Hahn (Scripps Research Institute, La Jolla, CA) and Linda van Aelst (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY), and were purified from *Escherichia coli*. To measure activation of Rac and Ras, EGF-stimulated cells were washed with cold PBS containing 1 mM sodium vanadate and lysed in magnesium lysis buffer (MLB) (25 mM HEPES, pH 7.5, 1% Igapal, 150 mM NaCl, 10 mM MgCl₂, 10% glycerol, 1 mM EDTA, 1 mM vanadate) containing protease inhibitors (1 mM PMSF, 1 µg/ml leupeptin, 1 µg/ml aprotinin). Immobilized GST-CRIB or RBD (8 µg per assay) was incubated with the cell lysates at 4°C for 30 minutes on a rotating wheel, washed three times with MLB and suspended in 50 µl Laemmli sample buffer. Proteins were separated by 14% SDS-PAGE, transferred to PVDF membranes and blotted with anti-Ras or anti-Rac antibodies.

GST-CRIB and GST-RBD immunostaining

EGF-stimulated cells were fixed and incubated with purified GST-CRIB or GST-RBD for 1 hour, followed by anti-GST antibody and secondary antibody.

siRNA and Q-RT-PCR analysis

Rac1 siRNA sequences were designed specifically against rat Rac1: 5'-AAAGAGAUCGGUGCUGUCAAAA-3'. Control siRNA duplexes were targeted against luciferase: 5'-CGTACGCGGAATACTTCGA-3' Rac1 was also knocked down using a pool of four siRNA duplexes (Smartpools, Dharmacon). For knockdown of Ras isoforms, pools of four siRNA duplexes for each Ras isoform (Smartpools, Dharmacon) were used. Cells were transfected with 10 nM siRNA using Lipofectamine 2000 (Invitrogen) for 48 hours. For analysis of Rac1 suppression, SDS-sample buffer was added directly to non-stimulated cells, and anti-Rac expression was analyzed by western blotting. For analysis of Rac and Ras isoform suppression, total mRNA from MTLn3 cells was isolated using the RNeasy mini kit (Qiagen). mRNA levels were analyzed by Q-RT-PCR as previously described (Chan et al., 2005; Wang et al., 2002). For signaling and motility experiments, siRNA-treated cells were plated on collagen-coated coverslips or acid-treated glass-bottom MatTek dishes (MatTek, Ashland, MA) at 24 hours after transfection.

Area extension analysis in EGF live upshift assays

For time-lapse experiments, quiescent cells were EGF stimulated on the microscope using a heated stage and an objective heater. Images were collected at 10-second intervals using 20× 0.4 NA or 40× 0.55 NA objectives with a video camera. Cell surface area was analyzed using NIH Image. Statistical significance was calculated using a two-tailed Student's *t*-test.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the Analytical Imaging Facility at Albert Einstein College of Medicine, especially to Michael Cammer for his valuable assistance in image analysis. This work was supported by National Institutes of Health grants PO1 CA 100324 (J.C., J.M.B.) and T32 DK 07513 (S.-C.Y.), and the Albert Einstein Cancer Center (NIC CA13330).

References

- Andrew N, Insall RH. Chemotaxis in shallow gradients is mediated independently of PtdIns 3-kinase by biased choices between random protrusions. *Nat Cell Biol.* 2007; 9:193–200. [PubMed: 17220879]
- Bailly M, Macaluso F, Cammer M, Chan A, Segall JE, Condeelis JS. Relationship between Arp2/3 complex and the barbed ends of actin filaments at the leading edge of carcinoma cells after epidermal growth factor stimulation. *J Cell Biol.* 1999; 145:331–345. [PubMed: 10209028]
- Bailly M, Ichetovkin I, Grant W, Zebda N, Machesky LM, Segall JE, Condeelis J. The F-actin side binding activity of the Arp2/3 complex is essential for actin nucleation and lamellipod extension. *Curr Biol.* 2001; 11:620–625. [PubMed: 11369208]
- Bar-Sagi D, Feramisco JR. Induction of membrane ruffling and fluid-phase pinocytosis in quiescent fibroblasts by ras proteins. *Science.* 1986; 233:1061–1068. [PubMed: 3090687]
- Bar-Sagi D, McCormick F, Milley RJ, Feramisco JR. Inhibition of cell surface ruffling and fluid-phase pinocytosis by microinjection of anti-ras antibodies into living cells. *J Cell Physiol Suppl.* 1987; 5:69–73. [PubMed: 3316255]
- Benard V, Bohl BP, Bokoch GM. Characterization of rac and cdc42 activation in chemoattractant-stimulated human neutrophils using a novel assay for active GTPases. *J Biol Chem.* 1999; 274:13198–13204. [PubMed: 10224076]
- Bokoch GM, Vlahos CJ, Wang Y, Knaus UG, Traynor-Kaplan AE. Rac GTPase interacts specifically with phosphatidylinositol 3-kinase. *Biochem J.* 1996; 315:775–779. [PubMed: 8645157]
- Bompard G, Caron E. Regulation of WASP/WAVE proteins: making a long story short. *J Cell Biol.* 2004; 166:957–962. [PubMed: 15452139]
- Bourne HR, Weiner O. A chemical compass. *Nature.* 2002; 419:21. [PubMed: 12214215]
- Bouzahzah B, Albanese C, Ahmed F, Pixley F, Lisanti MP, Segall JD, Condeelis J, Joyce D, Minden A, Der CJ, et al. Rho family GTPases regulate mammary epithelium cell growth and metastasis through distinguishable pathways. *Mol Med.* 2001; 7:816–830. [PubMed: 11844870]
- Burbelo PD, Snow DM, Bahou W, Spiegel S. MSE55, a Cdc42 effector protein, induces long cellular extensions in fibroblasts. *Proc Natl Acad Sci USA.* 1999; 96:9083–9088. [PubMed: 10430899]
- Burridge K, Wennerberg K. Rho and Rac take center stage. *Cell.* 2004; 116:167–179. [PubMed: 14744429]
- Chan AY, Raft S, Bailly M, Wyckoff JB, Segall JE, Condeelis JS. EGF stimulates an increase in actin nucleation and filament number at the leading edge of the lamellipod in mammary adenocarcinoma cells. *J Cell Sci.* 1998; 111:199–211. [PubMed: 9405304]
- Chan AY, Bailly M, Zebda N, Segall JE, Condeelis JS. Role of cofilin in epidermal growth factor-stimulated actin polymerization and lamellipod protrusion. *J Cell Biol.* 2000; 148:531–542. [PubMed: 10662778]
- Chan AY, Coniglio SJ, Chuang YY, Michaelson D, Knaus UG, Philips MR, Symons M. Roles of the Rac1 and Rac3 GTPases in human tumor cell invasion. *Oncogene.* 2005; 24:7821–7829. [PubMed: 16027728]
- Chen L, Janetopoulos C, Huang YE, Iijima M, Borleis J, Devreotes PN. Two phases of actin polymerization display different dependencies on PI(3,4,5)P3 accumulation and have unique roles during chemotaxis. *Mol Biol Cell.* 2003; 14:5028–5037. [PubMed: 14595116]
- Chiu VK, Bivona T, Hach A, Sajous JB, Silletti J, Wiener H, Johnson RL 2nd, Cox AD, Philips MR. Ras signalling on the endoplasmic reticulum and the Golgi. *Nat Cell Biol.* 2002; 4:343–350. [PubMed: 11988737]

- Condeelis J. How is actin polymerization nucleated in vivo? *Trends Cell Biol.* 2001; 11:288–293. [PubMed: 11413039]
- de Rooij J, Bos JL. Minimal Ras-binding domain of Raf1 can be used as an activation-specific probe for Ras. *Oncogene.* 1997; 14:623–625. [PubMed: 9053862]
- DeMali KA, Burridge K. Coupling membrane protrusion and cell adhesion. *J Cell Sci.* 2003; 116:2389–2397. [PubMed: 12766185]
- DeMali KA, Barlow CA, Burridge K. Recruitment of the Arp2/3 complex to vinculin: coupling membrane protrusion to matrix adhesion. *J Cell Biol.* 2002; 159:881–891. [PubMed: 12473693]
- DesMarais V, Ichetovkin I, Condeelis J, Hitchcock-DeGregori SE. Spatial regulation of actin dynamics: a tropomyosin-free, actin-rich compartment at the leading edge. *J Cell Sci.* 2002; 115:4649–4660. [PubMed: 12415009]
- DesMarais V, Macaluso F, Condeelis J, Bailly M. Synergistic interaction between the Arp2/3 complex and cofilin drives stimulated lamellipod extension. *J Cell Sci.* 2004; 117:3499–3510. [PubMed: 15252126]
- El-Sibai M, Nalbant P, Pang H, Flinn RJ, Sarmiento C, Macaluso F, Cammer M, Condeelis JS, Hahn KM, Backer JM. Cdc42 is required for EGF-stimulated protrusion and motility in MTLn3 carcinoma cells. *J Cell Sci.* 2007 in press.
- Ferguson GJ, Milne L, Kulkarni S, Sasaki T, Walker S, Andrews S, Crabbe T, Finan P, Jones G, Jackson S, et al. PI(3)Kgamma has an important context-dependent role in neutrophil chemokinesis. *Nat Cell Biol.* 2007; 9:86–91. [PubMed: 17173040]
- Funamoto S, Meili R, Lee S, Parry L, Firtel RA. Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell.* 2002; 109:611–623. [PubMed: 12062104]
- Gautreau A, Ho HY, Li J, Steen H, Gygi SP, Kirschner MW. Purification and architecture of the ubiquitous Wave complex. *Proc Natl Acad Sci USA.* 2004; 101:4379–4383. [PubMed: 15070726]
- Ghosh M, Song X, Mouneimne G, Sidani M, Lawrence DS, Condeelis JS. Cofilin promotes actin polymerization and defines the direction of cell motility. *Science.* 2004; 304:743–746. [PubMed: 15118165]
- Hancock JF. Ras proteins: different signals from different locations. *Nat Rev Mol Cell Biol.* 2003; 4:373–384. [PubMed: 12728271]
- Hawkins PT, Eguinoa A, Qiu RG, Stokoe D, Cooke FT, Walters R, Wennstrom S, Claesson-Welsh L, Evans T, Symons M, et al. PDGF stimulates an increase in GTP-Rac via activation of phosphoinositide 3-kinase. *Curr Biol.* 1995; 5:393–403. [PubMed: 7627555]
- Hill K, Welti S, Yu J, Murray JT, Yip SC, Condeelis JS, Segall JE, Backer JM. Specific requirement for the p85-p110alpha phosphatidylinositol 3-kinase during epidermal growth factor-stimulated actin nucleation in breast cancer cells. *J Biol Chem.* 2000; 275:3741–3744. [PubMed: 10660520]
- Hoeller O, Kay RR. Chemotaxis in the absence of PI(3,4,5)P3 gradients. *Curr Biol.* 2007; 17:813–817. [PubMed: 17462897]
- Holgado-Madruga M, Emler DR, Moscatello DK, Godwin AK, Wong AJ. A Grb2-associated docking protein in EGF- and insulin-receptor signalling. *Nature.* 1996; 379:560–564. [PubMed: 8596638]
- Ichetovkin I, Grant W, Condeelis J. Cofilin produces newly polymerized actin filaments that are preferred for dendritic nucleation by the Arp2/3 complex. *Curr Biol.* 2002; 12:79–84. [PubMed: 11790308]
- Insall RH, Borleis J, Devreotes PN. The aimless RasGEF is required for processing of chemotactic signals through G-protein-coupled receptors in Dictyostelium. *Curr Biol.* 1996; 6:719–729. [PubMed: 8793298]
- Jones BD, Paterson HF, Hall A, Falkow S. Salmonella typhimurium induces membrane ruffling by a growth factor-receptor-independent mechanism. *Proc Natl Acad Sci USA.* 1993; 90:10390–10394. [PubMed: 8234304]
- Kae H, Lim CJ, Spiegelman GB, Weeks G. Chemoattractant-induced Ras activation during Dictyostelium aggregation. *EMBO Rep.* 2004; 5:602–606. [PubMed: 15143344]
- Kiosses WB, Shattil SJ, Pampori N, Schwartz MA. Rac recruits high-affinity integrin alphavbeta3 to lamellipodia in endothelial cell migration. *Nat Cell Biol.* 2001; 3:316–320. [PubMed: 11231584]

- Kodaki T, Woscholski R, Hallberg B, Rodriguez-Viciana P, Downward J, Parker PJ. The activation of phosphatidylinositol 3-kinase by Ras. *Curr Biol*. 1994; 4:798–806. [PubMed: 7820549]
- Kundra V, Anand-Apte B, Feig LA, Zetter BR. The chemotactic response to PDGF-BB: evidence of a role for Ras. *J Cell Biol*. 1995; 130:725–731. [PubMed: 7622571]
- Liao J, Planchon SM, Wolfman JC, Wolfman A. Growth factor dependent AKT activation and cell migration requires the function of c-K(B)-Ras versus other cellular Ras isoforms. *J Biol Chem*. 2006; 281:29730–29738. [PubMed: 16908523]
- Machesky LM, Mullins RD, Higgs HN, Kaiser DA, Blanchoin L, May RC, Hall ME, Pollard TD. Scar, a WASP-related protein, activates nucleation of actin filaments by the Arp2/3 complex. *Proc Natl Acad Sci USA*. 1999; 96:3739–3744. [PubMed: 10097107]
- Miki H, Takenawa T. Regulation of actin dynamics by WASP family proteins. *J Biochem*. 2003; 134:309–313. [PubMed: 14561714]
- Miki H, Suetsugu S, Takenawa T. WAVE, a novel WASP-family protein involved in actin reorganization induced by Rac. *EMBO J*. 1998; 17:6932–6941. [PubMed: 9843499]
- Miki H, Yamaguchi H, Suetsugu S, Takenawa T. IRSp53 is an essential intermediate between Rac and WAVE in the regulation of membrane ruffling. *Nature*. 2000; 408:732–735. [PubMed: 11130076]
- Mochizuki N, Yamashita S, Kurokawa K, Ohba Y, Nagai T, Miyawaki A, Matsuda M. Spatio-temporal images of growth-factor-induced activation of Ras and Rap1. *Nature*. 2001; 411:1065–1068. [PubMed: 11429608]
- Mouneimne G, Soon L, DesMarais V, Sidani M, Song X, Yip SC, Ghosh M, Eddy R, Backer JM, Condeelis J. Phospholipase C and cofilin are required for carcinoma cell directionality in response to EGF stimulation. *J Cell Biol*. 2004; 166:697–708. [PubMed: 15337778]
- Nishio M, Watanabe K, Sasaki J, Taya C, Takasuga S, Iizuka R, Balla T, Yamazaki M, Watanabe H, Itoh R, et al. Control of cell polarity and motility by the PtdIns(3,4,5)P3 phosphatase SHIP1. *Nat Cell Biol*. 2007; 9:36–44. [PubMed: 17173042]
- O'Connor KL, Nguyen BK, Mercurio AM. RhoA function in lamellae formation and migration is regulated by the alpha6beta4 integrin and cAMP metabolism. *J Cell Biol*. 2000; 148:253–258. [PubMed: 10648558]
- Oikawa T, Yamaguchi H, Itoh T, Kato M, Ijuin T, Yamazaki D, Suetsugu S, Takenawa T. PtdIns(3,4,5)P3 binding is necessary for WAVE2-induced formation of lamellipodia. *Nat Cell Biol*. 2004; 6:420–426. [PubMed: 15107862]
- Pacold ME, Suire S, Perisic O, Lara-Gonzalez S, Davis CT, Walker EH, Hawkins PT, Stephens L, Eccleston JF, Williams RL. Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma. *Cell*. 2000; 103:931–943. [PubMed: 11136978]
- Park KC, Rivero F, Meili R, Lee S, Apone F, Firtel RA. Rac regulation of chemotaxis and morphogenesis in Dictyostelium. *EMBO J*. 2004; 23:4177–4189. [PubMed: 15470506]
- Price JT, Tiganis T, Agarwal A, Djakiew D, Thompson EW. Epidermal growth factor promotes MDA-MB-231 breast cancer cell migration through a phosphatidylinositol 3'-kinase and phospholipase C-dependent mechanism. *Cancer Res*. 1999; 59:5475–5478. [PubMed: 10554021]
- Ridley AJ. Microinjection of Rho and Rac into quiescent Swiss 3T3 cells. *Meth Enzymol*. 1995; 256:313–320. [PubMed: 7476447]
- Ridley AJ, Paterson HF, Johnston CL, Diekmann D, Hall A. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell*. 1992; 70:401–410. [PubMed: 1643658]
- Rodrigues GA, Falasca M, Zhang Z, Ong SH, Schlessinger J. A novel positive feedback loop mediated by the docking protein Gab1 and phosphatidylinositol 3-kinase in epidermal growth factor receptor signaling. *Mol Cell Biol*. 2000; 20:1448–1459. [PubMed: 10648629]
- Rodriguez-Viciana P, Warne PH, Dhand R, Vanhaesebroeck B, Gout I, Fry MJ, Waterfield MD, Downward J. Phosphatidylinositol-3-OH kinase as a direct target of Ras. *Nature*. 1994; 370:527–532. [PubMed: 8052307]
- Rodriguez-Viciana P, Warne PH, Khwaja A, Marte BM, Pappin D, Das P, Waterfield MD, Ridley A, Downward J. Role of phosphoinositide 3-OH kinase in cell transformation and control of the actin cytoskeleton by Ras. *Cell*. 1997; 89:457–467. [PubMed: 9150145]
- Sasaki AT, Chun C, Takeda K, Firtel RA. Localized Ras signaling at the leading edge regulates PI3K, cell polarity, and directional cell movement. *J Cell Biol*. 2004; 167:505–518. [PubMed: 15534002]

- Segall JE, Tyerech S, Boselli L, Masseling S, Helft J, Chan A, Jones J, Condeelis J. EGF stimulates lamellipod extension in metastatic mammary adenocarcinoma cells by an actin-dependent mechanism. *Clin Exp Metastasis*. 1996; 14:61–72. [PubMed: 8521618]
- Servant G, Weiner OD, Herzmark P, Balla T, Sedat JW, Bourne HR. Polarization of chemoattractant receptor signaling during neutrophil chemotaxis. *Science*. 2000; 287:1037–1040. [PubMed: 10669415]
- Spaargaren M, Bos JL. Rab5 induces Rac-independent lamellipodia formation and cell migration. *Mol Biol Cell*. 1999; 10:3239–3250. [PubMed: 10512863]
- Srinivasan S, Wang F, Glavas S, Ott A, Hofmann F, Aktories K, Kalman D, Bourne HR. Rac and Cdc42 play distinct roles in regulating PI(3,4,5)P3 and polarity during neutrophil chemotaxis. *J Cell Biol*. 2003; 160:375–385. [PubMed: 12551955]
- Tuxworth RI, Cheetham JL, Machesky LM, Spiegelmann GB, Weeks G, Insall RH. Dictyostelium RasG is required for normal motility and cytokinesis, but not growth. *J Cell Biol*. 1997; 138:605–614. [PubMed: 9245789]
- Vanhaesebroeck B, Leever SJ, Panayotou G, Waterfield MD. Phosphoinositide 3-kinases: a conserved family of signal transducers. *Trends Biochem Sci*. 1997; 22:267–272. [PubMed: 9255069]
- Wang W, Wyckoff JB, Frohlich VC, Oleynikov Y, Huttelmaier S, Zavadil J, Cermak L, Bottinger EP, Singer RH, White JG, et al. Single cell behavior in metastatic primary mammary tumors correlated with gene expression patterns revealed by molecular profiling. *Cancer Res*. 2002; 62:6278–6288. [PubMed: 12414658]
- Weaver AM, Young ME, Lee WL, Cooper JA. Integration of signals to the Arp2/3 complex. *Curr Opin Cell Biol*. 2003; 15:23–30. [PubMed: 12517700]
- Weiner OD, Neilsen PO, Prestwich GD, Kirschner MW, Cantley LC, Bourne HR. A PtdInsP(3)- and Rho GTPase-mediated positive feedback loop regulates neutrophil polarity. *Nat Cell Biol*. 2002; 4:509–513. [PubMed: 12080346]
- Weiner OD, Rentel MC, Ott A, Brown GE, Jedrychowski M, Yaffe MB, Gygi SP, Cantley LC, Bourne HR, Kirschner MW. Hem-1 complexes are essential for Rac activation, actin polymerization, and myosin regulation during neutrophil chemotaxis. *PLoS Biol*. 2006; 4:e38. [PubMed: 16417406]
- Welch HC, Coadwell WJ, Stephens LR, Hawkins PT. Phosphoinositide 3-kinase-dependent activation of Rac. *FEBS Lett*. 2003; 546:93–97. [PubMed: 12829242]
- West MA, Prescott AR, Eskelinen EL, Ridley AJ, Watts C. Rac is required for constitutive macropinocytosis by dendritic cells but does not control its downregulation. *Curr Biol*. 2000; 10:839–848. [PubMed: 10899002]
- Wheeler AP, Wells CM, Smith SD, Vega FM, Henderson RB, Tybulewicz VL, Ridley AJ. Rac1 and Rac2 regulate macrophage morphology but are not essential for migration. *J Cell Sci*. 2006; 119:2749–2757. [PubMed: 16772332]
- Yart A, Laffargue M, Mayeux P, Chretien S, Peres C, Tonks N, Roche S, Payrastra B, Chap H, Raynal P. A critical role for phosphoinositide 3-kinase upstream of Gab1 and SHP2 in the activation of ras and mitogen-activated protein kinases by epidermal growth factor. *J Biol Chem*. 2001; 276:8856–8864. [PubMed: 11134009]
- Yu J, Zhang Y, McIlroy J, Rordorf-Nikolic T, Orr GA, Backer JM. Regulation of the p85/p110 phosphatidylinositol 3'-kinase: stabilization and inhibition of the p110alpha catalytic subunit by the p85 regulatory subunit. *Mol Cell Biol*. 1998; 18:1379–1387. [PubMed: 9488453]
- Zheng Y, Bagrodia S, Cerione RA. Activation of phosphoinositide 3-kinase activity by Cdc42Hs binding to p85. *J Biol Chem*. 1994; 269:18727–18730. [PubMed: 8034624]




Fig. 1. EGF-stimulated PtdIns(3,4,5) P_3 production at the leading edge of lamellipod is rapid and sustained. Quiescent MTLn3 cells were stimulated with 5 nM EGF and fixed at various times. (A) Selected images of EGF-stimulated cells stained with anti-PtdIns(3,4,5) P_3 antibody. Bar, 10 μ m for all panels. (B) Quantification of leading-edge PtdIns(3,4,5) P_3 production following EGF stimulation. The average edge fluorescence (0-0.66 μ m from the perimeter) was measured as described in the Materials and Methods, normalized for the edge fluorescence of unstimulated cells and plotted as a function of time. The data are the mean \pm s.e.m. from four experiments. (C) Anti-phosphotyrosine immunoprecipitates from EGF-stimulated cells were assayed for PI 3-kinase activity. The data are the mean \pm s.e.m. from four experiments.




Fig. 2.

The kinetics of EGF-stimulated Ras and Rac activation and translocation. (A,B) GST-CRIB (from hPAK) or GST-RBD (from Raf 1) were immobilized on glutathione-Sepharose beads and incubated with EGF-stimulated cell lysates as described. Glutathione-Sepharose pull-downs were separated by 14% SDS-PAGE gel and blotted with anti-Rac or anti-Ras antibody. (A, upper panel) A representative anti-Rac immunoblot of a GST-CRIB pull-down from EGF-stimulated cells. (Lower panel) A representative anti-Ras immunoblot of a GST-RBD pull-down from EGF-stimulated cells. Where indicated, cells were treated with 100 nM wortmannin prior to EGF stimulation. Total Ras and Rac levels did not change during 5 minutes of EGF stimulation (data not shown). (B) Ras (closed diamonds) and Rac (open circles) pull-down assays were quantified by densitometry. Data were expressed as percentage of maximum activity, and show the mean \pm s.e.m. from nine experiments. (C) Representative fluorescent images of MTLn3 cells stimulated with EGF for various times and stained with anti-Rac antibodies. Bar, 10 μ m. (D) Representative images of EGF-stimulated MTLn3 cells stained with GST-RBD followed by anti-GST antibody, to visualize endogenous activated Ras. Bar, 10 μ m.





Fig. 3. Inhibition of Ras but not Rac reduces PtdIns(3,4,5)P₃ production at the leading edge of lamellipodia. (A,B) MTLn3 cells were transfected with control or Rac1 siRNA. (A) Representative images of control siRNA or Rac1 siRNA-treated cells stimulated with EGF for 3 minutes and stained with anti-PtdIns(3,4,5)P₃ antibody are shown. (B) Quantification of leading-edge PtdIns(3,4,5)P₃ in siRNA-treated cells. The data are mean \pm s.e.m. from four experiments. Bar, 10 μ m in all panels. (C,D) MTLn3 cells were transfected with control siRNA or pools of four siRNA duplexes specific for H-Ras, K-Ras or N-Ras. (C) Representative images of control siRNA or K-Ras siRNA-treated cells stimulated with EGF for 0, 1 or 3 minutes and stained with anti-PtdIns(3,4,5)P₃ antibody are shown. (D) Quantification of leading-edge PtdIns(3,4,5)P₃ in siRNA-treated cells. The data are mean \pm s.e.m. from 40 cells per condition, and are representative of two independent experiments.

**Fig. 4.**

Specific inhibition of Ras, but not Rac, inhibits EGF-stimulated protrusion. (A) MTLn3 cells were transfected with control or Rac1 siRNA. Time-lapse images of EGF-stimulated cells were recorded (20 seconds per frame) with a 20×0.4 NA objective, and surface areas were measured using NIH Image. The data show the mean ± s.e.m. from 41 cells per time point. (B) Data from A were normalized to the initial area of each cell. (C) MTLn3 cells were transfected with control siRNA or isoform-specific siRNA targeting H-Ras, K-Ras, N-Ras or both N-Ras and K-Ras. The surface area of control siRNA or Ras siRNA-treated cells was measured after various times of EGF stimulation. Cell areas were normalized to the initial area of each cell. Data show the mean ± s.e.m. from 9-14 cells per time point. (D) Pooled data from two experiments showing maximal EGF-stimulated protrusion in cells treated with control or anti-Ras siRNA. Data are the mean ± s.d.




Fig. 5. Specific inhibition of Ras, but not Rac, inhibits EGF-stimulated Arp2/3 recruitment. MTLn3 cells were transfected with control (left panels), Rac1 (middle panels) or K-Ras (right panels) siRNA. Cells were stimulated with EGF for 0, 1 or 3 minutes, fixed and stained with anti-Arp2/3 antibodies.




Fig. 6. Rac1 knockdown inhibits formation of focal adhesions. MTLn3 cells were transfected with control (luciferase, upper panels) or Rac1 (lower panels) siRNA. Cells were stimulated with EGF for 0, 1 or 3 minutes, fixed and stained with anti-paxillin antibodies.

Table 1
Rac is required for cell motility

	Net path (μm)
Control siRNA	19.4 \pm 2.0 ($n=30$)
Rac1 siRNA	5.4 \pm 0.6 ($n=37$)
Control siRNA	12.0 \pm 2.1 ($n=15$)
Rac1 siRNA (pooled oligonucleotides)	3.0 \pm 0.6 ($n=13$)
Control cells (DMSO)	30.9 \pm 1.7 ($n=20$)
NSC23766 (100 μM)	7.1 \pm 0.7 ($n=20$)

MTLn3 cells were transfected with control siRNA, a single Rac siRNA or a pool of four distinct Rac1 siRNAs. Alternatively, cells were treated with the Rac inhibitor NSC23766 or carrier (dimethyl sulfoxide, DMSO). Time-lapse phase images of cells in 5% serum were collected (1 frame/min). Net path was calculated using DIAS software (Solltech, Oakdale, CA).