

RESEARCH ARTICLE

The Shared Crosstalk of Multiple Pathways Involved in the Inflammation between Rheumatoid Arthritis and Coronary Artery Disease Based on a Digital Gene Expression Profile

Xuyan Niu^{1,9}, Cheng Lu^{1,9}, Cheng Xiao^{2,9}, Zhiguo Zhang³, Miao Jiang¹, Dan He⁴, Yanqin Bian⁴, Ge Zhang⁵, Zhaoxiang Bian⁵, Aiping Lu^{1,4,5*}

1. Institute of Basic Research in Clinical Medicine, China Academy of Chinese Medical Sciences, Beijing, 100700, China, 2. China-Japan Friendship Hospital, Beijing, 100029, China, 3. Institute of Basic Theory, China Academy of Chinese Medical Sciences, Beijing, 100700, China, 4. E-Institute of Chinese Traditional Internal Medicine, Shanghai Municipal Education Commission, Shanghai, 201203, China, 5. Institute for Advancing Translational Medicine in Bone & Joint Diseases, School of Chinese Medicine, Hong Kong Baptist University, Hong Kong, China

*lap64067611@126.com

These authors contributed equally to this work.



CrossMark
click for updates

 OPEN ACCESS

Citation: Niu X, Lu C, Xiao C, Zhang Z, Jiang M, et al. (2014) The Shared Crosstalk of Multiple Pathways Involved in the Inflammation between Rheumatoid Arthritis and Coronary Artery Disease Based on a Digital Gene Expression Profile. PLoS ONE 9(12): e113659. doi:10.1371/journal.pone.0113659

Editor: Michael Lipinski, Washington Hospital Center, United States of America

Received: August 7, 2014

Accepted: October 26, 2014

Published: December 16, 2014

Copyright: © 2014 Niu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper and its Supporting Information files.

Funding: This study was supported by (1) International cooperation project of Ministry of science and technology (Project ID: 2014DFA31490) to Cheng Xiao, (2) Interdisciplinary Research Matching Scheme (IRMS) of Hong Kong Baptist University (Project ID: RC-IRMS/12-13/02), Research Committee of Hong Kong Baptist University (Project ID: FRG2/12-13/027) to Ge Zhang, and (3) Science and Technology Projects for supervisors of Beijing outstanding doctorate dissertation (Project ID: 20118450201) to Aiping Lu.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Rheumatoid arthritis (RA) and coronary artery disease (CAD) are both complex inflammatory diseases, and an increased prevalence of CAD and a high rate of mortality have been observed in RA patients. But the molecular mechanism of inflammation that is shared between the two disorders is unclear. High-throughput techniques, such as transcriptome analysis, are becoming important tools for genetic biomarker discovery in highly complex biological samples, which is critical for the diagnosis, prognosis, and treatment of disease. In the present study, we reported one type of transcriptome analysis method: digital gene expression profiling of peripheral blood mononuclear cells of 10 RA patients, 10 CAD patients and 10 healthy people. In all, 213 and 152 differently expressed genes (DEGs) were identified in RA patients compared with normal controls (RA vs. normal) and CAD patients compared with normal controls (CAD vs. normal), respectively, with 73 shared DEGs between them. Using this technique in combination with Ingenuity Pathways Analysis software, the effects on inflammation of four shared canonical pathways, three shared activated predicted upstream regulators and three shared molecular interaction networks were identified and explored. These shared molecular mechanisms may provide the genetic basis and potential targets for

optimizing the application of current drugs to more effectively treat these diseases simultaneously and for preventing one when the other is diagnosed.

Introduction

Rheumatoid arthritis (RA) is a systemic, chronic and progressive inflammatory disease that affects mainly the synovial membrane of joints [1]. Systemic inflammatory diseases are associated with increased coronary artery disease (CAD) morbidity and mortality [2]. RA and CAD are both complex inflammatory diseases, and an increased prevalence of CAD and a high rate of mortality were observed in RA patients [3]. In addition, the relative risk of a myocardial infarction is three-fold higher among women with RA compared to controls [4]. Excessive cardiovascular events observed in individuals with RA are not fully explained by the traditional risk factors [5–7], and the underlying mechanism behind the high prevalence of CAD morbidity in RA is not completely understood [8].

There is an increasing interest in identifying ‘nontraditional’ novel risk factors such as genetic polymorphisms in the study of the molecular mechanisms of complex diseases [9, 10]. It has been speculated that genes that play an important role in the development and progression of RA may also play a role in comorbidities and mortality in this disease [11]. In clinical research, specific disease-modifying drugs (e.g., methotrexate and tumor necrosis factor (TNF) inhibitors) effectively control inflammation and frequently used in RA but not in prevention of CAD, also reduce CAD risk but with many limitations such as side effects, limited targets and absence of studies on effects and safety in depth [12]. For example, current clinical evidences suggested that abatacept, a novel CD80/86-CD28 T cell co-stimulation modulator, was effective on reducing inflammatory biomarkers in RA patients [13]. Coincidentally, it had also reported T-cell CD80/86-CD28 co-stimulation was vital for post-interventional accelerated atherosclerosis development, indicating promising clinical potential for prevention of post-interventional remodeling in CAD by abatacept [14]. Patrick H. Dessein *et al.* proved interleukin-6 (IL-6) concentrations independently contribute markedly to endothelial activation in RA more than other cardiovascular risk factors, which enlightened assessment of IL-6 concentrations might enhance cardiovascular risk stratification in RA [15]. Thus the shared novel molecular factors and pathways in RA and CAD may be considered new therapeutic or predicted targets [16, 17]. We therefore hypothesized that lots of commonly shared genes, molecules and pathways involved in chronic inflammation may exist in the form of networks both in RA and CAD and may serve as potential treatment targets.

Recently, several high-throughput techniques have been developed to study the expression of mRNAs, proteins, and metabolites [18], such as the next-generation

sequencing (NGS) platform [19–21]. Currently, one NGS protocol, 3'- tag digital gene expression (DGE) developed by Illumina (Illumina Inc., San Diego, CA, USA), has been widely used in transcriptome studies [22, 23]. Some reports have focused on the molecular mechanisms of pathophysiologic changes during RA or CAD independently by transcriptome or gene expression profiles technology [23–25], but few studies have focused on the associations and shared genes between RA and CAD by high-throughput techniques such as transcriptome analysis.

In the present study, we preformed the DGE method on peripheral blood mononuclear cells (PBMCs) of RA and CAD patients to review the transcriptome changes between them by identifying the DEGs compared with healthy volunteers. We analyzed the associations of pathways and networks related with those genes identified by Ingenuity Pathways Analysis (IPA, <http://www.ingenuity.com>) [26]. This research may provide the genetic basis for the shared prevention and treatment of these two complex related diseases, utilizing the inflammatory mechanisms to optimize the application of currently available drugs to more effectively treat them simultaneously.

Methods

Patients

RA and CAD patients and healthy volunteers were recruited at China-Japan Friendship Hospital in Beijing. The active RA sufferers were selected via their general practitioner from the outpatient rheumatology clinic, and their diagnosis was defined by the 1987 American College of Rheumatology revised criteria and the 2010 American College of Rheumatology/European League Against Rheumatism classification criteria for RA [27, 28]. Disease activity was assessed by the Disease Activity Score in 28 joints (DAS28) [29]. For the purposes of this study, the RA patients were considered to have active RA if they displayed the following symptoms for greater than 6 months: ① morning stiffness for more than 30 min; ② swelling in 3 or more joints, swelling of finger and/or wrist joints, symmetric swelling, rheumatoid nodules, positive rheumatoid factor, or radiographic erosions in the hand and/or wrist; ③ 6 or more tender joints; ④ $DAS28 > 3.2$ [29]; ⑤ erythrocyte sedimentation rate (ESR) ≥ 28 mm/h or C reactive protein (CRP) 20% higher than the normal value [27]. Because RA is two to three times more common in females than males, we chose female patients as the observed subjects in this study [30].

The CAD patients were diagnosed using diagnostic coronary arteriography to detect flow-limiting coronary artery stenosis, which was identified based on electrocardiogram (ECG) results (the inclusion criteria was shown below in detail) and adjudicated by two trained physicians on the basis of clinical examination indexes. Considering that the inflammatory response in CAD patients with acute coronary syndrome (ACS) is more obvious than in patients with stable angina [31], we chose CAD patients with ACS for inclusion in this study. ACS is a broad term that includes unstable angina (UA), non-ST elevation myocardial infarction

(NSTEMI) and ST elevation myocardial infarction (STEMI). Cases of ACS were determined by both formal (World Health Organization 1984 criteria for acute myocardial infarction (AMI), Braunwald criteria for unstable angina (UA)) and medical record criteria. Myocardial infarction (MI) was defined by the presence of symptoms consistent with cardiac ischemia within 24 hours of hospital presentation and troponin I levels above the 99th percentile with a test-specific coefficient of variation <10% [32]. STEMI was defined by the presence of criteria for MI plus one of the following: (a) persistent ST segment elevation of ≥ 1 mm in two contiguous ECG leads or (b) the presence of a new or presumably new left bundle branch block. NSTEMI was defined by the presence of criteria for MI but not for STEMI. The UA diagnosis required the presence of symptoms consistent with cardiac ischemia 24 hours prior to hospital admission, absence of MI criteria and at least one of the following: (a) history of coronary heart disease (CHD); (b) positive coronary disease stratification test (invasive or noninvasive); (c) transient ST segment changes ≥ 0.5 mm in two contiguous leads, new T-wave inversion of ≥ 1 mm and/or pseudonormalization of previously inverted T waves; (d) troponin >0.05 ng/mL; or (e) diagnostic concordance of two independent doctors.

Exclusion criteria included current or prior treatment with prednisone, systolic blood pressure <90 or >150 mm Hg, heart rate <50 or >130 bpm, diabetes mellitus, dyslipidemia, or a history of smoking. Patients with severe diseases of the lungs, liver, kidneys, and patients with mental or blood system disturbances, as well as women who were pregnant, breast-feeding or planning to become pregnant in the next 8 months were excluded from the study. The RA patients included were not diagnosed with any type of CAD, and the CAD patients did not have a concurrent RA diagnosis. The participants in the normal control group were considered healthy if their medical history did not reveal any chronic diseases, endemic infectious diseases or autoimmune diseases, and their physical examination and blood tests failed to show otherwise.

Peripheral blood samples were collected from 10 female RA patients, 10 female CAD patients, and 10 female healthy controls. And the detailed baseline characters of all participants are shown in [Table 1](#). Age- and sex-matched blood samples from healthy individuals were selected as the healthy controls. Peripheral blood samples of 3 mL were collected in the morning before breakfast from each individual participant in this study. The PBMCs were extracted immediately from each blood sample and then stored in Trizol under -80°C until analysis. All protocols involving human subjects were approved by the ethics committee of China-Japan Friendship Hospital (ethics ID: 2014–58), and informed consents were signed by all participants before commencing the study.

PBMC isolation and total RNA extraction

PBMCs were isolated by density gradient centrifugation over Ficoll-Hypaque gradient solution (Histopaque-1077, Sigma-Aldrich, USA), according to the manufacturer's instructions. Heparinized whole blood (3 mL) was diluted to

Table 1. Baseline characteristics of the RA, CAD patients and normal groups.

Characteristic	Normal	RA	CAD
Female Cases	10	10	10
Age, Mean years ± SD	45.8±3.5	54.5±7.1	59.2±4.2
BMI (kg/m²), mean ± SD	22.34±1.83	25.81±2.62	24.43±3.02
ESR (mm/h), mean ± SD	/	39.14±29.53	/
CRP (mg/L), mean ± SD	/	15.48±15.40	/
RF (IU/mL), mean ± SD	/	63.76±71.81	/
DAS28, mean ± SD	/	4.62±0.52	/
Troponin I (Tnl) (ng/mL), (cases in CAD group<0.05 ng/mL, cases in CAD group>0.05 ng/mL)	/	/	5, 5
CK-MB (ng/mL),(cases in CAD group <1.0 ng/mL, cases in CAD group>1.0 ng/mL)	/	/	5, 5

doi:10.1371/journal.pone.0113659.t001

6 mL with phosphate buffered saline (PBS, pH 7.4), layered on top of 3 mL Histopaque and centrifuged for 30 min at 400 × g. PBMCs were aspirated, washed twice, suspended in PBS and counted with a hemocytometer. PBMCs were lysed in Trizol reagent (1 mL/1 × 10⁷ PBMC) (Invitrogen, Karlsruhe, Germany; Carlsbad, CA) and stored at -80°C until further processing.

PBMC samples from RA, CAD patients and healthy normal controls were selected to isolate total RNA. Total RNA was isolated by the Trizol extraction method according to the manufacturer’s protocol. Total RNA was quantified with a NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific Inc., Marietta, OH, USA) followed by quality assessment with the 2100 Bioanalyzer using an RNA 6000 LabChip kit (Agilent Technologies, Palo Alto, CA) according to the manufacturer’s instructions. Acceptable quality values were in the 1.8–2.2 range for A260/A280 ratios, and for each total RNA sample, the RNA Integrity Number was >7.0.

DGE tag profiling and identification of differentially expressed genes

For each sample, 4 µg of total RNA was purified by adsorption of biotin oligo magnetic beads. After the binding of mRNA, cDNA synthesis was performed. Double strand cDNA was introduced into the cDNA fragment digested by *NlaIII* endonuclease and the binding fragments containing sequences of CATG sites and adjacent poly A tails in the 3’ end. After the precipitation of the 3’ cDNA fragment, an Illumina adaptor 1 was added to the 5’ end. Both the adaptor 1 and CATG sites can be recognized by *MmeI*, which cuts at a downstream CATG site and produces fragments of 17 bp tags with adaptor 1. Adaptor 2 was added to the 3’ end of these tags after eliminating the fragment with beads in the 3’ end. Next, these sequences were prepared for Solexa sequencing.

Clean-tags were obtained by filtering the adaptor sequences and removing low-quality sequences (containing ambiguous bases). Only the tags with perfect

matches or one mismatch were further considered and annotated based on the reference genes. The expression level of each gene was estimated by the frequency of clean tags and then normalized to TPM (number of transcripts per million clean tags) [22], which is a standard method and extensively used in DGE analysis [33]. TPM indicates reads per kilobase of transcript per million sequenced reads. The expression level of each gene was measured by the normalized number of matched clean tags.

The number of tags mapped to a given gene was considered to represent the expression level of this gene. Expression levels of a gene from two distinct samples were compared to give an expression difference. We identified the differentially expressed genes (DEGs) in the RA *vs.* the normal control group and the CAD *vs.* the normal control groups. Significance values for differences in expression were determined using a modified exact test, similar to *Fisher's exact* test. The *P*-values were adjusted using a *Benjamini* and Hochberg false discovery rate (FDR) of 1%. We classified the gene as differentially expressed only when the expression difference was more than 1.2-fold with *P*-value <0.01.

Pathway and network analysis by IPA

The Gene Bank accession numbers and matched identified DEGs in RA *vs.* normal and CAD *vs.* normal groups were set up as the identifiers of the two datasets. Each dataset was saved as an Excel file and uploaded into the Ingenuity Pathways Analysis system (IPA, Ingenuity Systems, <http://www.ingenuity.com>), which enabled the discovery, visualization and exploration of molecular interactions to identify the biological mechanisms, pathways and functions most relevant to the experimental datasets, genes, or proteins of interest. The proof-of-knowledge based IPA was performed to characterize the biomarkers confirmed by the pattern recognition analyses to evaluate the biomarkers based on their metabolic or signaling associations in canonical pathways and biological networks [34]. We utilized the IPA analysis system with the 'Core analysis' platform to analyze the identified DEGs of RA *vs.* normal control and CAD *vs.* normal control groups; comparisons between the canonical pathways and biological networks between the two groups were also carried out using the 'Comparison' platform in IPA. In this study, the score was -10 logarithms of *Fisher's exact* test *p*-values in canonical pathway analysis by IPA. Significances for biological functions were then assigned to each network by determining a *p*-value for the enrichment of the genes in the network for such functions compared with the whole Ingenuity Pathway Knowledge Base as a reference set.

Results

Gene expression profiles in RA vs. normal and CAD vs. normal groups

Differentially expressed genes (DEGs) in the RA or CAD groups compared with the normal control group were determined (>1.2 -fold change, false discovery rate (FDR) <0.05 ; **Tables S1, S2 in S1 File**). The number of DEGs is summarized in the Venn diagram (**Fig. 1A**) and the histogram (**Fig. 1B**). In comparison with the normal group, 213 and 152 DEGs were identified in the RA and CAD patients, respectively. It was striking that the majority of DEGs were up-regulated in the diseased states, both in RA and CAD. While 117 and 68 unique genes were up-regulated in RA and CAD, respectively, there were 23 and 11 significantly down-regulated genes in those two groups, respectively. **Figures 1A, B** show 73 shared DEGs between two states: 68 up-regulated and 5 down-regulated DEGs in both RA and CAD patients compared with normal controls. We also found the 73 shared DEGs in the two groups regulated with the same directions, respectively, compared to normal (**Table S3 in S1 File**).

Functions annotation comparison of DEGs in RA and CAD showing inflammation related functions shared in top

IPA was used to identify functions that were significantly over-represented by the DEGs in the comparison of RA and CAD. Next, we classified counting the number of DEGs involved in functions in three aspects: 'diseases and disorder', 'molecular and cellular functions' and 'physiological system development and function', shown in **Fig. 2**. Within the three aspects, there were several shared functions of DEGs in both RA and CAD compared with the normal controls, especially at diseases and disorder functions; most of the shared functions were focused on the inflammation related functions, such as inflammatory diseases, infectious disease, and the inflammatory response. This phenomenon indicated that inflammatory processes play an important role in both RA and CAD disease development. At the same time, among the shared molecular and cellular functions of DEGs in both RA and CAD, the functions of cell death and survival, cellular growth and proliferation, and cellular development were more important. In addition, in the shared physiological system development and functions, there were more DEGs that participated in the hematological system development and function and immune cell trafficking in both RA and CAD. Thus, the most important shared functions in which most DEGs participated were inflammation related functions within the diseases and disorder functions.

Network, function and canonical pathway analysis of shared DEGs in RA and CAD

The shared molecular network was built up from 95 shared DEGs in both RA vs. normal controls and CAD vs. normal controls, as shown in **Fig. 3A**. The function annotations of the shared DEGs molecular network indicated inflammatory




Fig. 1. Gene expression in groups of RA vs. normal, CAD vs. normal and shared DEGs. A: Venn diagram showing the number of uniquely up-regulated (red) or down-regulated (green) genes comparing individuals with RA and CAD to normal persons and shared DEGs; B: Bar diagram indicating number of DEGs in the RA vs. normal and the CAD vs. normal groups and the shared DEGs between them.

doi:10.1371/journal.pone.0113659.g001


response was most important, involving the most shared DEGs: 34 shared DEGs participated in this function, as shown in [Fig. 3C](#).

Based on the shared DEGs in [Table S3 in S1 File](#), we determined the top 4 shared biological pathways associated with inflammation in both RA and CAD using IPA ([Fig. 3B](#)). The calculation was either achieved according to the ratio (the number of genes from the data set that map to the canonical pathway in question divided by the total number of proteins that map to the same canonical pathway) or significance (*Fischer's* exact test was used to calculate a *p*-value determining the probability that the association between the proteins in the dataset and the canonical pathway was explained by chance alone). We chose the commonly shared pathways via score ($-\log(p\text{-value}) > 1.3$ (*p*-value of pathway < 0.05) in DEGs of RA vs. normal controls and CAD vs. normal controls as significant shared pathways in which the DEGs of the two diseases were both involved, as shown in [Table 2](#). Four shared signaling pathways related to inflammation between RA and CAD were found: IL-8 signaling, antigen presentation pathway, OX40 signaling pathway and TREM1 signaling. Additionally, 10 shared DEGs were also involved ([Table 2, Fig. 4](#)): CXCL1, CXCR1, CXCR2, BAX, DEFA1, MMP9, HLA-DRB4, HLA-DQA2, CCL2/MCP-1 and MPO, with same regulated directions. The shared DEGs play important roles in the 4 top shared signaling pathways and in the key nodes leading to inflammation.


The shared associated networks between RA and CAD

To further explore the commonly shared biological roots associated between RA and CAD, we identified molecular networks that were associated with DEGs in RA

Diseases and disorder



Molecular and cellular functions



Physiological system development and function




Fig. 2. The shared functions of DEGs in individuals with RA and CAD compared to normal controls. Radar diagrams show the shared functions of DEGs in RA and CAD compared to normal controls in aspects of 'Diseases and disorder', 'Molecular and cellular functions', and 'Physiological system development and function'. Blue areas represented the number of DEGs in RA vs. the normal number involved in the functions. Red areas represent the number of DEGs in CAD vs. normal controls involved in the functions.

doi:10.1371/journal.pone.0113659.g002

vs. normal controls and CAD vs. normal controls using IPA analysis. As shown in [Figs. 5A, 5B and 5C](#), there were 3 shared associated networks (SANs) that were found between RA and CAD: shared associated network (SAN) -1, SAN-2 and SAN-3. The functions of the associated merged networks between RA and CAD are shown in [Table 3](#). We found that the top functions of the three SANs were focused on immune cell trafficking, inflammatory response, inflammatory diseases, cell-to-cell signaling and interaction, and cell death and survival, most of which were related to biological processes of inflammation. Additionally, there were 11 shared DEGs involved in the three SANs: CTGF, EGR2, LTF, MMP9, NOV, RBP1, ARG, BMX, KCNJ2, HLA-DRB4, and SLPI.




Fig. 3. Network, molecular functions and canonical pathway analysis of shared DEGs in RA vs. normal and CAD vs. normal individuals. A: The molecular network of 95 shared DEGs in RA vs. normal and CAD vs. normal individuals; blue molecules represent the shared DEGs; B: the top 4 shared canonical pathways (with score $-\log(p\text{-value}) > 1.3$, $p\text{-value}$ of pathway < 0.05) related with inflammation which 95 shared DEGs in RA vs. normal and CAD vs. normal involved in: IL-8 signaling; antigen presentation pathway; OX40 signaling pathway; TREM1 signaling; C: the function classification of shared DEGs in both RA vs. normal and CAD vs. normal, the number in the bar diagram represented the DEGs numbers participated in the corresponding pathways.

doi:10.1371/journal.pone.0113659.g003

Table 2. The top 4 shared signaling pathways related to inflammation and shared DEGs in both RA vs. normal and CAD vs. normal subjects.

No.	Shared pathways	Score of pathways ($-\log(p\text{-value})$)	Shared DEGs involved in shared pathways	Fold of shared DEGs ($-\log(p\text{-value})$)	
				RA	CAD
1	IL-8 signaling	6.94	BAX	1.39	1.98
			CXCL1	1.82	1.47
			CXCR1	1.96	1.62
			CXCR2	1.41	1.32
			DEFA1	1.85	2.47
			MMP9	1.90	2.26
			MPO	1.23	1.43
2	Antigen presentation pathway	2.53	HLA-DRB4	1.71	1.99
			HLA-DQA2	1.30	1.51
3	OX40 signaling pathway	2.21	HLA-DRB4	1.71	1.99
			HLA-DQA2	1.30	1.51
4	TREM1 signaling	2.18	CCL2	2.07	1.43
			MPO	1.23	1.43

doi:10.1371/journal.pone.0113659.t002




Fig. 4. The significant commonly shared signaling canonical pathways between RA and CAD involved in inflammation. A: IL-8 signaling; B: antigen presentation pathway; C: OX40 signaling pathway; D: TREM1 signaling; E: picture legends. Purple circle: the same genes involved in the commonly shared pathways in RA and CAD.

doi:10.1371/journal.pone.0113659.g004

To understand the inflammatory related regulation shared by both RA and CAD, we predicted the upstream regulators of the DEGs in RA *vs.* normal controls and CAD *vs.* normal controls by IPA. The activation of three upstream regulators shared in both RA and CAD were predicted by the IPA platform: TNF, IL1 β and p38 MAPK. The networks of upstream regulators and target molecules in RA and CAD are drawn in [Fig. 5D and 5E](#) and [Table 4](#). Additionally, the shared downstream target DEGs were as follows: MMP9, EGR2, CTGF, HLA-DRB4 and RBP1 were found within both RA and CAD ([Table 4](#)). The top shared function of the predicted upstream regulator networks was also focused on the inflammatory response between RA and CAD ([Table 4](#)).




Fig 5. The shared molecular networks and upstream regulators networks of RA and CAD. A: shared associated network (SAN) -1; B: SAN-2; C: SAN-3; D: the network of upstream regulators and target molecules of DEGs in RA; E: the network of upstream regulators and target molecules of DEGs in CAD; F: the legends of predicted networks. In A, B and C, the molecules with orange circles stand for shared molecules between RA and CAD.

doi:10.1371/journal.pone.0113659.g005

Discussion

Both RA and CAD are reported to be chronic inflammatory disease, and usually occur sequentially or exist together, leading to high mortality rates [35, 36]. Sattar *et al.* suggested markers of systemic inflammation (such as interleukin (IL)-6, IL-1, and tumor necrosis factor (TNF)) have been correlated with an increased risk of cardiovascular death in patients with RA [37]. TNF- α and IL-6 have been reported

Table 3. The top functions of shared associated networks (SANs) between RA and CAD and the shared molecules involved (The molecules in italics are shared DEGs in both RA vs. normal and CAD vs. normal groups).

No.	Shared molecules in shared associated network (SAN)	Top functions of shared associated networks
1	<i>CTGF</i> , <i>EGR2</i> , ERK1/2, IgG, <i>LTF</i> , MAP2K1/2, <i>MMP9</i> , NF- κ B complex, <i>NOV</i> , p38 MAPK, <i>RBP1</i>	Immune cell trafficking, inflammatory response, cell-to-cell signaling and interaction.
2	<i>ARG</i> , <i>BMX</i> , IL-6, <i>KCNJ2</i> , TNFSTF14, TREM1	Cell death and survival, inflammatory diseases, inflammatory response.
3	<i>HLA-DRB4</i> , PRTN3, <i>SLPI</i> , SPHK1, TNF	Inflammatory response, immune cell trafficking, cell-to-cell signaling and interaction.

doi:10.1371/journal.pone.0113659.t003

Table 4. The top functions of predicted upstream regulators networks in RA vs. normal and CAD vs. normal and targets molecules involved in (The molecules in italics are shared predicted upstream regulators and targets in both RA vs. normal and CAD vs. normal groups).

Groups	Predicted upstream regulators	Targets of upstream regulators	Top functions of predicted upstream regulators networks
RA vs. normal	IgG, <i>p38 MAPK</i> , TGFB1, <i>TNF</i> , TREM1, EIF2AK2, IFNG, IFNL1, IL13, <i>IL1B</i> , IRF3	ADA, ADM, ARG1, BIRC5, C1QB, C1QC, CCL2/MCP-1, <i>CTGF</i> , CXCL1, EGR1, <i>EGR2</i> , FN1, GSTA1, <i>HLA-DRB4</i> , IFI27, IFI44L, IFIT1, IFIT3, IFITM3, ISG15, ITGB4, KCNJ2, KCNJ15, <i>MMP9</i> , NOV, <i>RBP1</i> , SNAI1, TNFSF11	Inflammatory response, connective tissue disorders, inflammatory disease.
CAD vs. normal	NF-κB (complex), <i>P38 MAPK</i> , <i>IL1B</i> , <i>TNF</i>	APOBEC3B, CAMP, CCL3, CCL3L1/CCL3L3, CHI3L1, <i>CTGF</i> , <i>EGR2</i> , <i>HLA-DRB4</i> , MMP8, <i>MMP9</i> , PTGES, <i>RBP1</i>	Inflammatory response, hypersensitivity response, cardiovascular disease.

doi:10.1371/journal.pone.0113659.t004

to be significantly associated with the severity of subclinical atherosclerosis, independent of Framingham risk score in RA [15, 38], and IL-6 was recently also documented to be very strongly and independently associated with surrogate markers of early atherogenesis in RA [15]. Most of those studies were based on hypothesis-driven candidate genes or inflammatory factors associated or performed in RA and CAD patients. However, how inflammatory modifications were made, the gene profile and the related pathways shared within the two disorders were not fully understood [39].

In the present study, we found 19 of the most important DEGs shared between RA and CAD. Based on the top shared functions, 4 shared canonical pathways, 3 shared associated networks, 3 upstream regulators predicted activation and the key DEGs involved in, we draw the summary path gram involved in inflammation in RA and CAD (Fig. 6), which may be considered as new targets or target networks for the prevention and treatment of patients with CAD and RA simultaneously.

The IL-8 signaling pathway was found to be the most important inflammatory signaling pathway shared both in RA and CAD, in which 7 identified DEGs were found to be commonly shared: BAX, CXCL1, CXCR1, CXCR2, DEFA1, MMP9 and MPO. Significant evidence implicates IL-8 as a major mediator of inflammation and joint destruction in rheumatoid arthritis [40]. The effects of IL-8 and its related ligands are mediated via two receptors, CXCR1 and CXCR2. Several studies have reported that CXCR1/2 mediate neutrophil migration and are involved in the cascade of events leading to inflammation. In addition to modifying fundamental pathological processes, non-competitive allosteric inhibitors of CXCR1/2 may have the additional benefit of providing partial relief for pain and, hence, may be a valid therapeutic target for the treatment of RA [41, 42]. In addition, we can also find reports about the increased levels of IL-8, CXCR1 and CXCR2 in CAD patients, which is supposed to be involved in the pathogenic mechanisms in heart failure [43, 44]. In our study, we acquired similar results to other reports: CXCR1 and CXCR2 can be up-regulated in both RA and CAD patients. Many MMPs are expressed at increased levels in RA tissues and in synoviocyte cultures in response to inflammatory cytokines including MMP9 [45]. Additionally, several mechanism and clinical studies support that MMP9 is a risk factor for CAD [46, 47]. MMP9 was up-regulated in both RA and CAD




Fig. 6. The summary view of shared pathways related with shared DEGs in RA and CAD involved in inflammation. The molecules with red color represent the shared DEGs in both RA vs. normal and CAD vs. normal; molecules with italics in pink shapes represent the shared upstream regulators in both RA vs. normal and CAD vs. normal.

doi:10.1371/journal.pone.0113659.g006

patients compared to the healthy controls in our study, which can lead to an inflammatory response through the regulation of endothelial cell migration during angiogenesis [48, 49]. Bax, a key node during apoptotic mechanisms participating in inflammation, was reported to be active in several studies of the pathogenesis of both RA and CAD [50, 51]. In response to inflammatory events, activated neutrophils release myeloperoxidase (MPO), which has been reported in both RA and CAD patients as well [38, 50]. In addition, we found the induction of the defensin gene: DEFA1/HNP-1 (defensin, alpha 1) was also interpreted as being up-regulated in peripheral blood in both RA and CAD. Bovin LF, *et al.* had confirmed that there was a significantly higher expression of DEFA1/HNP-1 in RA patients than in healthy controls [51], suggesting a role for defensins in the pathogenesis of RA. However, we did not find reports about the role that DEFA1 plays in the pathological process of CAD. Thus, DEFA1 may be considered a novel shared drug target in both RA and CAD.

The other shared signaling pathways, antigen presentation pathway, OX40 signaling pathway and TREM1 signaling, were found to participate in inflammatory process overlap or crossover with the IL-8 signaling, as shown in [Fig. 4](#), [Table 2](#). Chemokine (C-C motif) ligand 2 (CCL2) is referred to as monocyte chemoattractant protein-1 (MCP-1) and is an important risk factor through its role in recruiting circulating monocytes to sites of atherosclerotic lesions [[52](#), [53](#)]. In RA, MCP-1/CCL2 regulate up activation of normal T cell expression and secretion of (RANTES)/CCL5, growth-regulated oncogene α (GRO α)/CXCL1 and IL-8/CXCL8, which are potent chemotactic agents constitutively produced by RA synovial fibroblasts and further up-regulated upon cytokine stimulation [[54](#)]. Both HLA-DRB4 and HLA-DQA2 belong to class II histocompatibility antigens, which are expressed in antigen presenting cells (B lymphocytes, dendritic cells, macrophages) and are used to present antigenic peptides on the cell surface to be recognized by CD4 T-cells during inflammation, reported in the RA process in many papers [[55](#)]. In particular, the HLA-DRB1 gene was reported to be not only a risk factor for RA but also for CAD disease [[56](#)]. DRB4 is linked with allelic variants of DRB1, but is otherwise omitted and reported less frequently in RA or CAD. HLA-DQA2 was also associated with rheumatoid arthritis [[57](#)], but less frequently reported in CAD [[58](#)].

TNF, IL1 β and p38 MAPK were predicted to be activated as up-stream regulators shared in both RA and CAD in this study. Most of the shared DEGs were regulated by the three regulators leading to the inflammatory response in the end, such as CTGF, EGR2, MMP9, CXCL1, BAX, HLA-DRB4, CCL2/MCP-1 and MPO ([Fig. 6](#)). As we know, p38 MAPK, TNF and IL1 β play important roles in transducing inflammation, by which several transcription factors can be directly phosphorylated and activated to bring about pro-inflammatory factors in RA, CAD and other inflammatory diseases [[59](#), [60](#)]. Therefore, these regulators were also considered as key molecular targets for therapeutic intervention in chronic inflammatory diseases such as RA or CAD [[61](#), [62](#)]. For example, the inhibitors of p38 MAPK RWJ67657 and BIRB-796 have been tested in clinical studies in humans [[62](#)]. TNF inhibitors have proven their clinical efficacy in the treatment of RA, but the targeted regulation of TNF response for therapeutic benefits remains suboptimal [[63](#)]. In some reports, treatment of CAD patients with TNF- α antagonist did not show encouraging results and was even associated with worsening of cardiac events in some instances [[64](#)]. However, some others found that using TNF- α inhibitors was associated with a decreased risk for CAD in RA [[65](#)]. Oppositely, it was reported that neither treatment with, nor response to, anti-TNF therapy could be linked to any statistically significant decrease in the risk of acute coronary syndromes [[66](#)]. Perhaps tailored anti-TNF- α therapy in relation to the TNF- α genotype of CAD patients and targeting of the cytokine gene expression via signaling pathway inhibitors may have useful clinical implications [[64](#)]. The same time, preclinical and clinical studies of IL-1 β inhibition have shown efficacy in the treatment of several inflammatory disorders, suggesting that IL-1 β may be a novel therapeutic target for anti-inflammatory therapy such as CAD or RA [[67](#), [68](#)]. For example, Anakinra (Kineret), a

recombinant form of human interleukin-1 (IL-1) receptor antagonist, is approved for the treatment of RA in combination with methotrexate [68]. All of those regulator antagonists are single target drugs, only inhibiting certain pathogenetic targets to intervene in RA or CAD independently, and leading to several side effects at the same time. Thus, exploiting the combination drugs that can act on the shared multiple targets or network targets may be considered a novel therapeutic and preventive strategy for curing RA and CAD at the same time. For example, a drug that targets the p38 MAPK, TNF and IL1 β at the same time or that targets the shared inflammatory molecular network may be the next topic of research.

Limitations

Some potential limitations of the study should be considered. First, the patients participating in this study had either RA or CAD alone; patients with RA combined CAD were not included because of the complex situation of inflammation [69] and the lack of a relatively accepted inclusion criteria with integration of these two diseases. The verification of the shared inflammatory biomarker of the patients with both RA and CAD will be our future work. Additionally, the number of patients included in the study was small, partly because of funding constraints; a larger sample is needed to further verify the results of this study.

Conclusions

In summary, the present study gave a global understanding of the shared inflammatory characteristics in both RA and CAD. IL-8 signaling, antigen presentation, OX40 signaling and TREM1 signaling based on canonical pathways and the 19 shared DEGs involved were found to be the commonly shared inflammatory characteristics of RA and CAD. In particular, p38 MAPK, TNF and IL1 β were predicted to be commonly activated in RA and CAD. These shared pathways, key regulators and down-stream cytokines of inflammation may be considered to be novel targets or network targets for the treatment and prevention of RA and CAD at the same time. Further studies are needed to develop and confirm new drugs that can concurrently target the pathways of both RA and CAD or to predict when CAD will develop in RA patients in depth.

Supporting Information

S1 File. Supporting tables. Table S1. Differentially expressed genes of PBMCs in RA patients compared with normal control. Table S2. Differentially expressed genes of PBMCs in CAD patients compared with normal control. Table S3. The

shared 79 differentially expressed genes in both RA and CAD patients compared with normal control.

[doi:10.1371/journal.pone.0113659.s001](https://doi.org/10.1371/journal.pone.0113659.s001) (DOCX)

Author Contributions

Conceived and designed the experiments: AL GZ ZB. Performed the experiments: XN CL CX YB. Analyzed the data: XN CL ZZ YB. Contributed reagents/materials/analysis tools: CX MJ. Wrote the paper: XN CL MJ DH.

References

1. Lee DM, Weinblatt ME (2001) Rheumatoid arthritis. *Lancet* 358: 903–911.
2. Kriszbacher I, Koppan M, Bodis J (2005) Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 353: 429–430; author reply 429–430.
3. Gonzalez-Gay MA, Gonzalez-Juanatey C, Martin J (2005) Rheumatoid arthritis: a disease associated with accelerated atherogenesis. *Semin Arthritis Rheum* 35: 8–17.
4. Solomon DH, Karlson EW, Rimm EB, Cannuscio CC, Mandl LA, et al. (2003) Cardiovascular morbidity and mortality in women diagnosed with rheumatoid arthritis. *Circulation* 107: 1303–1307.
5. Dessein PH, Joffe BI, Veller MG, Stevens BA, Tobias M, et al. (2005) Traditional and nontraditional cardiovascular risk factors are associated with atherosclerosis in rheumatoid arthritis. *J Rheumatol* 32: 435–442.
6. Stavropoulos-Kalinoglou A, Metsios GS, Panoulas VF, Douglas KM, Nevill AM, et al. (2009) Associations of obesity with modifiable risk factors for the development of cardiovascular disease in patients with rheumatoid arthritis. *Ann Rheum Dis* 68: 242–245.
7. Kitas GD, Gabriel SE (2011) Cardiovascular disease in rheumatoid arthritis: state of the art and future perspectives. *Ann Rheum Dis* 70: 8–14.
8. del Rincon I, Freeman GL, Haas RW, O'Leary DH, Escalante A (2005) Relative contribution of cardiovascular risk factors and rheumatoid arthritis clinical manifestations to atherosclerosis. *Arthritis Rheum* 52: 3413–3423.
9. Sarzi-Puttini P, Atzeni F, Gerli R, Bartoloni E, Doria A, et al. (2010) Cardiac involvement in systemic rheumatic diseases: An update. *Autoimmun Rev* 9: 849–852.
10. Farzaneh-Far A, Roman MJ (2005) Accelerated atherosclerosis in rheumatoid arthritis and systemic lupus erythematosus. *Int J Clin Pract* 59: 823–824.
11. Matthey DL, Thomson W, Ollier WE, Batley M, Davies PG, et al. (2007) Association of DRB1 shared epitope genotypes with early mortality in rheumatoid arthritis: results of eighteen years of followup from the early rheumatoid arthritis study. *Arthritis Rheum* 56: 1408–1416.
12. Cynthia S. Crowson KPL, John M. Davis III, Daniel H. Solomon, Eric L. Matteson, Keith L. Knutson, Mark A. Hlatky, Sherine E. Gabriel (2013) Rheumatoid arthritis and cardiovascular disease. *American Heart Journal* 166: 622–628. e621.
13. Korhonen R, Moilanen E (2009) Abatacept, a novel CD80/86-CD28 T cell co-stimulation modulator, in the treatment of rheumatoid arthritis. *Basic Clin Pharmacol Toxicol* 104: 276–284.
14. Ewing MM, Karper JC, Abdul S, de Jong RC, Peters HA, et al. (2013) T-cell co-stimulation by CD28-CD80/86 and its negative regulator CTLA-4 strongly influence accelerated atherosclerosis development. *Int J Cardiol* 168: 1965–1974.
15. Dessein PH, Solomon A, Woodiwiss AJ, Norton GR, Tsang L, et al. (2013) Marked Independent Relationship between Circulating Interleukin-6 Concentrations and Endothelial Activation in Rheumatoid Arthritis. *Mediators Inflamm*.

16. **Genovese MC, Becker JC, Schiff M, Luggen M, Sherrer Y, et al.** (2005) Abatacept for rheumatoid arthritis refractory to tumor necrosis factor alpha inhibition. *N Engl J Med* 353: 1114–1123.
17. **Smolen JS, Beaulieu A, Rubbert-Roth A, Ramos-Remus C, Rovensky J, et al.** (2008) Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomised trial. *Lancet* 371: 987–997.
18. **Craddock N, Hurles ME, Cardin N, Pearson RD, Plagnol V, et al.** (2010) Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. *Nature* 464: 713–720.
19. **Mardis ER** (2008) The impact of next-generation sequencing technology on genetics. *Trends Genet* 24: 133–141.
20. **Oshlack A, Robinson MD, Young MD** (2010) From RNA-seq reads to differential expression results. *Genome Biol* 11: 220.
21. **Ioannidis JP, Allison DB, Ball CA, Coulibaly I, Cui X, et al.** (2009) Repeatability of published microarray gene expression analyses. *Nat Genet* 41: 149–155.
22. **t Hoen PA, Ariyurek Y, Thygesen HH, Vreugdenhil E, Vossen RH, et al.** (2008) Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms. *Nucleic Acids Res* 36: e141.
23. **Nuhrenberg TG, Langwieser N, Binder H, Kurz T, Stratz C, et al.** (2013) Transcriptome analysis in patients with progressive coronary artery disease: identification of differential gene expression in peripheral blood. *J Cardiovasc Transl Res* 6: 81–93.
24. **Teixeira VH, Olaso R, Martin-Magniette ML, Lasbleiz S, Jacq L, et al.** (2009) Transcriptome analysis describing new immunity and defense genes in peripheral blood mononuclear cells of rheumatoid arthritis patients. *PLoS One* 4: e6803.
25. **Joehanes R, Ying S, Huan T, Johnson AD, Raghavachari N, et al.** (2013) Gene expression signatures of coronary heart disease. *Arterioscler Thromb Vasc Biol* 33: 1418–1426.
26. **Calvano SE, Xiao W, Richards DR, Felciano RM, Baker HV, et al.** (2005) A network-based analysis of systemic inflammation in humans. *Nature* 437: 1032–1037.
27. **Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, et al.** (1988) The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 31: 315–324.
28. **Funovits J, Aletaha D, Bykerk V, Combe B, Dougados M, et al.** (2010) The 2010 American College of Rheumatology/European League Against Rheumatism classification criteria for rheumatoid arthritis: methodological report phase I. *Ann Rheum Dis* 69: 1589–1595.
29. **Prevo ML, van't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, et al.** (1995) Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 38: 44–48.
30. **Silman AJ, Pearson JE** (2002) Epidemiology and genetics of rheumatoid arthritis. *Arthritis Res* 4 Suppl 3: S265–272.
31. **Krintus M, Kozinski M, Kubica J, Sypniewska G** (2014) Critical appraisal of inflammatory markers in cardiovascular risk stratification. *Crit Rev Clin Lab Sci*: 1–17.
32. **Luepker RV, Apple FS, Christenson RH, Crow RS, Fortmann SP, et al.** (2003) Case definitions for acute coronary heart disease in epidemiology and clinical research studies: a statement from the AHA Council on Epidemiology and Prevention; AHA Statistics Committee; World Heart Federation Council on Epidemiology and Prevention; the European Society of Cardiology Working Group on Epidemiology and Prevention; Centers for Disease Control and Prevention; and the National Heart, Lung, and Blood Institute. *Circulation* 108: 2543–2549.
33. **Morrissy AS, Morin RD, Delaney A, Zeng T, McDonald H, et al.** (2009) Next-generation tag sequencing for cancer gene expression profiling. *Genome Res* 19: 1825–1835.
34. **Llorens F, Hummel M, Pastor X, Ferrer A, Pluvinet R, et al.** (2011) Multiple platform assessment of the EGF dependent transcriptome by microarray and deep tag sequencing analysis. *BMC Genomics* 12: 326.
35. **Scott DL, Wolfe F, Huizinga TW** (2010) Rheumatoid arthritis. *Lancet* 376: 1094–1108.

36. **Hansson GK** (2005) Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 352: 1685–1695.
37. **Sattar N, McCarey DW, Capell H, McInnes IB** (2003) Explaining how “high-grade” systemic inflammation accelerates vascular risk in rheumatoid arthritis. *Circulation* 108: 2957–2963.
38. **Rho YH, Chung CP, Oeser A, Solus J, Asanuma Y, et al.** (2009) Inflammatory mediators and premature coronary atherosclerosis in rheumatoid arthritis. *Arthritis Rheum* 61: 1580–1585.
39. **Liao KP, Solomon DH** (2013) Traditional cardiovascular risk factors, inflammation and cardiovascular risk in rheumatoid arthritis. *Rheumatology (Oxford)* 52: 45–52.
40. **Deleuran B, Lemche P, Kristensen M, Chu CQ, Field M, et al.** (1994) Localisation of interleukin 8 in the synovial membrane, cartilage-pannus junction and chondrocytes in rheumatoid arthritis. *Scand J Rheumatol* 23: 2–7.
41. **Min SH, Wang Y, Gonsiorek W, Anilkumar G, Kozlowski J, et al.** (2010) Pharmacological targeting reveals distinct roles for CXCR2/CXCR1 and CCR2 in a mouse model of arthritis. *Biochem Biophys Res Commun* 391: 1080–1086.
42. **Cunha TM, Barsante MM, Guerrero AT, Verri WA, Jr., Ferreira SH, et al.** (2008) Treatment with DF 2162, a non-competitive allosteric inhibitor of CXCR1/2, diminishes neutrophil influx and inflammatory hypernociception in mice. *Br J Pharmacol* 154: 460–470.
43. **Leonard DA, Merhige ME, Williams BA, Greene RS** (2011) Elevated expression of the interleukin-8 receptors CXCR1 and CXCR2 in peripheral blood cells in obstructive coronary artery disease. *Coron Artery Dis* 22: 491–496.
44. **Damas JK, Eiken HG, Oie E, Bjerkeli V, Yndestad A, et al.** (2000) Myocardial expression of CC- and CXC-chemokines and their receptors in human end-stage heart failure. *Cardiovascular Research* 47: 778–787.
45. **Rodriguez-Lopez J, Perez-Pampin E, Gomez-Reino JJ, Gonzalez A** (2006) Regulatory polymorphisms in extracellular matrix protease genes and susceptibility to rheumatoid arthritis: a case-control study. *Arthritis Res Ther* 8: R1.
46. **Li J, Lu H, Tao F, Zhou H, Feng G, et al.** (2013) Meta-analysis of MMP9-562C/T and the risk of coronary heart disease. *Cardiology* 124: 53–59.
47. **Niu W, Qi Y** (2012) Matrix metalloproteinase family gene polymorphisms and risk for coronary artery disease: systematic review and meta-analysis. *Heart* 98: 1483–1491.
48. **Wang X, Yu YY, Lieu S, Yang F, Lang J, et al.** (2013) MMP9 regulates the cellular response to inflammation after skeletal injury. *Bone* 52: 111–119.
49. **Mira E, Lacalle RA, Buesa JM, de Buitrago GG, Jimenez-Baranda S, et al.** (2004) Secreted MMP9 promotes angiogenesis more efficiently than constitutive active MMP9 bound to the tumor cell surface. *J Cell Sci* 117: 1847–1857.
50. **Yonezawa K, Morimoto N, Matsui K, Tenjin T, Yoneda M, et al.** (2012) Significance of the neutrophil myeloperoxidase index in patients with atherosclerotic diseases. *Kobe J Med Sci* 58: E128–137.
51. **Bovin LF, Rieneck K, Workman C, Nielsen H, Sorensen SF, et al.** (2004) Blood cell gene expression profiling in rheumatoid arthritis. Discriminative genes and effect of rheumatoid factor. *Immunol Lett* 93: 217–226.
52. **Charo IF, Taubman MB** (2004) Chemokines in the pathogenesis of vascular disease. *Circ Res* 95: 858–866.
53. **Tang W, Pankow JS, Carr JJ, Tracy RP, Bielinski SJ, et al.** (2007) Association of sICAM-1 and MCP-1 with coronary artery calcification in families enriched for coronary heart disease or hypertension: the NHLBI Family Heart Study. *BMC Cardiovasc Disord* 7: 30.
54. **Unemori EN, Amento EP, Bauer EA, Horuk R** (1993) Melanoma growth-stimulatory activity/GRO decreases collagen expression by human fibroblasts. Regulation by C-X-C but not C-C cytokines. *J Biol Chem* 268: 1338–1342.
55. **Nordang GB, Flam ST, Maehlen MT, Kvien TK, Viken MK, et al.** (2013) HLA-C alleles confer risk for anti-citrullinated peptide antibody-positive rheumatoid arthritis independent of HLA-DRB1 alleles. *Rheumatology (Oxford)* 52: 1973–1982.

56. **Gonzalez-Gay MA, Gonzalez-Juanatey C, Lopez-Diaz MJ, Pineiro A, Garcia-Porrua C, et al.** (2007) HLA-DRB1 and persistent chronic inflammation contribute to cardiovascular events and cardiovascular mortality in patients with rheumatoid arthritis. *Arthritis Rheum* 57: 125–132.
57. (2007) Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447: 661–678.
58. **Granada M, Wilk JB, Tuzova M, Strachan DP, Weidinger S, et al.** (2012) A genome-wide association study of plasma total IgE concentrations in the Framingham Heart Study. *J Allergy Clin Immunol* 129: 840–845 e821.
59. **Schett G, Zwerina J, Firestein G** (2008) The p38 mitogen-activated protein kinase (MAPK) pathway in rheumatoid arthritis. *Ann Rheum Dis* 67: 909–916.
60. **Ferrari R** (1999) The role of TNF in cardiovascular disease. *Pharmacol Res* 40: 97–105.
61. **Cohen P** (2002) Protein kinases—the major drug targets of the twenty-first century? *Nat Rev Drug Discov* 1: 309–315.
62. **Thalhamer T, McGrath MA, Harnett MM** (2008) MAPKs and their relevance to arthritis and inflammation. *Rheumatology* 47: 409–414.
63. **Nguyen CM, Bounthavong M, Mendes MA, Christopher ML, Tran JN, et al.** (2012) Cost utility of tumour necrosis factor-alpha inhibitors for rheumatoid arthritis: an application of Bayesian methods for evidence synthesis in a Markov model. *Pharmacoeconomics* 30: 575–593.
64. **Javed Q, Murtaza I** (2013) Therapeutic potential of tumour necrosis factor-alpha antagonists in patients with chronic heart failure. *Heart Lung Circ* 22: 323–327.
65. **Bili A, Tang X, Pranesh S, Bozaite R, Morris SJ, et al.** (2014) Tumor necrosis factor alpha inhibitor use and decreased risk for incident coronary events in rheumatoid arthritis. *Arthritis Care Res (Hoboken)* 66: 355–363.
66. **Ljung L, Simard JF, Jacobsson L, Rantapaa-Dahlqvist S, Askling J** (2012) Treatment with tumor necrosis factor inhibitors and the risk of acute coronary syndromes in early rheumatoid arthritis. *Arthritis Rheum* 64: 42–52.
67. **Qamar A, Rader DJ** (2012) Effect of interleukin 1beta inhibition in cardiovascular disease. *Curr Opin Lipidol* 23: 548–553.
68. **Kaiser C, Knight A, Nordstrom D, Pettersson T, Fransson J, et al.** (2012) Injection-site reactions upon Kineret (anakinra) administration: experiences and explanations. *Rheumatol Int* 32: 295–299.
69. **Crowson CS, Liao KP, Davis JM 3rd, Solomon DH, Matteson EL, et al.** (2013) Rheumatoid arthritis and cardiovascular disease. *Am Heart J* 166: 622–628 e621.