



Published in final edited form as:

Int J Obes (Lond). 2015 January ; 39(1): 69–74. doi:10.1038/ijo.2014.107.

Prenatal vitamin intake during pregnancy and offspring obesity

Marcelle M. Dougan, MPH¹, Walter C. Willett, MD, DrPH^{1,3,4}, and Karin B. Michels, ScD, PhD^{1,2,3}

¹Department of Epidemiology, Harvard School of Public Health, Boston, MA

²Obstetrics and Gynecology Epidemiology Center, Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

³Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

⁴Department of Nutrition, Harvard School of Public Health, Boston, MA

Abstract

Background/Objectives—In animal studies, exposure to multi-vitamins may be associated with obesity in the offspring; however, data in humans is sparse. We therefore examined the association between prenatal vitamin intake during pregnancy and offspring obesity.

Subjects/Methods—We investigated the association between prenatal vitamin intake and obesity among 29 160 mother-daughter dyads in the Nurses' Health Study II. Mothers of participants provided information on prenatal vitamin use during pregnancy with the nurse daughter. Information on body fatness at ages 5 and 10, body mass index (BMI) at age 18, weight in 1989 and 2009, waist circumference, and height was obtained from the daughter. Polytomous logistic regression was used to predict BMI in early adulthood and adulthood, and body fatness in childhood. Linear regression was used to predict waist circumference in adulthood.

Results—*In utero* exposure to prenatal vitamins was not associated with body fatness, either in childhood or adulthood. Women whose mothers took prenatal vitamins during pregnancy had a covariate-adjusted odds ratio of being obese in adulthood of 0.99 (95% CI 0.92 – 1.05, *P*-value = 0.68) compared to women whose mothers did not take prenatal vitamins. Women whose mothers took prenatal vitamins during pregnancy had a covariate-adjusted odds ratio of having the largest body shape at age 5 of 1.02 (95% CI 0.90 – 1.15, *P*-value = 0.78). In additional analyses, *in utero* exposure to prenatal vitamins was also unrelated to adult abdominal adiposity.

Conclusions—Exposure to prenatal vitamins was not associated with body fatness either in childhood or in adulthood.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding Author: Karin B. Michels, Obstetrics and Gynecology Epidemiology Center, Brigham and Women's Hospital, Harvard Medical School, 221 Longwood Avenue, Boston, MA 02115. kmichels@research.bwh.harvard.edu.

The authors have no conflicts of interest to disclose

Keywords

prenatal vitamins; body mass index; childhood obesity

1. Introduction

Over the past 30 years, the prevalence of obesity has increased dramatically across the United States and elsewhere. Although recent data suggest that the rise in the prevalence of obesity in adults and children appears to be slowing down and may even be leveling off, currently, 35.7% of adults and 18.4% of adolescents are obese (1), compared to 14.5% of adults (2) and 6.1% of adolescents (3) in 1971–1974. The increase in obesity is concerning, as it is well documented that obesity has serious consequences, including premature mortality, and elevated risks for diabetes, cardiovascular disease (4), some cancers (5–7), sub-fertility (8),(9), and depression (10). The economic burden associated with obesity is also quite significant: in one study, investigators reported that obese 45-year olds had a significantly reduced chance of surviving to age 65, and survivors incurred an approximately 40% higher lifetime Medicare costs, compared to normal weight 45-year olds (11).

In recent years, with growing acceptance that the intrauterine environment provides an important basis for future health outcomes (12), considerable progress has been made in examining this environment as a predictor of obesity later on in life. Ravelli *et al* (13) reported that men exposed to maternal starvation *in utero* during the first half of pregnancy had a significantly increased risk of being obese. In another study, exposure to maternal diabetes *in utero* and larger size for gestational age predicted obesity during childhood (14). In other studies, maternal obesity and gestational weight gain predicted childhood (15–16), (17) and later obesity (16).

Prenatal vitamin intake may increase obesity by increasing the amount of adipose tissue cells in the developing fetus. In animal studies, multivitamin supplementation was found to increase the risk of obesity among the offspring of Wistar rats who were fed an obesogenic diet (18). However, to the best of our knowledge, the role of prenatal vitamin supplementation during pregnancy in adult obesity in humans has not been examined.

We therefore examined the association between prenatal vitamin intake during pregnancy and obesity throughout life course among 29 160 participants of the Nurses' Health Study II (NHS II) whose mothers provided information on prenatal vitamin intake during pregnancy.

Materials/Subjects and Methods

Study subjects

Participants of this study are mother-daughter dyads from the Nurses' Health Study II (NHS II) and the Nurses' Mothers' Cohort Study. The NHS II was started in 1989 with the recruitment of 116 478 female registered nurses living in one of 15 US states, who were aged between 25 and 42 years. Participants were mailed a questionnaire about health and lifestyle factors in 1989 (baseline) and every 2 years thereafter. In 2001, participants of the

NHS II who were alive and free of cancer were asked if their mothers could participate in the Nurses' Mothers' Cohort Study, details of which have been previously published (19).

Assessment of Prenatal Vitamin Intake

Participants in the Nurses' Mothers' Cohort Study were asked whether they had taken prenatal vitamins during their pregnancy with the nurse daughter, and if so, whether they took the vitamins regularly. A total of 20 672 reported to have taken prenatal vitamins during pregnancy, of which 1 026 said they did not take the vitamins regularly. Because of the relatively low number of women who reported taking vitamins during pregnancy on an irregular basis, these were excluded from the analyses.

Assessment of Body Fatness

NHS II participants were asked at study enrollment to report their current height, current weight, and their weight at age 18. Current weight was updated on each biennial questionnaire. Body mass index (BMI) was calculated as weight in kg divided by the square of height in m². The validity of self-reported weight at age 18 and self-reported current height among 118 participants of this cohort was assessed in a validation study from records that were obtained from physical examinations conducted at college/nursing school entrance (20). Troy *et al* reported that the correlation between recalled and measured past weight was 0.87, although there was a slight under-reporting in weight at age 18. The correlation between self-reported height and measured height at age 18 was 0.94. Thus, the validity of recalled weight and self-reported height appears high in this cohort.

Childhood body fatness was determined by asking NHS II participants to identify their body size at age 5 and age 10, using a nine-level drawing which was developed by Stunkard (21) (Figure 1). The validity of long-term recall of childhood body fatness was examined during a follow up of the Third Harvard Growth Study, a longitudinal study of physical and mental growth which took place from 1922–1935 (22). More than 65 years later, using the same diagram described above, subjects who were then aged 71–76 years were asked to identify the level that best described their body size during childhood and adolescence. Among females, Pearson crude correlations between recalled body fatness and BMI at approximately the same ages were 0.60 for age 5, and 0.75 for age 10, which slightly attenuated, after adjusting for current BMI. Similar results have been observed in other studies (23–25), demonstrating that this type of recalled measure can provide fairly reliable information on early life body fatness.

In the 2005 questionnaire, participants of NHS II were also asked to provide measurements of their waist circumference. A total of 23 741 participants (81%) provided this information. The validity of measured waist circumference was assessed by Rimm *et al* (26) in a sample of 140 participants from a parallel cohort of older women. Self-reported data were compared with the average of measurements taken by two technicians, and the Pearson correlation between these two measures was 0.89, and the mean difference was 0.05 inches. Thus, although self-reported waist circumference may be underestimated, it is a reliable measure.

Assessment of Covariates

Information on possible risk factors for obesity was obtained from both the Nurses' Mothers' questionnaire and the NHS II questionnaire. Information on maternal age at birth of the daughter, birth order of the nurse, maternal education at time of birth, maternal diet during pregnancy, maternal physical activity level, maternal smoking during pregnancy, maternal domestic status, home ownership at time of birth, father's education at time of birth, father's profession at time of birth, preeclampsia, gestational diabetes, gestational weight gain, mother's BMI, utilization of prenatal care, and breastfeeding, was obtained from the Nurses' Mothers' Questionnaire. Age at menarche, age at first birth, smoking history, parity, alcohol consumption, menopausal status, husband's education, household income, and use of oral contraceptives were obtained from the NHS II questionnaire.

Exclusions

A total of 35,830 mothers of participants in the NHS II completed and returned Nurses' Mothers' Questionnaire. Nurses who were adopted or whose adoption status was unknown (n = 1 895), twin births (n = 587), missing information on age 5 body size (n = 583), age 10 body size (n = 47), BMI at age 18 (n = 269), body mass index in 2009 (n = 1 042), or whose mothers were missing information on prenatal vitamin intake (n = 1 221) or whose mothers took prenatal vitamins but not regularly (n = 1 026) were excluded from the analysis. Missing indicators were used for participants missing information on covariates. The final study population comprised 29 160 mother-daughter dyads.

Statistical Analysis

Follow-up for these analyses began in 1989 at NHS II study baseline, and ended in 2009, the most recent year for which complete information on the participants is available. Body mass index (BMI) in 2009 was categorized as < 18, 23– < 25 (reference), 25 – < 28, 28 – < 30, 30 – < 34, and ≥ 34 kg/m². Missing BMI in 2009 was substituted with BMI reported in 2007 for 1779 participants. BMI at age 18 was categorized as <18, 18–<20, 20–<22 (reference), 22–<23, 23–<25, ≥ 25 kg/m². We used polytomous logistic regression to estimate odds ratios of having being exposed to prenatal vitamins *in utero*, for each category of BMI relative to the reference group. We also modeled BMI in 2009 as a three-level categorical variable: <25, 25–<30, ≥ 30 kg/m². In additional analyses, BMI in 2009 and BMI at age 18 were modeled as continuous variables. Prenatal vitamin intake was coded as a dichotomous variable. Statistical models included potential predictors of obesity during childhood and adulthood: age of nurse at questionnaire return (continuous), maternal age at birth of nurse (< 20, 20 to < 25, 25 to < 30, 30 to < 35, 35 to < 40, ≥ 40 years), birth order of nurse (1, 2, 3, 4), mother's education (< 8 years, 8 years, 1 – 3 years high school, 4 years high school, 1 – 3 years college, ≥ 4 years college), maternal BMI (quintiles), consumption of dark leafy green vegetables during pregnancy (never, less than once a week, 1 – 6 times a week, once a day, twice or more a day), total activity level during pregnancy (highly active, active, mostly inactive/inactive), maternal smoking (non-smoker, quit during first trimester, quit after first trimester, smoked 1 – 15 cigarettes per day, smoked ≥ 15 cigarettes a day), living with nurse's father at time of birth (yes, no), owned a home at time of birth (yes, no), father's education (less than high school, high school, some college, college graduate), father a

professional (yes, no), preeclampsia (yes, no), gestational diabetes (yes, no), gestational weight gain (< 10, 10 – 14, 15 – 19, 20 – 29, 30 – 39, 40, lbs), utilization of prenatal care during pregnancy (yes, no), and ever breastfed (yes, no). Covariates pertaining to the nurse were age at menarche (< 11, 11, 12, 13, 14, 15 years), parity and age at first birth (nulliparous, 1–2 & age at first birth < 25, 1–2 & age at first birth 25–29, 1–2 and age at first birth 30+, 3–4 & age at first birth < 25, 3–4 & age at first birth 25–29, 3–4 & age at first birth 30+, 5 & age at first birth <25, 5 & age at first birth 25–29, and 5 children & age at first birth 30 years), alcohol consumption (non-drinkers, > 0 – 4.9, 5.0 – 9.9, 10.0 – 19.9, 20 g/day), smoking status (never, past, current), menopausal status (premenopausal, postmenopausal), husband's education (< high school, high school, 2 years college, 4 years college, graduate school), income in 2001, which is the most recent year for which income information was available (< \$30 000, 30 000 – 49 000, 50 000 – 74 000, 75 000 – 99 000, 100 000 – 149 000, 150 000), use of oral contraceptives (never, past, current), and physical activity level (< 3, 3 to < 9, 9 to < 18, 18 to < 27, 27 to < 42, 42, metabolic equivalents (METs) per week).

The association between body fatness during childhood (age 5 and age 10) was also analyzed using polytomous logistic regression. Because of sparse sample sizes at larger body types, we combined body size categories from level 5 – 9 into a single category.

Results

Among 29,160 mother-daughter dyads, 67% of the nurse mothers took prenatal vitamins during pregnancy with their nurse daughter whereas 33% did not. In 2009, the mean BMI of the adult nurse daughters was 27.3 kg/m², the median was 25.8 and the 5th and 95th percentiles were 19.9 and 39.5 respectively. At age 18, the mean BMI was 21.1 kg/m², the median was 20.6, and the 5th and 95th percentiles were 17.5 and 26.8 respectively. A total of 6434 participants reported being a Level 1 and 1843 reported being a Level 5 or higher body size at age 5.

Women whose mothers regularly took prenatal vitamins during pregnancy were slightly younger at baseline than women whose mothers did not take prenatal vitamins during pregnancy. Their mothers were also slightly younger at the time of the nurse's birth (Table 1).

The BMI in 2009 and at age 18, of nurses whose mothers took prenatal vitamins were almost the same as those whose mothers did not take prenatal vitamins. The proportion of participants reporting each level of body size was also similar in each group.

Compared to nurse mothers who did not take vitamins, nurse mothers who took prenatal vitamins during pregnancy were slightly younger. They were also more likely to consume green leafy vegetables during pregnancy and more likely to report a higher level of education (Table 1).

In the age-adjusted analysis, *in utero* exposure to prenatal vitamins was significantly associated with BMI in 2009 only for those with a BMI of 34 kg/m² or higher compared to the reference group of 23– < 25 kg/m². The age-adjusted odds ratio (95% confidence

interval, CI) for having a BMI of 34 or greater was 0.90 (95% CI 0.82–0.98) compared to those with a BMI of 23 – < 25 kg/m². After adjusting for other covariates related to the nurse mother this association was no longer significant (Table 2).

We also evaluated a separate model considering additional covariates related to the nurse, including age at menarche, age at first birth, smoking status, parity, and income (Table 2). Exposure to prenatal vitamins was not associated with BMI in 2009 after adjusting for these additional covariates.

Since birth weight may be in the causal pathway between prenatal vitamin intake and body size later in life, we assessed whether it may mediate the association. Associations remained unchanged when birth weight was added to the model.

Prenatal vitamin intake was also unrelated to BMI in 2009 when BMI was modeled as a three-level categorical variable. The age-adjusted OR for being overweight compared to normal weight was 1.00(95% CI 0.94–1.07), and for being obese compared to normal weight was 0.99(95% CI 0.92 –1.05). After adjusting for covariates related to the nurse mother, the OR for being overweight compared to normal weight was 0.99(95% CI 0.93–1.06), and the OR for being obese compared to normal weight was 0.99(95% CI 0.93–1.07).

In utero exposure to prenatal vitamins was significantly associated with BMI at age 18 in the age-adjusted analysis, only for those with a BMI of 25 kg/m² or higher. The age-adjusted OR was 0.85(95% CI 0.77–0.93) comparing those with a BMI of 25 kg/m² to the reference group of 20 – < 22 kg/m². After adjusting for additional covariates, this association was marginally significant (Table 2).

Body fatness at age 5 and age 10, comparing the highest (Type 5+) to the lowest (Type 1), was not significantly associated with exposure to prenatal vitamins before or after adjusting for covariates (Table 3).

In additional analyses, we also assessed the relation between pre-natal vitamin intake and BMI at age 18 and in 2009, modeling BMI as a continuous outcome. Prenatal vitamin intake was not associated with BMI, either at age 18 or in 2009. The average BMI at age 18 of women whose mothers took prenatal vitamins during pregnancy was approximately 0.07 kg/m² less than those whose mothers did not take prenatal vitamins during pregnancy (p-value 0.09), and the average BMI in 2009 of women whose mothers took prenatal vitamins during pregnancy was 0.03 kg/m² less than those whose mothers did not take prenatal vitamins during pregnancy (p-value 0.73).

Finally, we examined the association between prenatal vitamin intake and adult waist circumference in 2005. Prenatal vitamin intake was associated with waist circumference in the age-adjusted analysis: the mean waist circumference for women whose mothers took prenatal vitamins during pregnancy was 0.31 inches less than that of women whose mothers did not take prenatal vitamins during pregnancy (p-value < 0.01); however, after adjusting for covariates related to the mother and the nurse, the mean difference in waist circumference was 0.11 inches, and the association no longer persisted (p-value 0.16).

Discussion

In the Nurses' Health Study II, exposure to prenatal vitamins *in utero* was not associated with body mass index either at age 18 or in adulthood. Further, body size during childhood and waist circumference during adulthood was also not affected by prenatal vitamin intake.

To our knowledge, the literature on the association between *in utero* exposure to multivitamins and offspring obesity is limited. In animal models, multivitamin supplementation in Wistar rats fed an obesogenic diet was found to lead to an acceleration of obesity (18). Lewis *et al* (27) reported that maternal folate intake during pregnancy did not influence childhood body composition, consistent with our findings for body size at age 5 and age 10.

In utero exposure to prenatal vitamins may influence body fatness in the offspring via different mechanisms. For example, maternal malnutrition is believed to trigger excessive appetite in the offspring (28–29). In addition, fetal exposure to inadequate nutrition may increase the capacity of adipocytes to store lipid (30). These results suggest that the association between maternal nutrition and offspring obesity may depend on the nutritional status of the mother. Since our study was conducted in a relatively well-nourished population (United States, 1947–1964), the null association we found may be a consequence of this.

Our study has some limitations. More detailed information on prenatal vitamin intake may have enhanced our ability to detect differences in offspring body fatness. Furthermore, prenatal vitamin exposure was recalled by the mothers from several decades earlier, introducing the potential for recall bias, since mothers were aware of their daughters' body size. However, since an association between vitamin use during pregnancy and offspring overweight was not suspected, we do not expect that the recall would be differential by with respect to the body size in the daughter, and therefore expect any bias to be directed toward the null. Early case-control studies of peri-conceptional multi-vitamin use and the risk of neural tube defects (31–35) relied on recalled data for estimating maternal intake of prenatal vitamins up to 16 years prior to the study. In most of these studies, a significant protective effect of prenatal vitamin use was reported. These findings were later confirmed in subsequent randomized controlled trials (36–37), indicating that recalled vitamin intake can be a reliable way of assessing exposure.

A second limitation is that the timing of prenatal vitamin intake is unknown. It has been suggested in several studies of *in utero* exposures that the timing of the exposure may be a more important determinant of the outcome than the exposure itself. For example, Ravelli *et al* (13) found that the risk of obesity was significant in the offspring of women exposed to starvation in early pregnancy but not those exposed in the third trimester. In a study of folic acid and neural tube defects, Milunsky *et al* (38) reported that the critical exposure period during which folic acid was protective was between weeks 1 and 6 of conception. Folic acid after that period did not confer any protection. Therefore, more detailed information on when prenatal vitamins were actually taken may have been helpful in resolving this question.

Despite these limitations, our study has several strengths. We have a large study population with a high prevalence of prenatal vitamin supplement use. Furthermore, our measures of body fatness in adulthood, specifically weight and waist circumference have good validity (26).

In conclusion, we did not find any statistically significant association between exposure to prenatal vitamins *in utero*, and overweight or obesity either during childhood or during adulthood in this prospective study. Further studies on this subject should assess the timing and dose of prenatal vitamin intake. Although obesity continues to be an important public health problem in the US population, it is unlikely to be influenced by exposure to prenatal vitamins. Changes to current clinical recommendations of routine vitamin supplementation in pregnant women are not warranted based on these results.

Acknowledgements

All authors designed the study; M.M.D. performed statistical analysis and holds primary responsibility for the final content and drafted the manuscript; and all of the authors contributed intellectual content to the manuscript. All authors read and approved the final manuscript.

The Nurses' Mothers' Cohort Study was funded by the Intramural Research Program of the National Cancer Institute research contract N02-RC-17027, and by PO 263 MQ 411027 from the National Cancer Institute.

The Nurses' Health Study II is supported by Public Health Service grant CA50385 from the National Cancer Institute, National Institutes of Health, U.S. Department of Health and Human Services.


REFERENCES

- Ogden, CL.; Carroll, MD.; Kit, BK.; Flegal, KM. Prevalence of obesity in the United States, 2009–2010. Hyattsville, MD: National Center for Health Statistics; 2012. NCHS data brief, no 82. 2012
- National Center for Health Statistics. Prevalence of overweight, obesity and extreme obesity among adults: United States, trends 1976–80 through 2005–2006. 2008
- Ogden CL, Carroll MD. Prevalence of Obesity Among Children and Adolescents: United States, Trends 1963–1965 Through 2007–2008. 2010
- Flegal KM, Graubard BI, Williamson DF, Gail MH. Cause-Specific Excess Deaths Associated With Underweight, Overweight, and Obesity. *JAMA*. 2007 Nov 7; 298(17):2028–2037. 2007. [PubMed: 17986696]
- Ma Y, Yang Y, Wang F, Zhang P, Shi C, Zou Y, et al. Obesity and risk of colorectal cancer: a systematic review of prospective studies. *PLoS One*. 2013; 8(1)
- Fujihara S, Mori H, Kobara H, Nishiyama N, Kobayashi M, Oryu M, et al. Metabolic syndrome, obesity, and gastrointestinal cancer. *Gastroenterol Res Pract*. 2012
- Reeves KW, Carter GC, Rodabough RJ, Lane D, McNeeley SG, Stefanick ML, et al. Obesity in relation to endometrial cancer risk and disease characteristics in the Women's Health Initiative. *Gynecol Oncol*. 2011 May 1; 121(2):376–382. 2011. [PubMed: 21324514]
- Gesink Law DC, Maclehose RF, Longnecker MP. Obesity and time to pregnancy. *Human Reproduction*. 22(2):414–420. [PubMed: 17095518]
- Ramlau-Hansen CH, Thulstrup AM, Nohr EA, Bonde JP, Sørensen TIA, Olsen J. Subfecundity in overweight and obese couples. *Human Reproduction*. 2007 Jun 1; 22(6):1634–1637. 2007. [PubMed: 17344224]
- Chen Y, Jiang Y, Mao Y. Association between obesity and depression in Canadians. *Journal of Women's Health*. 2009; 18(10):1687–1692.
- Cai LP, Lubitz JM, Flegal KMP, Pamuk ERP. The Predicted Effects of Chronic Obesity in Middle Age on Medicare Costs and Mortality. *Medical Care*. 2010 Jun; 48(6):510–517. [PubMed: 20473195]

12. Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet*. 1986 May 10; 1(8489):1077–1081. [PubMed: 2871345]
13. Ravelli GP, Stein ZA, Susser MW. Obesity in young men after famine exposure in utero and early infancy. *N Engl J Med*. 1976; 295(7):349–353. [PubMed: 934222]
14. Lamb MM, Dabelea D, Yin X, Ogden LG, Klingensmith GJ, Rewers M, et al. Early-Life Predictors of Higher Body Mass Index in Healthy Children. *Annals of Nutrition and Metabolism*. 2010; 56(1):16–22. [PubMed: 19940472]
15. Stuebe AM, Forman M, Michels K. Maternal-recalled gestational weight gain, pre-pregnancy body mass index, and obesity in the daughter. *Int J Obes*. 2009 Jul; 33(7):743–752.
16. Schack-Nielsen L, Michaelsen KF, Gamborg M, Mortensen EL, Sorensen TIA. Gestational weight gain in relation to offspring body mass index and obesity from infancy through adulthood. *Int J Obes*. 2009; 34(1):67–74.
17. Heerwagen MJR, Miller MR, Barbour LA, Friedman JE. Maternal obesity and fetal metabolic programming: a fertile epigenetic soil. *Am J Physiol Regul Integr Comp Physiol*. 2010 Sep 1; 299(3):R711–R722. [PubMed: 20631295]
18. Szeto IMY, Das PJ, Aziz A, Anderson GH. Multivitamin supplementation of Wistar rats during pregnancy accelerates the development of obesity in offspring fed an obesogenic diet. *International Journal of Obesity*. 2009; 33(3):364–372. [Article]. [PubMed: 19153583]
19. Michels KB, Willett WC, Graubard BI, Vaidya RL, Cantwell MM, Sansbury LB, et al. A longitudinal study of infant feeding and obesity throughout life course. *Int J Obes*. 2007; 31(7): 1078–1085.
20. Troy LM, Hunter DJ, Manson JE, Colditz GA, Stampfer MJ, Willett WC. The validity of recalled weight among younger women. *Int J Obes Relat Metab Disord*. 1995 Aug; 19(8):570–572. [PubMed: 7489028]
21. Stunkard AJ, Sørensen T, Schulsinger F. Use of the Danish Adoption Register for the study of obesity and thinness. *Res Publ Assoc Res Nerv Ment Dis*. 1983; (60):115–120. [PubMed: 6823524]
22. Must A, Willett WC, Dietz WH. Remote recall of childhood height, weight, and body build by elderly subjects. *Am J Epidemiol*. 1993; 138(1):56–64. [PubMed: 8333427]
23. Koprowski C, Coates RJ, Bernstein L. Ability of women to recall past body size and age at menarche. *Obes Res*. 2001 Aug; 9(8):478–485. [PubMed: 11500528]
24. Muñoz KA, Ballard-Barbash R, Graubard BI, Swanson CA, Schairer C, Kahle LL. Recall of body weight and body size estimation in women enrolled in the breast cancer detection and demonstration project (BCDDP). *Int J Obes Relat Metab Disord* 1996. 1996 Sep; 20(9):854–859.
25. Must A, Phillips SM, Naumova EN, Blum M, Harris S, Dawson-Hughes B, et al. Recall of early menstrual history and menarcheal body size: after 30 years, how well do women remember? *Am J Epidemiol*. 2002 Apr 1; 155(7):672–679. [PubMed: 11914195]
26. Rimm EB, Stampfer MJ, Colditz GA, Chute CG, Litin LB, Willett WC. Validity of self-reported waist and hip circumferences in men and women. *Epidemiology*. 1990 Nov; 1(6):466–473. [PubMed: 2090285]
27. Lewis SJ, Leary S, Davey Smith G, Ness A. Body composition at age 9 years, maternal folate intake during pregnancy and methyltetrahydrofolate reductase (MTHFR) C677T genotype. *British Journal of Nutrition*. 2009; 102(04):493–496. [PubMed: 19660149]
28. Bouret SG. Role of early hormonal and nutritional experiences in shaping feeding behavior and hypothalamic development. *J Nutr*. 2010 Mar; 140(3):653–657. [PubMed: 20107150]
29. Taylor PD, Poston L. Developmental programming of obesity in mammals. *Exp Physiol*. 2007 Mar; 92(2):287–298. [PubMed: 17170060]
30. Muhlhausler B, Smith SR. Early-life origins of metabolic dysfunction: role of the adipocyte. *Trends Endocrinol Metab*. 2009 Mar; 20(2):51–57. [PubMed: 19095460]
31. Bower C, Stanley J. Dietary folate as a risk factor for neural-tube defects: evidence from a case-control study in Western Australia. *Med J Aust*. 1989 Jun 5; 150(11):613–619. [PubMed: 2725375]
32. Mills JL, Rhoads GG, Simpson JL, Cunningham GC, Conley MR, Lassman MR, et al. The absence of a relation between the periconceptional use of vitamins and neural-tube defects. *National*

- Institute of Child Health and Human Development Neural Tube Defects Study Group. N Engl J Med. 1989 Aug 17; 321(7):430–435. [PubMed: 2761577]
33. Mulinare J, Cordero JF, Erickson J, Berry RJ. Periconceptional use of multivitamins and the occurrence of neural tube defects. JAMA. 1988; 260(21):3141–3145. [PubMed: 3184392]
 34. Shaw GM, Schaffer D, Velie EM, Morland K, Harris JA. Periconceptional vitamin use, dietary folate, and the occurrence of neural tube defects. Epidemiology. 1995 May; 6(3):219–226. [PubMed: 7619926]
 35. Werler MM, Shapiro S, Mitchell AA. Periconceptional folic acid exposure and risk of occurrent neural tube defects. JAMA. 1993 Mar 10; 269(10):1257–1261. [PubMed: 8437302]
 36. Czeizel AE, Dudás I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. N Engl J Med. 1992 Dec 24; 327(26):1832–1835. [PubMed: 1307234]
 37. MRC Vitamin Study Research Group. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. Lancet. 1991 Jul 20; 338(8760):131–137. [PubMed: 1677062]
 38. Milunsky A, Jick H, Jick SS, Bruell CL, MacLaughlin DS, Rothman KJ, et al. Multivitamin/folic acid supplementation in early pregnancy reduces the prevalence of neural tube defects. JAMA. 1989 Nov 24; 262(20):2847–2852. [PubMed: 2478730]

44. Which diagram best depicts your outline at each age?



	1	2	3	4	5	6	7	8	9
Currently	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Age 5	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Age 10	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Age 20	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Age 30	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Age 40	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Figure 1.
Assessment of Childhood Body Fatness

Table 1

Age-standardized characteristics of daughters of the Nurses' Mothers' Cohort according to in utero exposure to prenatal vitamins at baseline (1989)

	Did you take prenatal vitamins during pregnancy?	
	No (n=9 514)	Yes, Regularly (n=19 646)
Nurse's age, years ¹	35.5(4.5)	32.9(4.5)
<i>From the Nurses' Mothers' Questionnaire:</i>		
Maternal age at time of nurse's birth, years	27.2(6.2)	25.6(5.3)
Mean maternal pre-pregnancy BMI, kg/m ²	21.5(2.8)	21.2(2.5)
Gestational weight gain, kg	23.5(8.9)	23.7(8.4)
Consumption of dark green leafy vegetables, %	58	63
Maternal smoking during pregnancy, %	24	25
Active/mostly active during pregnancy, %	91	91
Owned home at time of birth, %	51	46
Ever breastfed, %	45	47
Father a professional, %	27	32
Mother's education– some college or graduate, %	32	39
Father's education– some college or graduate, %	38	46
<i>From the Nurses' Health Study II Questionnaire:</i>		
Body size at age 5		
Level 1, n	2093	4322
Level 5+, n	666	1179
Body size at age 10		
Level 1, n	1617	3340
Level 5+, n	1142	2161
BMI ² at age 18, kg/m ²	21.2(3.2)	21.1(3.1)
BMI ² in 1989 (baseline), kg/m ²	23.8(4.8)	23.6(4.6)
BMI ² in 2009, kg/m ²	27.5(6.4)	27.3(6.2)
Waist circumference (2005), inches	34.6(6.0)	34.3(5.9)
Age at menarche, years	12.4(1.4)	12.4(1.4)
Nulliparous, %	36	31
Number of pregnancies over 6 months among parous women	2.0(0.8)	2.0(0.9)
Oral contraceptive use, %	82	84
Physical activity, METS/week	23.8(34.6)	24.5(34.6)
Any alcohol consumption, %	66	68
Alcohol consumption, grams/day ³	5.5(7.1)	5.4(7.1)
Ever a smoker, %	33	32
Husband's education– >= 4 yr college, %	55	56
Annual household income (2001)– \$100,000 or higher, %	34	36

Values are means(SD) or percentages and are standardized to the age distribution of the study population.

¹Value is not age adjusted.

²BMI: Body Mass Index.

³Alcohol consumption among drinkers.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2
Prenatal vitamin use and adult body mass index among NHS II daughters of the Nurses' Mothers' Cohort, 2009

Body Mass Index, kg/m ²	Prenatal Vitamin Intake		Age-adjusted OR ¹	Covariate-adjusted ² OR	Covariate-adjusted ³ OR
	No	Yes			
Current (2009) BMI ⁴					
< 18	2427	5431	0.99(0.92–1.08)	0.97(0.89–1.05)	0.98(0.90–1.06)
23 – < 25 (reference)	1514	3226	1.00	1.00	1.00
25 – < 28	1869	3825	0.98(0.90–1.06)	0.99(0.91–1.08)	0.99(0.90–1.08)
28 – < 30	983	1938	0.95(0.86–1.05)	0.96(0.87–1.07)	0.96(0.86–1.07)
30 – < 34	1294	2546	0.95(0.87–1.05)	0.99(0.90–1.09)	1.00(0.90–1.10)
34	1427	2680	0.90(0.82–0.98)	0.94(0.86–1.04)	0.96(0.87–1.07)
At Age 18					
< 18	779	1800	1.11(1.01–1.23)	1.08(0.97–1.19)	N/A ⁵
18 – < 20	2946	6222	1.01(0.95–1.08)	0.97(0.91–1.04)	
20 – < 22 (reference)	3042	6332	1.00	1.00	
22 – < 23	812	1662	0.97(0.88–1.06)	0.95(0.86–1.05)	
23 – < 25	984	1909	0.92(0.84–1.01)	0.93(0.85–1.03)	
25	951	1721	0.85(0.77–0.93)	0.90(0.82–0.99)	

¹ OR: odds ratio (95% confidence interval).

² Adjusted for maternal age, nurse's age in 2009, birth order of nurse, maternal education at time of birth, maternal diet during pregnancy, breastfeeding, total activity level, maternal smoking during pregnancy, maternal domestic status, home ownership at time of birth, father's education at time of birth, father's profession at time of birth, maternal preeclampsia, maternal gestational diabetes, maternal gestational weight gain, maternal pre-pregnancy BMI, and maternal utilization of prenatal care. (Model 1)

³ Model 1 + nurse's age at menarche, nurse's age at first birth, nurse's smoking status, parity of nurse, nurse's alcohol consumption, nurse's menopausal status, nurse's husband's education, nurse's household income in 2001, nurses physical activity in 2005, and nurse's use of oral contraceptives.

⁴ BMI: body mass index

⁵ Outcomes at Age 18 were not applicable to the model adjusted for nurse's covariates

Table 3

Prenatal vitamin use and body fatness at age 5 and age 10 among NHS II daughters of the Nurses' Mothers' Cohort, 2009

Childhood body size	Prenatal Vitamin Intake		Age-adjusted OR ¹	Covariate-adjusted ² OR
	No	Yes		
Age 5				
Level 1 (reference)	2130	4304	1.00	1.00
Level 2	3065	6592	1.01(0.94–1.08)	0.98(0.91–1.05)
Level 3	2398	5041	1.03(0.95–1.11)	1.00(0.92–1.07)
Level 4	1278	2509	0.98(0.90–1.06)	0.99(0.91–1.09)
Level 5+	643	1200	0.95(0.85–1.06)	1.02(0.90–1.15)
Age 10				
Level 1 (reference)	1636	3323	1.00	1.00
Level 2	2893	6298	1.00(0.93–1.08)	0.98(0.91–1.06)
Level 3	2250	4777	1.00(0.92–1.08)	0.97(0.90–1.06)
Level 4	1608	3106	0.93(0.85–1.01)	0.94(0.86–1.03)
Level 5+	1127	2142	0.95(0.86–1.04)	1.01(0.92–1.12)

¹OR: odds ratio (95% confidence interval).

²Adjusted for maternal age, nurse's age at 2009 questionnaire return, birth order of nurse, maternal education at time of birth, maternal diet during pregnancy, breastfeeding, total activity level, maternal smoking during pregnancy, maternal domestic status, home ownership at time of birth, father's education at time of birth, father's profession at time of birth, maternal preeclampsia, maternal gestational diabetes, maternal gestational weight gain, maternal BMI, and maternal utilization of prenatal care.