



Published in final edited form as:

*Anesthesiology*. 2014 September ; 121(3): 528–537. doi:10.1097/ALN.0000000000000316.

## General Anesthetic Isoflurane Modulates Inositol 1,4,5-Trisphosphate Receptor Calcium Channel Opening

J. Donald Joseph, Ph.D.<sup>\*,^</sup>, Yi Peng, M.D.<sup>\*,^</sup>, Don-On Daniel Mak, Ph.D.<sup>†</sup>, King-Ho Cheung, Ph.D.<sup>†,#</sup>, Horia Vais, Ph.D.<sup>†</sup>, J. Kevin Foskett, Ph.D.<sup>†</sup>, and Huafeng Wei, M.D., Ph.D.<sup>\*</sup>

<sup>\*</sup>Department of Anesthesiology and Critical Care, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania

<sup>†</sup>Department of Physiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania

### Abstract

**Background**—Pharmacological evidence suggests that inhalational general anesthetics induce neurodegeneration *in vitro* and *in vivo* through overactivation of inositol trisphosphate receptor (InsP<sub>3</sub>R) Ca<sup>2+</sup> release channels, but it is not clear whether these effects are due to direct modulation of channel activity by the anesthetics.

**Methods**—Using single-channel patch clamp electrophysiology, we examined the gating of rat recombinant type 3 inositol InsP<sub>3</sub>R (InsP<sub>3</sub>R-3) Ca<sup>2+</sup>-release channels in isolated nuclei (N = 3 to 15) from chicken lymphocytes modulated by isoflurane at clinically relevant concentrations in the absence and presence of physiological levels of the agonist InsP<sub>3</sub>. We also examined the effects of isoflurane on InsP<sub>3</sub>R-mediated Ca<sup>2+</sup> release from the endoplasmic reticulum, and changes in intracellular Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>).

**Results**—Clinically relevant concentrations (~ 1 Minimal Alveolar Concentration) of the commonly used general anesthetic, isoflurane, activated InsP<sub>3</sub>R-3 channels with open probability similar to channels activated by 1 μM InsP<sub>3</sub> (P<sub>o</sub> ≈ 0.2). This isoflurane modulation of InsP<sub>3</sub>R-3 P<sub>o</sub> depended biphasically on [Ca<sup>2+</sup>]<sub>i</sub>. Combination of isoflurane with subsaturating levels of InsP<sub>3</sub> in patch pipettes resulted in at least two-fold augmentations of InsP<sub>3</sub>R-3 channel P<sub>o</sub> compared to InsP<sub>3</sub> alone. These effects were not noted in the presence of saturating [InsP<sub>3</sub>]. Application of isoflurane to DT40 cells resulted in a 30% amplification of InsP<sub>3</sub>R-mediated [Ca<sup>2+</sup>]<sub>i</sub> oscillations, whereas InsP<sub>3</sub>-induced rise in [Ca<sup>2+</sup>]<sub>i</sub> and cleaved caspase-3 activity were enhanced by nearly 2.5 fold.

---

Corresponding author: Huafeng Wei, M.D., Ph.D., Department of Anesthesiology and Critical Care, Perelman School of Medicine, University of Pennsylvania, 305 John Morgan Building, 3620 Hamilton Walk, Philadelphia, PA 19104, Phone: 215-662-3193, Fax: 215-349-5078, weih@uphs.upenn.edu.

<sup>#</sup>Present address: Department of Physiology, University of Hong Kong, Hong Kong, China.

<sup>^</sup>These authors contributed equally to this project.

**Conflict of Interest:** The authors declare no competing interests.

Abstracts of this research work were presented at the Annual Meetings for Society for Neuroscience in Chicago of Illinois On October 21, 2009 (Program # 818.11/C4).

**Conclusion**—These results suggest that the InsP<sub>3</sub>R may be a direct molecular target of isoflurane and a player in the mechanisms of anesthetic-mediated pharmacological or neurotoxic effects.

## Introduction

The inositol trisphosphate receptor (InsP<sub>3</sub>R) is an intracellular Ca<sup>2+</sup> release channel found mostly on the membrane of the endoplasmic reticulum (ER). Activation of InsP<sub>3</sub>R by InsP<sub>3</sub> causes Ca<sup>2+</sup> release from the ER lumen into the cytoplasm, where it acts as a second messenger to regulate many physiological processes such as cell survival and neurogenesis.<sup>1–4</sup> InsP<sub>3</sub>R over-activation also regulates some pathological processes<sup>1;5</sup>, especially apoptosis and neurodegeneration.<sup>6–8</sup> InsP<sub>3</sub>R channel activity can be regulated by a diverse array of interacting proteins,<sup>4</sup> including neurodegenerative-associated proteins such as Huntingtin-associated protein 1 and Alzheimer's disease mutant presenilin 1 and presenilin 2.<sup>9;10</sup>

More than 260 million patients worldwide receive surgeries under general anesthesia each year. The diversity of the molecular mechanisms of general anesthetics is still not clear. Activation of InsP<sub>3</sub>Rs may play important roles in anesthetic-mediated regulation of intracellular Ca<sup>2+</sup> homeostasis and some physiological and pathological processes.<sup>11–14</sup> General anesthetics, especially isoflurane, may precondition cells and provide cytoprotection by moderately activating InsP<sub>3</sub>Rs and Ca<sup>2+</sup> release from the ER.<sup>12</sup> Isoflurane may also cause apoptosis and neurodegeneration by causing abnormal Ca<sup>2+</sup> release from the ER *via* overactivation of InsP<sub>3</sub>R.<sup>3;11;13;15–17</sup> Sensitized InsP<sub>3</sub>R activity in Alzheimer's and Huntington diseases appears to render neurons vulnerable to isoflurane-induced Ca<sup>2+</sup> release from the ER and subsequent apoptosis.<sup>15;17;18</sup> Isoflurane increases levels of β-site amyloid β precursor protein-cleaving enzyme<sup>16</sup> and aggregation of mutated Huntingtin proteins<sup>17</sup> *via* activation of InsP<sub>3</sub>R. General anesthetics may cause neuronal apoptosis by disruption of intracellular Ca<sup>2+</sup> homeostasis.<sup>18–21</sup> Despite these strong associations of anesthetic exposure and InsP<sub>3</sub>R activation in normal and pathological conditions, there is a lack of evidence to support direct activation of the InsP<sub>3</sub>R by isoflurane.

In this study, we report for the first time that clinically relevant concentrations of isoflurane directly modulate the activity of the InsP<sub>3</sub>R channel and sensitize the channel to basal levels of InsP<sub>3</sub>, resulting in InsP<sub>3</sub>R mediated Ca<sup>2+</sup> release from the ER and amplification of InsP<sub>3</sub>-induced [Ca<sup>2+</sup>]<sub>i</sub> signals, and induction cell apoptosis. These results suggest that the InsP<sub>3</sub>R may be one of the molecular mechanisms of anesthetic-mediated pharmacological and toxic effects in neurodegeneration.

## Materials and Methods

### Cell Culture

**DT40 chicken lymphocytes**—DT40 cells lacking the genes for all 3 isoforms of InsP<sub>3</sub>Rs (DT40-KO, RIKEN Cell Bank No. RCB 1467, Ibaraki, Japan) and DT40-KO cells stably over-expressing the rat type 3 InsP<sub>3</sub>R (DT40-R3)<sup>22</sup> were used in this study. While the work was conducted in a chicken lymphocyte cell line, the channel studied is the rat recombinant

InsP<sub>3</sub>R channel. We elected to use InsP<sub>3</sub>R-3 instead of other InsP<sub>3</sub>R subtypes because InsP<sub>3</sub>R-3 has been shown to have relatively higher opening probability activated by InsP<sub>3</sub>R agonists and a useful *in vitro* model to examine the InsP<sub>3</sub>R activation by its agonists.<sup>22</sup> Given the high level of primary sequence homology among mammalian InsP<sub>3</sub>Rs, and the substantial similarities in the major features of the regulation of endogenous and recombinant InsP<sub>3</sub>R channels by cytoplasmic InsP<sub>3</sub> and Ca<sup>2+</sup> (biphasic activation and inhibition by Ca<sup>2+</sup>; and monotonic, saturable activation by InsP<sub>3</sub>),<sup>4</sup> we expect that our findings that isoflurane affects Ca<sup>2+</sup> and InsP<sub>3</sub> regulation of InsP<sub>3</sub>R channels have relevance for similar regulations of neuronal InsP<sub>3</sub>R channels, and thereby have relevance for Ca<sup>2+</sup> homeostasis in neurons. Cells were maintained in suspension culture in Roswell Park Memorial Institute medium 1640 media containing 10% fetal calf serum, 1% chicken serum, penicillin (100 units/ml), streptomycin (100 µg/ml), and glutamine (2 mM) at 37°C in a 95% air and 5% CO<sub>2</sub> humidified incubator.

### Electrophysiology

There is no way to directly observe single-channel activity of the InsP<sub>3</sub>R in the native ER membrane since performing patch-clamp electrophysiology on the ER directly has not been achieved. Since the outer nuclear membrane is continuous with the ER membrane topologically, and since there is no evidence to suggest that the biochemical properties of the outer nuclear membrane are distinct from those of the rest of the ER, nuclear patch clamping of the ER membrane is a tool to record ER-localized ion channel activities, including the InsP<sub>3</sub>R.<sup>4</sup> We have examined the effects of isoflurane on InsP<sub>3</sub>Rs located on the outer nuclear membrane. Preparation of isolated nuclei from DT40 cells was performed as described.<sup>10;22–26</sup> Cells were washed twice with PBS and suspended in a nuclear isolation solution containing 150 mM KCl, 250 mM sucrose, 1.5 mM 2-mercapoethanol, 10 mM Tris-HCl, 0.05 mM PMSF, and protease inhibitor mixture (Complete; Roche Molecular Biochemicals, Indianapolis, IN), adjusted to pH 7.3 with KOH. Isolated nuclei were placed in a recording chamber containing a standard bath solution: 140 mM KCl, 10 mM Hepes, and 0.5 mM BAPTA, ≈200 nM free-Ca<sup>2+</sup>, adjusted to pH 7.3 with KOH. During the nuclear isolation protocol and the introduction of nuclei onto the stage of the patch clamp microscope, any endogenous IP<sub>3</sub> is washed away. Selected intact nuclei were patch-clamped and single channel activities were recorded in the on-nucleus configuration. Recording pipettes with resistances of 8–10 MΩ were used. The pipette solution contained 140 mM KCl, 0.5 mM ATP, 10 mM Hepes (pH 7.3), with varying concentrations of isoflurane, InsP<sub>3</sub> and Ca<sup>2+</sup>. The free Ca<sup>2+</sup> concentration was varied by addition of an appropriate Ca<sup>2+</sup> chelator, as previously described.<sup>27</sup> Single-channel currents were amplified using an Axopatch-200B amplifier (Molecular Devices, Downingtown, PA), filtered at 1 kHz, and digitized at 5 kHz with an ITC-16 interface (Instrutech, Port Washington, NY) and Pulse+ Pulse Fit software (HEKA Elektronik, Farmingdale, NY). All recordings were performed at room temperature with the pipette electrode at –40 mV relative to the reference bath electrode. Single channel-analyses were performed on recordings exhibiting only a single channel using QuB (University of Buffalo, Buffalo, New York) and Igor Pro (WaveMetrics, Lake Oswego, OR) software. Figures were generated using Igor Pro software and Adobe Illustrator (Adobe System Incorporated, San Jose, CA).

### Measurement of cytoplasmic $\text{Ca}^{2+}$ concentration ( $[\text{Ca}^{2+}]_i$ )

DT40-KO or DT40-R3 cells were seeded onto glass coverslips coated with 0.01% poly-L-ornithine for 1 hr before measurements. Cells were loaded with 2.5  $\mu\text{M}$  Fura-2AM (Molecular Probes, Grand Island, NY) for 30 min at room temperature in Hanks' balanced salt solution (HBSS, Sigma, Pittsburgh, PA) containing 1.8 mM  $\text{CaCl}_2$  and 0.8 mM  $\text{MgCl}_2$ , pH 7.4. Coverslips were then placed in a sealed perfusion chamber (Warner Instruments, Hamden, CT) and continuously perfused at room temperature with HBSS. Fura-2AM was alternately excited at 340 and 380 nm, and emitted fluorescence (510 nm) was collected and recorded using a CCD-based imaging system running IPLab v3.7 software (Biovision Technologies, Exton, PA). The data are presented as the ratio of fluorescence intensities recorded at 340 nm and 380 nm excitations (F340/F380) during baseline and isoflurane application. F340/F380 ratios were recorded from > 30 cells in at least three separate experiments. For analyses of  $[\text{Ca}^{2+}]_i$  oscillations, ratio images were collected from > 60 cells over 20 min in at least four separate experiments. The percentage of cells with obvious  $[\text{Ca}^{2+}]_i$  responses (a single transient rise or sustained oscillations), peak  $[\text{Ca}^{2+}]_i$ , area under the  $[\text{Ca}^{2+}]_i$  *versus* time curve, and oscillation frequency were determined and analyzed as described previously.<sup>28</sup> HBSS samples were collected from inflow and outflow tubes of the recording chamber to determine the anesthetic concentration to which the cells were exposed. High performance liquid chromatography measurements showed that isoflurane was consistently maintained at 0.4 mM (not shown). This concentration corresponds to a minimum alveolar concentration of ~1.

### Detection of caspase-3 activity

Proteolytic activation of caspase-3 was measured in lysates from DT40 wild type (DT40-wt), DT40-KO and DT40-R3 cells following exposure to isoflurane using previously described protocols.<sup>13</sup> Caspase-3 activity was measured using Ac-DEVD-AFC hydrolysis kit (Caspase-3, Calbiochem, Billerica, MA). Briefly, DT40-KO or DT40-R3 cells grown in six well plates were exposed to isoflurane (2.4%) for 24 h which resulted in 0.8 mM isoflurane in the culture medium<sup>29</sup> and reliably produced isoflurane toxicity in this anesthesia exposure cell model,<sup>13</sup> harvested *via* trypsinization, and washed with phosphate buffered saline. The cell pellet was gently resuspended in CelLytic™ M lysis buffer with protease inhibitor cocktail (Sigma). Lysate was centrifuged and the resultant supernatant was used for the assay. Ac-DEVD-AFC, the caspase substrate, was added at a final concentration of 50  $\mu\text{M}$  and the samples were incubated for 45 min at 37°C. Ac-DEVD-AFC hydrolysis was monitored by fluorescence emission of the released AFC (excitation, 400 nm; emission, 500 nm) using a multi-wavelength excitation dual wavelength-emission fluorimeter (Delta RAM; photon Technology International, Edison, NJ). Under these excitation and emission conditions, Ac-DEVD-AFC hydrolysis produced a yellow-green fluorescence as suggested by the manufacturer (Calbiochem).

### Analysis and Statistics

Data and statistical analyses were done with IGOR Pro software (Wavemetrics). All data are presented as mean  $\pm$  SEM. Sample size for the electrophysiological and  $\text{Ca}^{2+}$  imaging experiments were used based on previous experiences<sup>27;30</sup>. Experimenters were not blinded

to conditions. To avoid error introduced by subjectivity, all electrophysiological experiments in which nuclear membrane patches were successfully isolated were recorded and analyzed.

Statistical comparisons between data points were done using two-tailed tests. All statistically significant differences between data points stated were established using unpaired *t*-tests (with Bonferroni correction for multiple comparisons within a family of inferences). All lack of statistically significant differences between data points stated were established by ANOVA. Unless a more stringent *P*-value is stated,  $P < 0.05$  was used for rejecting the null hypothesis.

## Results

### Activation of InsP<sub>3</sub>R by isoflurane

Previous studies indicated that InsP<sub>3</sub>Rs are involved in isoflurane-mediated increases in  $[Ca^{2+}]_i$  in PC12 cells, DT40 chicken B lymphocytes, Huntington's striatal neuronal cell lines and primary cortical neurons.<sup>11;13;16;31</sup> To test whether isoflurane might directly modulate the activity of the InsP<sub>3</sub>R, we measured single InsP<sub>3</sub>R channel currents in native ER membranes by nuclear patch clamp electrophysiology.<sup>4;10;23;24;26;28</sup> Homo-tetrameric InsP<sub>3</sub>R-3 channel activities were recorded by patch-clamping outer membranes of nuclei isolated from DT40-KO cells stably expressing the rat type 3 InsP<sub>3</sub>R.<sup>10;22;32</sup> This isoform has ligand regulation and permeation properties similar to other InsP<sub>3</sub>R isoforms, but with robust gating that provides sensitive detection of modulation of channel activity in patch clamp electrophysiology. InsP<sub>3</sub>R-3 channel activity was observed with 2  $\mu$ M  $Ca^{2+}$  and either 1 or 10  $\mu$ M InsP<sub>3</sub> in the pipette solution (fig. 1A–C) in 90% of patches (table 1; 18/20 and 31/34 for 1 and 10  $\mu$ M InsP<sub>3</sub>, respectively). Surprisingly, with 400  $\mu$ M isoflurane in the pipette solution, InsP<sub>3</sub>R channel activity could be elicited in the absence of InsP<sub>3</sub> (fig. 1D), albeit with low open probability ( $P_o$ ) and in < 20% of patches (table 1; 15/85). The channels activated by isoflurane were identified as InsP<sub>3</sub>R by their sensitivity to the InsP<sub>3</sub>R antagonists heparin (100  $\mu$ g/ml) or xestospongin C (1  $\mu$ M) (fig. 1E and F; table 1). The mean open probability, open duration ( $t_o$ ) and closed time ( $t_c$ ) of the InsP<sub>3</sub>R-3 channels activated by 400  $\mu$ M isoflurane were similar to those of channels activated by subsaturating 1  $\mu$ M InsP<sub>3</sub> (fig. 1G–I), suggesting that isoflurane activates InsP<sub>3</sub>R-3 channels with similar kinetics to the endogenous agonist InsP<sub>3</sub> but with lower efficacy. Interestingly, the concentration dependence of the modification of InsP<sub>3</sub>R-3 channel activity by isoflurane was biphasic: at low [isoflurane] (< 400  $\mu$ M), increases in [isoflurane] increased channel  $P_o$ , but channel  $P_o$  decreased as [isoflurane] was increased beyond 400  $\mu$ M. (fig. 2). Thus, InsP<sub>3</sub>R-3 channels are activated within a narrow range of [isoflurane]. This is very different from InsP<sub>3</sub> activation of the channel, in which channel  $P_o$  increases with [InsP<sub>3</sub>] until the saturating [InsP<sub>3</sub>] is reached. Beyond that, further increase in [InsP<sub>3</sub>] does not enhance channel activity any more.<sup>32</sup>

### $[Ca^{2+}]_i$ dependence of isoflurane-induced InsP<sub>3</sub>R channel activity

Gating of InsP<sub>3</sub>Rs is under complex allosteric regulation by InsP<sub>3</sub> and  $[Ca^{2+}]_i$ . In general, InsP<sub>3</sub>R channel activity is biphasically regulated by  $[Ca^{2+}]_i$  with maximum channel  $P_o$  observed over a broad range of  $[Ca^{2+}]_i$  in the presence of saturating (10  $\mu$ M) InsP<sub>3</sub>. In the

presence of subsaturating  $[\text{InsP}_3]$ , sensitivity of the channel to inhibition by high  $[\text{Ca}^{2+}]_i$  is enhanced, resulting in a narrower  $P_o$  versus  $[\text{Ca}^{2+}]_i$  dependence.<sup>4,32</sup> Thus, we investigated the  $[\text{InsP}_3]$  and  $[\text{Ca}^{2+}]_i$  dependencies of isoflurane-activated  $\text{InsP}_3\text{R}$  channel activity. At low (0.4 or 0.9  $\mu\text{M}$ ) or high (6 and 10  $\mu\text{M}$ )  $[\text{Ca}^{2+}]_i$ , 400  $\mu\text{M}$  isoflurane-activated channel  $P_o$  was substantially lower ( $P_o < 0.006$ ) than that observed in 2  $\mu\text{M}$   $\text{Ca}^{2+}_i$ , ( $P_o = 0.2$ ; fig. 3), resulting in a narrow biphasic  $[\text{Ca}^{2+}]_i$  dependence.

### Isoflurane modulates $\text{InsP}_3\text{R}$ channel sensitivity to $\text{InsP}_3$

Although isoflurane activated  $\text{InsP}_3\text{R}$ -3 channel activity in the absence of  $\text{InsP}_3$  in the pipette solution, the sensitivity of isoflurane activation of the channel to the competitive inhibitor heparin suggests that isoflurane may activate the  $\text{InsP}_3\text{R}$ -3 by sensitizing the channel to low  $[\text{InsP}_3]$  generated locally in the patched nuclear membrane. To test this, we recorded  $\text{InsP}_3\text{R}$  channel activity in DT40-R3 cells with 400  $\mu\text{M}$  isoflurane, 2  $\mu\text{M}$   $\text{Ca}^{2+}$  and a range of  $[\text{InsP}_3]$  in the pipette solution. As expected,  $\text{InsP}_3$  enhanced channel  $P_o$  in a dose-dependent manner (fig. 4A–D). At very low (0.1 and 0.5  $\mu\text{M}$ )  $[\text{InsP}_3]$ , 400  $\mu\text{M}$  isoflurane potentiated channel activity to a level greater than either agonist alone (fig. 4E). In contrast, at higher (1  $\mu\text{M}$ )  $[\text{InsP}_3]$ , 400  $\mu\text{M}$  isoflurane no longer potentiated channel activity significantly (fig. 4E). In saturating (10  $\mu\text{M}$ )  $[\text{InsP}_3]$ , 400  $\mu\text{M}$  isoflurane did not enhance channel activity measurably (fig. 4E). These results suggest that isoflurane increases the functional sensitivity of the  $\text{InsP}_3\text{R}$ -3 to  $\text{InsP}_3$  only at low, subsaturating  $[\text{InsP}_3]$  ( $< 0.1$   $\mu\text{M}$ ).

### Isoflurane modulates $\text{InsP}_3\text{R}$ -mediated $[\text{Ca}^{2+}]_i$ signaling and apoptosis

Our single channel recordings indicate that isoflurane at clinically relevant concentrations potentiates the activity of the  $\text{InsP}_3\text{R}$ -3 in low levels of  $[\text{InsP}_3]$  that can exist in unstimulated cells in basal conditions. To determine whether this effect influences intracellular  $\text{Ca}^{2+}$  signaling, we measured  $[\text{Ca}^{2+}]_i$  in individual DT40-KO or DT40-R3 cells kept in complete growth medium. Application of 400  $\mu\text{M}$  isoflurane resulted in a significant transient increases in  $[\text{Ca}^{2+}]_i$  in the DT40-R3 cells that were absent in DT40-KO cells lacking  $\text{InsP}_3\text{R}$  expression (fig. 5A–C).

To further study the potentiating effects of isoflurane on intracellular  $\text{Ca}^{2+}$  signaling in weakly stimulated cells, the cells were perfused with complete growth medium containing a low concentration (50 ng/ml) of anti-IgM antibody to weakly stimulate the B-cell receptor to generate low, but higher than basal, levels of  $\text{InsP}_3$ .<sup>33</sup> 82  $\pm$  4% of DT40-R3 cells responded to anti-IgM antibody with either a single transient elevation  $[\text{Ca}^{2+}]_i$  or with  $[\text{Ca}^{2+}]_i$  oscillations (fig. 5D and E). Whereas the addition of 400  $\mu\text{M}$  isoflurane did not change the percentage of cells responding to anti-IgM antibody (fig. 5D and E), it enhanced the percentage of cells that responded with a single prolonged  $[\text{Ca}^{2+}]_i$  peak (fig. 5E), consistent with a stronger  $\text{InsP}_3\text{R}$  response. Because of the variable responses of cells (oscillations vs. single-peak responses), we quantified the  $[\text{Ca}^{2+}]_i$  responses in all cells by the areas under the  $[\text{Ca}^{2+}]_i$ -versus-time curves (AUC). The presence of isoflurane significantly increased the AUC for cells responding with a single-peak transient (fig. 5F;  $P < 0.001$ ) or with  $[\text{Ca}^{2+}]_i$  oscillations (fig. 5G;  $P < 0.01$ ). These results suggest that isoflurane, at a clinically used concentration, potentiates low level  $\text{InsP}_3$ -mediated  $[\text{Ca}^{2+}]_i$  signaling.

Prolonged exposure to isoflurane is associated with widespread cell death in diverse *in vivo* and *in vitro* systems.<sup>13;16</sup> Because activation of InsP<sub>3</sub>R-mediated Ca<sup>2+</sup> signaling has been linked to apoptosis,<sup>34;35</sup> and considering the effects of isoflurane on InsP<sub>3</sub>R gating observed here, we assessed the role of this effect in isoflurane-mediated apoptosis. Caspases are a family of cysteine proteases that play crucial roles in apoptosis. Activation of caspase-3 is a central event in the progression of programmed cell death. We therefore monitored caspase-3 cleavage as a measure of apoptosis. In agreement with our previous observations of cells expressing endogenous InsP<sub>3</sub>R,<sup>13;16</sup> isoflurane triggered apoptosis in the DT40-R3 cells, but not in the InsP<sub>3</sub>R deficient cells (fig. 5H). These observations support a role for exaggerated activation of InsP<sub>3</sub>R in isoflurane-induced apoptosis, consistent with previous observations.<sup>11;13;16</sup>

## Discussion

We have, for the first time, demonstrated that isoflurane at clinically relevant concentrations modulates the activity of the InsP<sub>3</sub>R Ca<sup>2+</sup> release channel at the single channel level. The modulating effects of isoflurane on the InsP<sub>3</sub>R regulate Ca<sup>2+</sup> release from the ER that results in exaggerated [Ca<sup>2+</sup>]<sub>i</sub> signaling. We have also demonstrated, for the first time at the molecular level, that isoflurane causes Ca<sup>2+</sup> release from the ER *via* this activation of InsP<sub>3</sub>R, and can therefore affect intracellular Ca<sup>2+</sup> homeostasis, regulation of cytosolic Ca<sup>2+</sup> oscillations and cell survival. These results provide novel insights into possible molecular mechanisms of anesthesia-mediated effects on neurodegeneration and cognitive function.

### Inhalational anesthetic modulate InsP<sub>3</sub>R activation

It was previously demonstrated that both isoflurane and halothane may elevate [Ca<sup>2+</sup>]<sub>i</sub> primarily by inducing Ca<sup>2+</sup> release from the intracellular Ca<sup>2+</sup> stores in neurons.<sup>36</sup> Our results provide the first evidence that an inhalational anesthetic can modulate activation of InsP<sub>3</sub>R channels. Interestingly, isoflurane activates InsP<sub>3</sub>R channels with a biphasic dose response with optimal concentrations around 0.4 – 1mM, close to clinically used concentrations for general anesthesia. This activation of InsP<sub>3</sub>R channels by isoflurane showed a strong Ca<sup>2+</sup> concentration dependence, with the optimum [Ca<sup>2+</sup>]<sub>i</sub> of 2 μM, qualitatively similar to the [Ca<sup>2+</sup>] dependence of InsP<sub>3</sub>-activated channel activity. The InsP<sub>3</sub> competitive antagonist, heparin, blocked the ability of isoflurane to activate InsP<sub>3</sub>Rs. Whether isoflurane requires InsP<sub>3</sub> for it to activate the channel, or isoflurane activation is direct but also sensitive to heparin inhibition is unclear. The former may be likely since isoflurane (0.4 mM) potentiated activation of the InsP<sub>3</sub>R by low concentrations of InsP<sub>3</sub>, but failed to further enhance InsP<sub>3</sub>R-3 activity at saturating [InsP<sub>3</sub>]. Nevertheless, further studies are needed to investigate possible biochemical interactions between isoflurane and the InsP<sub>3</sub>R. Together, these results establish the InsP<sub>3</sub>R as a novel target of isoflurane and perhaps other inhalational or intravenous general anesthetics.

### Isoflurane enhances Ca<sup>2+</sup> signals and cell apoptosis *via* activation of InsP<sub>3</sub>R

The biphasic effects of [Ca<sup>2+</sup>]<sub>i</sub> on InsP<sub>3</sub>R activation play important roles on intracellular Ca<sup>2+</sup> oscillations, waves, and spreading of global Ca<sup>2+</sup> signals.<sup>37</sup> Our finding that isoflurane activates InsP<sub>3</sub>R-3 channels raised the possibility that it could enhance [Ca<sup>2+</sup>]<sub>i</sub> signals.

Application of 400  $\mu\text{M}$  isoflurane resulted in a transient increase in  $[\text{Ca}^{2+}]_i$  in DT-40 cells expressing  $\text{InsP}_3\text{R-3}$  but not in DT-40 cells lacking the channels. In addition,  $[\text{Ca}^{2+}]_i$  signals generated by IgM stimulation of DT40-R3 cells, both single-peaks and oscillations, were more prominent in the presence of isoflurane. (fig. 5F and G).

We previously showed in *in vitro* and *in vivo* models that exposure to isoflurane for prolonged durations significantly induced apoptosis that required activation of  $\text{InsP}_3\text{Rs}$ .<sup>11;13</sup> Our results here suggest that this is mediated at least in part by isoflurane activation of  $\text{InsP}_3\text{R}$  channel gating. The effects of isoflurane on changes in  $[\text{Ca}^{2+}]_i$  are in good agreement with the conclusions reached from the single channel recordings of the effects of isoflurane on  $\text{InsP}_3\text{R}$  activity observed in the current study. It has long been known that anesthetics including halothane activate the other major ER  $\text{Ca}^{2+}$  release channels, the RyR channel complex,<sup>38</sup> and this is thought to be the basis for malignant hyperthermia.<sup>39</sup> Like  $\text{InsP}_3\text{Rs}$ , RyRs are expressed throughout the nervous system and play important roles in both normal cell functions<sup>40</sup> and in various neurodegenerative diseases.<sup>2;41-43</sup> Our previous study indicated a role of RyR activation in isoflurane-induced apoptosis in neuronal tissue cultures.<sup>29</sup> Both  $\text{InsP}_3\text{R}$  and RyR contribute to regulation of intracellular  $\text{Ca}^{2+}$  homeostasis and may interact with each other through  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release in a common pathway in normal neuronal function and neurodegeneration. Excessive  $\text{Ca}^{2+}$  release from the ER *via* these release channels could cause  $\text{Ca}^{2+}$  overload in mitochondria and depletion of ER  $\text{Ca}^{2+}$ , both of which can contribute to apoptosis.<sup>6;34;44</sup> In addition, mitochondrial  $\text{Ca}^{2+}$  overload causes cytochrome C release, activating caspase-3, which can cleave the  $\text{InsP}_3\text{R}$ , resulting in a constituent  $\text{Ca}^{2+}$  leak from the ER.<sup>45</sup> Thus, excessive or prolonged activation of  $\text{InsP}_3\text{R}$  by isoflurane may set in motion a cascade of events resulting in apoptosis in different tissue culture cells including neurons,<sup>3;11;13;15-17</sup> and in developing brains.<sup>16</sup> Our data suggest that the  $\text{InsP}_3\text{R}$  may represent an ideal target for prevention of the harmful side effects of inhalational anesthetics. Inhibition of excessive activation of the  $\text{InsP}_3\text{R}$  may ameliorate or prevent anesthesia-mediated neurodegeneration as demonstrated in animal models.<sup>16</sup> This is especially relevant in pediatric<sup>46;47</sup> and aged patients,<sup>48</sup> who appear to be the most vulnerable to the harmful side effects of anesthetics. Unfortunately, there is no good pharmacology for the  $\text{InsP}_3\text{R}$ . Although heparin is an  $\text{InsP}_3\text{R}$  antagonist, its poor penetration across cell membranes limits its use for protection against anesthetic neurotoxicity in animals or patients. It should be noted that anesthetics have also been shown to be protective against various stresses, also by activation of  $\text{InsP}_3\text{Rs}$  in different tissue culture models.<sup>3;12;14;49;50</sup> As demonstrated here (fig. 6), mild or moderate activation of  $\text{InsP}_3\text{R}$  and moderate  $\text{Ca}^{2+}$  release from the ER by isoflurane provides cytoprotection, possibly *via* physiological  $\text{Ca}^{2+}$  uptake into mitochondria and stimulation of ATP production,<sup>25</sup> or by activation of AKT and MAPK/ERK cytoprotective pathways.<sup>12;14;50</sup> Nevertheless, it is prudent to minimize the use of general anesthetics as much as possible, so that their beneficial effects can be utilized and the harmful effects be minimized.

In summary, our results indicate that the commonly used inhalational anesthetic isoflurane modulates gating of the  $\text{InsP}_3\text{R}$   $\text{Ca}^{2+}$  release channel, enhancing ER  $\text{Ca}^{2+}$  release and contributing to isoflurane-mediated apoptosis. These results suggest that  $\text{InsP}_3\text{Rs}$  are molecular targets of general anesthetics and that these receptors may provide the basis for



some pharmacologic effects of general anesthetics and therapeutic interventions for anesthesia-mediated cell death by apoptosis.

## Acknowledgments

**Source of Financial Support:** This work was supported by: National Institutes of Health, Baltimore, Maryland, awards GM084979 and GM084979-02S1 to H.W., GM056328 and MH059937 to J.K.F., and K08-GM073224 to H.W.; March of Dimes Birth Defects Foundation Research Grant 12-FY08-167 to H.W., White Plains, New York; and the Research Fund at the Department of Anesthesiology at the University of Pennsylvania, Philadelphia, Pennsylvania.


The authors thank Qingcheng Meng, Ph.D., Department of Anesthesiology, University of Pennsylvania, Philadelphia, Pennsylvania for assistance in measuring isoflurane concentrations used in the electrophysiology and  $\text{Ca}^{2+}$  measurement experiments. We appreciate the valuable discussions and support from Roderic Eckenhoff, M.D., Maryellen Eckenhoff, Ph.D., and Lee Fleisher, M.D., Department of Anesthesiology and Critical Care, University of Pennsylvania, Philadelphia, Pennsylvania.

## References


1. Stutzmann GE. Calcium dysregulation, IP3 signaling, and Alzheimer's disease. *Neuroscientist*. 2005; 11:110–5. [PubMed: 15746379]
2. Banerjee S, Hasan G. The InsP3 receptor: Its role in neuronal physiology and neurodegeneration. *Bioessays*. 2005; 27:1035–47. [PubMed: 16163728]
3. Zhao X, Yang Z, Liang G, Wu Z, Peng Y, Joseph DJ, Inan S, Wei H. Dual effects of isoflurane on proliferation, differentiation, and survival in human neuroprogenitor cells. *Anesthesiology*. 2013; 118:537–49. [PubMed: 23314167]
4. Foskett JK, White C, Cheung KH, Mak DO. Inositol trisphosphate receptor  $\text{Ca}^{2+}$  release channels. *Physiol Rev*. 2007; 87:593–658. [PubMed: 17429043]
5. Criollo A, Vicencio JM, Tasdemir E, Maiuri MC, Lavandro S, Kroemer G. The inositol trisphosphate receptor in the control of autophagy. *Autophagy*. 2007; 3:350–3. [PubMed: 17404493]
6. Hanson CJ, Bootman MD, Roderick HL. Cell signalling: IP3 receptors channel calcium into cell death. *Curr Biol*. 2004; 14:R933–5. [PubMed: 15530388]
7. Joseph SK, Hajnoczky G. IP3 receptors in cell survival and apoptosis:  $\text{Ca}^{2+}$  release and beyond. *Apoptosis*. 2007; 12:951–68. [PubMed: 17294082]
8. Decuyper JP, Monaco G, Bultynck G, Missiaen L, De SH, Parys JB. The IP(3) receptor-mitochondria connection in apoptosis and autophagy. *Biochimica et Biophysica Acta*. 2011; 1813:1003–13. [PubMed: 21146562]
9. Tang TS, Tu H, Chan EY, Maximov A, Wang Z, Wellington CL, Hayden MR, Bezprozvanny I. Huntingtin and huntingtin-associated protein 1 influence neuronal calcium signaling mediated by inositol-(1,4,5) triphosphate receptor type 1. *Neuron*. 2003; 39:227–39. [PubMed: 12873381]
10. Cheung KH, Shineman D, Muller M, Cardenas C, Mei LJ, Yang J, Tomita T, Iwatsubo T, Lee VMY, Foskett JK. Mechanism of  $\text{Ca}^{2+}$  disruption in Alzheimer's disease by presenilin regulation of InsP(3) receptor channel gating. *Neuron*. 2008; 58:871–83. [PubMed: 18579078]
11. Yang H, Liang G, Hawkins BJ, Madesh M, Pierwola A, Wei HF. Inhalational anesthetics induce cell damage by disruption of intracellular calcium homeostasis with different potencies. *Anesthesiology*. 2008; 109:243–50. [PubMed: 18648233]
12. Bickler PE, Zhan X, Fahlman CS. Isoflurane preconditions hippocampal neurons against oxygen-glucose deprivation: Role of intracellular  $\text{Ca}^{2+}$  and mitogen-activated protein kinase signaling. *Anesthesiology*. 2005; 103:532–9. [PubMed: 16129978]
13. Wei HF, Liang G, Yang H, Wang QJ, Hawkins B, Madesh M, Wang SP, Eckenhoff RG. The common inhalational anesthetic isoflurane induces apoptosis *via* activation of inositol 1,4,5-trisphosphate receptors. *Anesthesiology*. 2008; 108:251–60. [PubMed: 18212570]

14. Bickler PE, Fahlman CS. The inhaled anesthetic, isoflurane, enhances Ca<sup>2+</sup>-dependent survival signaling in cortical neurons and modulates MAP kinases, apoptosis proteins and transcription factors during hypoxia. *Anesth Analg*. 2006; 103:419–29. [PubMed: 16861427]
15. Liang G, Wang QJ, Li Y, Kang B, Eckenhoff MF, Eckenhoff RG, Wei HF. A presenilin-1 mutation renders neurons vulnerable to isoflurane toxicity. *Anesth Analg*. 2008; 106:492–500. [PubMed: 18227305]
16. Zhao Y, Liang G, Chen Q, Joseph DJ, Meng Q, Eckenhoff RG, Eckenhoff MF, Wei H. Anesthetic-induced neurodegeneration mediated *via* inositol 1,4,5-trisphosphate receptors. *J Pharmacol Exp Ther*. 2010; 333:14–22. [PubMed: 20086058]
17. Wang Q, Liang G, Yang H, Wang S, Eckenhoff MF, Wei H. The common inhaled anesthetic isoflurane increases aggregation of huntingtin and alters calcium homeostasis in a cell model of Huntington's disease. *Toxicol Appl Pharmacol*. 2011; 250:291–8. [PubMed: 21059370]
18. Wei H. The role of calcium dysregulation in anesthetic-mediated neurotoxicity. *Anesth Analg*. 2011; 113:972–4. [PubMed: 22021793]
19. Zhang GH, Dong YL, Zhang B, Ichinose F, Wu X, Culley DJ, Crosby G, Tanzi RE, Xie ZC. Isoflurane-induced caspase-3 activation is dependent on cytosolic calcium and can be attenuated by memantine. *J Neurosci*. 2008; 28:4551–60. [PubMed: 18434534]
20. Zhao YL, Xiang Q, Shi QY, Li SY, Tan L, Wang JT, Jin XG, Luo AL. GABAergic excitotoxicity injury of the immature hippocampal pyramidal neurons' exposure to isoflurane. *Anesth Analg*. 2011; 113:1152–60. [PubMed: 21918167]
21. Sinner B, Friedrich O, Zink W, Zausig Y, Graf BM. The toxic effects of s(+)-ketamine on differentiating neurons *in vitro* as a consequence of suppressed neuronal Ca<sup>2+</sup> oscillations. *Anesth Analg*. 2011; 113:1161–9. [PubMed: 21788311]
22. Li C, Wang XL, Vais H, Thompson CB, Foskett JK, White C. Apoptosis regulation by Bcl-x(L) modulation of mammalian inositol 1,4,5-trisphosphate receptor channel isoform gating. *Proc Natl Acad Sci U S A*. 2007; 104:12565–70. [PubMed: 17636122]
23. Mak DO, Vais H, Cheung KH, Foskett JK. Patch-clamp electrophysiology of intracellular Ca<sup>2+</sup> channels. *Cold Spring Harb Protoc*. 2013; 2013:787–97. [PubMed: 24003191]
24. Mak DO, Vais H, Cheung KH, Foskett JK. Isolating nuclei from cultured cells for patch-clamp electrophysiology of intracellular Ca(2+) channels. *Cold Spring Harb Protoc*. 2013; 2013:880–4. [PubMed: 24003193]
25. Cardenas C, Miller RA, Smith I, Bui T, Molgo J, Muller M, Vais H, Cheung KH, Yang J, Parker I, Thompson CB, Birnbaum MJ, Hallows KR, Foskett JK. Essential regulation of cell bioenergetics by constitutive InsP3 receptor Ca<sup>2+</sup> transfer to mitochondria. *Cell*. 2010; 142:270–83. [PubMed: 20655468]
26. Mak DO, Vais H, Cheung KH, Foskett JK. Nuclear patch-clamp electrophysiology of Ca<sup>2+</sup> channels. *Cold Spring Harb Protoc*. 2013; 2013:885–91. [PubMed: 24003194]
27. Mak DO, McBride S, Foskett JK. Inositol 1,4,5-trisphosphate [correction of tris-phosphate] activation of inositol trisphosphate [correction of tris-phosphate] receptor Ca<sup>2+</sup> channel by ligand tuning of Ca<sup>2+</sup> inhibition. *Proc Natl Acad Sci U S A*. 1998; 95:15821–5. [PubMed: 9861054]
28. Cheung KH, Mei L, Mak DO, Hayashi I, Iwatsubo T, Kang DE, Foskett JK. Gain-of-function enhancement of IP3 receptor modal gating by familial Alzheimer's disease-linked presenilin mutants in human cells and mouse neurons. *Sci Signal*. 2010; 3:ra22. [PubMed: 20332427]
29. Wei H, Kang B, Wei W, Liang G, Meng QC, Li Y, Eckenhoff RG. Isoflurane and sevoflurane affect cell survival and BCL-2/BAX ratio differently. *Brain Res*. 2005; 1037:139–47. [PubMed: 1577762]
30. Vais H, Siebert AP, Ma Z, Fernandez-Mongil M, Foskett JK, Mak DO. Redox-regulated heterogeneous thresholds for ligand recruitment among InsP3R Ca<sup>2+</sup>-release channels. *Biophys J*. 2010; 99:407–16. [PubMed: 20643058]
31. Wang Q, Liang G, Yang H, Wang S, Eckenhoff MF, Wei H. The common inhaled anesthetic isoflurane increases aggregation of huntingtin and alters calcium homeostasis in a cell model of Huntington's disease. *Toxicol Appl Pharmacol*. 2010; 250:291–8. [PubMed: 21059370]

32. Vais H, Foskett JK, Ullah G, Pearson JE, Daniel Mak DO. Permeant calcium ion feed-through regulation of single inositol 1,4,5-trisphosphate receptor channel gating. *J Gen Physiol.* 2012; 140:697–716. [PubMed: 23148262]
33. Huang R, Cioffi J, Kimberly R, Edberg J, Mayer L. B cell differentiation factor-induced human B cell maturation: Stimulation of intracellular calcium release. *Cell Immunol.* 1995; 164:227–33. [PubMed: 7656331]
34. Inan S, Wei H. The cytoprotective effects of dantrolene: A ryanodine receptor antagonist. *Anesth Analg.* 2010; 111:1400–10. [PubMed: 20861418]
35. Bezprozvanny I. Calcium signaling and neurodegenerative diseases. *Trends Mol Med.* 2009; 15:89–100. [PubMed: 19230774]
36. Kindler CH, Eilers H, Donohoe P, Ozer S, Bickler PE. Volatile anesthetics increase intracellular calcium in cerebrocortical and hippocampal neurons. *Anesthesiology.* 1999; 90:1137–45. [PubMed: 10201687]
37. Berridge MJ. Inositol trisphosphate and calcium signalling. *Nature.* 1993; 361:315–25. [PubMed: 8381210]
38. Akata T, Nakashima M, Izumi K. Comparison of volatile anesthetic actions on intracellular calcium stores of vascular smooth muscle: Investigation in isolated systemic resistance arteries. *Anesthesiology.* 2001; 94:840–50. [PubMed: 11388536]
39. Denborough M. Malignant hyperthermia. *Lancet.* 1998; 352:1131–6. [PubMed: 9798607]
40. Bardo S, Cavazzini MG, Emptage N. The role of the endoplasmic reticulum Ca<sup>2+</sup> store in the plasticity of central neurons. *Trends Pharmacol Sci.* 2006; 27:78–84. [PubMed: 16412523]
41. Paschen W, Frandsen A. Endoplasmic reticulum dysfunction - a common denominator for cell injury in acute and degenerative diseases of the brain? *J Neurochem.* 2001; 79:719–25. [PubMed: 11723164]
42. Berridge MJ. Calcium signalling and Alzheimer's disease. *Neurochem Res.* 2011; 36:1149–56. [PubMed: 21184278]
43. Marambaud P, Dreses-Werringloer U, Vingdeux V. Calcium signaling in neurodegeneration. *Mol Neurodegener.* 2009; 4:20. [PubMed: 19419557]
44. Zhang Y, Dong Y, Wu X, Lu Y, Xu Z, Knapp A, Yue Y, Xu T, Xie Z. The mitochondrial pathway of anesthetic isoflurane-induced apoptosis. *J Biol Chem.* 2010; 285:4025–37. [PubMed: 20007710]
45. Assefa Z, Bultynck G, Szlufcik K, Nadif KN, Vermassen E, Goris J, Missiaen L, Callewaert G, Parys JB, De Smedt H. Caspase-3-induced truncation of type 1 inositol trisphosphate receptor accelerates apoptotic cell death and induces inositol trisphosphate-independent calcium release during apoptosis. *J Biol Chem.* 2004; 279:43227–36. [PubMed: 15284241]
46. Kalkman CJ, Peelen L, Moons KG, Veenhuizen M, Bruens M, Sinnema G, de Jong TP. Behavior and development in children and age at the time of first anesthetic exposure. *Anesthesiology.* 2009; 110:805–12. [PubMed: 19293699]
47. Wilder RT, Flick RP, Sprung J, Katusic SK, Barbaresi WJ, Mickelson C, Gleich SJ, Schroeder DR, Weaver AL, Warner DO. Early exposure to anesthesia and learning disabilities in a population-based birth cohort. *Anesthesiology.* 2009; 110:796–804. [PubMed: 19293700]
48. Moller JT, Cluitmans P, Rasmussen LS, Houx P, Rasmussen H, Canet J, Rabbitt P, Jolles J, Larsen K, Hanning CD, Langeron O, Johnson T, Lauven PM, Kristensen PA, Biedler A, van Beem H, Fraidakis O, Silverstein JH, Beneken JE, Gravenstein JS. Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study. ISPOCD investigators. International Study of Post-Operative Cognitive Dysfunction. *Lancet.* 1998; 351:857–61. [PubMed: 9525362]
49. Wei H, Inan S. Dual effects of neuroprotection and neurotoxicity by general anesthetics: Role of intracellular calcium homeostasis. *Prog Neuropsychopharmacol Biol Psychiatry.* 2013; 47:156–61. [PubMed: 23721657]
50. Gray JJ, Bickler PE, Fahlman CS, Zhan X, Schuyler JA. Isoflurane neuroprotection in hypoxic hippocampal slice cultures involves increases in intracellular Ca<sup>2+</sup> and mitogen-activated protein kinases. *Anesthesiology.* 2005; 102:606–15. [PubMed: 15731600]




**Figure 1. Isoflurane activates single inositol trisphosphate receptor ( $InsP_3R$ ) channel gating** (A–F) Typical single-channel current traces in membrane patches from nuclei isolated from DT40-KO cells stably expressing rate type 3  $InsP_3R$ . Concentrations of  $InsP_3$ , isoflurane and  $InsP_3R$  inhibitors used are as indicated. In this and all subsequent figures, currents shown were recorded at room temperature with applied potential ( $V_{app}$ ) =  $-40$  mV, and arrows indicate the closed channel current level. Concentration of free  $Ca^{2+}$  ( $[Ca^{2+}]_i$ ) =  $2 \mu M$  in all the experiments unless stated otherwise. (G–I) Bar graphs showing  $InsP_3R$  gating characteristics: mean open probability ( $P_o$ ), open duration ( $t_o$ ) and closed time ( $t_c$ ), respectively, in various inositol trisphosphate ( $InsP_3$ ) or isoflurane (ISO) concentrations. Error bars indicate standard error of the mean, and number of experiments analyzed is tabulated. \*\* and \* indicate  $P$  ( $t$ -test with Bonferroni correction)  $< 0.005$  and  $0.05$ , respectively, for the quantities connected by the brackets.




**Figure 2. Dependence of inositol trisphosphate receptor (InsP<sub>3</sub>R) channel activity on isoflurane concentration**

(A–D) Typical single-channel current traces of type 3 (InsP<sub>3</sub>R-3) channel activity in various isoflurane concentrations as indicated. Concentration of free Ca<sup>2+</sup> ( $[Ca^{2+}]_i$ ) = 2  $\mu M$ . See figure 1D for single InsP<sub>3</sub>R channel current trace in 400  $\mu M$  isoflurane. (E) [isoflurane] dependence of InsP<sub>3</sub>R-3 channel open probability  $P_o$ . Error bars indicate standard error of the mean, and number of experiments analyzed is tabulated.



**Figure 3. Ca<sup>2+</sup> dependence of isoflurane-activated type 3 inositol trisphosphate receptor (InsP<sub>3</sub>R-3) channel**

(A) Typical single channel current traces of InsP<sub>3</sub>R channels activated by 400 μM isoflurane in free Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>) of 900 nM. (B) [Ca<sup>2+</sup>]<sub>i</sub> dependence of InsP<sub>3</sub>R-3 channel open probability (P<sub>o</sub>) in 400 μM isoflurane. Error bars indicate standard error of the mean, and number of experiments analyzed is tabulated. Note the logarithmic channel P<sub>o</sub> axis. Low channel P<sub>o</sub> (<0.01, marked with §) were observed in all [Ca<sup>2+</sup>]<sub>i</sub> examined other than 2 μM. There is no statistically significant difference among these low channels P<sub>o</sub> (P > 0.05, ANOVA test). These channel P<sub>o</sub> were so low that the number of active channel(s) (N) in the membrane patch cannot be accurately determined. Thus, the data plotted are effectively the NP<sub>o</sub> value, which may be an over-estimate of the actual P<sub>o</sub> value.




**Figure 4. Effects of 400  $\mu M$  isoflurane on the activity of inositol trisphosphate receptor (InsP<sub>3</sub>R) channel in various  $[InsP_3]$**

(A–D) Typical single-channel current traces of InsP<sub>3</sub>R channels in various  $[InsP_3]$  in the absence and presence of 400  $\mu M$  isoflurane, as indicated. See figure 1A–C for InsP<sub>3</sub>R channel current traces in 0, 1, and 10  $\mu M$  InsP<sub>3</sub> with no isoflurane, and figure 1D for InsP<sub>3</sub>R channel current trace in 400  $\mu M$  isoflurane only. (E) Statistically similar InsP<sub>3</sub>R channel activities ( $P > 0.05$ , ANOVA test) were observed in 100 nM  $[InsP_3]$ –1  $\mu M$  in the absence of isoflurane. With 400  $\mu M$  isoflurane, statistically similar InsP<sub>3</sub>R channel activities ( $P > 0.05$ , ANOVA test) were also observed in 0  $[InsP_3]$ –1  $\mu M$ . Importantly, 400  $\mu M$  isoflurane significantly increased InsP<sub>3</sub>R channel activity at low  $[InsP_3]$  (0, 100, and 500 nM). Error bars indicate standard error of the mean, and number of experiments analyzed is tabulated. \*\* and \* indicate  $P$  ( $t$ -test)  $< 0.005$  and  $0.05$ , respectively, for the quantities connected by the brackets. InsP<sub>3</sub>R channel  $P_o$  in saturating 10  $\mu M$  InsP<sub>3</sub> were significantly


higher ( $P < 0.05$ ,  $t$ -test with Bonferroni correction) than those in subsaturating [InsP<sub>3</sub>], in the presence or absence of 400  $\mu$ M isoflurane. However, InsP<sub>3</sub>R channel activity in saturating (10  $\mu$ M) InsP<sub>3</sub> was not enhanced by 400  $\mu$ M isoflurane ( $P > 0.05$ ,  $t$ -test).





**Figure 5. Isoflurane amplifies cytosolic  $Ca^{2+}$  signals in DT40 cells by sensitizing inositol trisphosphate receptor (InsP<sub>3</sub>R) activation and induces cell apoptosis**

(A) Representative single cell  $Ca^{2+}$  responses to 400  $\mu$ M isoflurane (Iso) in DT-40 cells with triple knock out of InsP<sub>3</sub>R (DT40-KO) or DT40-KO transfected with rat recombinant type 3 InsP<sub>3</sub>R (DT40-R3) cells. Arrows mark the addition of iso to the complete growth medium. (B) Peak amplitudes and (C) area under curve (AUC) for DT40-KO and DT40-R3 cells responding to isoflurane. (D) Representative single cell  $Ca^{2+}$  responses and oscillations induced by addition (marked by arrows) of 50 ng/ml anti-IgM antibody ( $\alpha$ -IgM) or 50 ng/ml  $\alpha$ -IgM plus isoflurane in Dulbecco's Modified Eagle Medium. (E) Fraction of cells with various kind of response to stimulation by  $\alpha$ -IgM or  $\alpha$ -IgM+iso in the medium. (F) AUC of cells with single  $Ca^{2+}$  peak response triggered by  $\alpha$ -IgM or  $\alpha$ -IgM+iso. (G) AUC of cells with  $Ca^{2+}$  oscillations in response to stimulation by  $\alpha$ -IgM or  $\alpha$ -IgM+iso. (H) Isoflurane significantly elevated caspase-3 activity only in DT40 wild type (DT40-wt) cells or in DT40 expressing only InsP<sub>3</sub>R (DT40-R3), but not in DT40 cells with triple knock out of InsP<sub>3</sub>R (DT40-KO). All calcium measurement data are summary of at least 141 cells (N = 141, fig. 5 B and C) or 212 cells (N = 212, fig. 5 E, F, and G) from 4 separate experiments. The N values for caspase-3 activity are the average from three separate experiments (N = 3). Error bars indicate standard error of the mean, and number of experiments analyzed is tabulated. \*, \*\*, and \*\*\* indicate significant difference ( $P < 0.05$ ,  $P < 0.01$  and  $P < 0.001$ , respectively,  $t$  test). \* indicate significant difference ( $P < 0.05$ ,  $t$ -test) between quantities connected by the brackets in the bar graphs.



**Figure 6. Modulation of inositol trisphosphate (InsP<sub>3</sub>)receptor (InsP<sub>3</sub>R) activity by isoflurane and the effects on cell survival**

Isoflurane can activate InsP<sub>3</sub>R or potentiate the activation of InsP<sub>3</sub>R by endogenous agonist InsP<sub>3</sub> generated by various extracellular stimulation including antibody for IgM (IgM Ab). These effects can be inhibited by InsP<sub>3</sub>R antagonists xestospongin and heparin. Moderate activation of InsP<sub>3</sub>R by isoflurane at low concentration for short duration cause moderate Ca<sup>2+</sup> release from the endoplasmic reticulum (ER) and elevation of cytosolic Ca<sup>2+</sup>, which in turn, inhibit cell death by inducing endogenous cytoprotective mechanisms (left side), such as activation of phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)- protein kinase B (AKT) and microtubule-associated protein kinase (MAPK)/extracellular-signal-regulated kinases (ERK) pathways. Excessive activation of InsP<sub>3</sub>R by isoflurane at high concentration for prolonged duration cause excessive Ca<sup>2+</sup> release from the endoplasmic reticulum and abnormal elevation of cytosolic Ca<sup>2+</sup>, resulting in induction of cell death *via* apoptosis directly (right side), through rerelease of cytochrome C (Cyc C) from mitochondria into cytosolic space. ATP = adenosine triphosphate.

Table 1

Channel Detection Rate (Pd) of Isoflurane-activated Type 3 InsP<sub>3</sub>R

	Isoflurane (0.4 mM)	InsP <sub>3</sub> (μM)	Heparin (100 μg/ml)	Xestospongins C (μM)	Channel Detection Rate (Pd)	Patches (n)
	-	0	-	-	0/14(0)	14
	+	0	-	-	15/85(0.18)	85
Type-3 InsP <sub>3</sub> R	-	1	-	-	18/20(0.9)	20
	-	10	-	-	31/34(0.92)	34
	+	-	+	+	0/40(0)	40
	+	-	-	+	0/40(0)	40
InsP <sub>3</sub> R Knockout	-	10	-	-	0/10(0)	10
	+	0	-	-	0/40(0)	40

InsP<sub>3</sub>R – inositol 1,4,5-trisphosphate receptors.