

A long noncoding RNA connects c-Myc to tumor metabolism

Chiu-Lien Hung^{a,1}, Ling-Yu Wang^{a,1}, Yen-Ling Yu^b, Hong-Wu Chen^a, Shiv Srivastava^c, Gyorgy Petrovics^c, and Hsing-Jien Kung^{a,b,2}

^aDepartment of Biochemistry and Molecular Medicine, Comprehensive Cancer Center, University of California at Davis, Sacramento, CA 95817; ^bInstitute of Molecular and Genomic Medicine, National Health Research Institutes, Zhunan Town, Miaoli County 350, Taiwan; and ^cDepartment of Surgery, Center for Prostate Disease Research, Uniformed Services University of the Health Sciences, Bethesda, MD 20814-4799

Edited* by Michael G. Rosenfeld, University of California, San Diego, La Jolla, CA, and approved November 18, 2014 (received for review August 13, 2014)

Long noncoding RNAs (lncRNAs) have been implicated in a variety of physiological and pathological processes, including cancer. In prostate cancer, prostate cancer gene expression marker 1 (PCGEM1) is an androgen-induced prostate-specific lncRNA whose overexpression is highly associated with prostate tumors. PCGEM1's tumorigenic potential has been recently shown to be in part due to its ability to activate androgen receptor (AR). Here, we report a novel function of PCGEM1 that provides growth advantages for cancer cells by regulating tumor metabolism via c-Myc activation. PCGEM1 promotes glucose uptake for aerobic glycolysis, coupling with the pentose phosphate shunt to facilitate biosynthesis of nucleotide and lipid, and generates NADPH for redox homeostasis. We show that PCGEM1 regulates metabolism at a transcriptional level that affects multiple metabolic pathways, including glucose and glutamine metabolism, the pentose phosphate pathway, nucleotide and fatty acid biosynthesis, and the tricarboxylic acid cycle. The PCGEM1-mediated gene regulation takes place in part through AR activation, but predominantly through c-Myc activation, regardless of hormone or AR status. Significantly, PCGEM1 binds directly to target promoters, physically interacts with c-Myc, promotes chromatin recruitment of c-Myc, and enhances its transactivation activity. We also identified a c-Myc binding domain on PCGEM1 that contributes to the PCGEM1-dependent c-Myc activation and target induction. Together, our data uncover PCGEM1 as a key transcriptional regulator of central metabolic pathways in prostate cancer cells. By being a coactivator for both c-Myc and AR, PCGEM1 reprograms the androgen network and the central metabolism in a tumor-specific way, making it a promising target for therapeutic intervention.

lncRNA | tumor metabolism | c-Myc coactivator | prostate cancer

Long noncoding RNAs (lncRNAs) have recently drawn increasing attention as important players in physiological and pathological processes. In cancer, aberrant expression and mutations of lncRNAs can contribute to tumor development and progression by promoting proliferation, invasion, metastasis, and survival (1–3). lncRNAs thus may serve as diagnostic biomarkers and therapeutic targets for cancer. lncRNAs function at several levels of cellular processes, and the majority thus far studied are involved in gene regulation either at the transcriptional or posttranscriptional level (4). At the transcriptional level, lncRNA can serve as a chaperon or scaffold to deliver transcriptional factor to the chromatin site, to modulate the chromatin conformation by recruiting histone-modifying complexes, and to connect distal gene-regulatory elements together to effectively modulate the transcription of the targeted loci (4).

Prostate cancer gene expression marker 1 (PCGEM1) is a prostate tissue-specific lncRNA highly associated with prostate cancer (5). Over 80% of patient specimens show elevated levels (5), and the occurrence of PCGEM1 overexpression seems to be significantly higher in African-American patients, whose population has the highest prostate-cancer incidence in the world (6). The clinical evidence thus strongly indicates the tumorigenic potential of PCGEM1 in prostate-cancer development. The oncogenic property of PCGEM1 is further demonstrated by its ability to promote cell

proliferation and increase colony formation upon overexpression (6), as well as conferring resistance to doxorubicin-induced apoptosis via attenuation of p53 and p21 responses (7). Despite being a hormone-regulated malignancy at early stages, the majority of prostate cancer develops into hormone independence during progression, resulting in disease relapse, and makes the current hormone-deprivation therapy ineffective. Recent research has therefore emphasized mechanisms underlying cancer progression to hormone independence or castration resistance. In a systematic transcriptome analysis using a human prostate cancer cell line (LNCaP) mouse xenograft model to identify genes differentially expressed during tumor progression, PCGEM1 was found significantly up-regulated in the castration-recurrent stage, implicating its role in the development of hormone-refractory cancer (8). The recent work of Yang et al. (9) elegantly demonstrated that PCGEM1 is associated with and activates androgen receptor (AR), which contributes to the development of castration-resistant prostate cancer. Although a recent report does not support the role of PCGEM1 or PRNCR1 in AR activation (10), evidence from other studies described above suggests PCGEM1 as a potentially useful biomarker as well as a therapeutic target for prostate cancer. In the present study, we define a previously unidentified role of PCGEM1 in prostate carcinogenesis. It functions as a master regulator of tumor metabolism that facilitates the biosynthesis of cellular building materials, providing proliferative advantages for cancer cells. PCGEM1 regulates metabolic programming by enhancing activation of c-Myc and AR, which, in turn, control the

Significance

Long noncoding RNA (lncRNA) has been implicated in carcinogenesis and regarded as an emerging alternative target for cancer therapy. This study provides, to our knowledge, the first case of lncRNA that serves as a coactivator of c-Myc and a master regulator of tumor metabolism. This lncRNA, prostate cancer gene expression marker 1 (PCGEM1), is overexpressed in prostate cancer and implicated in castration resistance. It also forms an independent complex with androgen receptor (AR). Being a dual coactivator for AR and c-Myc, PCGEM1 reprograms the transcriptional network of metabolic genes and androgen-responsive genes, making it an ideal therapeutic target for prostate cancer. The present study provides significant insights into lncRNA's role in c-Myc activation, tumor metabolism, and prostate carcinogenesis.

Author contributions: C.-L.H., L.-Y.W., and H.-J.K. designed research; C.-L.H., L.-Y.W., and Y.-L.Y. performed research; C.-L.H., L.-Y.W., S.S., and G.P. contributed new reagents/analytic tools; C.-L.H., L.-Y.W., H.-W.C., and H.-J.K. analyzed data; and C.-L.H., L.-Y.W., and H.-J.K. wrote the paper.

The authors declare no conflict of interest.

*This Direct Submission article had a prearranged editor.

Freely available online through the PNAS open access option.

¹C.-L.H. and L.-Y.W. contributed equally to this work.

²To whom correspondence should be addressed. Email: hkung@ucdavis.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1415669112/-DCSupplemental.

expression profiles of multiple key metabolic pathways. We found that *PCGEM1* directly binds c-Myc, enhances c-Myc transactivation potency, and facilitates the recruitment of c-Myc to the chromatin target sites. These functions are independent of its association with AR. However, being a coactivator of both c-Myc and AR, *PCGEM1* reprograms both metabolic and AR genes and represents an ideal target for therapy.

Results

PCGEM1 Regulates Prostate Cancer Cell Growth and Tumor Metabolism.

PCGEM1 is an androgen-inducible gene whose expression is reported to associate exclusively with AR-positive cell lines (5). To further explore its physiological role, we used the androgen-responsive, androgen-dependent LNCaP prostate cancer cell line and developed derivatives with either constitutive overexpression of *PCGEM1* (LNCaP/*PCGEM1*) or with doxycycline (DOX)-inducible knockdown of *PCGEM1* (LNCaP/sh*PCGEM1*). Consistent with a previous report (6), *PCGEM1* overexpression led to accelerated cell growth (Fig. 1A). By contrast, knockdown of *PCGEM1* by DOX treatment to LNCaP/sh*PCGEM1* cells resulted in retarded proliferation (Fig. 1B) and G1 arrest (Fig. 1C). Prolonged *PCGEM1* knockdown also gave rise to increased caspase 3/7 activity, indicating induced apoptosis in the knockdown cells (Fig. 1D). Given that tumor cells often develop altered metabolism to cope with the demand of cell-mass increase during growth, we next examined whether the *PCGEM1*-dependent proliferation involves metabolic reprogramming. Indeed, LNCaP/*PCGEM1* showed increased glucose uptake and lactate production, indicating elevated glycolysis (Fig. 1A). *PCGEM1* overexpression also increased the activity of glucose-6-phosphate dehydrogenase (G6PD), the rate-limiting enzyme of pentose phosphate pathway (PPP) shunting the carbon flow from glucose to ribose-5-phosphate, through which generating the reducing agent NADPH that is essential for maintaining cellular redox status. The subsequent increase of ribose-5-phosphate presumably further accelerated nucleotide synthesis. Citrate, the intermediate for fatty-acid synthesis, was also increased in LNCaP/*PCGEM1* cells (Fig. 1A). These results suggest that *PCGEM1* overexpression leads to increased glucose uptake and glycolysis that facilitate macromolecule biosynthesis and ensures the supply of intracellular reducing energy. Knockdown of *PCGEM1* in contrast, resulted in decreased glucose consumption and lactate production, as well as decreased G6PD activity and citrate level,

indicating reduced glycolysis and anabolism (Fig. 1B). Together, our data suggest that *PCGEM1* plays an important role in regulating metabolism essential for cell-cycle progression, proliferation, and survival of LNCaP cells.

***PCGEM1* Regulates Metabolic Genes.** To investigate how *PCGEM1* regulates cellular metabolism, we surveyed expression of metabolic enzymes in *PCGEM1* knockdown cells (LNCaP/sh*PCGEM1* + DOX) and found that the *PCGEM1*-mediated metabolic alterations take place at the transcriptional level. Several enzymes involved in glucose uptake, glycolysis, PPP, lipid synthesis, glutamine metabolism, and the TCA cycle were prominently down-regulated in *PCGEM1* knockdown cells (Fig. 2). These data strongly suggest that *PCGEM1* functions as a key regulator of multiple metabolic genes, whose expression alterations in turn lead to metabolic outcomes beneficial to tumor growth.

PCGEM1 Regulates Metabolic Genes in both AR-Dependent and -Independent Manners.

In prostate-cancer cells, androgen signaling and AR play an important role in regulating cellular metabolism (11). Consistent with the report of Massie et al. (11), our data showed that androgen (DHT) significantly induced genes in multiple metabolic pathways, including glucose uptake, glycolysis, PPP, lipid, and nucleotide synthesis. Additionally, we found that several enzymes in glutamine metabolism and the TCA cycle were also induced upon DHT treatment (Fig. 3A). Strikingly, when knocking down *PCGEM1* (LNCaP/sh*PCGEM1* + DOX), the DHT-dependent gene induction was drastically compromised, suggesting that *PCGEM1* is a key regulator for androgen-dependent metabolic gene expression (Fig. 3A). *PCGEM1* has been recently shown to be a coactivator of AR (9). In agreement with Yang et al. (9), we independently confirmed the association of AR with *PCGEM1* and extended the results to demonstrate *PCGEM1*'s ability to transactivate the prostate-specific antigen (PSA) enhancer and FK506 binding protein 5 promoter, two well-known AR targets (Fig. S1A and B). We also showed that AR was recruited to a subset of the *PCGEM1*-regulated gene promoters and that the chromatin recruitment was partially reduced in *PCGEM1* knockdown cells (Fig. S1C). These results indicate that *PCGEM1* may facilitate AR binding to some of the metabolic gene promoters. However, given the limited reduction of AR chromatin occupancy in the knockdown cell observed, we




Fig. 1. *PCGEM1* regulates proliferation and metabolism. (A and B) MTT proliferation assay and metabolic profiles of *PCGEM1*-overexpressing cells (A) and DOX-inducible *PCGEM1* knockdown cells (B). Cells were collected for metabolic enzyme reactions on day 3 after lentiviral transduction or DOX treatment and were monitored for proliferation to 10 d. The measurement of glucose consumption, lactate production, G6PD activity, and citrate and NADPH levels were normalized by cell number. SD was derived from biological triplicates. Inset graphs illustrate *PCGEM1* expression level. (C) The DNA content of LNCaP/sh*PCGEM1* cells was detected by propidium iodide staining and flow-cytometry analysis. Population of each cell-cycle stage is quantified and plotted, as a percentage of the total cell population, in the stacked bar graph. (D) Caspase 3/7 activity was measured in LNCaP cells transduced with pLKO.1 or sh*PCGEM1* for the indicated days.




Fig. 2. *PCGEM1* regulates expression of metabolic enzymes in multiple pathways. Expression of the metabolic genes in *PCGEM1* knockdown LNCaP cell was examined by qRT-PCR. Positive and negative values indicate up and down-regulation compared with control cell, respectively.

anticipated that the *PCGEM1*-mediated AR activation and target transcription may rely on other mechanisms, such as the formation of androgen-dependent chromatin looping reported by Yang et al. (9). Together, our results suggest that *PCGEM1* plays an essential role in the androgen-induced metabolic gene regulation via AR.

Interestingly, in addition to its role in androgen response, *PCGEM1* seemed to be essential for the tumor metabolic regulation in hormone-deprived conditions as well. Under hormone deprivation (CDT), knockdown of *PCGEM1* caused down-regulation of multiple metabolic genes (Fig. 3A, +DOX CDT) and further resulted in significant metabolic alterations indicative of down-regulated glycolysis and macromolecule biosynthesis (Fig. 3B). These results revealed *PCGEM1*'s critical role in tumor metabolism independent of androgen signaling and suggest that, in addition to AR, another transcription factor may be involved in *PCGEM1*'s metabolic regulatory role. This assumption was strongly supported by the observation that, in AR-negative prostate cancer cell PC3, whose growth and survival do not require androgen, overexpression of *PCGEM1* was capable of widely inducing the metabolic genes (Fig. 3C). Our data support the notion that, as an androgen-inducibile gene, *PCGEM1* enhances AR activity and contributes to the androgen-induced metabolic reprogramming. Moreover, we also revealed *PCGEM1*'s critical role in metabolic regulation independent of androgen and AR.

***PCGEM1* Functions as a Coactivator of c-Myc.** Several transcription factors, such as c-Myc, p53 and HIF-1 α , are implicated in regulating cancer-cell metabolism (12, 13). To identify the transcription factors involved in *PCGEM1*-mediated metabolic regulation, we performed RNA immunoprecipitation (RIP) to examine whether they form complexes with *PCGEM1*. As shown in Fig. 4A, we found that, in addition to AR, ectopically expressed *PCGEM1* predominantly interacted with c-Myc whereas no significant association with p53 or HIF-1 α was detected. The *PCGEM1* and c-Myc interaction was also detected at the endogenous level, and the specificity was confirmed by c-Myc exclusively binding with *PCGEM1* but not *PRNCR1*, another prostate cancer-associated lncRNA (Fig. S2). Consistently, in RNA pull-down assays using purified recombinant c-Myc or LNCaP cell lysate, we showed that the in vitro transcribed biotinylated *PCGEM1* associated with c-Myc through direct binding (Fig. 4B). Given the evidence of physical interaction, we further tested whether *PCGEM1* directly enhances c-Myc transactivation potency. Using a Myc-responsive luciferase construct, we showed that overexpression of *PCGEM1* itself was capable of inducing c-Myc transactivation activity whereas the combination of *PCGEM1* and c-Myc overexpression synergistically enhanced the promoter activity (Fig. 4C). In a reciprocal experiment, the c-Myc activity was compromised when knocking down endogenous *PCGEM1* (+DOX), indicating that *PCGEM1* functions as a natural coactivator of c-Myc in prostate cancer cells (Fig. 4D). In conclusion, we identified c-Myc as a novel binding partner of *PCGEM1*, which positively regulates the transactivity of c-Myc.

c-Myc Is the Major Effector of *PCGEM1*-Dependent Metabolism Regulation. Global metabolic programming is one of the most prominent c-Myc properties in tumorigenesis (14). Our findings that *PCGEM1* formed complexes with and coactivated c-Myc strongly suggest c-Myc as a key effector of *PCGEM1* in metabolic gene regulation. This hypothesis was supported by the following evidence. First, several metabolic enzymes showing more than twofold expression alteration upon *PCGEM1* knockdown have been documented as c-Myc targets (Fig. 2; summarized in Fig. S3) (14, 15). Second, chromatin isolation by RNA purification (ChIRP) analysis showed that *PCGEM1* physically associated to a subset of the metabolic gene promoters (Fig. 5A) that are c-Myc binding sites identified in The Encyclopedia of DNA Elements (ENCODE) ChIP-seq data (also confirmed by our c-Myc ChIP assay) (Fig. 5B). Therefore, the DNA loci that were detected in the captured *PCGEM1* complex suggest overlapped chromatin occupancy of *PCGEM1* with c-Myc, indicating lncRNA-transcriptional complex formation on these targets. The few c-Myc binding loci failing to show *PCGEM1* enrichment could be due to various reasons, including the assay stringency or the




Fig. 3. Metabolic gene regulation in androgen-dependent and -independent manners. (A) Metabolic expression profiles of LNCaP/shPCGEM1 cells cultured in hormone-deprived (CDT) or androgen-treated (DHT) conditions. The expression levels of control cells in CDT medium (-DOX CDT) were used as baseline to compare with other treatment. Relative expression fold changes are illustrated by heat map (red, up-regulation; blue, down-regulation). Color scale indicates the 95th percentile of either up- or down-regulated expression range. (B) Metabolic profiles of the *PCGEM1* knockdown cell under hormone deprivation. The procedures for *PCGEM1* knockdown and enzymatic reactions are as described in Fig. 1B. (C) Metabolic gene expression in PC3 cells overexpressing control vector or *PCGEM1* was detected by qRT-PCR. The normalized expression levels were compared using vector control as the baseline (shown by fold, $P < 0.05$).




Fig. 4. *PCGEM1* functions as a c-Myc coactivator. (A) Association of ectopically expressed *PCGEM1* with HA-tagged c-Myc, AR, p53, or HIF-1 α was detected by RIP assay. The relative levels of coimmunoprecipitated *PCGEM1* were calculated as fold difference compared with vector control (* P < 0.05, ** P < 0.01). (B) RNA pull-down of the in vitro transcribed biotinylated *PCGEM1* incubated with LNCaP cell lysate (Left) or purified recombinant c-Myc protein (Right). The biotinylated GFP mRNA served as a negative control. (C) Myc responsive luciferase was cotransfected with empty vector, c-Myc, *PCGEM1*, or both into PC3 cells for the reporter assay. Coexpression of *PCGEM1* and c-Myc significantly enhanced the luciferase activity. (D) The luciferase activity (same as C) was measured in LNCaP/sh*PCGEM1* cell cultured with or without DOX treatment. Knockdown of the endogenous *PCGEM1* significantly reduced c-Myc transactivity. The relative luciferase activity was calculated by normalization against *Renilla-Luc* activity (* P < 0.05, ** P < 0.01).

possibility that *PCGEM1* associates with the target at regions distal to the c-Myc binding site.

To further understand the mechanism of how *PCGEM1* regulates c-Myc, we first tested whether it increases the protein stability of c-Myc. We found that no significant changes in the protein turnover rate were detected (Fig. S4A) and that no detectable difference of total c-Myc protein level was observed in the *PCGEM1* knockdown cell either (Fig. 5B, Inset). We next tested whether *PCGEM1* promotes c-Myc loading to the chromatin. Significantly, the ChIP assay confirmed c-Myc occupancy on a significant number of *PCGEM1*-dependent metabolic gene promoters and showed that its recruitment was drastically impaired by the knockdown of *PCGEM1* (+DOX) (Fig. 5B). Concurrently, the recruitment of RNA polymerase II (PolII) was generally diminished in *PCGEM1* knockdown cells (Fig. 5C), indicating a suppressed transcriptional status of these metabolic genes. c-Myc-mediated transcription in vivo involves several histone modification events at the target promoters (16). In particular, once bound to the target, c-Myc induces local hyperacetylation of both histone H3 and H4, further maintaining the active chromatin status (16, 17). Given the reduced c-Myc occupancy upon *PCGEM1* knockdown, we examined whether the histone hyperacetylation on the target loci is decreased as well. Consistently, we found that *PCGEM1* knockdown (+DOX) resulted in decreased H3 and H4 acetylation on most of the targets (Fig. 5D and E), indicating inactive chromatin and compromised transcriptional status. Together, these data suggest that *PCGEM1* forms complexes with c-Myc on their target loci and functions in promoting the recruitment of c-Myc and presumably other transcriptional regulators to enhance c-Myc transcriptional activity.

Structural and Functional Domain Mapping of *PCGEM1*. To map the c-Myc binding domain on *PCGEM1*, various in vitro transcribed and biotinylated *PCGEM1* truncates were incubated with recombinant c-Myc protein for RNA pull-down assays. The results showed that, compared with others, the RNA fragment

containing nucleotides 481–878 associated to c-Myc with the highest affinity (Fig. 6A). Subsequently, we used synthesized tiling oligos that are complementary to the *PCGEM1* transcript to probe the c-Myc-bound *PCGEM1* fragments. The fourteenth probe complementary to nucleotides 781–840 exhibited the strongest intensity (Fig. 6B). This region falls within the fragment 481–878 identified in Fig. 6A. Conversely, both RIP assays and in vitro RNA pull-down assays showed significantly reduced association of c-Myc with the mutant *PCGEM1* Δ 761–849 (Fig. 6A and C), strongly suggesting that nucleotides 781–840 represent the major c-Myc binding site on *PCGEM1*. Importantly, the c-Myc binding site seemed to be essential for *PCGEM1*'s role in c-Myc transactivation. We found that the synergistic effect of *PCGEM1* on c-Myc transactivation was significantly diminished when expressing the mutant *PCGEM1* (Δ 761–849) (Fig. 6D) whereas its efficacy on AR transactivation was not affected (Fig. 6E). Given that AR binds to *PCGEM1* on a distinct site locating on nucleotides 421–480 (9) (Fig. S1D), our data suggest that the c-Myc binding domain on *PCGEM1* is functionally distinct from that of AR. Finally, using the AR-negative PC3 cell, we tested whether the c-Myc binding domain on *PCGEM1* is essential for the downstream target induction. As shown in Fig. 6F, compared with full-length *PCGEM1*, the ability of the Δ 761–849 mutant to




Fig. 5. *PCGEM1* associates with the target chromatin and promotes c-Myc recruitment and active transcription. (A) Chromatin occupancy of *PCGEM1* to the target loci was detected by ChIP assay. The *PCGEM1*-associated metabolic gene promoters were detected by qPCR, and the recovered DNA level was estimated as the percentage of input chromatin. *Actin* promoter was used as negative control. The Inset graph illustrates the efficiency and specificity of RNA retrieved from streptavidin-bound probes. *GAPDH* mRNA was used to evaluate nonspecific binding of the biotinylated probes. (B–E) ChIP analysis of c-Myc (B) and PolII (C) recruitment, and acetylated-histone H3 (D), -histone H4 (E) on metabolic gene promoters. Recovered DNA in the precipitated complexes was analyzed by qPCR, and the DNA levels were normalized against nonspecific IgG-bound DNA to obtain the enrichment folds, indicated by fold of IgG. Significantly reduced chromatin enrichment in *PCGEM1* knockdown cell is indicated by asterisks (* P < 0.05, ** P < 0.01). The Inset Western blots show similar c-Myc protein level in –DOX and +DOX cells.

induce metabolic genes was considerably weakened. A similar gene regulation pattern was observed when overexpressing the full-length and deletion mutant *PCGEM1* in LNCaP cells although the presence of AR and the endogenous *PCGEM1* may have masked some of the c-Myc effects (Fig. S5). Together, we identified the c-Myc binding domain on *PCGEM1* and showed that the structural domain for c-Myc binding is consistent as the functional domain for c-Myc target-gene regulation.

Discussion

Distinct from normal cells, cancer cells acquire alterations in central metabolic pathways to fulfill their high demands on biomass and energy production, while maintaining appropriate redox. These metabolic changes are critical for cancer cells to sustain rapid proliferation and adapt to a dynamic tumor microenvironment (12, 18, 19). In this report, we identified the lncRNA *PCGEM1* as a key regulator of metabolic pathways that provide proliferating advantages for prostate-cancer cells. We found that *PCGEM1*-overexpressing cells showed significantly enhanced glucose uptake and lactate production, indicative of an increased glycolysis rate, as well as an increased cellular level of citrate, G6PD activity, and NADPH, indicating elevated biosynthesis of fatty acid and nucleotide and redox control. Conversely, knockdown of *PCGEM1* resulted in opposite metabolic outcomes that further arrested cell-cycle progression and induced apoptosis. These findings indicate an essential role of *PCGEM1* in tumor metabolic regulation, critical for cancer-cell proliferation and survival. By enhancing aerobic glycolysis, the most common metabolic phenotype, known as the Warburg effect in cancer (20), overexpression of *PCGEM1* facilitates anabolism to produce cellular building materials (21, 22) and consequently drives the cell to higher proliferation potential.

Our study has uncovered *PCGEM1* lncRNA as a key transcriptional regulator of metabolic genes that profoundly affects the gene expression profiles in several pathways linked to tumor metabolism. The *PCGEM1*-mediated tumor metabolic regulation can largely be attributed to its dual role as c-Myc and AR coactivators. In prostate-cancer cells, AR has been shown to regulate metabolic genes involved in glycolysis, PPP, fatty acid, and nucleotide synthesis (11). On the other hand, as one of the

most highly amplified oncogenes in many cancers (23), c-Myc emerges as a master regulator of global metabolism, including those regulated by AR, as well as glutamine metabolism and the TCA cycle (14, 15). We showed, in both cases, that *PCGEM1* associated with the transcription factors, enhanced their trans-activation activities, enriched their recruitments to the target promoters, and up-regulated the target genes in all of the pathways described above. Our data indicate that the complex formation of *PCGEM1* with c-Myc and AR is mutually independent. First, *PCGEM1* associated with c-Myc in the absence of AR in vitro, and vice versa. Moreover, the *PCGEM1*-c-Myc association was detected in AR-negative cells. Second, c-Myc did not interact with lncRNA *PRNCR1*, which is required for AR and *PCGEM1* complex formation (9). Third, *PCGEM1* enhanced c-Myc activity and up-regulated c-Myc-targeted metabolic genes in cells without AR expression. Finally, c-Myc and AR bind to *PCGEM1* on distinct domains, and deletion of the c-Myc binding site did not affect the *PCGEM1*-induced AR transcriptional activity. Therefore, we propose that the *PCGEM1*-mediated transactivation of c-Myc and AR is functionally and structurally distinct. We cannot completely rule out that, under certain conditions, c-Myc and AR may be embedded in the same complex and coregulate certain targets. We have used ChIP-reChIP to test whether there is *PCGEM1*-mediated corecruitment of c-Myc on the AR-targeted loci. At least for *ENO1*, *HK2*, *GLS*, *GSR*, and *LDHA* promoters, such corecruitment was not observed (Fig. S6A), indicating the formation of two independent *PCGEM1* complexes. On the other hand, even without coexisting in the same complex, *PCGEM1*-dependent activation of c-Myc and AR can enhance the cross talk between these two transcriptional programs. It has previously been reported that AR is able to induce the expression of c-Myc, in turn reinforcing amplification of the AR-transcriptional signals (24). The c-Myc induction by AR can also confer hormone-independent growth of prostate cancer cells (25, 26). Our data showed that, upon androgen stimulation, the metabolic gene induction was drastically diminished when knocking down *PCGEM1* (Fig. 3A) or c-Myc (Fig. S6B), indicating that the androgen-induced metabolic reprogramming is dependent on the action of *PCGEM1* and c-Myc. Together, we propose a model for the androgen-stimulated




Fig. 6. c-Myc binding domain on *PCGEM1*. (A) In vitro binding of c-Myc recombinant protein with full-length and various *PCGEM1* constructs. Equal amount of biotinylated *PCGEM1* RNA was incubated with c-Myc in the pull-down assay. The streptavidin-RNA-bound c-Myc was analyzed by Western blotting. (B) The recombinant c-Myc and *PCGEM1* RNA pull-down assay was coupled with a dot blot assay. Twenty-seven probes, each containing 60 bases of the complementary *PCGEM1* sequence, were dotted on membrane for c-Myc binding domain detection. Probe 14 complementary to *PCGEM1* 781–840 exhibited the strongest signal. The biotinylated full-length *PCGEM1* was dotted at right lower corner as a positive control for streptavidin-HRP chemiluminescent reaction. (C) Association of HA-tagged c-Myc with full-length (FL) or mutant (Δ761–849) *PCGEM1* was detected by RIP assay. (D) Myc-responsive element reporter and (E) PSA enhancer reporter assays were carried out to determine the transactivities of c-Myc and AR in *PCGEM1* (FL and Δ761–849) overexpressing PC3 cell, as described in Fig. 4C. (F) Metabolic gene expression in PC3 cells overexpressing control vector, full-length, or mutant (Δ761–849) *PCGEM1*. The relative expression levels were calculated as described in Fig. 3C. Asterisks indicate statistical significance (* $P < 0.05$, ** $P < 0.01$).

metabolic regulation that is through AR-dependent transcriptional up-regulation of the key regulators *PCGEM1* and c-Myc, as well as the formation of a dual *PCGEM1* complex with AR and c-Myc (Fig. S6C).

Regarding the mechanisms as to how *PCGEM1* serves as a coactivator of transcriptional factors, Yang et al. demonstrated that the *PCGEM1*-induced AR activation involves looping of the chromatin enhancer to the promoter for subsequent transcriptional activation (9). The data from Prensner et al. (10), however, indicated otherwise. In our system, we were able to detect *PCGEM1*-enhanced AR activation, but we also identified an AR-independent function of *PCGEM1* through c-Myc. At present, we know little about how *PCGEM1* enhances the transcriptional potential of c-Myc. We observed that the chromatin recruitment of c-Myc is compromised when knocking down *PCGEM1*, indicating that *PCGEM1* facilitates binding of c-Myc to the targeted promoters tested. Because heterodimerization of c-Myc and Max is sufficient to bind the E-box containing DNA in vitro (27, 28), it is unlikely that the DNA binding of c-Myc per se in prostate cancer cell requires *PCGEM1*. In fact, we found that *PCGEM1* did not affect the interaction of c-Myc with Max or c-Myc protein stability (Fig. S4), indicating a mechanism independent of facilitating active Myc–Max dimerization or increasing the c-Myc protein level for enhanced transactivation. On the other hand, because the target recognition and binding for c-Myc in vivo depends on preexisting chromatin modification, such as histone methylation and acetylation (29), it is possible that *PCGEM1* binds to epigenetic modifiers or specific histone marks of active chromatin, and in turn, promotes and stabilizes c-Myc enrichment on the target chromatin. The detailed mechanism requires further investigation.

Given the kinship of lncRNA to gene expression, the roles of lncRNAs in regulating metabolism are anticipated but have not been extensively studied. Recently, it was shown that *lncRNA-p21* associates with HIF-1 α and modulates glycolysis under hypoxia (30). As to c-Myc regulation, two recent reports identified lncRNA *CCAT1-L* and *GHET1* involved in modulating transcription and RNA stability of c-Myc (31, 32). To our knowledge, the present study is the first report of an lncRNA that binds c-Myc and

functions as a coactivator of c-Myc that modulates metabolic programming. Targeting tumor metabolism and its key regulators has emerged as an alternative strategy to complement the conventional genotoxic stress-based cancer therapy (19, 33, 34). One rationale is that “starvation” therapy may use a different cellular death mechanism to overcome the resistance to apoptosis often developed after cancer therapy (35). However, despite some success, the challenge of targeting individual metabolic enzymes in tumor, while avoiding toxic effects on normal proliferating cells, remains due to their essential housekeeping roles. Because *PCGEM1* is prostate tissue-specific (5) and is overexpressed in cancer cells (6), it may serve as a unique target of metabolic regulation for prostate-cancer therapy. The specific role of *PCGEM1* in prostate cancer metabolism and AR activation (ref. 9 and this study) makes it an ideal therapeutic target for prostate cancer.

Materials and Methods

Detailed materials and methods are provided in *SI Materials and Methods*. The sequences of all primers and probes used in this study are listed in *Tables S1* and *S2*.

LNCaP cells stably expressing Tet repressor (TR) were used to generate the inducible *PCGEM1* knockdown cell line (LNCaP/sh*PCGEM1*). Briefly, lentiviral particles carrying the sh*PCGEM1* construct that was driven by Tet-operating H1 promoter were generated according to the manufacturer’s instructions (Invitrogen) and were subsequently infected to LNCaP/TR cells. After 48 h of transduction, drug-resistant clones were selected by zeocin. The obtained stable cell lines were cultured and maintained in blasticidin (10 ng/mL) and zeocin (100 ng/mL; Invitrogen). *PCGEM1* expression levels in the isolated zeocin-resistant clones were monitored by quantitative RT (qRT)-PCR to confirm knockdown efficiency. To induce knockdown, doxycycline (DOX) (100 ng/mL) was added to culture medium for 3 d. *PCGEM1*-overexpressing LNCaP and PC3 cells were also generated by the lentiviral system, following the same procedure.

ACKNOWLEDGMENTS. This work was supported by National Science Council of Taiwan Grant NSC102-2320-B-400-018. We also acknowledge US National Institutes of Health Grants CA114575, CA165263, and DK065977, as well as National Health Research Institutes of Taiwan Grant O3A1-MGPP18-014.

1. Tsai MC, Spitale RC, Chang HY (2011) Long intergenic noncoding RNAs: New links in cancer progression. *Cancer Res* 71(1):3–7.
2. Gibb EA, Brown CJ, Lam WL (2011) The functional role of long non-coding RNA in human carcinomas. *Mol Cancer* 10:38.
3. Spizzo R, Almeida MI, Colombatti A, Calin GA (2012) Long non-coding RNAs and cancer: A new frontier of translational research? *Oncogene* 31(43):4577–4587.
4. Geisler S, Coller J (2013) RNA in unexpected places: Long non-coding RNA functions in diverse cellular contexts. *Nat Rev Mol Cell Biol* 14(11):699–712.
5. Srikantan V, et al. (2000) *PCGEM1*, a prostate-specific gene, is overexpressed in prostate cancer. *Proc Natl Acad Sci USA* 97(22):12216–12221.
6. Petrovics G, et al. (2004) Elevated expression of *PCGEM1*, a prostate-specific gene with cell growth-promoting function, is associated with high-risk prostate cancer patients. *Oncogene* 23(2):605–611.
7. Fu X, Ravindranath L, Tran N, Petrovics G, Srivastava S (2006) Regulation of apoptosis by a prostate-specific and prostate cancer-associated noncoding gene, *PCGEM1*. *DNA Cell Biol* 25(3):135–141.
8. Romanuik TL, et al. (2010) LNCaP Atlas: Gene expression associated with in vivo progression to castration-recurrent prostate cancer. *BMC Med Genomics* 3:43.
9. Yang L, et al. (2013) lncRNA-dependent mechanisms of androgen-receptor-regulated gene activation programs. *Nature* 500(7464):598–602.
10. Prensner JR, et al. (2014) The lncRNAs *PCGEM1* and *PRNCR1* are not implicated in castration resistant prostate cancer. *Oncotarget* 5(6):1434–1438.
11. Massie CE, et al. (2011) The androgen receptor fuels prostate cancer by regulating central metabolism and biosynthesis. *EMBO J* 30(13):2719–2733.
12. Cairns RA, Harris IS, Mak TW (2011) Regulation of cancer cell metabolism. *Nat Rev Cancer* 11(2):85–95.
13. Ward PS, Thompson CB (2012) Metabolic reprogramming: A cancer hallmark even Warburg did not anticipate. *Cancer Cell* 21(3):297–308.
14. Dang CV (2013) MYC, metabolism, cell growth, and tumorigenesis. *Cold Spring Harb Perspect Med* 3(8):a014217.
15. Morrish F, Isern N, Sadilek M, Jeffrey M, Hockenbery DM (2009) c-Myc activates multiple metabolic networks to generate substrates for cell-cycle entry. *Oncogene* 28(27):2485–2491.
16. Martinato F, Cesaroni M, Amati B, Guccione E (2008) Analysis of Myc-induced histone modifications on target chromatin. *PLoS ONE* 3(11):e3650.
17. Frank SR, Schroeder M, Fernandez P, Taubert S, Amati B (2001) Binding of c-Myc to chromatin mediates mitogen-induced acetylation of histone H4 and gene activation. *Genes Dev* 15(16):2069–2082.
18. Dang CV (2012) Links between metabolism and cancer. *Genes Dev* 26(9):877–890.
19. Schulze A, Harris AL (2012) How cancer metabolism is tuned for proliferation and vulnerable to disruption. *Nature* 491(7424):364–373.
20. Warburg O (1956) On the origin of cancer cells. *Science* 123(3191):309–314.
21. Newsholme EA, Crabtree B, Ardawi MS (1985) The role of high rates of glycolysis and glutamine utilization in rapidly dividing cells. *Biosci Rep* 5(5):393–400.
22. Vander Heiden MG, Cantley LC, Thompson CB (2009) Understanding the Warburg effect: The metabolic requirements of cell proliferation. *Science* 324(5930):1029–1033.
23. Beroukhi R, et al. (2010) The landscape of somatic copy-number alteration across human cancers. *Nature* 463(7283):899–905.
24. Ni M, et al. (2013) Amplitude modulation of androgen signaling by c-MYC. *Genes Dev* 27(7):734–748.
25. Bernard D, Pourtier-Manzanedo A, Gil J, Beach DH (2003) Myc confers androgen-independent prostate cancer cell growth. *J Clin Invest* 112(11):1724–1731.
26. Gao L, et al. (2013) Androgen receptor promotes ligand-independent prostate cancer progression through c-Myc upregulation. *PLoS ONE* 8(5):e63563.
27. Papoulas O, Williams NG, Kingston RE (1992) DNA binding activities of c-Myc purified from eukaryotic cells. *J Biol Chem* 267(15):10470–10480.
28. Littlewood TD, Amati B, Land H, Evan GI (1992) Max and c-Myc/Max DNA-binding activities in cell extracts. *Oncogene* 7(9):1783–1792.
29. Guccione E, et al. (2006) Myc-binding-site recognition in the human genome is determined by chromatin context. *Nat Cell Biol* 8(7):764–770.
30. Yang F, Zhang H, Mei Y, Wu M (2014) Reciprocal regulation of HIF-1 α and lncRNA-p21 modulates the Warburg effect. *Mol Cell* 53(1):88–100.
31. Xiang JF, et al. (2014) Human colorectal cancer-specific *CCAT1-L* lncRNA regulates long-range chromatin interactions at the MYC locus. *Cell Res* 24(5):513–531.
32. Yang F, et al. (2014) Long non-coding RNA *GHET1* promotes gastric carcinoma cell proliferation by increasing c-Myc mRNA stability. *FEBS J* 281(3):802–813.
33. Tennant DA, Durán RV, Gottlieb E (2010) Targeting metabolic transformation for cancer therapy. *Nat Rev Cancer* 10(4):267–277.
34. Vander Heiden MG (2011) Targeting cancer metabolism: A therapeutic window opens. *Nat Rev Drug Discov* 10(9):671–684.
35. Kim RH, Bold RJ, Kung HJ (2009) ADI, autophagy and apoptosis: Metabolic stress as a therapeutic option for prostate cancer. *Autophagy* 5(4):567–568.