

Detection of *Salmonella* in Blood by PCR using *iroB* gene

VITHIYA GANESAN¹, BELGODE NARASIMHA HARISH², GODFRED ANTONY MENEZES³, SUBASH CHANDRA PARIJA⁴

ABSTRACT

Background: *Salmonella*, a genus of more than 2500 serological variants (serovars), includes many organisms that can cause human disease. Enteric fever remains an important public health problem in developing countries. Non typhoidal *Salmonella* generally produce a self limited gastroenteritis in healthy individuals whereas in extremes of age and immunocompromised cause severe fatal disease. The protean manifestations make this disease a true diagnostic challenge.

Aim: The present study was carried out to optimize PCR for detecting the *Salmonella* genus using *iroB* gene and evaluate its use in the rapid diagnosis of typhoid and non typhoidal salmonellosis.

Materials and Methods: The study was carried out between August 2009 and July 2011 on blood samples from patients

attending JIPMER hospital, Pondicherry, India with clinical suspicion of enteric fever and salmonellosis. Whole blood was used DNA extraction and conventional PCR done with *iroB* and *fliC* primers. Blood culture and Widal test were performed for all the patients.

Statistical Analysis: Performed using Fischer's exact test with Graphpad InStat 3.

Results: PCR results were compared with blood culture. Sensitivity and specificity of PCR with *fliC* gene are 95.6% and 93.3% respectively. Sensitivity and specificity of PCR with *iroB* gene are 96.6% and 93.3% respectively.

Conclusion: With *iroB* gene, additional cases of *Salmonella* Paratyphi A and non typhoidal *Salmonella* were detected when compared to *fliC* gene.

Keywords: Non typhoidal *Salmonella*, *iroB* gene, *fliC* gene

INTRODUCTION

Salmonellae are facultative intracellular pathogens. *Salmonella enterica* serovar Typhi is exclusively host-restricted to humans, and early intracellular replication within reticuloendothelial mononuclear cells is critical for pathogenesis [1]. In contrast, non-typhoidal *Salmonella* (NTS) serovars have a broad host range and usually cause self-limiting mucosal diarrheal disease and carriage in immunocompetent humans [2].

Culturing blood specimens has become much faster and easier, since the advent of automated blood culture instruments such as the BACTEC 9000 systems (Becton Dickinson, Cockeysville, MD, USA) and BacTAlert (BioMerieux, Durham, NC, USA). However, it still takes several days to detect and identify *Salmonella* [3,4]. Due to the time required for blood culture identification alternative methods of identification of *Salmonella* are being sought. In particular, DNA detection methods such as the polymerase chain reaction (PCR) have been investigated [5]. PCR-based typing techniques are becoming increasingly popular in developing countries. PCR can be useful in salmonellosis especially in cases when empirical antibiotic is started and bacterial load is low.

Many of the molecular methods done for detection of *Salmonella* in blood were based on using *fliC* gene which is specific for *Salmonella* typhi. The flagellum of *S. typhi* like all other bacterial flagella is composed of pre-distinct portion, basal body, hook and filament. The filament is made up of repeating subunits of a single protein, flagellin. *Salmonella* typhi typically has only phase I flagellar antigen, d. This d antigen however is the phase I antigen in a number of *Salmonella* species like *Salmonella muenchen*. A unique sequence present in the hypervariable region VI of *S. typhi* distinguishes it from all other *Salmonella* spp especially *S. muenchen*. PCR for *S. typhi* utilizes primers designed to detect this region of the flagellin gene. Studies have shown a good specificity with these primers [6]. On the other hand, the incidence of enteric fever caused by *S. Paratyphi A* has been increasing in Asia. It is estimated to have more than 5 million cases of paratyphoid fever compared to more than 25 million cases of typhoid fever [7]. Typhoid is a highly adapted invasive

disease that is restricted to humans and shows little association with the immunocompromised. In contrast, non-typhoidal salmonellosis have a broad vertebrate host range, an epidemiology that often involves food and animals, and a dramatically more severe and invasive presentation in immunocompromised adults, in particular in those with HIV. The prevalence of non-typhoidal *Salmonella* (NTS) bacteraemia has risen in many countries and is probably related to the increase in HIV infection. Although invasive disease caused by NTS has been recently reported from many African and Asian countries, the infection is relatively unknown in India.

So, in the current study the primer is chosen such that it detects other typhoidal as well as NTS. Analysis of the evolution of the genus *Salmonella* revealed that it is of monophyletic origin and that *Salmonella* serotypes last had a common ancestor with close relatives, such as *Escherichia coli*, some 100 to 150 million years ago [8]. During their separate evolution, *Salmonella* serotypes acquired many genes by phage or plasmid-mediated horizontal transfer. These genes now distinguish this genus from other bacteria and are thus obvious candidates for the development of DNA-based methods for identification of *Salmonella* serotypes. Two genetic regions on the *S. enterica* chromosome, *Salmonella* pathogenicity island 2 and the *iroBC* operon, indeed show this phylogenetic distribution [9].

The first genes of the *iroA* locus, designated *iroBC*, were identified in *S. enterica* serotype Typhi during a genetic screen for genes that are regulated by the iron response regulator Fur (Ferric Uptake Regulator). Regulation by Fur results in expression of *iroBC* under iron-limited growth conditions. In contrast, during growth under iron sufficiency expression is prevented by binding of the Fur-Fe²⁺ repressor complex to a Fur DNA binding site in the *iroB* promoter region. However, the *iroBC* gene products show homology to proteins that have so far not been associated with iron uptake or defense against oxidative stress in other bacteria. *IroB* shows homology with bacterial glycosyl transferases, and *IroC* is a member of the ATP binding cassette (ABC) family of transport proteins [10].

Primers	Nucleotide sequence
<i>fljC</i> -F	5'- ACTGCTAAAACCACTACT -3'
<i>fljC</i> -R	5'-TGGAGACTTCGGTTGCGTAG-3'
<i>iroB</i> -F	5'-TGC GTATTCTGTTTGTCCGGTCC-3'
<i>iroB</i> -R	5'-TACGTTCCCACCATTCTTCCC-3'

[Table/Fig-1]: Primers used for PCR amplification of *fljC* and *iroB* genes

Age	Number(%)	Salmonella in blood culture
1-10	5(8.3)	4
11-20	20(33.3)	13
21-30	16(26.6)	8
31-40	11(18.3)	2
41-50	2(3.3)	1
>50	6(10)	2

[Table/Fig-2]: Age details of the study population

	Blood culture	<i>fljC</i> gene positive	<i>iroB</i> gene positive
<i>Salmonella</i> Typhi	24	23	23
<i>Salmonella</i> Paratyphi A	1	0	1
Group B <i>Salmonella</i>	3	0	3
<i>Salmonella</i> spp	2	0	2

[Table/Fig-3]: Comparison of results between blood culture, *fljC* and *iroB* gene

AIM

To optimize PCR for *iroB* gene of *S. enterica* and evaluate its usefulness in the rapid diagnosis of typhoidal and non typhoidal salmonellosis.

OBJECTIVES


To optimize PCR for *iroB* gene of *Salmonella enterica*. and to compare *iroB* gene detection with that of *fljC* gene in diagnosis of typhoid fever.

MATERIALS AND METHODS

The study was approved by Institute Ethics Committee, JIPMER, Pondicherry. This study was carried out between August 2009 and July 2011 on blood samples from 60 patients attending JIPMER Hospital, Pondicherry with clinical suspicion of enteric fever and non typhoidal salmonellosis. Two groups of 30 patients each were included in the study. Members of both sexes and all age groups were included. Group I consisted of blood samples from 30 patients whose blood culture grew *Salmonella* spp. Group II consisted of blood samples from 30 patients with clinical suspicion of Salmonellosis but blood cultures negative for *Salmonella* spp.

Patients with clinical suspicion of enteric fever who tested positive by Standard Agglutination test for Brucellosis, leptospirosis by IgM ELISA, rickettsial infections by Weil Felix test, malarial parasites by peripheral smear and QBC were excluded from the study. Three to five ml of blood were collected from patients coming to microbiology outpatient department in EDTA coated containers with informed consent. Pure culture of *Salmonella* Typhi H901 was used for standardisation of PCR.

DNA extraction: Extraction of DNA from whole blood was done as per procedure described by Haque et al., [11] with several modifications. Blood samples collected in EDTA containers inoculated into BHI broth and incubated for 6 h. One ml of supernatant was centrifuged at 10,000 rpm for 5 min. One ml of lysis buffer (0.2% Triton X100 in Tris HCl pH 8.0) was added to the pellet. The mixture was gently aspirated several times to effect hemolysis. The tube was centrifuged at 12,000 rpm for 6 min, the supernatant was discarded, and the procedure was repeated several times till

[Table/Fig-4]: Gel image after PCR for *iroB* gene of *S. enterica*

a clear white pellet was observed. The pellet was washed twice with distilled water. After the removal of the supernatant, the pellet was resuspended in 20-30 μ l of distilled water. The tubes were sealed, kept in boiling water for 20 min, and brought back to room temperature before being used as a template for PCR.

PCR Conditions: Primers targeting the flagellin gene of *Salmonella* Typhi were used after verifying against the database in the GenBank {GenBank accession no (AC): L21912} oligonucleotides ST3 (5'- ACTGCTAAAACCACTACT -3') and ST4 (5'- TGGAGACTTCGGTTGCGTAG-3') were used to amplify a 366 bp fragment corresponding to nucleotide 1060-1077 and 1407-1426 {(AC): L21912} [6]. Forward and reverse primers of *iroB* gene include oligonucleotides (5'-TGC GTATTCTGTTTGTCCGGTCC-3') and (5'-TACGTTCCCACCATTCTTCCC-3'). With these primers, a 606-bp DNA fragment could be amplified from all *S. enterica* isolates [10]. A 25 μ l amplification mixture containing 10 μ l of master mix (Bangalore Genei, Bangalore) 7 μ l of MilliQ water, 1 μ l *fljC*-F, 1 μ l *fljC*-R, 1 μ l *iroB*-F, 1 μ l *iroB*-R with 4 μ l of the extracted DNA were used. Primer details are in [Table/Fig-1]. The reaction was carried out in an Eppendorf Mastercycler (Eppendorf, Germany). The reaction mixture was submitted to touch down PCR of 25 cycles of 2 min denaturation at 94 $^{\circ}$ C, annealing at 68 $^{\circ}$ C for 1 min 15 sec and elongation at 72 $^{\circ}$ C for 2 min followed by 20 cycles of initial denaturation of 94 $^{\circ}$ C of 2 min, annealing at 55 $^{\circ}$ C for 1 min and elongation at 72 $^{\circ}$ C for 2 min. A final elongation of 10 min duration was done at 72 $^{\circ}$ C.

Detection of PCR products: A reaction mixture of 5 μ l was fractionated electrophoretically at 80V for 45 min in a 1.5% agarose gel containing 0.5 μ g of ethidium bromide. A positive control (Amplification mixture with DNA extracted from standard strain of *Salmonella* Typhi) and a negative control (Amplification mixture without DNA) were also included in each lot. The bands obtained were documented using a gel documentation system. With *iroB* primers, a 606-bp DNA fragment could be amplified from all *S. enterica* isolates. The *fljC* primers amplified a 366 bp fragment, in the flagellin gene of *S. typhi*.

Blood culture: Under aseptic precautions, 5-10 ml of blood from adults and 1-3ml of blood from paediatric age group was collected and inoculated into a biphasic medium consisting of Brain Heart Infusion (BHI) agar and BHI broth (Hi Media Laboratories Ltd, Mumbai). Bottles were checked daily for growth on the BHI slant or any turbidity in the BHI broth. If no visible growth occurred, blind subculture was performed. Any colony obtained on subculture was identified by biochemical tests as per standard recommendations and confirmed by specific antisera. (Denka Seiken Co. Ltd, Japan).

RESULTS

In this study, the age of the patients ranged from 6 to 60 yrs with a mean of 26 yrs. Age distribution of the study population described

in [Table/Fig-2]. Of the 60 patients included in the study, 30 grew *Salmonella* isolates in the blood culture. 24 isolates of *Salmonella* Typhi, 2 *Salmonella* spp, 3 Group B *Salmonella* and 1 *Salmonella* Paratyphi A. Of which 23 were positive for *fljC* gene by PCR. Over all 29 samples were positive for *iroB* gene [Table/Fig-3 & 4]. One sample which has grown *S.typhi* in blood culture and negative for *fljC* and *iroB* by PCR. This could be due to inhibitors in blood, mainly heme and leucocyte DNA, or added anticoagulants like EDTA. Two samples which were negative by blood culture were positive for *fljC* and *iroB* by PCR. PCR was also carried out on 10 blood samples which yielded *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter* spp. in culture. All these samples were negative for *fljC* and *iroB* genes by PCR.

DISCUSSION

Salmonellosis remains a major cause of morbidity and mortality in developing countries. Early diagnosis is not only necessary for detection of the cases but also to prevent fatal complications like intestinal perforation in case of enteric fever and focal infections like meningitis, osteomyelitis, septic arthritis and pneumonia in case of NTS. Age distribution of the population is needed to define the optimum age of immunization and choice of vaccines for public health programmes in developing countries. Highest attack rates occur in school going children and young adults. Invasive nontyphoidal salmonellosis shows a markedly contrasting age distribution showing a bimodal age distribution. In this study, though the number of cases of NTS is too small to extrapolate the pattern, they followed the same bimodal age distribution.

Generally isolation from blood culture is poor due to factors like prior antibiotic therapy, low bacterial count as low as 1CFU/ml [12]. Blood culture is usually negative in 30-65% of typhoid fever patients [13]. After standardization of PCR for flagellin gene of *S.Typhi*, it was evaluated against the results of blood culture in the absence of a better diagnostic test. PCR was able to detect flagellin gene in 25 out of the 60 patients (41.6%). The sensitivity of PCR was determined to be 96%. Several studies have reported high sensitivity of PCR of blood for *S.Typhi* [9,14]. PCR detected *iroB* gene in 31 out of the 60 blood samples (51.6%). The sensitivity of PCR was determined to be 96.6%.

However, initial research indicated that PCR has similar sensitivity to blood culture and lower specificity [15]. A prospective study of the concentration of *S. Typhi* in blood of typhoid fever patients showed a median value of 0.3 (range of 0.1 to 399) CFU/ml, well below current PCR-based detection limits [16]. Gordon et al [2] have shown that NTS in bacteremic patients are present at a similarly low concentration (1 CFU/ml).

To overcome the difficulties caused by the low numbers of *Salmonella* bacteria present in typhoid patient blood samples, pre-enrichment of bacteria is necessary prior to PCR detection. A recent study demonstrated that the 5 hour broth culture enrichment improved PCR sensitivity by 10 times for spiked blood, and 100 times for spiked stool samples [17]. In another study, ox bile tryptone soy broth was optimized for blood culture which caused lysis of cells to release the intracellular bacteria without inhibiting growth of

S.Typhi, and then detected by PCR targeting *fljC* of *S.Typhi* [18]. In the current study, 6 hour BHI broth culture enrichment done prior to DNA extraction and by using *iroB* gene typhoidal and non typhoidal *Salmonellae* are detected which is advantageous over the more commonly targeted gene, *fljC*.

CONCLUSION

With *iroB* gene, additional cases of *S.Paratyphi A* and non typhoidal *Salmonella* were detected when compared to *fljC* gene. DNA extraction results were good after six hours of BHI enrichment of blood samples rather than directly from blood. In PCR positive and culture negative cases, antibiotic sensitivity results will not be known. Primers targeting drug resistance genes will serve the purpose.

REFERENCES

- [1] Bhan MK, Bahl R, Bhatnagar S. Typhoid and paratyphoid fever. *Lancet*. 2005;366(9487):749-62.
- [2] Gordon MA, Kankwatira AMK, Mwafurirwa G, Walsh AL, Hopkins MJ, Parry CM. Invasive Non-typhoid *Salmonellae* Establish Systemic Intracellular Infection in HIV-Infected Adults: An Emerging Disease Pathogenesis. *Clin Infect Dis*. 2010;50(7):953-62.
- [3] Reisner BS, Woods GL. Times to Detection of Bacteria and Yeasts in BACTEC 9240 Blood Culture Bottles. *J Clin Microbiol*. 1999;37(6):2024-26.
- [4] Durmaz G, Us T, Aydinli A, Kiremitci A, Kiraz N, Akgun Y. Optimum Detection Times for Bacteria and Yeast Species with the BACTEC 9120 Aerobic Blood Culture System: Evaluation for a 5-Year Period in a Turkish University Hospital. *J Clin Microbiol*. 2003;41(2):819-21.
- [5] Wain J, Hosoglu S. The laboratory diagnosis of enteric fever. *J Infect Dev Ctries*. 2008;2(6):421-25.
- [6] Khan S, Harish BN, Menezes GA, Acharya NS, Parija SC. Early diagnosis of typhoid fever by nested PCR for flagellin gene of *Salmonella enterica* serotype Typhi. *Indian J Med Res*. 2012;136: 850-54.
- [7] Crump JA, Luby SP, Mintz ED. The global burden of typhoid fever. *Bull. World Health Organ*. 2004; 82(5):346-53.
- [8] Ochman H, Wilson AC. Evolution in bacteria: Evidence for a universal substitution rate in cellular genomes. *J Mol Evol*. 1987;26(1-2):74-86.
- [9] Baumler AJ, Tsolis RM, van der Velden AW, Stojiljkovic I, Anic S, Heffron F. Identification of a new iron regulated locus of *Salmonella typhi*. *Gene*. 1996;183(1-2):207-13.
- [10] Baumler AJ, Heffron F, Reissbrodt R. Rapid detection of *Salmonella enterica* with primers specific for *iroB*. *J Clin Microbiol*. 1997;35:1224-30.
- [11] Haque A, Ahmed J, Qureshi JA. Early detection of typhoid by polymerase chain reaction. *Ann Saudi Med*. 1999;19(4):337-40.
- [12] Wain J, Diep TS, Ho VA, Walsh AM, Hoa NTT, Parry CM. Quantitation of Bacteria in Blood of Typhoid Fever Patients and Relationship between Counts and Clinical Features, Transmissibility, and Antibiotic Resistance. *J Clin Microbiol*. 1998;36(6):1683-87.
- [13] Chaudhry R, Laxmi BV, Nisar N, Ray K, Kumar D. Standardisation of polymerase chain reaction for the detection of *Salmonella typhi* in typhoid fever. *J Clin Pathol*. 1997;50(5):437-39.
- [14] Sánchez-Jiménez MM, Cardona-Castro N. Validation of a PCR for diagnosis of typhoid fever and salmonellosis by amplification of the *hilA* gene in clinical samples from Colombian patients. *J Med Microbiol*. 2004;53:875-78.
- [15] Levy H, Diallo S, Tennant SM, Livio S, Sow SO, Tapia M. PCR method to identify *Salmonella enterica* serovars Typhi, Paratyphi A, and Paratyphi B among *Salmonella* Isolates from the blood of patients with clinical enteric fever. *J Clin Microbiol*. 2008;46(5):1861-66.
- [16] Wain J, Pham VB, Ha V, Nguyen NM, To SD, Walsh AL. Quantitation of bacteria in bone marrow from patients with typhoid fever: relationship between counts and clinical features. *J Clin Microbiol*. 2001;39(4):1571-76.
- [17] Bhutta ZA. Current concepts in the diagnosis and treatment of typhoid fever. *BMJ*. 2006; 333(7558):78-82.
- [18] Zhou L, Pollard AJ. A fast and highly sensitive blood culture PCR method for clinical detection of *Salmonella enterica* serovar Typhi. *Ann Clin Microbiol Antimicrob*. 2010; 19(9):14.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Microbiology, Velammal Medical College Hospital and Research Institute, Madurai, India.
2. Head of the Department, Department of Microbiology, Jawaharlal Institute of Postgraduate Medical Science and Research, Puducherry, India.
3. Assistant Professor, College of Applied Medical Sciences, University of Ha'il, Ha'il, Kingdom of Saudi Arabia (KSA).
4. Dean, Jawaharlal Institute of Postgraduate Medical Science and Research, Puducherry, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. G. Vithiya,
4/437, Babu Nagar Main Road, Anupanadi, Madurai, India.
Phone : 9486312483, E-mail : vithiya.md@gmail.com

Date of Submission: **Mar 09, 2014**
Date of Peer Review: **Jul 02, 2014**
Date of Acceptance: **Sep 10, 2014**
Date of Publishing: **Nov 20, 2014**

FINANCIAL OR OTHER COMPETING INTERESTS: None.