



Published in final edited form as:  
*J Am Chem Soc.* 2013 ; 135(29): 10878.

## Isolation of notoamide E, a key precursor in the biosynthesis of prenylated indole alkaloids in a marine-derived fungus, *Aspergillus* sp

**Sachiko Tsukamoto,**

Kumamoto University, Graduate School of Pharmaceutical Sciences Kato, Hikaru

**Hikaru Kato,**

Kumamoto University, Graduate School of Pharmaceutical Sciences, Greshock, Thomas

**Thomas J. Greshock,**

Colorado State University, Chemistry Hirota, Hiroshi

**Hiroshi Hirota,**

Riken Advanced Science Institute, Ohta, Tomihisa

**Tomihisa Ohta,** and

Kanazawa University, Faculty of Pharmaceutical Sciences Williams, Robert

**Robert M. Williams**

Colorado State University, Department of Chemistry

---

In Figure 1, the structures of notoamide C (**3**) and 3-*epi*-notoamide C (**7**) should be corrected as shown: This error was caused by the configuration of notoamide C, which was initially assigned as 3*R* based on biogenetic considerations.<sup>1</sup> Recently, the absolute configuration of notoamide C was corrected to be 3*S* based on the biochemical conversion of notoamide E into notoamide C by recombinant NotB.<sup>2</sup> Recently, the relative and absolute configuration of notoamide C was independently reported by Chen, et al. by single crystal X-ray analysis of notoamide C isolated from *Aspergillus* sp. XS-20090066.<sup>3</sup> Accordingly the configuration of **7** was corrected to be 3*R*.

## REFERENCES

1. Kato H, Yoshida T, Tokue T, Nojiri Y, Hirota H, Ohta T, Williams RM, Tsukamoto S. *Angew. Chem., Int. Ed.* 2007; 46:2254.
2. Li S, Finefield JM, Sunderhaus JD, McAfoos TJ, Williams RM, Sherman DH. *J. Am. Chem. Soc.* 2012; 134:788. [PubMed: 22188465]
3. Chen M, Shao C-L, Fu X-M, Xu R-F, Zheng J-J, Zhao D-L, She Z-G, Wang C-Y. *J. Nat. Prod.* 2013; 76:547–553. [PubMed: 23527875]

