

ORIGINAL ARTICLE

Adaptation of an abundant *Roseobacter* RCA organism to pelagic systems revealed by genomic and transcriptomic analyses

Sonja Voget¹, Bernd Wemheuer¹, Thorsten Brinkhoff², John Vollmers¹, Sascha Dietrich¹, Helge-Ansgar Giebel², Christine Beardsley², Carla Sardemann², Insa Bakenhus², Sara Billerbeck², Rolf Daniel¹ and Meinhard Simon²

¹Institute of Microbiology and Genetics, Genomic and Applied Microbiology and Göttingen Genomics Laboratory, University of Göttingen, Göttingen, Germany and ²Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, Germany

The RCA (*Roseobacter* clade affiliated) cluster, with an internal 16S rRNA gene sequence similarity of >98%, is the largest cluster of the marine *Roseobacter* clade and most abundant in temperate to (sub)polar oceans, constituting up to 35% of total bacterioplankton. The genome analysis of the first described species of the RCA cluster, *Planktomarina temperata* RCA23, revealed that this phylogenetic lineage is deeply branching within the *Roseobacter* clade. It shares not >65.7% of homologous genes with any other organism of this clade. The genome is the smallest of all closed genomes of the *Roseobacter* clade, exhibits various features of genome streamlining and encompasses genes for aerobic anoxygenic photosynthesis (AAP) and CO oxidation. In order to assess the biogeochemical significance of the RCA cluster we investigated a phytoplankton spring bloom in the North Sea. This cluster constituted 5.1% of the total, but 10–31% (mean 18.5%) of the active bacterioplankton. A metatranscriptomic analysis showed that the genome of *P. temperata* RCA23 was transcribed to 94% in the bloom with some variations during day and night. The genome of *P. temperata* RCA23 was also retrieved to 84% from metagenomic data sets from a Norwegian fjord and to 82% from stations of the Global Ocean Sampling expedition in the northwestern Atlantic. In this region, up to 6.5% of the total reads mapped on the genome of *P. temperata* RCA23. This abundant taxon appears to be a major player in ocean biogeochemistry.

The ISME Journal (2015) 9, 371–384; doi:10.1038/ismej.2014.134; published online 1 August 2014

Introduction

Our understanding of the role of abundant individual taxa in ocean biogeochemistry is hampered by the fact that so far very few of such taxa have been isolated and are available to genomic and postgenomic analyses (Yooshep *et al.*, 2010). It is extremely difficult to obtain representative isolates of the major players and only novel approaches made this effort more successful (Giovannoni and Stingl, 2007). However, still today detailed information on the role of abundant individual taxa in oceanic cycling of matter is available only for *Prochlorococcus* and for *Cand. Pelagibacter ubique* of the SAR11 clade. *Pelagibacter ubique* of the SAR11 clade (Giovannoni *et al.*, 2005; Tripp *et al.*, 2008; Sowell *et al.*, 2009; Thompson *et al.*, 2011). Isolates of these taxa are available, their genomes

have been sequenced and thus form a basis for postgenomic studies and relating metagenomic and, more importantly, metatranscriptomic and metaproteomic information to individual taxa.

The *Roseobacter* and SAR11 clades are the most prominent subdivisions of Alphaproteobacteria in the ocean's near surface waters (Giovannoni and Stingl, 2005). The RCA (*Roseobacter* clade affiliated) cluster with an internal sequence similarity of the 16S rRNA gene of at least 98%, constitutes up to 35% of total bacterioplankton and is most abundant in temperate to (sub)polar oceans, but absent in tropical and subtropical regions (Selje *et al.*, 2004; Giebel *et al.*, 2009; Giebel *et al.*, 2011). It is divided into a subcluster with sequences from temperate regions and one with sequences of subpolar and polar origin (Giebel *et al.*, 2011). Because of the high abundance and distinct biogeography of the RCA cluster, there is great interest to elucidate its functional role and biogeochemical significance. However, no information on the genomic data and features of organisms of this important cluster are yet available because RCA organisms defy isolation and cultivation greatly.

Correspondence: M Simon, Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, C v Ossietzky Str 9-11, Oldenburg D-26111, Germany.

E-mail: m.simon@icbm.de

Received 24 April 2014; revised 17 June 2014; accepted 21 June 2014; published online 1 August 2014

An isolate of this cluster, retrieved from the southern North Sea, became recently available and was characterized as type species of the first described species of the genus *Planktomarina*, *Planktomarina temperata* RCA23 (Giebel *et al.*, 2013). In the North Sea, *P. temperata* RCA23 represents the most abundant ribotype of the RCA cluster that comprises persistently between 2% and 20% of total bacterioplankton (Selje *et al.*, 2004; Giebel *et al.*, 2011; Teeling *et al.*, 2012) and is a major representative of the *Roseobacter* clade in the active bacterioplankton (Wemheuer *et al.*, 2014). Based on these observations we hypothesized that the genome of this organism exhibits distinct features of its adaptation to life in the nutrient-poor pelagic environment with respect to genome organization and metabolic potential. Further, we hypothesized that the genome of *P. temperata* RCA23 is well represented in metagenomic and metatranscriptomic data in the North Sea because of its high abundance and activity. Therefore, we sequenced the genome of *P. temperata* RCA23 to elucidate its metabolic potential. In addition, we assessed its significance and functional role during a phytoplankton spring bloom in the southern North Sea by applying a metatranscriptomic approach. Finally, metagenomic data sets of pelagic marine systems were mined for the presence, abundance and genomic features of this organism.

Materials and methods

Origin and growth of P. temperata RCA23

P. temperata RCA23 was originally isolated from a water sample collected in the southern North Sea (Giebel *et al.*, 2011, 2013). It was grown in liquid culture of autoclaved sea water amended with marine broth (40% of peptone and yeast extract; Giebel *et al.*, 2013) to an optical density of 0.2. Biomass was harvested by 20 min centrifugation (Beckman JA10, Krefeld, Germany) at 7500 r.p.m. and 4 °C and DNA was extracted using the MasterPure DNA Purification Kit (Epicentre, Madison, Wisconsin, USA) according to the manufacturer's protocol.

Sequence determination, gene annotation and phylogenetic analyses

To sequence the genome of *P. temperata* RCA23 a pyrosequencing run was performed using a Roche GS-FLX 454 sequencer (Branford, CT, USA) with Titanium chemistry. All sequencing steps were performed according to the manufacturer's protocols and recommendations. In total 411 932 reads were generated and assembled to 78 contigs bigger than 500 bp with a 26-fold coverage. Furthermore, 576 fosmid Sanger-sequences were added to the data set to identify the contig order. Gap closure and polishing were carried out using the Staden software package (Staden, 1996) and PCR-based techniques on genomic DNA. Open reading frames were

identified using YACOP (Tech and Merkl, 2003) and GLIMMER (Delcher *et al.*, 2007). The open reading frame finding was inspected manually with Artemis and open reading frames were corrected by checking the GC frame plot, ribosomal binding site and blast hits against the NCBI nr database.

Orthologous protein sequences were identified using reciprocal BLASTp-analysis combined with global alignments based on the Needleman–Wunsch algorithm. Only hits with e -values $< 1e - 20$ were considered and additionally filtered, based on sequence identity cutoffs of the respective global alignments. A cutoff value of 30% sequence identity was chosen to identify orthologs. For multilocus sequence analysis, protein sequences of 162 genes, for which one ortholog but no paralog was found in every comparison strain, were concatenated. The multilocus sequence analysis tree was constructed using ARB v5.1 (Ludwig *et al.*, 2004) and the evolutionary history was inferred using the neighbor-joining method and recovered reproducibly with the maximum-likelihood method (Figure 1).

Study area, sample collection and chlorophyll

The significance of the RCA cluster was investigated during a phytoplankton spring bloom in the southern North Sea. Samples were collected at 11 stations at 2 m depth between 25 and 31 May 2010 on board RV Heincke by 41 Niskin bottles mounted on a CTD rosette (Sea-Bird, Bellevue, WA, USA). For pyrosequencing, metagenomic and metatranscriptomic analyses, 50 l of sea water were prefiltered through a 10- μ m nylon net and a filter sandwich consisting of a precombusted glass fiber and 3- μ m polycarbonate filter (47 mm diameter). Bacterioplankton was harvested from a prefiltered 1-liter sample on a filter sandwich consisting of a glass fiber and 0.2- μ m polycarbonate filter (47 mm). One filter sandwich was used for RNA extraction. For gene expression analysis and metagenome sequencing, at least four filter sandwiches were subjected to RNA and DNA extraction. Chlorophyll *a* concentrations were determined as described (Giebel *et al.*, 2011).

Enumeration of bacteria, CARD-FISH and BrdU-FISH

Bacterial cell numbers were determined by flow cytometry after staining of subsamples with SybrGreen I (Sigma-Aldrich, Munich, Germany), preserved with glutardialdehyde (final concentration 1%) and stored at -80 °C until further analysis as described Giebel *et al.* (2011). The abundance of RCA cells was determined by catalyzed reporter deposition fluorescence *in situ* hybridization (CARD-FISH) and the proportion of DNA-synthesizing RCA cells by incorporation of BrdU (4 h incubation) according to Pernthaler *et al.* (2002) and applying an RCA-specific probe set (RCA996, C Beardsley and I Bakenhus, unpublished data). Cells were counted via epifluorescence microscopy using $\times 1000$ magnification and suitable filter sets for DAPI-stained total cells, Alexa₄₈₈-stained RCA cells,




Figure 1 Neighbor-joining tree based on multilocus sequence analysis (MLSA) of genome-sequenced organisms of the *Roseobacter* clade. The tree was constructed using ARB v5.1 and is based on a similar tree of Newton *et al.* (2010) but includes additional genome sequences and an extended gene set. Filled circles indicate nodes also recovered reproducibly with maximum-likelihood calculation. Numbers at the nodes are bootstrap values (only >50% are shown) from 1000 replicates. Subclades of the *Roseobacter* clade are marked by different colors. (T) indicates type strains. *Escherichia coli* K12 MG1655 was used as outgroup.

and Cy3-stained (Perkin Elmer, Waltham, MA, USA) BrdU-positive cells, respectively.

16S rRNA gene amplicons, metagenomics and transcriptomics

Environmental DNA and RNA were coextracted and the composition of the active bacterial community was assessed by 16S rRNA PCRs as described before Wemheuer *et al.* (2014). For metagenomic and metatranscriptomic analyses DNA and cDNA were sequenced on an Illumina/Solexa GAIIX system (San Diego, CA, USA). In total, 54 334 282 paired-end sequences of 100 bp were generated for the metagenomic and 78 042 122 single-read sequences of 75–100 bp for the metatranscriptomic data sets, respectively (Supplementary Table S1). The sequences were quality trimmed and the Illumina adapter sequences were removed with Trimmomatic v0.30 (Bolger *et al.*, 2014) using the following parameters: adapter:2:40:15 leading:3 trailing:3

slidingwindow:4:15 minlen:50. Reads derived from ribosomal RNA gene fragments were filtered with SortMeRNA (Kopylova *et al.*, 2012). The remaining sequences were mapped with Bowtie 2 (Langmead and Salzberg, 2012) using the implemented end-to-end mode, which requires that the entire read align from one end to the other.

To compare the RNA-Seq results, the read counts were normalized to remove biases like the length of the transcript and the sequencing depth of a sample. We used the Nucleotide activity Per Kilobase of exon model per million Mapped reads (NPKM), a derivative of RPKM (reads per kilo base per million), as a normalized read count value (Wiegand *et al.*, 2013).

Data deposition

The genome sequence of *P. temperata* RCA23 has been deposited at the NCBI GenBank database with the accession number CP003984. The metagenomic

and metatranscriptomic data sets have been deposited at MG-Rast with the accession numbers: 4548721.3 (Station 3 gDNA), 4550305.3 (Station 3 cDNA), 4548722.3 (Station 9 gDNA), 4550303.3 (Station 9 cDNA), 4548723.3 (Station 13 gDNA) and 4550304.3 (Station 13 cDNA).

Results and discussion

Genomic features of *P. temperata* RCA23

P. temperata RCA23 is deeply branching within the *Roseobacter* clade, not affiliated to any of the known subclades, as shown by multilocus sequence analysis (Figure 1). The genome of *P. temperata* RCA23 encompasses 3.29 Mbp and carries 3101 genes of which 3054 encode proteins (Supplementary Table S2). It is the smallest of all closed and the third smallest of all genomes of the *Roseobacter* clade and carries the second lowest number of genes of all organisms of this clade (Figure 2a and Supplementary Table S2). The only organism of this clade with a significantly smaller genome is

Rhodobacterales bacterium HTCC2255 (Luo *et al.*, 2013). Further, partial genomes of two other organisms of the *Roseobacter* clade with estimated genome sizes of <3 Mbp were recently reported from sequence analyses of single amplified genomes retrieved from coastal and oceanic surface waters (Swan *et al.*, 2013; Luo *et al.*, 2014). Members of the *Roseobacter* clade with a genome size of <3.8 Mbp are exclusively of pelagic origin (Figure 2a and Supplementary Table S2). The mean genome sizes of the pelagic members of this clade is 4.06 Mbp and 0.5 Mbp smaller than that of members associated to other organisms, surfaces or sediments (Figure 2b and Supplementary Table S3). The relatively small genome of *P. temperata* RCA23 and other pelagic members of the *Roseobacter* clade and the reduction in genomic traits is consistent with genomic streamlining and adaptation to nutrient-poor pelagic marine ecosystems (see below; Giovannoni *et al.*, 2005; Luo *et al.*, 2013; Swan *et al.*, 2013). The genomic difference between both groups of the *Roseobacter* clade obviously reflects the divergent evolutionary history of this clade with genomic




Figure 2 Genomic traits of *P. temperata* RCA23 in comparison to other organisms of the *Roseobacter* clade. Relation between genome size and number of genes (a). Box-Whisker plot of the GC content (b), percentage coding genes (c) and CDS (coding DNA sequences) count (d) of 14 members of the *Roseobacter* clade associated to other organisms, surfaces or sediments and of 24 members with a pelagic life style listed in Supplementary Table S2. Data include 12 closed and 26 draft genomes. Blue circle and +: *P. temperata* RCA23. The boxes show the median (solid line), mean (dashed line), the 25th and 75th percentile and the whiskers the 10th and 90th percentiles. •: Outlayers. For further details see Supplementary Tables S2 and S3.

streamlining of at least several pelagic members and gained genomic content by the associated members (Luo *et al.*, 2013). The much larger genome size of *P. temperata* RCA23 and other pelagic members of the *Roseobacter* clade, however, harbour genomes much larger than highly streamlined marine pelagic bacteria such as *Cand. P. ubique* of the SAR11 clade. This feature points out that, besides streamlining, distinct aspects of their life history and adaptation to their niche in the marine pelagic realm were also important in shaping the evolution of these pelagic members of the *Roseobacter* clade (Giovannoni *et al.*, 2014). These authors have identified a low number of σ -factors as a feature to distinguish between streamlined and non-streamlined genomes of bacteria with a more complex life style. The genome of *P. temperata* RCA23 harbors six σ -factors, a value much lower than that of other members of the *Roseobacter* clade and in the same range as that of many typical streamlined genomes of a size of <2 Mb (Giovannoni *et al.*, 2014).

The GC content of 54% of *P. temperata* RCA23 is lower than that of all associated (mean = $60.07 \pm 3.58\%$, $N = 14$) and the fifth lowest of the 24 pelagic members of the *Roseobacter* clade included in this analysis (mean = $59.28 \pm 6.82\%$; Supplementary Tables S2 and S3). However it is substantially higher than the genomic GC content of many other marine bacteria, ranging below 40%, such as *Rhodobacterales* bacterium HTCC2255 (Supplementary Table S2), *Cand. P. ubique* and pelagic bacteria subjected to single cell genome sequencing (Swan *et al.*, 2013; Luo *et al.*, 2014). It has been speculated that a reduced genomic GC content may be an adaptation to nitrogen limitation in marine systems (Swan *et al.*, 2013), but clear-cut evidence is still missing, also because it is still unclear whether pelagic heterotrophic bacteria are carbon- or nitrogen-limited. The percentage of coding DNA of *P. temperata* RCA23 is very similar to the mean of all pelagic members of the *Roseobacter* clade which, however, is significantly higher than the mean of the organisms of this clade associated to other organisms, surfaces or sediment (Figure 2c and Supplementary Table S3).

The genome of *P. temperata* RCA23 shares not >65.7% of homologous genes (2009) with other members of the *Roseobacter* clade (Supplementary Tables S2 and S4). It harbors no prophage, which is consistent with the fact that prophage induction using mitomycin C and UV light was unsuccessful (H-A Giebel, unpublished data). Further, the genome carries no plasmid, no complete GTA (gene transfer agent) cluster (Supplementary Table S4) and no CRISPR (clustered regularly interspaced small palindromic repeats). CRISPR are uncommon in the genomes of most organisms of the *Roseobacter* clade and were so far detected in only two members of this clade, *Dinoroseobacter shibae* and *Maritimibacter alkaliphilus* HTCC2654. The lack of plasmids in the genome of *P. temperata* RCA23 and other pelagic

members of this clade with a genome size of <3.5 Mb (Supplementary Table S2) appears to be an adaptation to the pelagic life style and exhibits another feature of streamlining not considered yet in other analyses. In contrast, plasmids are common genomic elements of all members of the *Roseobacter* clade associated to other organisms, surfaces and sediments and constitute between 2% and 33% of the genomic content (Supplementary Table S2; Pradella *et al.*, 2010).

GTAs have been detected in Alphaproteobacteria and all genomes of the *Roseobacter* clade except in *Rhodobacterales* bacteria HTCC2083 and HTCC2255 (Zhao *et al.*, 2009; Newton *et al.*, 2010). GTA-related gene transfer was suggested as a potential adaptation mechanism of these bacteria to maintain the metabolic flexibility in the dynamic marine environment (Biers *et al.*, 2008). The fact that the genome of *P. temperata* RCA23 encodes only three putative GTA-related genes (c18030, c18040 and c18050; Supplementary Table S4) implies that this mode of gene transfer was discarded during adaptation to the pelagic life style, presumably because of little benefits by this type of genetic exchange in the nutrient-poor pelagic environment. Hence, the mentioned missing genomic features, the absence of plasmids and GTA in the genome of *P. temperata* RCA23 indicate a less diversified genomic organization as compared to other organisms of the *Roseobacter* clade and suggests reduced genetic exchange with other members of this clade.

In comparison to typical groups of other *Roseobacter* clade organisms (Newton *et al.*, 2010) 10 genomic islands (GIs), harboring 22.6% of all genes, are present in the genome of *P. temperata* RCA23 (Supplementary Figure S1). This bacterium encodes genes for chemotaxis, possesses a monotrichous flagellum and is motile (Table 1; Giebel *et al.*, 2013). Phylogenetic analysis of the flagella synthesis cluster present in GI 3 shows that the flagellar gene sets of the *Roseobacter* clade are divided into two distinct groups (Supplementary Figure S2). The flagella genes of the majority of the *Roseobacter* clade organisms fall into group I, but those of *P. temperata* RCA23 into group II. This group contains relatively few sequences of *Roseobacter* clade members but additionally includes strains of the SAR116 clade and of *Rhodobacter sphaeroides*. The existence of these two groups of flagella gene clusters obviously reflects the evolutionary history of the *Roseobacter* clade with substantial lateral gene transfer (Luo *et al.*, 2013). GI 5 harbours the genes for the group I CO dehydrogenase (*coxI*; Supplementary Table S4). The existence of the *coxI* gene cluster enables members of the *Roseobacter* clade to oxidize CO (Cunliffe, 2011).

The genome of *P. temperata* RCA23 encodes all basic metabolic functions. Carbohydrates are metabolized via the Entner–Doudoroff pathway that appears typical for the *Roseobacter* clade and also for the SAR11 clade (Giovannoni *et al.*, 2005; Fuerch

Table 1 Comparison of selected biosynthetic/catabolic genes and pathways of representative members of the *Roseobacter* clade

		<i>P. temperata</i> RCA23	<i>P. inhibens</i> DSM17395	<i>R. pomeroyi</i> DSS-3	<i>Sulfitobacter</i> sp. NAS-14.1	<i>Sulfitobacter</i> sp. GA1101	<i>R. litoralis</i> Och 149	<i>Rhodobacteraceae bacterium</i> HTCC2083	<i>Roseovarius</i> sp. TM1035	<i>Roseovarius</i> sp. 217	<i>S. stellata</i> E-37	<i>O. batsensis</i> HTCC2597	<i>L. vestfoldensis</i> SKA53	<i>O. arcticus</i> 238	<i>O. antarcticus</i> 307	<i>D. shibae</i> DFL-12	<i>Jannaschia</i> sp. CCS1	<i>M. alkaliphilus</i> HTCC2654	<i>Rhodobacterales bacterium</i> HTCC2150	<i>Rhodobacterales bacterium</i> HTCC2255
Trophic strategy	photosynthesis cluster	■					■	■	■	■		■				■				
	rhodopsin												■	■						■
	H ₂ oxidation								■		■									
	gene transfer agents		■	■	■	■		■	■	■	■	■	■	■	■	■	■	■	■	■
Aromatic degradation	B-ketoadipate	■	■	■	■	■		■	■	■	■	■	■			■	■	■	■	■
	gentisate pathway				■	■										■	■	■	■	■
	benzoate	■			■	■					■					■	■	■	■	■
	phenylacetic acid	■	■	■	■	■	■				■					■	■	■	■	■
	homoprotocatechuate	■	■	■	■	■	■				■					■	■	■	■	■
	homogentisate pathway				■	■	■	■				■				■	■	■	■	■
Carbon monoxide utilization	group I CO DH	■	■		■	■		■	■	■	■	■	■	■	■	■	■	■	■	■
	group II CO DH	■	■		■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
C1 compound utilization	C1 incorp (serine)	■	■					■	■	■	■					■	■	■	■	■
	MeOH oxidation					■		■	■	■	■					■	■	■	■	■
	TMA oxidation					■	■	■	■	■	■		■	■	■	■	■	■	■	■
	formaldehyde oxidation	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
	formate oxidation	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
C2 compound utilization	ethylmalonyl pathway	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
	glyoxylate shunt																			■
Motility, sensing, and attachment	chemotactic ability	■	■		■	■	■	■	■	■	■	■	■			■	■	■	■	■
	motility	■	■	■	■	■	■	■	■	■	■	■	■			■	■	■	■	■
	flp pilus (type IV)				■	■	■	■	■	■	■	■	■	■	■	■	■	■	■	■
	lux quorum sensing		■	■	■	■	■	■	■	■	■	■	■			■	■	■	■	■
	<i>virB</i> syst (type IV SS)		■		■			■	■	■	■	■	■	■	■	■	■	■	■	■
Secondary metabolite	antibiotic production (TDA)		■																	

Abbreviations: CO DH, carbon monoxide dehydrogenase; DMSP, dimethylsulphonium propionate; NRPS, non-ribosomal peptide synthase; PKS, polyketide synthetase; TDA, tropodithietic acid; TMA, trimethylamine. Colored boxes represent the presence of a gene or pathway within a genome. Data for *P. temperata* RCA23 were generated in this study; the data for representative members of the *Roseobacter* clade are taken from Newton *et al.* (2010) and colored according to the different subclasses in Figure 1.

the pelagic members of the *Roseobacter* clade (Supplementary Figure S4A and B; Supplementary Tables S2 and S3). A total of 208 transport proteins of *P. temperata* RCA23 belongs to the ATP-binding cassette (ABC) family (Supplementary Figure S4B and C; Supplementary Table S2). This number is lower than the mean of the pelagic members of the *Roseobacter* clade, whereas the proportion of ABC transporters of *P. temperata* RCA23 (72.5%) is very close to and the number of ABC transporters per Mb of *P. temperata* RCA23 (63.2) identical to the mean of all pelagic members (Supplementary Table S2). Interestingly, the other two deeply branching pelagic members of the *Roseobacter* clade, *Rhodobacterales* bacteria HTCC2255 and HTCC 2150, have the lowest numbers of ABC transporters per Mb of all roseobacters, 47.6 and 50.2, values close to that of *Cand. P. ubique*, 51.1 (Supplementary Table S2). The mean of the absolute numbers of ABC transporters of the pelagic members is significantly lower than that of the associated *Roseobacter* clade members (Student's *t*-test; $P=0.014$; Supplementary Table S3), presumably a result of the different evolutionary history of both groups of this clade with respect to gene acquisition and loss (Luo *et al.*, 2013). A high proportion of ABC transporters, which exhibit high substrate affinities, was interpreted as an adaptation of pelagic bacteria to the nutrient-poor oceanic environment (Giovannoni *et al.*, 2005; Lauro *et al.*, 2009). As all members of the *Roseobacter* clade exhibit a rather high percentage of ABC transporters, this feature may not be an adaptation to the nutrient-poor oceanic environment. It may also reflect the generally highly dynamic nutrient supply to members of this clade, considered as opportunistic marine bacteria and being able to adapt to a variety of nutrient conditions (Brinkhoff *et al.*, 2008; Teeling *et al.*, 2012; Luo *et al.*, 2013). The number of tripartite ATP-independent periplasmic transporters of *P. temperata* RCA23 is lower than the mean of the pelagic as well as associated members of the *Roseobacter* clade (Figure 3d and Supplementary Table S3). The great majority of tripartite ATP-independent periplasmic transporters of *P. temperata* RCA23 mediates uptake of dicarboxylic acids (Supplementary Table S4). Physiological tests showed that *P. temperata* RCA23 is able to grow on a large variety of organic substrates including amino acids, monosaccharides and short-chain fatty acids (Giebel *et al.*, 2013). This high versatility appears to reflect the high number of transporter proteins.

Even though quite a few genomic features are similar to those of other members of the *Roseobacter* clade, the genome of *P. temperata* RCA23 is distinct in lacking genes encoding the Flp pilus (type IV) for attachment, the VirB system for discharge of genetic material and protein, and genes encoding enzymes for quorum sensing and synthesizing other secondary metabolites (Table 1). The only other organism of the *Roseobacter* clade also missing the genes encoding attachment properties is *Rhodobacterales*

bacterium HTCC2255 (Table 1). In *Octadecabacter arcticus* these genes are present but fragmented into three partial clusters and not arranged homologously to the other organisms of this clade (Table 1; Vollmers *et al.*, 2013). The lacking feature for attachment in *P. temperata* RCA23 is consistent with the observation that the RCA cluster has not been detected in the fraction of particle-associated bacteria in the North Sea (Giebel *et al.*, 2011) and that growth on agar plates is weak and unreliable (Giebel *et al.*, 2013). This lacking attachment ability, however, is in contrast to the life style of another RCA cluster isolate, strain LE17, retrieved from Californian coastal waters that lives associated with a dinoflagellate (Mayali *et al.*, 2008).

In order to distinguish between genomic features of copiotrophic and oligotrophic bacteria Lauro *et al.* (2009) compared *Photobacterium angustum* S14 (copiotrophic; genome size 5.10 Mbp), and *Sphingopyxis alaskensis* RB 2256 (oligotrophic; genome size 3.37 Mbp). These authors found that categories of clusters of orthologous groups (COG) for motility (N), transcription (K), defense mechanisms (V) and signal transduction (T) constitute lower and COG categories for lipid transport and metabolism (I) and secondary metabolites, biosynthesis, transport and catabolism (Q) higher fractions in the oligotrophic as compared to the copiotrophic bacterium. The fractions of COG categories K, V and T in the genome of *P. temperata* RCA23 constitute even smaller fractions and COG categories I and Q higher fractions than in *S. alaskensis* (Supplementary Table S5), confirming the finding by Lauro *et al.* (2009) and emphasizing that *P. temperata* RCA23 is a truly oligotrophic bacterium in these respects. Interestingly, COG category I (lipid transport and metabolism) of the pelagic members of the *Roseobacter* clade constitutes a significantly higher fraction relative to the associated members (Supplementary Table S5) and *P. temperata* RCA23 has the third highest fraction of this COG category of all pelagic roseobacters (Supplementary Table S5). Genes affiliated to this COG category include short-chain fatty acid dehydrogenases, hydratases, dehydratases and acetyltransferases (Supplementary Table S4). The COG category of cell motility (N) of *P. temperata* RCA23 has a fraction not as low as that of *S. alaskensis*, suggesting that this feature is of adaptive significance for the life style of *P. temperata* RCA23 in the pelagic environment. The respective means of the associated and pelagic groups of the *Roseobacter* clade do not exhibit such a clear-cut difference between these COG categories and thus may reflect that members of this clade in general are more adapted to nutrient-poor environments than *P. angustum* (Supplementary Table S5).

Occurrence of the RCA cluster and *P. temperata* RCA23 in the North Sea

In order to assess the significance of the RCA cluster and *P. temperata* RCA23 among the active microbial




Figure 3 (a) Bathymetry of the southern North Sea (German Bight) and sampling stations in May 2010. Arrows indicate stations where the samples for the metagenomic and transcriptomic analyses were collected. (b) Chlorophyll *a*, bacterial abundance, salinity and temperature at 2 m depth. (c) Percentages of the RCA cluster as detected by CARD-FISH with an RCA-specific probe, of DNA-synthesizing RCA cells (BrdU positive) and of active RCA (cDNA). Samples in (b) and (c) are ordered as inside and outside the bloom except station 1, which was a separate bloom in the southernmost area. *: Missing data.

players in the southern North Sea, we investigated bacterioplankton abundance, composition and activity in and outside a diatom-dominated phytoplankton spring bloom in the German Bight (Figure 3). The RCA cluster dominated the *Roseobacter* clade by >90% (Wemheuer *et al.*, 2014), constituted 3.0–6.5% (mean = $5.1 \pm 1.2\%$) of total bacterioplankton and the active bacterioplankton community to even higher proportions, to 10–31.3% (mean = $18.5 \pm 7.7\%$), as determined by the cDNA derived from the 16S rRNA (Figure 3c; Wemheuer *et al.*, 2014). Between 11.9% and 49.6% (mean = $34.0 \pm 10.7\%$) of the RCA cells were actively dividing as determined by incorporation of bromodeoxyuridine (BrdU; Figure 3c).

One sample collected outside (station 3) and two samples collected inside the bloom, during day (1100 hours; station 13) and at night (0330 hours; station 9), were subjected to a metagenomic and metatranscriptomic analysis. The genome of *P. temperata* RCA23 was retrieved from the metagenomes to >96% and accounted for 7.8–15.4% of total reads (Table 2; Figure 4). It was also retrieved from the metatranscriptomes, to 17.3% outside and to at least 93.4% inside the bloom (Table 2; Figure 4). Inside the bloom it accounted for 6.7% of total metatranscriptomic reads during night and for 8% during day (Table 2). These data, in line with the other above-mentioned data, show that *P. temperata* RCA23 was an abundant and in the

bloom also highly active member of the bacterioplankton in the southern North Sea.

For comparison and to validate our genome mapping approach, we carried out similar genomic mappings in the metagenomes and transcriptomes for organisms of the SAR11 clade and SAR92 cluster, phylogenetic lineages also abundant at these stations (Wemheuer *et al.*, 2014). The genomes of *Cand. P. ubiquus* HTCC1062 of the SAR11 clade and of the Gammaproteobacterium HTCC2207 of the SAR92 cluster were retrieved to 92.7% and 44.6% in the metagenome outside the bloom, and to 50.2% and 95.6% inside the bloom, with some differences between day and night (Table 2). Both genomes accounted for not >1.7% of total metagenomic reads. The genome of *Cand. P. ubiquus* HTCC1062 had already been retrieved to 96% from metagenomic data of a North Sea phytoplankton spring bloom, but only from the pooled data of all combined samples (Teeling *et al.*, 2012). We retrieved the genomes of *Cand. P. ubiquus* HTCC1062 and of Gammaproteobacterium HTCC2207 also from metatranscriptomic reads, in the bloom sample collected during the day to 40.9% and 89.1%, respectively, and at night to 42.6% and 34.1%, respectively. These data show little variations between day and night for the transcriptome of *Cand. P. ubiquus* HTCC1062, but a much lower fraction of the genome of HTCC2207 transcribed at night. The transcriptomic reads of both organisms

Table 2 Days of sampling, mapped reads, percentage of genomic coverage and percentage of total reads of *P. temperata* RCA23, *Cand. Pelagibacter ubique* HTCC1062 of the SAR11 clade and of Gammaproteobacterium HTCC2207 of the SAR92 cluster of the metagenome (DNA) and metatranscriptome (RNA) retrieved at stations outside (non-bloom) and inside the phytoplankton spring bloom during day (1100 hours) and night (0330 hours)

Station	Target	No. of reads	Organism (lineage)	Mapped reads	Percentage of genome coverage	Percentage of total reads
3						
Non-bloom 26 May 2010	DNA	23 381 812	<i>P. temperata</i> RCA23	1 819 024	97.1	7.8
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	407 611	92.7	1.7
			HTCC2207 (SAR92)	50 361	44.6	0.2
	RNA	2 223 804	<i>P. temperata</i> RCA23	19 435	17.3	0.9
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	16 947	17.9	0.8
			HTCC2207 (SAR92)	4021	2.6	0.2
9						
bloom, night 29 May 2010	DNA	12 671 944	<i>P. temperata</i> RCA23	1 783 951	96.4	14.1
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	112 728	89.0	0.9
			HTCC2207 (SAR92)	45 926	50.2	0.4
	RNA	8 101 537	<i>P. temperata</i> RCA23	543 596	93.4	6.7
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	35 853	42.6	0.4
			HTCC2207 (SAR92)	35 853	34.1	0.3
13						
bloom, day 29 May 2010	DNA	16 156 091	<i>P. temperata</i> RCA23	2 480 608	96.0	15.4
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	23 806	66.3	0.1
			HTCC2207 (SAR92)	238 054	95.5	1.5
	RNA	15 349 574	<i>P. temperata</i> RCA23	1 222 858	94.6	8.0
			<i>Cand. P. ubique</i> HTCC1062 (SAR11)	27 026	40.9	0.2
			HTCC2207 (SAR92)	179 429	89.1	1.2
GOS02-13 Norwegian fjord	DNA	1 083 836	<i>P. temperata</i> RCA23	12 254	81.9	1.1
	DNA	863 687	<i>P. temperata</i> RCA23	54 895	83.9	6.4
	RNA	256 537		4176	23.7	1.6

For exact locations of the stations see Figure 3a.

accounted for not >1.2% of total metatranscriptomic reads, indicating that these organisms were far less abundant and active than *P. temperata* RCA23.

The fact that at least 93% of the genome of *P. temperata* RCA23 were transcribed in the bloom is remarkable because it reflects that the population of this abundant organism was highly active and used basically the entire potential of its metabolic properties. The largest fraction of the remaining unmapped regions refers to hypothetical proteins mostly located in GIs (Figure 4). To the best of our knowledge this is the first report of such a high fraction of the genome of an abundant marine bacterium transcribed and retrieved from a metatranscriptome. A similarly high genome coverage in a metatranscriptome analysis was recently found in a sediment-derived microbial community degrading hexadecane under anoxic conditions. The genome of *Smithella spec.*, largely dominating this community, was transcribed to 94% (Embree *et al.*, 2014). Transcribed genomic fractions of 52–95% have been reported from pure culture experiments with various bacteria such as *Helicobacter pylori* (Sharma *et al.*, 2010), *Bacillus anthracis* (Passalacqua *et al.*, 2009), *Bacillus subtilis* (Nicolas *et al.*, 2012) and *Prochlorococcus MED4* (Wang *et al.*, 2014). Hence, the high fraction of the genome of *P. temperata* RCA23 transcribed is in line with these reports but may indicate that the ambient population of this organism consisted of cells in different metabolic

stages with respect to growth and substrate utilization.

A more detailed analysis of the transcriptomic patterns of *P. temperata* RCA23 shows that transcripts of COG category E (amino-acid transport and metabolism) exhibited the highest NPKM-normalized reads of all COG categories, followed by COG category R (general function) and J (translation, ribosomal structure and biogenesis) (Table 3, Supplementary Table S6). The high values of normalized transcripts of amino-acid and carbohydrate transport and metabolism (categories E and G) appear to reflect the significance of these substrates for growth of *P. temperata* RCA23, and is consistent with the broad substrate spectrum of this organism (Giebel *et al.*, 2013). Pronounced differences between the day and night transcriptome occurred in COG categories for translation, ribosomal structure and biogenesis (category J) and for cell motility, but not for chemotaxis (both category N), with higher fractions at night than at day (Supplementary Table S6). Further, the genes c29660, c29670 and c29760, encoding the light-harvesting protein B-870 beta chain PufB and alpha chain PufA and cytochrome c551, respectively, were highly overexpressed at night (Supplementary Figure S5 and Supplementary Table S6). This observation is in line with the well-known fact that bacteriochlorophyll *a* is only produced during night in AAP bacteria (Wagner-Döbler and Biebl, 2006).




Figure 4 Circular plot of metagenome and transcriptome read mappings onto the genome of *P. temperata* RCA23. Read counts from the Bowtie 2 mappings are shown. From innermost to outer circle: GC content, CDS counts, stations 3, 9 and 13 (DNA and cDNA), unmappable regions and genomic islands.

Genes encoding small heat-shock protein IbpA (c1240, c20690), RNA polymerase σ -32 factors RpoH1 and RpoH2 (c9960, c21310), chaperonin GroEL (c20740), GroS (c20750) and chaperone protein DnaK (c30090) were also highly overexpressed at night. The small heat-shock protein DnaK and chaperonin GroS, GroEL are important in preventing stress-induced aggregation of various proteins in the cytosol of bacteria (Mogk *et al.*, 2003; Stolyar *et al.*, 2007; Ting *et al.*, 2010). Hence, *P. temperata* RCA23 obviously undergoes some stress during night. So far we have no clue about a more specific type of stress that upregulates the transcription of these genes during night.

As a conclusion from this analysis the population of *P. temperata* RCA23 undergoes intense metabolic reconstruction during night including protein synthesis and stress response, but also enhances the synthesis of flagella proteins. This intense metabolic reconstruction may be a result of the different modes of energy conservation of this AAP

bacterium during day and night. It has been shown that *Roseobacter litoralis*, another AAP bacterium of the *Roseobacter* clade, changes its proteome drastically, mainly by downregulating protein synthesis, when shifted from dark to light conditions (Zong and Zhao, 2012). Hence the mode of energy conservation appears to have a pronounced effect on the global regulation of metabolic networks in AAP bacteria of the *Roseobacter* clade, and presumably beyond, with implications of their diurnal participation in turnover of organic matter in marine ecosystems.

Global and biogeochemical significance of the RCA cluster and P. temperata RCA23

Because of the known global distribution of the RCA cluster (Selje *et al.*, 2004) we examined publicly available metagenomic data sets for the presence of genomic features of *P. temperata* RCA23. The photosynthetic operon was present in an

Table 3 NPKM-normalized reads in the COG categories with type of metabolism of the genome of *P. temperata* RCA23 retrieved in the metatranscriptome at stations 9 and 13 in the German Bight of the North Sea

COG category	Type of metabolism	Station 9 (night) NPKM	Station 13 (day) NPKM
B	Chromatin structure and dynamics	661	954
C	Energy production and conversion	61 408	57 017
D	Cell cycle control, cell division and chromosome partitioning	7190	7533
E	Amino acid transport and metabolism	101 802	107 628
F	Nucleotide transport and metabolism	24 455	26 003
G	Carbohydrate transport and metabolism	59 872	59 448
H	Coenzyme transport and metabolism	36 820	42 486
I	Lipid transport and metabolism	45 455	49 575
J	Translation, ribosomal structure and biogenesis	71 558	59 835
K	Transcription	44 710	41 837
L	Replication, recombination and repair	36 761	39 159
M	Cell wall/membrane/envelope biogenesis	37 299	39 404
N	Cell motility	8425	4975
O	Post-translational modification, protein turnover and chaperones	50 598	45 863
P	Inorganic ion transport and metabolism	32 016	32 078
Q	Secondary metabolites biosynthesis, transport and catabolism	16 285	16 089
R	General function prediction only	84 440	90 432
S	Function unknown	68 057	68 212
T	Signal transduction mechanisms	20 797	19 875
U	Intracellular trafficking, secretion and vesicular transport	7431	7254
V	Defense mechanisms	8258	8799

Abbreviations: COG, clusters of orthologous groups; NPKM, Nucleotide activity Per Kilobase of exon model per million Mapped reads.

environmental bacterial artificial chromosome clone derived from Californian coastal waters (Supplementary Figure S3; Béjà *et al.*, 2002; Yutin and Béjà, 2005) and in metagenomic data sets from a Norwegian fjord, the western English Channel, the western coastal Atlantic visited during the Global Ocean Sampling expedition (GOS), and in the eastern (Monterey Bay) and southwestern Pacific in Australian waters (Rusch *et al.*, 2007; Yutin *et al.*, 2007; Gilbert *et al.*, 2008, 2010; Thomas *et al.*, 2010; Rich *et al.*, 2011).

The genome of *P. temperata* RCA23 was retrieved from metagenomic and metatranscriptomic data of a Norwegian fjord to 83.9% and 23.7%, respectively (Gilbert *et al.*, 2008; Table 2). At the GOS stations, the genome of *P. temperata* RCA23 was retrieved from the metagenomic data to 81.9% (Table 2) and accounted for 0.7% to 6.5% of the mapped reads (Supplementary Figure S6B). This finding, together with the observation that the structure of the photosynthetic operon of the AAP bacteria at these stations was similar to that of *P. temperata* RCA23 (see above), is consistent with the assumption that this organism constituted largely the AAP communities at these stations. In the GOS and the Norwegian fjord data sets, 80% and 81%, respectively, of the genes of *P. temperata* RCA23 showed a coverage of $\geq 80\%$. In contrast, 62.5% of the genes of the GIs were not mapped or showed a low coverage (Supplementary Table S7).

Conclusion

This study, on the basis of the sequenced genome of *P. temperata* RCA23 and the North Sea metagenome

and metatranscriptome, sheds new light on the significance of the RCA cluster for biogeochemical processes in marine pelagic systems, considering the high abundances of this organism and the entire RCA cluster in temperate to polar oceans (Selje *et al.*, 2004; West *et al.*, 2008; Giebel *et al.*, 2009, 2011; Wemheuer *et al.*, 2014). It appears to carry out a life style well adapted to the nutrient- and energy-poor marine pelagic realm. The entire population of this organism, and possibly even single cells, simultaneously transcribe almost the entire genome during phytoplankton bloom situations and convert complementary energy by harvesting light. This versatile physiology together with the wealth of high-affinity transporter proteins, despite the absence of distinct genomic and metabolic traits typical for the *Roseobacter* clade, and a streamlined genome relative to other *Roseobacter* clade organisms, helps to explain the success of this abundant taxon in temperate and presumably also (sub)polar marine systems. Our findings illustrate how important it is to dissect the role of individual taxa for better understanding their participation in processing organic and inorganic matter in pelagic marine systems.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

We thank Renate Gahl-Janssen for growing *P. temperata* RCA23 and harvesting biomass for DNA extraction, Melissa Büngener for technical assistance in sequence analyses, the crew of RV Heincke and M Wolterink for

helping with sample collection and processing during the North Sea cruise and the prophage induction experiment. This work was supported by grants from the Volkswagen Foundation in Lower Saxony, Germany, and by TRR51 of Deutsche Forschungsgemeinschaft.

References

- Béjà O, Suzuki MT, Heidelberg JF, Nelson WC, Preston CM, Hamada T *et al.* (2002). Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature* **415**: 630–633.
- Biers EJ, Wang K, Pennington C, Belas R, Chen F, Moran MA. (2008). Occurrence and expression of gene transfer agent (GTA) genes in marine bacterioplankton. *Appl Environ Microbiol* **74**: 2933–2939.
- Bolger AM, Lohse M, Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*; e-pub ahead of print 28 April 2014; doi:10.1093/bioinformatics/btu170.
- Brinkhoff T, Giebel HA, Simon M. (2008). Diversity, ecology, and genomics of the Roseobacter clade: a short overview. *Arch Microbiol* **189**: 531–539.
- Cunliffe M. (2011). Correlating carbon monoxide oxidation with *cox* genes in the abundant Marine Roseobacter Clade. *ISME J* **5**: 685–691.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. (2007). Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* **23**: 673–679.
- Embree M, Nagarajan H, Movahedi N, Chitsaz H, Zengler K. (2014). Single-cell genome and metatranscriptome sequencing reveal metabolic interactions of an alkane-degrading methanogenic community. *ISME J* **8**: 757–767.
- Fuerch T, Preusse M, Tomasch J, Zech H, Wagner-Döbler I, Rabus R *et al.* (2009). Metabolic fluxes in the central carbon metabolism of *Dinoroseobacter shibae* and *Phaeobacter gallaeciensis*, two members of the marine Roseobacter clade. *BMC Microbiol* **9**: 209.
- Giebel HA, Brinkhoff T, Zwisler W, Selje N, Simon M. (2009). Distribution of Roseobacter RCA and SAR11 lineages and distinct bacterial communities from the subtropics to the Southern Ocean. *Environ Microbiol* **11**: 2164–2178.
- Giebel HA, Kalhoefer D, Lemke A, Thole S, Gahl-Janssen R, Simon M *et al.* (2011). Distribution of Roseobacter RCA and SAR11 lineages in the North Sea and characteristics of an abundant RCA isolate. *ISME J* **5**: 8–19.
- Giebel HA, Kalhoefer D, Gahl-Janssen R, Choo YJ, Lee K, Cho JC *et al.* (2013). *Planktomarina temperata* gen. nov., sp. nov., belonging to the globally distributed RCA cluster of the marine Roseobacter clade, isolated from the German Wadden Sea. *Int J Syst Evol Microbiol* **63**: 4207–4217.
- Gilbert JA, Field D, Huang Y, Edwards R, Weizhong L, Gilna R *et al.* (2008). Detection of large numbers of novel sequences in the metatranscriptomes of complex marine microbial communities. *PLoS One* **3**: e3042.
- Gilbert JA, Field D, Swift P, Thomas S, Cummings D, Temperton B *et al.* (2010). The taxonomic and functional diversity of microbes at a temperate coastal site: a ‘multi-omic’ study of seasonal and diel temporal variation. *PLoS One* **5**: e15545.
- Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D *et al.* (2005). Genome streamlining in a cosmopolitan oceanic bacterium. *Science* **309**: 1242–1245.
- Giovannoni SJ, Stingl U. (2007). The importance of culturing bacterioplankton in the ‘omics’ age. *Nat Rev Microbiol* **5**: 820–826.
- Giovannoni SJ, Stingl U. (2005). Molecular diversity and ecology of microbial plankton. *Nature* **437**: 343–348.
- Giovannoni SJ, Cameron Thrash J, Temperton B. (2014). Implications of streamlining theory for microbial ecology. *ISME J* **8**: 1553–1565.
- Kopylova E, Noé L, Touzet H. (2012). SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**: 3211–3217.
- Langmead B, Salzberg S. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.
- Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S *et al.* (2009). The genomic basis of trophic strategy in marine bacteria. *Proc Natl Acad Sci USA* **106**: 15527–15533.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar *et al.* (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–1371.
- Luo H, Csu M, Hughes A, Moran MA. (2013). Evolution of divergent life history strategies in marine Alphaproteobacteria. *mbio* **4**: e00373–13.
- Luo H, Swan BK, Stepanauskas R, Hughes A, Moran MA. (2014). Evolutionary analysis of a streamlined lineage of surface ocean Roseobacters. *ISME J* **8**: 1428–1439.
- Mayali X, Franks PS, Azam F. (2008). Cultivation and ecosystem role of a marine Roseobacter clade-affiliated cluster bacterium. *Appl Environ Microbiol* **74**: 2595–2603.
- Mogk A, Deuerling E, Vorderwülbecke S, Vierling E, Bukau B. (2003). Small heat shock proteins, ClpB and the DnaK system form a functional triade in reversing protein aggregation. *Mol Microbiol* **50**: 585–595.
- Newton RJ, Griffin LE, Bowles KM, Meile C, Gifford S, Givens CE *et al.* (2010). Genome characteristics of a generalist marine bacterial lineage. *ISME J* **4**: 784–798.
- Nicolas P, Mäder U, Dervyn E, Rochat T, Léduc A *et al.* (2012). Condition-dependent transcriptome reveals high-level regulatory architecture in *Bacillus subtilis*. *Science* **335**: 1103–1106.
- Passalacqua KD, Varadarajan A, Ondov BD, Okou DT, Zwick ME, Bergman NH. (2009). Structure and complexity of a bacterial transcriptome. *J Bacteriol* **191**: 3203–3211.
- Pernthaler A, Pernthaler J, Schattenhofer M, Amann R. (2002). Identification of DNA-synthesizing bacterial cells in coastal North Sea plankton. *Appl Environ Microbiol* **68**: 5728–5736.
- Pradella S, Päucker O, Petersen J. (2010). Genome organization of the marine Roseobacter clade member *Marinovum algicola*. *Arch Microbiol* **192**: 115–126.
- Rich VI, Pham VD, Eppley JM, Shi Y, DeLong EF. (2011). Time-series analyses of Monterey Bay coastal microbial picoplankton using a ‘genome proxy’ microarray. *Environ Microbiol* **13**: 116–134.
- Rusch DB, Halpern A, Sutton G, Heidelberg KB, Williamson S, Yooseph S *et al.* (2007). The Sorcerer II Global Ocean Sampling expedition: northwest Atlantic through eastern tropical Pacific. *PLoS Biol* **5**: e77.

- Selje N, Simon M, Brinkhoff T. (2004). A newly discovered Roseobacter cluster in temperate and polar oceans. *Nature* **427**: 445–448.
- Sharma CM, Hoffmann S, Darfeuille F, Reignier J, Findeiss S, Sittka A *et al.* (2010). The primary transcriptome of the major human pathogen *Helicobacter pylori*. *Nature* **460**: 250–254.
- Sowell SM, Wilhelm LJ, Norbeck AD, Lipton LS, Nicora CD, Barofsky DF *et al.* (2009). Transport functions dominate the SAR11 metaproteome at low-nutrient extremes in the Sargasso Sea. *ISME J* **3**: 93–105.
- Staden R. (1996). The Staden sequence analysis package. *Mol Biotech* **5**: 233–241.
- Stolyar S, He Q, Joachimiak MP, He ZL, Yang ZK, Borglin SE *et al.* (2007). Response of *Desulfovibrio vulgaris* to alkaline stress. *J Bacteriol* **189**: 8944–8952.
- Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, González JM *et al.* (2013). Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in the surface ocean. *Proc Natl Acad Sci USA* **110**: 11463–11468.
- Tech M, Merkl R. (2003). YACOP: Enhanced gene prediction obtained by a combination of existing methods. *In Silico Biology* **3**: 441–451.
- Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM *et al.* (2012). Substrate-controlled succession of marine bacterioplankton populations induced by a phytoplankton bloom. *Science* **336**: 608–611.
- Thomas T, Rusch D, DeMaere MZ, Yung PY, Lewis M, Halpern A *et al.* (2010). Functional genomic signatures of sponge bacteria reveal unique and shared features of symbiosis. *ISME J* **4**: 1557–1567.
- Thompson AW, Huang K, Saito MA, Chisholm SW. (2011). Transcriptome response of high- and low-light-adapted *Prochlorococcus* strains to changing iron availability. *ISME J* **5**: 1580–1594.
- Ting L, Williams TJ, Cowley MJ, Lauro FM, Guilhaus M, Raftery MJ *et al.* (2010). Cold adaptation in the marine bacterium, *Sphingopyxis alaskensis*, assessed using quantitative proteomics. *Environ Microbiol* **12**: 2658–2676.
- Tripp HJ, Kitner JB, Schwalbach MS, Dacey JWH, Wilhelm LJ, Giovannoni SJ. (2008). SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature* **452**: 741–744.
- Vollmers J, Voget S, Dietrich S, Gollnow K, Smits M, Meyer K *et al.* (2013). Poles apart: extreme genome plasticity and a new xanthorhodopsin-like gene family in the genomes of *Octadecabacter arcticus* 238 and *Octadecabacter antarcticus* 307. *PLoS One* **8**: e63422.
- Wagner-Döbler I, Biebl H. (2006). Environmental biology of the marine Roseobacter lineage. *Ann Rev Microbiol* **60**: 255–280.
- Wang B, Lu L, Lv H, Jiang H, Qu G, Tian C *et al.* (2014). The transcriptome landscape of *Prochlorococcus* MED4 and the factors for stabilizing the core genome. *BMC Microbiol* **14**: 11.
- Wemheuer B, Güllert S, Billerbeck S, Giebel H-A, Voget S, Simon M *et al.* (2014). Impact of a phytoplankton bloom on the diversity of the active bacterial community in the southern North Sea as revealed by metatranscriptomic approaches. *FEMS Microbiol Ecol* **87**: 378–389.
- West NJ, Obernosterer I, Zemb Lebaron P. (2008). Major differences of bacterial diversity and activity inside and outside of a natural iron-fertilized phytoplankton bloom in the Southern Ocean. *Environ Microbiol* **10**: 738–756.
- Wiegand S, Dietrich S, Hertel R, Bongaerts J, Evers S, Volland S *et al.* (2013). RNA-Seq of *Bacillus licheniformis*: active regulatory RNA features expressed within a productive fermentation. *BMC Genomics* **14**: 667.
- Yooshef S, Nealson KH, Rusch DB, McCrown JP, Dupont CL, Kim M *et al.* (2010). Genomic and functional adaptation in surface ocean planktonic prokaryotes. *Nature* **468**: 60–67.
- Yutin N, Béjà O. (2005). Putative novel photosynthetic reaction centre organizations in marine aerobic anoxygenic photosynthetic bacteria: insights from metagenomics and environmental genomics. *Environ Microbiol* **7**: 2027–2033.
- Yutin N, Suzuki MT, Teeling H, Weber M, Venter JC, Rusch DB *et al.* (2007). Assessing diversity and biogeography of aerobic anoxygenic phototrophic bacteria in surface waters of the Atlantic and Pacific Oceans using the Global Ocean Sampling expedition metagenomes. *Environ Microbiol* **9**: 1464–1475.
- Zhao Y, Wang K, Budinoff C, Buchan A, Lang A, Jiao N *et al.* (2009). Gene transfer agent (GTA) genes reveal diverse and dynamic Roseobacter and Rhodospirillum populations in the Chesapeake Bay. *ISME J* **3**: 364–373.
- Zong R, Zhao N. (2012). Proteomic responses of *Roseobacter litoralis* OCh149 to starvation and light regimen. *Microbes Environ* **27**: 430–442.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)