



HHS Public Access

Author manuscript

Peptides. Author manuscript; available in PMC 2016 June 01.

Published in final edited form as:

Peptides. 2015 June ; 68: 140–147. doi:10.1016/j.peptides.2014.07.015.

Aedes* cadherin mediates the *in vivo* toxicity of the Cry11Aa toxin to *Aedes aegypti

Su-Bum Lee^{#1}, Jianwu Chen^{#2}, Karlygash G. Aimanova², and Sarjeet S. Gill^{1,2,*}

¹Environmental Toxicology graduate program, University of California, Riverside, CA, 92521, USA

²Department of Cell Biology and Neuroscience, University of California, Riverside, CA, 92521, USA

These authors contributed equally to this work.

Abstract

Cadherin plays an important role in the toxicity of *Bacillus thuringiensis* Cry proteins. We previously cloned a full-length cadherin from *Aedes aegypti* larvae and reported this protein binds Cry11Aa toxin from *B. thuringiensis* subsp. *israelensis* with high affinity, ≈ 16.7 nM. Based on these results, we investigated if *Aedes* cadherin is involved in the *in vivo* toxicity of Cry11Aa toxin to *Ae. aegypti*. We established a mosquito cell line stably expressing the full-length *Aedes* cadherin and transgenic mosquitoes with silenced *Aedes* cadherin expression. Cells expressing the *Aedes* cadherin showed increased sensitivity to Cry11Aa toxin. Cry11Aa toxin at 400 nM killed approximately 37% of the cells in 3 h. Otherwise, transgenic mosquitoes with silenced *Aedes* cadherin expression showed increased tolerance to Cry11Aa toxin. Furthermore, cells expressing *Aedes* cadherin triggered Cry11Aa oligomerization. These results show the *Aedes* cadherin plays a pivotal role in Cry11Aa toxicity to *Ae. aegypti* larvae by mediating Cry11Aa oligomerization. However, since high toxicity was not obtained in cadherin-expressing cells, an additional receptor may be needed for manifestation of full toxicity. Moreover, cells expressing *Aedes* cadherin were sensitive to Cry4Aa and Cry11Ba but not Cry4Ba. However transgenic mosquitoes with silenced *Aedes* cadherin expression showed no tolerance to Cry4Aa, Cry4Ba, and Cry11Ba toxins. These results suggest that while *Aedes* cadherin may mediate Cry4Aa and Cry11Ba toxicity, this cadherin but is not the main receptor of Cry4Aa, Cry4Ba and Cry11Ba toxin in *Ae. aegypti*.

Keywords

Bacillus thuringiensis; Cry11Aa toxin; cadherin; C6/36 cell; transgenic mosquito; oligomerization; cytotoxicity

© 2014 Elsevier Inc. All rights reserved.

*Corresponding author: sarjeet.gill@ucr.edu. Tel: 951-827-4621/3547.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Unlike other members of the *Bacillus cereus* group, *B. thuringiensis* is pathogenic to insects by producing insecticidal proteins, which consists of one or more proteins, called Cry or Cyt toxins [21]. As these proteins are highly selective to the target insect, and harmless to humans and vertebrates, this species has been used for the control of insect pests in agriculture and public health [14].

One subsp., *B. thuringiensis* subsp. *israelensis* (Bti), has been used for the control of insect vectors of human diseases, including *Simulium damnosum*, a vector of onchocerciasis, and *Aedes* and *Culex* mosquito species, that can be vectors for dengue fever, chikungunya and yellow fever, and filariasis and West Nile fever, respectively [33]. These control programs are possible because Bti, while having high insecticidal activity, has low toxicity to non-target organisms. Consequently, it is an important alternative and environmental-friendly method for control of mosquito and black fly populations.

While Bti has high activity, its mechanism of action is still poorly understood. The bacterium has a megaplasmid, pBtoxis, which encodes a number of toxins (Cry4Aa, Cry4Ba, Cry10Aa, Cry11Aa, Cyt1Aa, Cyt1Ca and Cyt2Ba) [4]. Among them, Cry11Aa is one of the more active toxins towards *Ae. aegypti* [11]. Fernandez et al. [19] reported domain II of Cry11Aa is important in receptor recognition and binding. Domain II contains four putative loop regions, α -8, 1, 2 and 3. Using competitive binding assays, peptide-displaying phages and mutagenesis, it was revealed that loop α -8 in Cry11Aa was involved in toxicity and receptor binding.

Many putative Cry toxin receptors have been identified in mosquitoes [31]. An aminopeptidase N (APN) from *Anopheles quadrimaculatus* bound Cry11Ba, and a cadherin receptor from *An. gambiae* was identified and bound Cry4Ba [1, 26, 39]. In *Ae. aegypti*, Cry11Aa bound four proteins (200, 100, 65 and 62 kDa) in brush border membrane vesicles (BBMV) isolated from *Ae. aegypti* midgut epithelia [18]. Among them, the 65 kDa protein was identified as a glycosylphosphatidylinositol (GPI)-anchored alkaline phosphatase (ALP) and was a functional receptor of Cry11Aa toxin in *Ae. aegypti* midgut cells. The 100 kDa protein was identified as an APN and two of these have been characterized [9, 10]. In a previous study, we showed that the *Aedes* cadherin (AAEL007478 and AAEL007488), which is homologous to the lepidopteran Bt-R₁ and mediates Cry1A toxicity in Lepidoptera, bound Cry11Aa with high affinity [8]. This finding suggests that the cadherin is associated with the insecticidal activity of the Cry11Aa toxin.

Based on these results, we investigated further whether *Aedes* cadherin mediates Cry11Aa toxicity *in vivo*. We established a cell line expressing *Aedes* cadherin and a transgenic mosquito line that silences *Aedes* cadherin. We determined the ability of the Cry11Aa protein to cause cytotoxicity with cells expressing *Aedes* cadherin, and showed that the transgenic mosquitoes showed increased tolerance to Cry11Aa toxicity.

2. Materials and Methods

2.1 Cell culture

C6/36 (*Aedes albopictus*) cells were grown and maintained in L15 medium (Gibco, Grand Island, NY) supplemented with 10% FBS (Gibco), 1% L-glutamine (Gibco) and 1% penicillin-streptomycin (Gibco) at 27°C. The cells were grown as a monolayer in T-25 culture flasks (BD Falcon, Franklin Lakes, NJ), 6-well tissue culture plates, or 96-well tissue culture plates (Corning, Tewksbury, MA).

2.2 Construction of cadherin in pACTIN.SV

The pACTIN.SV and pIE1.SV vectors used, including enhanced GFP gene ORF (*EGFP*) (Fig. 1B), were obtained from Drs. Huynh and Zieler (National Institutes of Health, Bethesda, MD) [27].

A full-length *Aedes* cadherin cDNA (AaeCad) cloned into pCR2.1 vector was previously reported [8]. To remove the 5' and 3' UTRs, partial AaeCad fragments (5EM and 3EM) were prepared with a set of primers (Supplementary Table S1). The 5' end modified fragment (5EM) was amplified using a sense primer (5EM-S), which contained the restriction enzyme sites NotI and StuI, a Kozak sequence (CCACC) and a start codon, and an antisense primer (5EM-A) with a restriction enzyme site, Bstz17I. The 3' end modified fragment (3EM) was amplified using a sense primer (3EM-S), containing a BlnI restriction enzyme site and an antisense primer (3EMA), which had a HA-tag (TACCCATACGACGTCCTCCAGACTACGCT), a stop codon, and the restriction sites NheI, PmeI and SacI. All PCR products were cloned into the pCR2.1 vector (Invitrogen, Grand Island, NY) and fully sequenced (Institute for Integrative Genome Biology, University of California, Riverside, CA). To construct the modified AaeCad, the 5EM and AaeCad were each digested with NotI and Bstz17I (New England Biolabs, Ipswich, MA), the fragments then gel-purified and ligated. Next, this construct and the 3EM were digested with BlnI and SacI, gel-purified, and ligated. To construct an expression vector, the modified AaeCad in pCR2.1 vector and pACTIN.SV vector were separately digested with XbaI and SacI, gel-purified and ligated (Fig. 1A).

The blasticidin-resistance gene (*bla*) from pCoBlast vector (Invitrogen) was amplified using a sense primer (BLA-S), with a restriction enzyme site (BglII), a Kozak sequence, and a start codon, and an antisense primer (BLA-A), which had a stop codon and a restriction enzyme site (PmlI). To construct an antibiotic selection vector to be used for co-expression, the *Bla* gene and pIE1.SV were separately digested with BglII and PmlI, gel-purified, and then ligated (Fig. 1C).

2.3 Cell transfection

To construct a stable cell line expressing *Aedes* cadherin, pACTIN.SV containing the cloned AaeCad (5 µg) together with pIE1.SV containing a blasticidin-resistance gene (1 µg) for selection, were co-transfected into C6/36 cells using the FuGENE6 transfection reagent (Roche Applied Science, Madison, WI) following the manufacturer's protocol. Another stable C6/36 cell line expressing GFP was made using pACTIN.SV containing an EGFP

gene ORF (5 µg) together with pIE1.SV containing a blasticidin-resistance gene (1 µg), using identical conditions. In brief, 1.5×10^5 C6/36 cells/well in a 6-well plate were incubated overnight at 27°C. To these cells the plasmid transfection mixtures in FuGENE 6 transfection reagent were added and the cells incubated for an additional 3 days at 27°C. Media was removed and replaced with fresh medium containing blasticidin (15 µg), and a week later, the media was replaced with media with additional blasticidin (7.5 µg). The selecting media was replaced every 3-4 days until cell colonies were observed. Colonies were picked, diluted, and transferred to 96-well plates in order to obtain a single cell in each well. These single cells were grown and maintained in the fresh medium containing blasticidin (7.5 µg) at 27°C. Homogeneous cell lines were analyzed by western blotting to find a cell line highly expressing AaeCad.

2.4 Western blotting with cells

Cells expressing AaeCad or EGFP were harvested, washed with phosphate buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 2 mM KH₂PO₄, pH7.4) twice and resuspended in SDS-PAGE sample buffer. The collected cells were boiled for 10 min and centrifuged at 10,000×g for 10 min to remove insoluble material. The supernatants were loaded in SDS polyacrylamide gel (10%), and then electrotransferred to nitrocellulose membrane. The membrane was blocked with blocking solution (PBS, 5% skim milk and 0.1% Tween-20) for 1 h at room temperature, and washed with PBST (PBS and 0.1% Tween-20). The blocked membrane was incubated with an anti-HA antibody (a rabbit polyclonal IgG, Santa Cruz Biotechnology, Dallas, TX) or an anti-cadherin fragment antibody detecting AaeCad CR7-11 [8] (1:3,000) overnight at 4°C. The membrane was washed with PBST, and then subsequently incubated with anti-rabbit horseradish peroxidase (HRP, 1:5,000) secondary antibody (Sigma, St. Louis, MO) for 90 min at room temperature. After washing with PBST, the HRP activity was revealed with a luminal substrate (Thermo Scientific, Lafayette, CO) and exposed to an X-ray film in a darkroom.

2.5 Immunolocalization of AaeCad

C6/36 cells (1×10^5 cells) were plated in a slide chamber (Fisher, Hampton, NH) and incubated overnight at 27°C, then washed three times with PBS, and then fixed in 4% paraformaldehyde in PBS for 1 h at room temperature. The fixed cells were treated with a blocking solution (PBS, 2% BSA) for 1 h at room temperature. The cells were then incubated with anti-cadherin fragment antibody (1:200 dilution) in PBS containing 1% BSA for 1 h at room temperature, washed three times with washing buffer (PBS, 0.1% BSA, and 2% goat serum), and incubated with Cy3-conjugated goat anti-rabbit IgG (1:1000, Jackson Immuno Research, West Grove, PA) in PBS containing 0.1% BSA and 2% goat serum for 1 h at room temperature in dark condition. The fluorescence was observed using Zeiss 510 confocal microscope (Institute for Integrative Genome Biology).

2.6 Purification of Cry toxins

For crude Cry toxin production, *B. thuringiensis* strains producing Cry4Aa, Cry4Ba, Cry11Aa, or Cry11Ba [7, 15], were grown in nutrient broth sporulation medium containing erythromycin (25 µg/ml) at 30°C for 4-5 days for cell autolysis to occur [30]. Spores and

crystal inclusions were harvested 10,000xg for 10 min at 4°C, washed twice with sterilized water, and stored in water at –80°C until used.

For purified Cry toxins, the inclusion bodies for Cry toxins were isolated as previous reported [13]. The purified inclusions were washed, solubilized in 50 mM Na₂CO₃ pH 10.5 buffer, and then activated by trypsin (1:20 w/w) at 37°C. Solubilized Cry toxins were quantified using the BCA assay (Pierce, Rockford, IL). Activated Cry toxins were filtered using 0.2 µm syringe filter and stored at –80°C until needed.

2.7 Cytotoxicity test with Cry toxins

The toxicity of Cry toxins to C6/36 cells expressing AaeCad or EGFP was analyzed by using a MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) cytotoxicity assay [12]. In brief, 6×10^4 cells were plated in each well of a 96-well cell culture dish and incubated overnight at 27°C. The plate was centrifuged at 1,000xg for 5 min. The medium was removed and replaced with the fresh medium containing the activated Cry toxins (final concentration 50 – 400 nM) and the cells were incubated for 3 h at 27°C. After centrifugation at 1,000xg for 5 min, the medium was removed and replaced with a mixture of 200 µl fresh medium and 20 µl MTT solution (5 mg/ml MTT in PBS), and the cells were incubated for 2 h at 27°C. The plate was centrifuged at 1,000xg for 5 min and the medium was removed. Isopropanol-HCl-SDS solution (120 µl) was added to each well and the absorbance was read at 570 nm with a microplate reader (Molecular Devices, Sunnyvale, CA).

2.8 Cry11Aa toxin oligomerization

The oligomerization of Cry11Aa was performed by incubation with cells expressing AaeCad or EGFP [34, 35]. The activated Cry11Aa toxin (final concentration 400 nM) or the mixture of Cry11Aa protoxin (final concentration 1,400 nM) and midgut juice from *Ae. aegypti* larvae were incubated with 1×10^6 cells in 100 µl L15 medium for 1 h at 37°C. The mixture of Cry11Aa and cells was harvested at 500xg for 10 min and washed with PBS twice. The pellets were resuspended in SDS-PAGE sample buffer, boiled for 10 min, and centrifuged at 10,000xg for 10 min. The supernatants were loaded in SDS polyacrylamide gel (10%) and immunoblotted with anti-Cry11Aa antibody as noted above.

2.9 Construction of transformation vector and transgenic mosquitoes

The pBAC[3xP3-EGFP afm] transformation vector containing 3xP3 (eye-specific promoter), TATA box, and enhanced GFP [24] and the phsp-pBac helper plasmid [23] were obtained from Drs. Alexander Raikhel and Vladimir Kokoza, Department of Entomology, University of California, Riverside, CA. The hairpin dsRNA construct of AaeCad was made by using an *Aedes* heat shock promoter (689 bp), a cadherin (700 bp, AAEL007478) cadherin invert repeat having an APN intron (140 bp, AAEL008155), and SV40 (250 bp), and cloned into pBAC[3xP3-EGFP afm] transformation vector in GenScript company, Piscataway, NJ (Fig. 6).

The mixture of pBAC[3xP3-EGFP afm, AaeCad dsRNA] and phsp-pBac helper plasmid were injected into *Ae. aegypti* embryos (Insect Transformation Facility, University of

Maryland, Rockville, MD). G0 eggs were hatched and reared with a mixture of dog food and yeast (3:1) in deoxygenated tap water at 29°C, 8:12 h light:dark, and 50% humidity. Adult G0 were backcrossed with wild-type females or males and G1 progeny were screened for green fluorescent eye using fluorescence microscopy (Nikon SMZ1500). Homozygous larvae were obtained by subsequently mating individual male and female showing green-fluorescent eyes until all progeny larvae contained green-fluorescent eyes. To express the hairpin dsRNA of AaeCad regulated by the *Aedes* heat shock promoter in larvae, 2nd instar wild-type and transgenic larvae were incubated for 1 h at 37°C.

Bioassays were performed with crude Cry4Aa, Cry4Ba, Cry11Aa, and Cry11Ba. In brief, 20 early fourth-instar larvae were transferred to plastic cups containing 200 ml tap water and then fed Cry toxins at different concentration for 24 h. Bioassays were performed twice and the dose-response values were analyzed by probit (EPA) or Origin program (Origin Lab, Northampton, MA).

2.10 Quantitative Real-time PCR (qPCR)

Total RNA was extracted from wild-type or transgenic larvae using TRIzol. cDNA was synthesized from total RNA of each sample with SuperScript III (Invitrogen), diluted, and 5- μ l aliquots were used as template for qPCR. Respective primers specific to AaeCad (AaeCad-S and AaeCad-A, AAEL007478) and Actin (Actin-S and Actin-A, AAEL011197) or 40S ribosomal protein S7 (S7-S and S7-A, AAEL009496) as reference genes for quantification were designed to have similar properties in terms of nucleotide length and %GC content (Supplementary Table S1). Our microarray data showed the actin expression (AAEL011197) and 40S ribosomal protein S7 (AAEL009496) were not changed (-0.05 , -0.06 , or -0.07 / 0.17 , -0.08 , or 0.09 fold) in Cry11Aa-treated larvae midgut at LC₁₀, LC₅₀, LC₉₀ compared to untreated larvae midgut. PCR conditions, including the template cDNA, primer concentrations and annealing temperatures, were adjusted for amplification efficiencies (Efficiency 90 – 110%) for all genes. Optimized PCR master mix (20 μ l) contained the following components: 10 μ l iQ SYBR Green supermix (Bio-Rad, Irvine, CA), 5 μ l cDNA, 2 μ l 10 μ M of each primer, and 1 μ l water. The qPCR was performed using CFX Real-time PCR (Bio-Rad). Optimized thermal program consisted of: one cycle of 95°C/1 min and 40 cycles of 95°C/1 min, 62°C/1 min, and 72°C/1 min, followed by a final extension of one cycle 72°C /5 min. Following qPCR, the homogeneity of the PCR product was confirmed by melting curve analysis. To check for gDNA contamination in RNA preparation, the melting curve of 40S ribosomal protein S7 using primers that spanned a 114-bp intron was analyzed on every run. Quantification of the transcript levels or relative copy number of the genes was conducted according to the Pfaffl method [37]. Quantitative PCR was performed three times using independently prepared whole body cDNA.

3. Results

3.1 AaeCad are stably expressed in cells

A stable cell line expressing the *Aedes* cadherin under control of the actin promoter was obtained using a co-expressed selection vector. Analysis of these cells by western using anti-HA antibody and anti-cadherin fragment antibodies showed the presence of a band around

200 kDa in AaeCad-transfected cells. This size band was not detected in control cells expressing only EGFP (Fig. 2). Since the cadherin protein in the midgut of *Aedes* is also of this size [8] the full length cadherin is expressed in these cells (Fig. 2C).

To determine the sub-cellular location of AaeCad the cells were analyzed using an anti-cadherin fragment antibody and a secondary antibody conjugated with Cy3. Cells expressing AaeCad and control cells expressing EGFP were stained without permeabilization to detect AaeCad localized in the plasma membrane. Control cells did not show any specific immunofluorescence staining (Fig. 3A). However, red immunofluorescence was observed in the plasma membrane of AaeCad expressing cells (Fig. 3B).

3.2 Cells expressing AaeCad are more sensitive to Cry11Aa, Cry4Aa, and Cry11Ba toxins

Since *Aedes* cadherin binds the Cry11Aa and Cry11Ba toxins and hence should mediate Cry toxicity [8, 32], the toxicity of Cry proteins to AaeCad expressing cells was analyzed at final concentrations of 50 - 400 nM. Control cells expressing the EGFP protein were insensitive to all Cry toxins tested even at concentrations of up to 400 nM. However, cells expressing AaeCad showed significant sensitivity to the Cry11Aa toxin from 200 nM (29% death). Cry11Aa toxin at 400 nM killed approximately 37% of the cells in 3 h (Fig. 4). Moreover, these cadherin-expressing cells showed significant sensitivity to Cry4Aa (37.2% death) and Cry11Ba (25.7%) toxins, but were insensitive to Cry4Ba toxin (Table 1).

3.3 Cells expressing AaeCad trigger Cry11Aa oligomerization

It has been previously demonstrated that the toxicity of Cry toxins depends on the ability of these toxins to oligomerize in the presence of its receptor [22, 28]. Therefore, to determine whether the full-length AaeCad expressed in the C6/36 cells triggers Cry11Aa oligomerization within a cellular context, trypsin activated Cry11Aa at concentrations that causes cytotoxicity was analyzed. Alternatively, a higher concentration of the Cry11Aa protoxin activated with *Aedes* midgut juice was used as reported [36]. Both of these toxins were incubated with control cells or cells expressing AaeCad. In Cry11Aa incubations with cells expressing AaeCad, 140 kDa (intermediate) and 200 kDa (oligomer) bands from both activated toxin and protoxin were detected (Fig. 5, lane 2 and 4). However, only the Cry11Aa protoxin activated with midgut juice showed a 36 kDa (monomer) band (Fig. 5, lane 3 and 4) in both control and AaeCad expressing cells. Potentially this could be detected in lanes 3 and 4 because of the higher toxin concentrations used. Nevertheless, these data demonstrates that Cry11Aa oligomers are formed in the membranes of viable *Aedes* cells by both the trypsin activated Cry11Aa and midgut juice activated Cry11Aa protoxin.

3.4 Transgenic mosquitoes with silenced AaeCad are tolerant to Cry11Aa toxin

As noted above the *Aedes* cadherin has been previously shown to bind mosquitoicidal Cry toxins [8] and our data above shows that *Aedes* cells expressing cadherin increased susceptibility to these toxins. To analyze whether the cadherin gene plays a role in the *in vivo* toxicity of the Cry toxins, we silenced AaeCad expression in *Aedes* larvae. A hairpin dsRNA construct of the AaeCad was produced under control of the *Aedes* heat shock promoter and transgenic mosquitoes expressing this construct were obtained using pBAC transformation (Fig. 6, S1).

To silence AaeCad expression, wild-type and transgenic larvae were heat-shocked at 2nd instar for 1 h at 37°C. After this heat shock, the larvae were allowed to develop to the 4th instar. The transcript levels of AaeCad in whole larvae and in the midgut were significantly reduced in transgenic larvae compared to wild-type larvae (Table 2). Transgenic larvae with silenced AaeCad showed increased tolerance to Cry11Aa toxicity (Fig. 7C). However, silencing AaeCad did not show any difference in toxicity to Cry4Aa, Cry4Ba, and Cry11Ba (Fig. 7A, 7B, 7D).

4. Discussion

In Lepidoptera, binding of Cry1 toxins to cadherin causes a mild denaturation of Cry toxin and proteolytic cleavage of helix α -1 which results in a toxin conformational change [22, 28]. This conformational change is thought to cause the formation of oligomers that bind to a secondary receptor, which can be APN or ALP anchored to the membrane by a GPI anchor [6, 17]. Similarly, in *Aedes* mosquitoes, either Cry11Aa protoxin in the presence of BBMV or proteolytically activated Cry11Aa toxin incubated with AaeCad receptor fragment (CR7-11) alone can trigger toxin oligomerization [36, 38]. Here we also showed that C6/36 cells expressing full-length AaeCad can cause Cry11Aa oligomerization by the use of trypsin-activated Cry11Aa toxin directly or Cry11Aa protoxin in the presence of midgut juice (Fig. 5). There are some differences in the oligomeric forms obtained with these two toxins. Further, oligomerization of the Cry11Aa toxin is observed in cells at toxin concentration levels at which cytotoxicity is observed. Also, it is reported that the partial *Tenebrio molitor* cadherin (TmCad1) can promote the coleopteran-specific Cry3A toxin oligomerization [16]. Therefore, various cadherin receptors from three major insect pest orders have a conserved function leading to toxin oligomerization, which has been demonstrated as an important step for Cry toxicity.

In this study, AaeCad expressed in cells can not only trigger Cry11Aa oligomerization mentioned afore, but also the cells expressing AaeCad are susceptible to Cry11Aa, Cry4Aa, and Cry11Ba toxin (Fig. 4 and Table 1). Cry11Aa, Cry4Aa, and Cry11Ba toxins at 400 nM can cause about 37%, 37.2% and 25.7% cytotoxicity and Cry4Ba shows no cytotoxicity at the same dose. Our previous toxin binding competition assays showed Cry11Ba readily competes with Cry11Aa binding to AaeCad, while Cry4Aa slightly competes, and Cry4Ba does not compete [8]. In this study, Cry11Aa and Cry11Ba also showed significant cytotoxicity and Cry4Ba caused no cytotoxicity. However, Cry4Aa cytotoxicity is higher than expected since its binding with AaeCad is much weaker than Cry11Aa and Cry11Ba [8]. We continue to investigate this difference between the toxin binding assay and cytotoxicity for Cry4Aa.

In a similar approach, *Heliothis* cadherin transiently expressed in *Drosophila* S2 cells showed low cytotoxicity, maximum 20% and 5% cytotoxicity were caused by Cry1Ac and Cry1Ab, respectively at 330 nM [29]. However, only 180 nM Cry1Ab toxin is required to kill most of *Trichoplusia* High Five cells stably expressing *Manduca* cadherin [40, 41]. Notably, High Five cells apparently have a much higher level of cadherin expression compared with the AaeCad in C6/36 cells, and with *Heliothis* cadherin in S2 cells. Thus, it is very likely that the different cadherin expression levels could contribute to the different

sensitivities of cell lines to the corresponding toxins. On the other hand, it appears that Cry11Aa and Cry1A toxins shared the same mechanism of action because Cry11Aa can also bind to the same kinds of protein receptors as Cry1A toxin, for example, cadherin, APN and ALP. According to sequential binding model [5], in addition to cadherin, the secondary receptors are essential for the full toxicity *in vivo*. Thus it is understandable that C6/36 cells expressing only AaeCad show only limited sensitivity to mosquitocidal toxins. To obtain more sensitive cell lines, secondary receptors may have to be introduced into these cells expressing AaeCad.

Although AaeCad RNAi-mediated silencing had been done in *Aedes* mosquitoes by dsRNA feeding [38], a transgenic *Aedes* mosquito line with silenced AaeCad was established in order to obtain more homogeneous animals that can be used to screen a number of other toxins. Importantly, these transgenic mosquitoes revealed a higher tolerance to Cry11Aa (about 13 fold) at LC₅₀ value than *Aedes* larvae fed AaeCad dsRNA, where only a 50% reduction in Cry11Aa toxicity was obtained compared to wild-type mosquitoes [38].

Our bioassay data suggests AaeCad is involved in the toxicity of Cry11Aa but not in the toxicity of Cry4Aa, Cry4Ba, and Cry11Ba (Fig. 7). These bioassay data for Cry11Aa and Cry4Ba are consistent with toxin binding competition assay performed by Chen et al, 2009 [8] and cytotoxicity data from C6/36 cells expressing AaeCad in this study. Thus, while cadherin is critical for Cry11Aa toxicity it is not important for *in vivo* Cry4Ba toxicity. However, Cry4Aa and Cry11Ba *in vivo* toxicity in cadherin-silenced mosquitoes is quite different from the *in vitro* binding assays performed in earlier studies [8, 32] and the cytotoxicity test data reported here. These different results suggest that receptors other than AaeCad could be involved and may be more important for the toxicity of Cry4Aa and Cry11Ba, than they are for Cry11Aa.

Collectively, our data suggest that AaeCad clearly mediates the *in vivo* Cry11Aa toxicity to *Ae. aegypti* mosquitoes, and that AaeCad is not involved in Cry4Ba toxicity. In *Anopheles gambiae*, two *Anopheles* cadherins, AgCad1 and AgCad2, were identified as putative receptors for Cry4Ba and Cry11Ba toxins, respectively [25, 26]. It is possible a cadherin receptor other than AaeCad is involved in Cry4A or Cry11Ba toxicity. In contrast, Cry4Ba can form oligomer spontaneously in the presence of small unilamellar vesicles [38], and thus it is quite likely that no cadherin receptor is required for Cry4Ba cytotoxicity. Thus, Cry4Ba is a unique toxin and its mechanism of action appears to differ from that of Cry11Aa. Recently an ATP-binding cassette (ABC) transporter was shown to play a vital role in Cry1A toxicity and toxin resistance in lepidopteran insects [2, 3, 20]. It is still unknown if any ABC transporter is also involved in the mechanism of action of mosquitocidal toxins, including Cry4Aa, Cry4Ba and Cry11Ba. We believe novel RNAi technology and genome editing tools could be used for further elucidating the mechanism of action of these toxins

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The technical assistance of Maria Ramirez and Jacqueline Ledezma is gratefully acknowledged. This research was funded in part through grants from the National Institutes of Health, 1R01AI066014 and the University of California Agricultural Experiment Station.

6. References

- [1]. Abdullah MA, Valaitis AP, Dean DH. Identification of a *Bacillus thuringiensis* Cry11Ba toxin-binding aminopeptidase from the mosquito, *Anopheles quadrimaculatus*. *BMC Biochem.* 2006; 7:16. [PubMed: 16716213]
- [2]. Atsumi S, Miyamoto K, Yamamoto K, Narukawa J, Kawai S, Sezutsu H, et al. Single amino acid mutation in an ATP-binding cassette transporter gene causes resistance to Bt toxin Cry1Ab in the silkworm, *Bombyx mori*. *Proc Natl Acad Sci.* 2012; 109:E1591–E8. [PubMed: 22635270]
- [3]. Baxter SW, Badenes-Pérez FR, Morrison A, Vogel H, Crickmore N, Kain W, et al. Parallel evolution of *Bacillus thuringiensis* toxin resistance in Lepidoptera. *Genetics.* 2011; 189:675–9. [PubMed: 21840855]
- [4]. Berry C, O’Neil S, Ben-Dov E, Jones AF, Murphy L, Quail MA, et al. Complete sequence and organization of pBtoxis, the toxin-coding plasmid of *Bacillus thuringiensis* subsp. *israelensis*. *Appl Environ Microbiol.* 2002; 68:5082–95. [PubMed: 12324359]
- [5]. Bravo, A.; Gill, SS.; Soberon, M. *Bacillus thuringiensis*: Mechanisms and Use. In: Gilbert, LI.; Kostas, I.; Gill, SS., editors. *Comprehensive Molecular Insect Science*. Elsevier; 2005. p. 175-205.
- [6]. Bravo A, Gomez I, Conde J, Munoz-Garay C, Sanchez J, Miranda R, et al. Oligomerization triggers binding of a *Bacillus thuringiensis* Cry1Ab pore-forming toxin to aminopeptidase N receptor leading to insertion into membrane microdomains. *Biochim Biophys Acta.* 2004; 1667:38–46. [PubMed: 15533304]
- [7]. Chang C, Yu YM, Dai SM, Law SK, Gill SS. High-level cryIVD and cytA gene expression in *Bacillus thuringiensis* does not require the 20-kilodalton protein, and the coexpressed gene products are synergistic in their toxicity to mosquitoes. *Appl Environ Microbiol.* 1993; 59:815–21. [PubMed: 8481007]
- [8]. Chen J, Aimanova KG, Fernandez LE, Bravo A, Soberon M, Gill SS. *Aedes aegypti* cadherin serves as a putative receptor of the Cry11Aa toxin from *Bacillus thuringiensis* subsp. *israelensis*. *Biochem J.* 2009; 424:191–200. [PubMed: 19732034]
- [9]. Chen J, Aimanova KG, Pan S, Gill SS. Identification and characterization of *Aedes aegypti* aminopeptidase N as a putative receptor of *Bacillus thuringiensis* Cry11A toxin. *Insect Biochem Mol Biol.* 2009; 39:688–96. [PubMed: 19698787]
- [10]. Chen J, Likitvivanavong S, Aimanova KG, Gill SS. A 104 kDa *Aedes aegypti* aminopeptidase N is a putative receptor for the Cry11Aa toxin from *Bacillus thuringiensis* subsp. *israelensis*. *Insect Biochem Mol Biol.* 2013; 43:1201–8. [PubMed: 24128608]
- [11]. Chilcott C, Ellar DJ. Comparative Toxicity of *Bacillus thuringiensis* var. *israelensis* Crystal Proteins *in vivo* and *in vitro*. *J Gen Microbiol.* 1988; 134:2551–8. [PubMed: 3254944]
- [12]. Chow E, Gill S. A rapid colorimetric assay to evaluate the effects of *Bacillus thuringiensis* toxins on cultured insect cells. *J Tissue Cult Methods.* 1989; 12:39–42.
- [13]. Cowles EA, Yunovitz H, Charles JF, Gill SS. Comparison of toxin overlay and solid-phase binding assays to identify diverse CryIA(c) toxin-binding proteins in *Heliothis virescens* midgut. *Appl Environ Microbiol.* 1995; 61:2738–44. [PubMed: 7618886]
- [14]. Crickmore N, Zeigler DR, Feitelson J, Schnepf E, Van Rie J, Lereclus D, et al. Revision of the nomenclature for the *Bacillus thuringiensis* pesticidal crystal proteins. *Microbiol Mol Biol Rev.* 1998; 62:807–13. [PubMed: 9729610]
- [15]. Delecluse A, Rosso ML, Ragni A. Cloning and expression of a novel toxin gene from *Bacillus thuringiensis* subsp. *jegathesan* encoding a highly mosquitocidal protein. *Appl Environ Microbiol.* 1995; 61:4230–5. [PubMed: 8534090]

- [16]. Fabrick JA, Forlow Jech L, Henneberry TJ. Novel pink bollworm resistance to the Bt toxin Cry 1Ac: effects on mating, oviposition, larval development and survival. *J Insect Sci.* 2009; 9:24. [PubMed: 19613847]
- [17]. Fernandez LE, Aimanova KG, Gill SS, Bravo A, Soberon M. A GPI-anchored alkaline phosphatase is a functional midgut receptor of Cry11Aa toxin in *Aedes aegypti* larvae. *Biochemical J.* 2006; 394:77–84.
- [18]. Fernandez LE, Martinez-Anaya C, Lira E, Chen J, Evans A, Hernandez-Martinez S, et al. Cloning and epitope mapping of Cry11Aa-binding sites in the Cry11Aa-receptor alkaline phosphatase from *Aedes aegypti*. *Biochemistry.* 2009; 48:8899–907. [PubMed: 19697959]
- [19]. Fernandez LE, Perez C, Segovia L, Todriguez MH, Gill SS, Bravo A, et al. Cry11Aa toxin from *Bacillus thuringiensis* binds its receptor in *Aedes aegypti* mosquito larvae through loop α -8 of domain II. *FEBS Lett.* 2005; 579:3508–14. [PubMed: 15963509]
- [20]. Gahan LJ, Pauchet Y, Vogel H, Heckel DG. An ABC transporter mutation is correlated with insect resistance to *Bacillus thuringiensis* Cry1Ac toxin. *PLoS Genet.* 2010; 6:e1001248. [PubMed: 21187898]
- [21]. Gill SS, Cowles EA, Pietrantonio PV. The mode of action of *Bacillus thuringiensis* endotoxins. *Annu Rev Entomol.* 1992; 37:615–36. [PubMed: 1311541]
- [22]. Gomez I, Sanchez J, Miranda R, Bravo A, Soberon M. Cadherin-like receptor binding facilitates proteolytic cleavage of helix α -1 in domain I and oligomer pre-pore formation of *Bacillus thuringiensis* Cry1Ab toxin. *FEBS Lett.* 2002; 513:242–6. [PubMed: 11904158]
- [23]. Handler AM, Harrell RA 2nd. Germline transformation of *Drosophila melanogaster* with the piggyBac transposon vector. *Insect Mol Biol.* 1999; 8:449–57. [PubMed: 10634970]
- [24]. Horn C, Wimmer EA. A versatile vector set for animal transgenesis. *Dev Genes Evol.* 2000; 210:630–7. [PubMed: 11151300]
- [25]. Hua G, Zhang Q, Zhang R, Abdullah AM, Linser PJ, Adang MJ. AgCad2 cadherin in *Anopheles gambiae* larvae is a putative receptor of Cry11Ba toxin of *Bacillus thuringiensis* subsp. *jegathesan*. *Insect Biochem Mol Biol.* 2013; 43:153–61. [PubMed: 23231770]
- [26]. Hua G, Zhang R, Abdullah MA, Adang MJ. *Anopheles gambiae* cadherin AgCad1 binds the Cry4Ba toxin of *Bacillus thuringiensis israelensis* and a fragment of AgCad1 synergizes toxicity. *Biochemistry.* 2008; 47:5101–10. [PubMed: 18407662]
- [27]. Huynh CQ, Zieler H. Construction of modular and versatile plasmid vectors for the high-level expression of single or multiple genes in insects and insect cell lines. *J Mol Biol.* 1999; 288:13–20. [PubMed: 10329122]
- [28]. Jimenez-Juarez N, Munoz-Garay C, Gomez I, Saab-Rincon G, Damian-Almazo JY, Gill SS, et al. *Bacillus thuringiensis* Cry1Ab mutants affecting oligomer formation are non-toxic to *Manduca sexta* larvae. *J Biol Chem.* 2007; 282:21222–9. [PubMed: 17537728]
- [29]. Jurat-Fuentes JL, Adang MJ. The *Heliothis virescens* cadherin protein expressed in *Drosophila* S2 cells functions as a receptor for *Bacillus thuringiensis* Cry1A but not Cry1Fa toxins. *Biochemistry.* 2006; 45:9688–95. [PubMed: 16893170]
- [30]. Lereclus D, Agaisse H, Gominet M, Chaufaux J. Overproduction of encapsulated insecticidal crystal proteins in a *Bacillus thuringiensis* spo0A mutant. *Biotechnology (N Y).* 1995; 13:67–71. [PubMed: 9634751]
- [31]. Likirvivanavong S, Chen J, Evans AM, Bravo A, Soberon M, Gill SS. Multiple receptors as targets of Cry toxins in mosquitoes. *J Agric Food Chem.* 2011; 59:2829–38. [PubMed: 21210704]
- [32]. Likirvivanavong SCJ, Bravo A, Soberón M, Gill SS. Cadherin, alkaline phosphatase, and aminopeptidase N as receptors of Cry11Ba toxin from *Bacillus thuringiensis* subsp. *jegathesan* in *Aedes aegypti*. *Appl Environ Microbiol.* 2011; 77:24–31. [PubMed: 21037295]
- [33]. Margalith, Y.; Ben-Dov, E. Biological Control by *Bacillus thuringiensis* subsp. *israelensis*. In: Rechicgl, JE.; Rechicgl, NA., editors. *Insect Pest Management: Techniques for Environmental Protection.* CRC Press; 2000. p. 243-301.
- [34]. Munoz-Garay C, Rodriguez-Almazan C, Aguilar JN, Portugal L, Gomez I, Saab-Rincon G, et al. Oligomerization of Cry11Aa from *Bacillus thuringiensis* has an important role in toxicity against *Aedes aegypti*. *Appl Environ Microbiol.* 2009; 75:7548–50. [PubMed: 19820153]

- [35]. Obata F, Kitami M, Inoue Y, Atsumi S, Yoshizawa Y, Sato R. Analysis of the region for receptor binding and triggering of oligomerization on *Bacillus thuringiensis* Cry1Aa toxin. *Febs J.* 2009; 276:5949–59. [PubMed: 19740106]
- [36]. Perez C, Munoz-Garay C, Portugal LC, Sanchez J, Gill SS, Soberon M, et al. *Bacillus thuringiensis* ssp. *israelensis* Cyt1Aa enhances activity of Cry11Aa toxin by facilitating the formation of a pre-pore oligomeric structure. *Cell Microbiol.* 2007; 9:2931–7. [PubMed: 17672866]
- [37]. Pfaffl M. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 2001; 29:e45. [PubMed: 11328886]
- [38]. Rodríguez-Almazán C, Reyes EZ, Zúñiga-Navarrete F, Muñoz-Garay C, Gómez I, Evans AM, et al. Cadherin binding is not a limiting step for *Bacillus thuringiensis* subsp. *israelensis* Cry4Ba toxicity to *Aedes aegypti* larvae. *Biochem J.* 2012; 443:711–7. [PubMed: 22329749]
- [39]. Zhang R, Hua G, Urbauer JL, Adang MJ. Synergistic and inhibitory effects of aminopeptidase peptides on *Bacillus thuringiensis* Cry11Ba toxicity in the mosquito *Anopheles gambiae*. *Biochemistry.* 2010; 49:8512–9. [PubMed: 20809561]
- [40]. Zhang X, Candas M, Griko NB, Rose-Young L, Bulla LA Jr. Cytotoxicity of *Bacillus thuringiensis* Cry1Ab toxin depends on specific binding of the toxin to the cadherin receptor BT-R₁ expressed in insect cells. *Cell Death Differ.* 2005; 12:1407–16. [PubMed: 15920532]
- [41]. Zhang X, Candas M, Griko NB, Taussig R, Bulla LAJ. A mechanism of cell death involving an adenylyl cyclase/PKA signaling pathway is induced by the Cry1Ab toxin of *Bacillus thuringiensis*. *Proc Natl Acad Sci USA.* 2006; 103:9897–902. [PubMed: 16788061]

Highlights

Aedes cadherins were stably expressed in a mosquito cell line and silenced in transgenic mosquitoes.

Cells expressing *Aedes* cadherin are more sensitive to Cry11Aa, Cry4Aa, and Cry11Ba toxins, but not Cry4Ba.

Cells expressing *Aedes* cadherin trigger Cry11Aa oligomerization just as with lepidopteran cadherins.

Transgenic mosquitoes with silenced *Aedes* cadherin show increased tolerance to Cry11Aa toxin




Fig. 1.

Construction of expression vectors. (A) pACTIN.SV-AaeCad. Modified *Aedes* cadherin (AaeCad) cDNA from *Ae. aegypti* larvae of 5301 bp in length was cloned into pACTIN.SV vector. (B) pACTIN.SV-EGFP. The pACTIN.SV vector, including the *EGFP* gene, was used for selection of a control cell line. (C) pIE1.SV-Bla. A blasticidin-resistance sequence (*bla* gene) from the pCoBlast vector was cloned into pIE1.SV vector. To make a stable cell line, pACTIN.SV-AaeCad and pIE1.SV-Bla / pACTIN.SV-EGFP and pIE1.SV-Bla were co-transfected into C6/36 (*Aedes albopictus*) cells and then cells were grown and maintained in fresh medium containing blasticidin.




Fig. 2.

AaeCad is stably expressed in AaeCad-transfected cells. To determine the expression of *Aedes* cadherin cloned into pACTIN.SV, non-soluble proteins were extracted and detected with an anti-HA antibody (1: EGFP-transfected cells, 2: AaeCad-transfected cells) or an anti-cadherin fragment antibody (3: EGFP-transfected cells, 4: AaeCad-transfected cells, 5: Brush border membrane vesicles from *Ae. aegypti* midgut). In all cases, we detected a band around 200 kDa in AaeCad-transfected cells. The 200-kDa protein corresponds to the full-length cadherin.




Fig. 3. *Aedes* cadherin is localized in the plasma membrane. To determine the sub-cellular location of *Aedes* cadherin, the expression was determined by immunofluorescence staining under confocal microscopy. Red immunofluorescence staining without permeabilization showed that *Aedes* cadherin were localized in the plasma membrane (B). (A) Immunostaining of control cells using anti-cadherin fragment antibody. (B) Immunostaining of cells expressing *Aedes* cadherin using anti-cadherin fragment antibody.




Fig. 4. Cells expressing *Aedes* cadherin are more sensitive to Cry11Aa toxin. The toxicity of Cry11Aa to these cells was analyzed in a MTT cytotoxicity assay using activated Cry11Aa toxin at final concentrations of 50 - 400 nM. After cells were incubated with activated Cry11Aa for 3 hours, live cells were analyzed by measuring reduced MTT. Control cells expressing the EGFP protein (black) were insensitive to Cry11Aa toxin up to 400 nM. However, cells expressing AaeCad (red) showed sensitivity to the toxin and Cry11Aa toxin at 400 nM killed approximately 37% of the cells in 3 hours (n=3, t-test, $p < 0.05$).




Fig. 5. Cells expressing *Aedes* cadherin trigger Cry11Aa oligomerization. Control cells expressing EGFP and cells expressing *Aedes* cadherin were incubated with activated Cry11Aa (final concentration 400 nM) and with trypsin or Cry11Aa protoxin (final concentration 1,400 nM) with midgut juice. After 1 h at 37°C, cells were washed, boiled for 10 min, and separated in 10% SDS-PAGE gel. Cry11Aa oligomerization was detected with an anti-Cry11Aa antibody. Lane 1: EGFP-transfected cells and active Cry11Aa, 2: AaeCad-transfected cells and active Cry11Aa, 3: EGFP-transfected cells and Cry11Aa protoxin with midgut juice, 4: AaeCad-transfected cells and Cry11Aa protoxin with midgut juice.




Fig. 6. Construction of the mosquito transformation vector. The hairpin dsRNA construct of *Aedes* cadherin was made by arranging *Aedes* heat shock promoter, *Aedes* cadherin, APN, *Aedes* cadherin inverted repeat (IR), and SV40. This construct was cloned into pBAC[3xP3-EGFP afm] transformation vector containing 3xP3 (eye-specific promoter), TATA box, and enhanced GFP.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 7. Transgenic mosquitoes with silenced *Aedes* cadherin expression show increased tolerance against Cry11Aa toxin. Bioassay was performed with early 4th instar larvae heat shocked at 2nd instar. Wild-type larvae (■) and transgenic larvae ▲ were incubated with Cry4Aa (A), Cry4Ba (B), Cry11Aa (C), and Cry11Ba (D) and mortalities were analyzed in Probit and Origin.

Table 1

Cells expressing *Aedes* cadherin are more sensitive to Cry11Aa, Cry4Aa and Cry11Ba toxins.

	EGFP-transfected cell (% survival)	AaeCad-transfected cell (% survival)
Cry11Aa	99.7±11.4	79.0±6.70 *
Cry4Aa	94.4±3.68	62.8±6.32 *
Cry4Ba	114±8.76	107±5.67
Cry11Ba	114±16.5	74.3±11.5 *

The toxicities of Cry4Aa, Cry4Ba, and Cry11Ba to these cells were analyzed in a MTT cytotoxicity assay using activated toxins at a final concentration of 400 nM. After cells were incubated with activated toxins for 3 h, live cells were analyzed by measuring reduced MTT. Control cells expressing the EGFP protein were insensitive to all toxins at 400 nM (94.4 – 114% survival compared to non-treated control cells). However, cells expressing *Aedes* cadherin showed significantly sensitivity to Cry4Aa (37.2% death) and Cry11Ba (25.7% death) toxins compared to non-treated cells expressing *Aedes* cadherin (n=3, t-test, $p < 0.05$). Values given are relative to the Cry11Aa toxicity to EGFP-transfected cells.

Table 2

Transgenic mosquitoes down-regulate *Aedes* cadherin expression in whole body.

	Transcript change in whole body	Transcript change in midgut
AaeCad	0.20 ± 0.23	0.49 ± 0.31

The larval AaeCad transcript levels were determined by qPCR. Transcript levels are relative to the transcript levels in wild-type larvae where the value is 1.0.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript