



Published in final edited form as:

Ann Biomed Eng. 2014 February ; 42(2): 445–459. doi:10.1007/s10439-013-0902-7.

Creating living machines

Roger D. Kamm¹ and Rashid Bashir²

Roger D. Kamm: rdkamm@mit.edu; Rashid Bashir: rbashir@illinois.edu

¹Massachusetts Institute of Technology, Cambridge, MA

²University of Illinois at Urbana-Champaign, Urbana, IL

Abstract

Development of increasingly complex integrated cellular systems will be a major challenge for the next decade and beyond, as we apply the knowledge gained from the sub-disciplines of tissue engineering, synthetic biology, micro-fabrication and nanotechnology, systems biology, and developmental biology. In this prospective, we describe the current state-of-the-art in the context of differentiating source cells from more primitive, pluripotent cells, and organizing these cells into populations of a single cell type to produce the components or building blocks of higher order systems and finally, combining multiple cell types, possibly in combination with scaffolds possessing specific physical or chemical properties, to produce greater functionality. As these “living machines” increase in capabilities, exhibit emergent behavior and potentially reveal the ability for self-assembly, self-repair, and even self-replication, questions arise regarding the ethical implications of this work. Future prospects as well as ways of addressing these complex ethical questions will be addressed.

Keywords

tissue engineering; systems biology; synthetic biology; biobots; vascular networks; neuromuscular junctions; biological machines

A. Introduction

The past century has witnessed tremendous advances in science and technology, but in no other field are these more evident than in the biological sciences. New discoveries in synthetic biology, tissue engineering, systems biology, and developmental biology, which have accelerated in the past 10 to 20 years, are nothing short of phenomenal. This new appreciation of fundamental biological processes is changing our lives in numerous ways ranging from new healthcare technologies to alternative energy sources to environmental protection.

But the potential for even greater and more significant advances lies ahead. Until recently, biological research has led primarily to a detailed yet largely qualitative understanding of fundamental phenomena at the molecular and cellular scales. These qualitative concepts are

being increasingly cast in quantitative form through new programs at the intersection of engineering and biology. The synthesis of ideas and approaches from these diverse disciplines has been termed *convergence* in a recent study from the National Research Council that lays out a visionary plan for future biological research⁴⁹. While many of the specifics remain ill defined, it is becoming increasingly evident that the advances in molecular and cellular biology of the 20th century are certain to translate into entirely new technologies in the 21st.

One of the greatest opportunities lies in the potential to understand populations of multiple cell types and their interactions. To a large degree, this has been one of the driving forces behind developmental biology, tissue engineering, and systems biology. Yet, it can be argued that a tremendous gap exists between understanding processes at the level of a single cell and the behavior of large-scale tissues, i.e. how the local rules of interaction result in global functionalities and diverse phenotypes. This is an issue involving complex systems of multiple interacting components that could fruitfully draw upon considerable advances in the engineering realms of forward engineering, forward design, and manufacturing of large, complex systems.

B. The Foundational Disciplines

Creation of living machines is critically dependent upon developments drawn from a range of existing disciplines, a unique and thoughtful fusion of which can result in a new discipline where engineers are designing machines and systems with biological components and cells at various length scales (Figure 1). We first review some of the significant advances, and discuss them in the context of the present state-of-the-art (see Figure 2 for a collection of representative research nuggets). Progress in creating living machines relies on the fusion and convergence of these different fields that have to be seamlessly integrated to result in the pedagogical foundations of a new discipline dealing with design and realization of engineered biological machines from cells.

Synthetic biology is the field that perhaps most closely parallels what we propose. Among the major advances that have been made is a detailed and systematic process for mathematically modeling gene regulatory networks¹¹². For this purpose, a systems objective has been adopted that is capable of describing and analyzing how changes in the genetic code of the cell affect cell behavior. More than that, it also provides the framework for engineering modifications in cell behavior that are desired in order to confer new and useful functionality. This, more than any other advance, lays the groundwork for engineering cell-cell interactions.

As significant as these advances are, much of the work thus far has been directed toward engineering individual strains of bacteria or yeast systems. This is a logical approach for many reasons, but places constraints on the types of functionality that cells might ultimately achieve by interactions with multiple different cell types. Mammalian cells, in particular, offer unique capabilities, developed over millions of years of evolutionary pressure, and exhibiting a wide range of coordinated behavior. It is not coincidental that mammals are considered the highest form of life, since the collective behavior of an organism created

from a large variety of different cell types has far more potential to attain high-level function, which would, in fact, be impossible with a single phenotype/genotype. Therefore, although not all biological machines will be based on mammalian cells, the ability to apply the lessons learned from synthetic biology to all types of cells, acting in a coordinated manner, will be essential in creating machines that are both multifunctional, and ecologically stable in the sense that no one cell type overtakes the others.

Tissue engineering and its sister field, regenerative medicine, also have much to contribute to the creation of living machines. Notably, tissue engineering has achieved some major technological advances over the years. Artificial skin was the first engineered tissue to reach the market. In 1982, the first composite living skin was produced which led to a commercial product⁴⁵. Other tissues such as bladder and cartilage are either available commercially or soon to come onto the market (see table 1 in⁵). Still others have been the subject of intense investigation, and the technologies are rapidly improving. Products are at various stages of development for liver¹³¹, bladder³, pancreas²⁶, heart tissue¹⁸, and others⁵, but fundamental barriers persist, such as the need for a vascular system to meet the metabolic needs of the tissue construct.

It might be argued that one of the factors that has prevented tissue engineering from having attained broader success is the lack of a fundamental understanding of the complex processes that lead to tissue formation, complexity that comes in many forms such as mechanistic, biological, or systems. Systems biology has been building the tools that will enable this deeper understanding, through the development of computational approaches to simulate biological processes at multiple length scales²³¹⁰. These range from models based on first principles to those that are purely data-driven, and all have been fruitfully employed. But whatever their approach, they collectively offer the potential to serve as a receptacle of the knowledge gained about a particular system or process, and allow us to capture and simulate the complex interactions between systems and between scales that might otherwise be incomprehensible.

Much of our fundamental understanding of biological systems and the methods that might be used to grow them comes from developmental biology. Since the study of developmental biology most closely mimics the 'bottom-up' or emergence driven behavior of cellular clusters, foundational principles from this very broad and widely researched field will play a critical role in creation of the proposed biological machines. Since these studies are challenging to do in mammalian based systems due to the longer time spans and the ethical considerations involved, model systems such as zebra fish are being studied in great detail¹⁷. Genetic manipulation and engineering is an important tool to vary the genetic programming in these model systems to examine the resultant phenotypes and correlate genes to phenotype and function. Many of the morphogens that enable cell-cell signaling during development have been identified and the process of development can be manipulated by controlling morphogen gradients or localized cell depolarization. For example, introduction of ion channels that alter the local transmembrane potential are also being used to study the resulting development of model organisms such as planarians or tadpoles to form organisms with two heads or multiple organs with stably formed and reproducing organisms⁴.

Cells used in biological machines ultimately need to be compatible with each other in terms of inter-cellular signaling and maintaining long-term stability, and this argues for the use of cells from the same species if not the same organism. And since it is likely that some of the cells will need to be genetically modified, one logical solution would be to derive cells from a pluripotent source. Therefore, stem cell biology is another pillar in the discipline of biological machines, and here again, we are able to draw upon a wealth of prior work in support of these efforts. Protocols already exist for the differentiation of many of the cell types needed for the various functionalities needed in these machines from stem cells or other precursors. And while not every cell type can be derived, the methods developed and collection of known morphogenic agents serve as a useful starting point.

The basic idea of design and realization of living machines mimics that of forward engineering design, where precise models and specifications exist for the components. Using these, the design of more complex machines can be achieved. This is similar to the goals of synthetic biology where libraries and specifications for parts are used to create newer functions. Similarly, systems design and top-down decomposition of a complex system such as a biological machines and the design specification of the sub components will be critical for the design of different machines using those subcomponents.

Many advances in microfluidics and nanotechnology over the past decade have now allowed the development of this enabling field of technology with applications in biology, medicine, electronics, materials, energy and other areas. The technologies for fabrication of soft polymers such as PDMS, hydrogels, and biomaterials can especially have major implications in the development of biological machines. BioMEMS (bio-Micro Electro Mechanical Systems) and Microfluidic devices are being used a point of care devices⁴¹, in-vitro cell culture devices for drug screening, and implantable hybrid devices for innovative solutions to organic-inorganic interfaces²². 3-D fabrication and printing of cells and polymer scaffolds can be used to place cells with spatial control on or in scaffolds to realize new physical designs for biological machines. Techniques such as photo-polymerization using stereolithography, 3-D rapid prototyping, and 3-D printing can be especially useful in this context⁹⁴⁸.

Many examples exist of developments in each of these important foundational areas where the fusion and synthesis of these concepts can contribute to the development of cellular and biological machines (Figure 2). This ‘forward-engineering’ of biological components is certainly an important element of what is called the ‘New Biology’, which envisions the integration of multiple disciplines, leading ultimately to enhanced understanding of fundamental principles in biology, and ultimately novel approaches to the problems that face society (Figure 3). Yet another perspective emerged from discussions at a recent *Keck Futures Initiative* that focused on the extension of synthetic biology into more advanced cell types, such as mammalian cells, and moving toward poly-culture systems, potentially using cells from multiple species to generate new and unique behaviors. Thus, what we present here is an idea that is increasingly being explored, accepted, and embraced by the research community. Ours is one perspective on the future, in particular, a future that envisions living machines that are engineered to perform specific tasks that differ from or perhaps combine or enhance the capabilities of existing organisms. We summarize here the current state-of-

the-art in the critical foundational disciplines, recent work that makes the case that many of the needed elements are already existent, and attempt to identify future needs in terms of where efforts are needed in order to achieve these ambitious goals and develop a systematic process for creating living machines.

C. Biological Machine and Essential Elements

Can we truly “engineer” or “orchestrate” the growth of entire biological systems? To do so will require that we overcome a number of significant barriers. Biological systems are exquisitely complex, even at the scale of a single cell, and the level of complexity escalates precipitously when multi-cell interactions need to be considered. One embodiment of the conceptual framework leading to a biological machine is shown in Figure 4; an increasing level of complexity is expected as one progresses through cells, modules, and eventually a machine (Figure 5). Take, for example, the case of simple interactions between neural-controlled muscle of a scale that requires a vascular supply, as needed for a biological robot (“biobot”) of millimeter or greater scale. Certainly the individual cell types needed – neuronal, muscular, vascular – can be isolated or even derived from pluripotent cells, and these can be co-cultured in systems that enable some level of interaction. Inducing these cells to combine into a functional, stable muscle actuator for use in a biobot is an enormous step, however, and will require significant advances on many fronts.

Given the enormous complexity of living systems, it seems implausible that one could engineer and control the position of each cell and its interactive functionality with its neighbors. So, although we can and must specify the design parameters of the machine, we can only hope to assemble clusters of cells into the approximately correct arrangement, and rely on natural processes to establish the functional complexity needed for its operation. As one example, we can locate motor neuron clusters in close proximity to an engineered muscle strip, but the formation of axons with functional synapses to muscle must be accomplished by the cells themselves, possibly via release of chemotactic agents, cell migration towards these chemicals, and subsequent formation of the synapse itself. In a sense, the cells must participate in the fabrication process of the machine, and in doing so, we must rely on biological processes (e.g., those that occur during development, regeneration or wound repair) to help. We consider this “emergent behavior” to be an intrinsic competence of the cells, and must at some level rely on it to proceed naturally. Here, the concepts associated with developmental biology, as mentioned earlier, come into play. So we view the manufacture of a living machine as requiring both top-down design specification and a process for “assembling the parts”, as well as a reliance on emergent processes that are programmed into the genetic code of the cells, and involve various forms of cell-cell and cell-matrix interactions.

D. What is Possible Today?

While this all may seem futuristic and to some, unrealistic, an argument can be made that the creation of living machines is not only possible, but that some simple machines could even be on the near-term horizon. We construct this argument by working up in complexity. We first consider homotypic cell clusters; that is, collections of cells of a single type that

function in some unified manner. Examples considered here include muscle, neuronal networks, and vascular networks. Then we consider initial efforts to combine multiple cells types (heterotypic cell clusters) into a more complex functional unit.

D1. Homotypic Cell Clusters

Muscle strips—Many of the machines one might envision, require actuation or motility, and for this, will need to incorporate muscle or some form of contractile cell. Due to their intrinsic tendency to rhythmically contract, cardiomyocytes have been the cell of choice in many of the early motile machines. One method that garnered considerable attention was the use of temperature-responsive polymers functionalized with cell-adhesive ligands to culture cell sheets of cardiomyocytes. Uniformly grafted sheets of poly(N-isopropylacrylamide) (PIPAAM) were formed on polystyrene culture dishes by irradiation with an electron beam to form the cell sheets³². Kitamori and colleagues first demonstrated a micropump powered by a cell sheet of cardiomyocytes achieving flow rates of 2 nL min^{-1} ³⁶. The same group recently improved on their method by wrapping a sheet of functional primary cardiomyocytes around a hollow PDMS sphere with inlet and outlet capillary tubes to engineer a bio-artificial hybrid pump³⁷. The fluid oscillating frequency measured at $37 \text{ }^\circ\text{C}$ was 0.4 Hz and the maximum observed linear displacement of tracking particles was $70 \text{ }\mu\text{m}$. The expected flow rate was 47 nL min^{-1} , which was an improvement over the previous design.

Cardiac muscle cells are self-paced and hence lack the potential for control offered by skeletal muscle cells that contract on demand by external activation. Recently, methods have been developed to produce contractile skeletal muscle strips, tethered to compliant posts as shown in Figure 6, both to support the tension needed for proper myotube formation and to allow for direct inference of the contractile stress³⁹. These methods produce highly stratified muscle, with clearly delineated sarcomeric structure, however, the levels of stress generated, even taking into account the amount of matrix material incorporated into the muscle construct, ($\sim 200 \text{ Pa}$) are more than 100-fold lower than what can be generated in vivo ($>10^5 \text{ Pa}$) (Sakar et al., 2012). Although the reasons for this considerable difference in contractile strength are unclear, it could be related either to the lack of capillary blood flow to the muscle or the mechanism of activation, which is often by electrical stimulation rather than synaptic activation by motor neurons.

Neuronal Networks—Neurons are a natural source of cells that can gather and process data, and then direct another cell population to perform a desired function. For example, sensory cells detecting a toxin might transmit a signal to a neuronal network that would direct secretory cells to synthesize and release a neutralizing agent, or muscle cells to move in the direction of the stimulus to initiate other actions. In short, neuronal clusters could be called upon to receive a signal and direct a coordinated response from multiple other cell types. It should be recognized that for some applications, the sensing and responding cells may be one in the same; beta cells, for example, sense glucose levels and secrete appropriate levels of insulin in response. Often, however, complex sensing/response problems involve some degree of processing by the central nervous system.

Progress in developing functional neuronal networks has progressed, but slowly. An essential feature of such networks is the ability to communicate, and in the case of neurons, communication occurs via signal transmission, either bi-directionally via electrical signals passed through gap junctions, or in a uni-directional fashion mediated by neurotransmitters released at a synapse. Synaptic connections have been extensively studied, and can readily be established between cultured neurons of various types derived from embryonic stem cells¹². Moreover, functional networks produced with such neurons have been demonstrated¹⁹, although additional reports and detailed characterization are still lacking. Creation of neuronal networks that perform specific functions, however, has been proceeding at a slow pace, and relies generally on the ability to place the neurons in a particular geometrical arrangement and precisely control connections. These constructs are often referred to as a “brain on a chip”⁴². And while these experimental advances coupled with computational models of function are significant, many challenges lie ahead before these constructs can be designed and constructed to perform a specific processing function.

Vascular Networks—For years, one of the barriers to progress in tissue-engineered organs was the inability to generate a perfusable microvascular network that could provide adequate gas and nutrient exchange to the regenerated tissue. Recent work in several labs has produced several methods that now overcome this constraint. The earliest work demonstrated that microvascular networks could be patterned onto a PDMS substrate, lined with endothelial cells, and perfused¹⁵. Others²⁴ demonstrated that straight channels could be cast in gel (collagen in this instance), lined with endothelial cells, and perfused, showing excellent cell viability and wall permeabilities that are approaching *in vivo* values. These later led to methods to form networks that could be cast in 3-dimensional gel matrices, the casting material removed, and endothelial cells seeded on the walls of the channels produced²⁷.

One disadvantage of all these methods involving casting channels in gels, is that the smallest channels tend to be >100 μm in diameter, so still roughly 10 \times larger than the real microvascular bed. While casting methods might yet achieve these dimensions, an alternative has been developed during the past several years that shows considerable promise. When endothelial cells are plated onto gel surfaces, they can be induced to sprout into the gel, recreating the process of angiogenesis that occurs during, for example, wound healing or cancer. These vessels form capillary-sized vessels with a more natural branching pattern^{28,46,7}, and can now be grown so that they span a region of matrix of up to several mm, and can be perfused, representing an important step toward *in vitro* vascularized organs. Several methods have been demonstrated to induce network growth, including the addition of growth factors such as VEGF, either in uniform concentration or in a gradient, choice of an appropriate matrix material (fibrin or a fibrin-collagen mix appear most conducive), co-culture with various stromal cell types, and the application of physical factors such as interstitial flow across an endothelial monolayer^{33,40} or through a cell-seeded 3D matrix²⁸. And while many issues remain to be addressed such as functionality, control of network morphology, and long-term phenotypic stability, the recent advances bode well for ultimate success of these methods, essentially using the natural ability of endothelial cells to generate microvascular networks.

D2. Heterotypic Cell Clusters and Living Machines

Neuromuscular Junctions—Among the most basic cell-cell communication systems is one that allows neurons to communicate with other neurons, or motoneurons to activate skeletal muscle. As an example of heterotypic systems, we consider here the latter case. In the preceding section, we discussed recent work that has led to the growth in vitro of muscle strips that are on the scale of 100 μm in diameter and mms in length. Many of these systems have been grown from cardiomyocytes, so possess the capability of cyclic activation, ideal for first generation biobots. In order to develop the ability to control the motions, however, will likely require skeletal muscle along with a means of activation. One promising approach has been to express in the muscle cells a light sensitive calcium channel (e.g., channelrhodopsin (Sakar et al., 2012)). But the more natural means would be to use motoneurons that activate the myotubes by direct synaptic excitation. And while significant advances have been made in the generation of skeletal muscle strips in vitro, there has been very limited success in the formation of functional synapses³⁸.

Co-cultures of Vascular Cells—Vascular networks, by themselves, are of little value, since their purpose is to provide nutrients and gas exchange for other cells in a tissue. Some recent experiments attest to the viability of doing so in an in vitro setting. For example, Yeon et al.⁴⁶ have used fibroblasts as a “feeder cell” in 3D culture to help create a vascular network (Figure 8(A)), and Chan et al.⁷ have created networks through gels in the presence of agarose beads (Figure 8(B)) which they show could contain tumor cells or various types of stromal cells. Other co-culture systems with fibroblasts have also been demonstrated in which the vascular network is cast into a hydrogel, which can also be seeded with other cell types²⁴²⁷. Note that most of these publications have appeared during the past one or two years, so this area of research is likely to grow considerably.

Biobots—Among the various living machines envisioned, biological robots, or “biobots” are perhaps the most advanced. These are relatively simple systems capable of moving under the rhythmic contraction of cardiac myocytes appropriately seeded onto a flexible substrate. Montemagno and colleagues developed a microdevice using a silicon backbone with self-assembled cardiomyocytes grown on a chromium/gold layer⁴⁴. They used photolithographic techniques to fabricate a micron-sized, self-assembled, and self-actuated walking bio-microactuator powered by cardiomyocyte muscular tissue, achieving a maximum speed of 38 $\mu\text{m}/\text{s}$. Park and colleagues established a swimming microrobot by micromolding PDMS²⁰. Parker and colleagues assembled cardiomyocytes on various PDMS thin films with proteins to create muscular thin films (MTFs)¹⁴ to reverse-engineer jellyfish-like constructs, dubbed “medusoids”²⁹. The Medusoid propulsion was like that of a jellyfish and was externally-driven by electrically-paced power and was able to replicate the momentum transport and body lengths traveled per swimming stroke of the natural system (Figure 9a). Most recently, a 3D printer has also been used for the assembly of “bio-bots” with poly(ethylene glycol) (PEG) hydrogels and neonatal rat cardiomyocytes (Figure 9b)⁸. The bio-bots consisted of a ‘biological bimorph’ cantilever structure as the actuator to power the bio-bot, and the elastic properties of the bio-bots were tuned similar to that of neonatal rat cardiomyocytes to maximize their contractile force ($\sim 5 \mu\text{N}$). The maximum recorded velocity of the bio-bot was $\sim 236 \mu\text{m}\cdot\text{s}^{-1}$.

While current biobots have limited capabilities (e.g., they can only function in cell culture medium and their movement is uncontrolled) they are useful in that they constitute an experimental platform on which new functionalities can be incrementally added. As mentioned above, optogenetic muscle cells can be used to gain control over the rate of contraction, hence the speed of motion. Using these same optogenetically modified cells, the direction of movement can be controlled by selectively activating muscles positioned to produce movement in multiple directions. In a further extension, motor neurons could be used to produce more efficient contraction via synaptic control, and these, too, could be activated by light.

Organs on a Chip—Last year, the FDA, NIH and DARPA took the unprecedented step of introducing joint programs aimed at developing a disruptive change in the technologies used by the pharmaceutical industries to discover new drugs. These programs called for the development of “organs-on-a-chip” that, by extrapolating into the future, could replicate human organs and their interactions with sufficient fidelity to be used to screen for new therapies and their potential off-target complications that often block drugs that have reached the stage of clinical testing. Some small steps have been made in the development of microfluidic systems that can replicate certain aspects of organ function (Figure 10). The grand challenge of the new programs is to further this development, and produce a “body-on-a-chip” technology that not only could model the response of a single organ to a new compound, but also anticipate the off-target effects of that same drug as it interacts with multiple other subsystems or organs. While it might be some time before such systems are capable of screening the millions of compounds in a particular library, in the nearer term, these might be useful as secondary screens, taking hits identified by the more conventional multi-well systems and refining these to a smaller number that can be moved up the ladder for further testing. At some stage, these technologies might replace animal testing, realizing significant savings in resources and time, but also providing a more realistic test bed by using human cells. Longer-term, such systems might be produced based on cells for a particular patient, providing the ultimate in identifying patient-specific treatment protocols.

E. More Complex Living Machines on the Horizon

With but a little imagination, it is not difficult to envision how these nascent technologies might be developed into higher level machines with a wide spectrum of functionalities. Here we describe a few examples merely to provide stimulus for others to expand our horizons in this critical new field.

Smart Plants

Society faces continuing challenges to feed our growing populations and despite dramatic improvements in food production, we continue to face shortages. In addition, problems remain in addressing the increasingly dramatic swings in the yearly cycles that periodically devastate a particular crop or food supply. What if plants could sense their surroundings, process that information, and enter into dormant or active growing periods based on that input? That is, what if plants had the ability to sense, “think”, and act in a manner that would enhance their ability to survive under adverse conditions? Higher-level organisms have

developed neurons for this purpose, but plants, too, could benefit from these capabilities. Either by genetically modifying plant cells so that they can form logic circuits and process information, or by introducing a mammalian-derived neuron cell type into plants, such capabilities might be realized.

What if we were able to enable plants to not only sense their condition, but process that information and respond appropriately? One example would be if they were to shut down all vital functions – to enter into a state of dormancy – during periods of draught. Another example might be that plants experiencing inadequate water, other nutrients or sunlight might send a signal requesting more.

Hyper-organs

While much attention has been focused on the growth of organ systems that currently exist in order to replace a dysfunctional liver, kidney, heart, etc., one can also envision implantable systems (“organs”) that perform other functions, not currently accomplished with any existing system. For example, one might envision a drug delivery system for chronic illnesses that senses the concentration of a desired drug, then directs other cells to synthesize and secrete the drug into the circulation. Another system might consist of a simple elastic reservoir connected to the circulation via a sphincter controlled by cells that sense the concentration of some cytokine in the blood. Sensing drives the relaxation of sphincter contraction, allowing for the release of needed factors into the body. Additional possibilities include blood vessels that pump relieving the load on the heart, or cell based sensors to measure the increase in pressure from vascular occlusions and release anti-thrombotic factors produced by cell based factories embedded within the blood vessels.

Emergent Manufacturing

One of the unique potential advantages of biological machines is their inherent capability for growth, self-assembly, and self repair. Living organisms require no external guidance to develop into mature systems, and one might argue that living machines should be no different. As was discussed earlier, it might be sufficient to place the machine “parts” in proper relative proximity to each other, and leave the final steps of assembly to biological processes.

But it clearly is not that simple, as developmental biologist have worked for decades to understand the emergent behaviors that lead to the maturation of a living organism. At its most fundamental level, all steps of development can be reduced to the “initial condition” or pre-programming intrinsic to the embryonic stem cells of particular machine/organism, the signaling that occurs (both intracellular intercellular, and between the cells and their local extracellular matrix), and perhaps with global signals generated by the external environment. While enormously complex in its entirety, the individual steps leading to emergence can be understood and, in principle, applied. So, while emergent manufacturing may be on the distant horizon, the tools and fundamental understanding needed to make it a reality largely exist today.

Bio-based Surveillance Systems

Despite tremendous advances, living systems still outperform non-living ones in the critical areas of sensing – we still rely on dogs for detecting drugs or explosives in luggage, and voice recognition by computers continues to lag far behind human capabilities. Among these, smell is the one sense that is least dependent upon information processing by the central nervous system, and therefore, represents a logical first target for applications in surveillance or detection. Some advances have been made in the development of systems that attempt to replicate cellular sensing, but the potential of using mammalian cells in a sensing device remains largely unexplored.

F. Potential Benefits and Dangers of Living Machines

The potential benefits of a fully biological machine are numerous and transformative. One might envision machines that can self-assemble, repair themselves, and even self-replicate under appropriate controls. But this raises numerous important ethical questions, in addition to the scientific/engineering barriers to progress. At what stage does a biological machine become a living being? What ethical issues does this kind of research raise? How can cell based systems have a positive impact on the world around us and how can we prevent harmful outcomes of this research? At what level of complexity does a biological machine become a living organism, and what features distinguish one from the other? And how far can or should we go in “engineering” cellular systems that resemble in form or function a living entity that can perform a specific function? A machine that can sense its surroundings, process the information that it gathers, and perform some function based on that decision process possesses many of the same qualities that we often attribute exclusively to life and natural living beings. Technologies developed to protect our borders might also be used to collect information on unsuspecting citizens, and machines designed to self-replicate or evolve pose enormous potential risks. Obviously, the ethical issues are of tremendous importance and the time for discussion is now, not once the technologies have already been developed.

G. Conclusions

Synthetic biology brought us the prospect of engineering single cells to perform entirely new functions or to exhibit characteristics different from their natural counterparts. Other disciplines have strived to create and understand through modeling and experiments the behavior of complex, multi-cellular systems. Through this emerging technology, we have already witnessed advances in mono-culture constructs (e.g., muscle strips, microvascular networks, neuronal circuits, myocyte-driven biobots) as well as heterotypic multi-culture systems (neuromuscular junctions, organs-on-a-chip, etc.). In this prospective, we have sought to present the case that even greater potential exists in the use of multiple cell types, each performing different functions in a coordinated manner, to produce higher-order forms of living machines. While we can only speculate regarding the future of these endeavors, the groundwork is now being laid through advances in a number of related fields. We propose here that integrated cellular systems be recognized as an emerging discipline, and that efforts be undertaken to develop the nascent ideas presented in this prospective piece to

promote related research and also initiate discussions of the critical ethical questions that this research raises.

Acknowledgments

The authors acknowledge support from the National Science Foundation (NSF) Science and Technology Center (STC) Emergent Behavior of Integrated Cellular Systems (EBICS) Grant CBET-0939511, as well as valuable discussions with Robert Nerem, Todd McDevitt, Ron Weiss, Linda Griffith, Hyunjoon Kong, Taher Saif, K. Jimmy Hsia, and many other colleagues from the EBICS STC.

Bibliography

1. Allen JW, Bhatia SN. Engineering liver therapies for the future. *Tissue engineering*. 2002; 8:725–37. [PubMed: 12459052]
2. Andrianantoandro E, Basu S, Karig DK, Weiss R. Synthetic biology: new engineering rules for an emerging discipline. *Molecular systems biology*. 2006; 2:2006.0028. [PubMed: 16738572]
3. Atala A, Bauer SB, Soker S, Yoo JJ, Retik AB. Tissue-engineered autologous bladders for patients needing cystoplasty. *Lancet*. 2006; 367:1241–6. [PubMed: 16631879]
4. Beane WS, Morokuma J, Lemire JM, Levin M. Bioelectric signaling regulates head and organ size during planarian regeneration. *Development (Cambridge, England)*. 2013; 140:313–22.
5. Berthiaume F, Maguire TJ, Yarmush ML. Tissue engineering and regenerative medicine: history, progress, and challenges. *Annual review of chemical and biomolecular engineering*. 2011; 2:403–30.
6. Booth R, Kim H. Characterization of a microfluidic in vitro model of the blood-brain barrier (μ BBB). *Lab on a chip*. 2012; 12:1784–92. [PubMed: 22422217]
7. Chan JM, Zervantonakis IK, Rimchala T, Polacheck WJ, Whisler J, Kamm RD. Engineering of in vitro 3D capillary beds by self-directed angiogenic sprouting. *PloS one*. 2012; 7:e50582. [PubMed: 23226527]
8. Chan V, Park K, Collens MB, Kong H, Saif TA, Bashir R. Development of miniaturized walking biological machines. *Scientific reports*. 2012; 2:857. [PubMed: 23155480]
9. Chan V, Zorlutuna P, Jeong JH, Kong H, Bashir R. Three-dimensional photopatterning of hydrogels using stereolithography for long-term cell encapsulation. *Lab on a chip*. 2010; 10:2062–70. [PubMed: 20603661]
10. Chuang H-Y, Hofree M, Ideker T. A decade of systems biology. *Annual review of cell and developmental biology*. 2010; 26:721–44.
11. Davidson E, Levin M. Gene regulatory networks. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102:4935. [PubMed: 15809445]
12. Dodla MC, Mumaw J, Stice SL. Role of astrocytes, soluble factors, cells adhesion molecules and neurotrophins in functional synapse formation: implications for human embryonic stem cell derived neurons. *Current stem cell research & therapy*. 2010; 5:251–60. [PubMed: 20214556]
13. Domansky K, Inman W, Serdy J, Dash A, Lim MHM, Griffith LG. Perfused multiwell plate for 3D liver tissue engineering. *Lab on a chip*. 2010; 10:51–8. [PubMed: 20024050]
14. Feinberg AW, Feigel A, Shevkoplyas SS, Sheehy S, Whitesides GM, Parker KK. Muscular thin films for building actuators and powering devices. *Science (New York, NY)*. 2007; 317:1366–70.
15. Fidkowski C, Kaazempur-Mofrad MR, Borenstein J, Vacanti JP, Langer R, Wang Y. Endothelialized microvasculature based on a biodegradable elastomer. *Tissue engineering*. 2005; 11:302–9. [PubMed: 15738683]
16. Grosberg A, Alford PW, McCain ML, Parker KK. Ensembles of engineered cardiac tissues for physiological and pharmacological study: heart on a chip. *Lab on a chip*. 2011; 11:4165–73. [PubMed: 22072288]
17. Grunwald DJ, Eisen JS. Headwaters of the zebrafish -- emergence of a new model vertebrate. *Nature reviews. Genetics*. 2002; 3:717–24.

18. Hansen A, Eder A, Bönstrup M, Flato M, Mewe M, Schaaf S, Aksehirlioglu B, Schwoerer AP, Schwörer A, Uebeler J, Eschenhagen T. Development of a drug screening platform based on engineered heart tissue. *Circulation research*. 2010; 107:35–44. [PubMed: 20448218]
19. Heikkilä TJ, Ylä-Outinen L, Tanskanen JMA, Lappalainen RS, Skottman H, Suuronen R, Mikkonen JE, Hyttinen JAK, Narkilahti S. Human embryonic stem cell-derived neuronal cells form spontaneously active neuronal networks in vitro. *Experimental neurology*. 2009; 218:109–16. [PubMed: 19393237]
20. Kim J, Park J, Yang S, Baek J, Kim B, Lee SH, Yoon E-S, Chun K, Park S. Establishment of a fabrication method for a long-term actuated hybrid cell robot. *Lab on a chip*. 2007; 7:1504–8. [PubMed: 17960278]
21. Kim S, Lee H, Chung M, Jeon NL. Engineering of functional, perfusable 3D microvascular networks on a chip. *Lab on a chip*. 2013:1489–1500.10.1039/c3lc41320a [PubMed: 23440068]
22. Kim, T-i; McCall, JG.; Jung, YH.; Huang, X.; Siuda, ER.; Li, Y.; Song, J.; Song, YM.; Pao, HA.; Kim, R-H.; Lu, C.; Lee, SD.; Song, I-S.; Shin, G.; Al-Hasani, R.; Kim, S.; Tan, MP.; Huang, Y.; Omenetto, FG.; Rogers, JA.; Bruchas, MR. Injectable, Cellular-Scale Optoelectronics with Applications for Wireless Optogenetics. *Science*. 2013; 340:211–216. [PubMed: 23580530]
23. Lauffenburger, Da. The multiple dimensions of Integrative Biology. *Integrative biology: quantitative biosciences from nano to macro*. 2012; 4:9. [PubMed: 22158999]
24. Leung AD, Wong KHK, Tien J. Plasma expanders stabilize human microvessels in microfluidic scaffolds. *Journal of biomedical materials research. Part A*. 2012; 100:1815–22. [PubMed: 22489049]
25. Long C, Finch C, Esch M, Anderson W, Shuler M, Hickman J. Design optimization of liquid-phase flow patterns for microfabricated lung on a chip. *Annals of biomedical engineering*. 2012; 40:1255–67. [PubMed: 22271245]
26. Lysy PA, Weir GC, Bonner-Weir S. Concise review: pancreas regeneration: recent advances and perspectives. *Stem cells translational medicine*. 2012; 1:150–9. [PubMed: 23197762]
27. Miller JS, Stevens KR, Yang MT, Baker BM, Nguyen D-HT, Cohen DM, Toro E, Chen AA, Galie PA, Yu X, Chaturvedi R, Bhatia SN, Chen CS. Rapid casting of patterned vascular networks for perfusable engineered three-dimensional tissues. *Nature materials*. 2012; 11:768–74.
28. Moya ML, Hsu Y, Lee AP, Hughes CCW, George SC. In Vitro Perfused Human Capillary Networks. *Tissue engineering. Part C, Methods*. 2013; 19
29. Nawroth JC, Lee H, Feinberg AW, Ripplinger CM, McCain ML, Grosberg A, Dabiri JO, Parker KK. A tissue-engineered jellyfish with biomimetic propulsion. *Nature biotechnology*. 2012; 30:792–7.
30. Sakar MS, Neal D, Boudou T, Borochin Ma, Li Y, Weiss R, Kamm RD, Chen CS, Asada HH. Formation and optogenetic control of engineered 3D skeletal muscle bioactuators. *Lab on a chip*. 2012; 12:4976–85. [PubMed: 22976544]
31. Sakar MS, Neal DM, Boudou T, Borochin MA, Li Y, Weiss R, Kamm R, Chen CS, Asada HH. Formation and optogenetic control of engineered 3D skeletal muscle bioactuators. *Lab on a Chip*. 2012
32. Shimizu T, Yamato M, Kikuchi A, Okano T. Cell sheet engineering for myocardial tissue reconstruction. *Biomaterials*. 2003; 24:2309–16. [PubMed: 12699668]
33. Song JW, Munn LL. Fluid forces control endothelial sprouting. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:15342–7. [PubMed: 21876168]
34. Sung JH, Esch MB, Prot J-M, Long CJ, Smith A, Hickman JJ, Shuler ML. Microfabricated mammalian organ systems and their integration into models of whole animals and humans. *Lab on a chip*. 2013:1201–1212.10.1039/c3lc41017j [PubMed: 23388858]
35. Sung JH, Yu J, Luo D, Shuler ML, March JC. Microscale 3-D hydrogel scaffold for biomimetic gastrointestinal (GI) tract model. *Lab on a chip*. 2011; 11:389–92. [PubMed: 21157619]
36. Tanaka Y, Morishima K, Shimizu T, Kikuchi A, Yamato M, Okano T, Kitamori T. An actuated pump on-chip powered by cultured cardiomyocytes. *Lab on a chip*. 2006; 6:362–8. [PubMed: 16511618]
37. Tanaka Y, Sato K, Shimizu T, Yamato M, Okano T, Kitamori T. A micro-spherical heart pump powered by cultured cardiomyocytes. *Lab on a chip*. 2007; 7:207–12. [PubMed: 17268623]

38. Umbach, Ja; Adams, KL.; Gundersen, CB.; Novitch, BG. Functional neuromuscular junctions formed by embryonic stem cell-derived motor neurons. *PloS one*. 2012; 7:e36049. [PubMed: 22574134]
39. Vandenburgh H. High-content drug screening with engineered musculoskeletal tissues. *Tissue engineering. Part B, Reviews*. 2010; 16:55–64. [PubMed: 19728786]
40. Vickerman V, Kamm RD. Mechanism of a flow-gated angiogenesis switch: early signaling events at cell-matrix and cell-cell junctions. *Integrative biology: quantitative biosciences from nano to macro*. 2012; 4:863–74. [PubMed: 22673733]
41. Watkins N, Irimia D, Toner M, Bashir R. On a chip. *IEEE pulse*. 2011; 2:19–27. [PubMed: 22147065]
42. Wheeler BC, Brewer GJ. Designing Neural Networks in Culture: Experiments are described for controlled growth, of nerve cells taken from rats, in predesigned geometrical patterns on laboratory culture dishes. *Proceedings of the IEEE Institute of Electrical and Electronics Engineers*. 2010; 98:398–406. [PubMed: 21625406]
43. Wilson K, Das M, Wahl KJ, Colton RJ, Hickman J. Measurement of contractile stress generated by cultured rat muscle on silicon cantilevers for toxin detection and muscle performance enhancement. *PloS one*. 2010; 5:e11042. [PubMed: 20548775]
44. Xi J, Schmidt JJ, Montemagno CD. Self-assembled microdevices driven by muscle. *Nature materials*. 2005; 4:180–4.
45. Yannas, Burke JF, Orgill DP, Skrabut EM. Wound tissue can utilize a polymeric template to synthesize a functional extension of skin. *Science (New York, NY)*. 1982; 215:174–6.
46. Yeon JH, Ryu HR, Chung M, Hu QP, Jeon NL. In vitro formation and characterization of a perfusable three-dimensional tubular capillary network in microfluidic devices. *Lab on a chip*. 2012; 12:2815–22. [PubMed: 22767334]
47. Zheng Y, Chen J, Craven M, Choi NW, Totorica S, Diaz-Santana A, Kermani P, Hempstead B, Fischbach-Teschl C, López JA, Stroock AD. In vitro microvessels for the study of angiogenesis and thrombosis. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109:9342–7. [PubMed: 22645376]
48. Zorlutuna P, Jeong JH, Kong H, Bashir R. Stereolithography-Based Hydrogel Microenvironments to Examine Cellular Interactions. *Advanced Functional Materials*. 2011; 21:3642–3651.
49. *A New Biology for the 21st Century*. Washington, DC: National Academies Press; 2009.




Figure 1.
The fusion of different disciplines and specialties needed to develop living machines.




Figure 2.

Synthesis and examples of the different disciplines: (a) gene networks in bacteria can be programmed to result in different patterns, (b) the “Vacanti mouse” as an example of tissue engineering and control of biological phenotype, (c) systems biology depicted in terms of increasing complexity of mechanistic, biological, and systems understanding, (d) developmental biology examples such as growth of a zebrafish embryo (left) and examples of phenotypic control by altering the electrical polarization of adult stem cells to generate a 2nd head on a planarian, or to develop a second working eye induced on tadpole gut, (e) stem cell differentiation to produce neurons, muscle or endothelial cells, (f) systems design and top-down decomposition of complex systems such as an airliner and design specification of the sub components, and (g) examples in microfluidics, lab on chip, and 3-D fabrication using stereolithographic printing of cells and polymers for tissue engineering and 3-D soft systems




Figure 3. Conceptual schematic of the interdisciplinary foundations of the New Biology with implications for many solutions to societal problems. Reproduced from *A New Biology for the 21st Century*.




Figure 4. Conceptual schematic of a Biological Machine with cells, scaffolds and physical or chemical cues to result in machines that exhibit specific functionalities.




Figure 5.
Increasing levels of complexity to go from cells, modules, and machines.




Figure 6. Formation of muscle strips anchored to flexible posts. A) Sequential images of myoblasts in a collagen/Matrigel solution after seeding into a rectangular well with two compliant posts. B) Schematic of the resulting muscle strip formed around the two compliant posts. C) Fluorescent image showing cell membranes (green) and nuclei (red). D) Muscle strip as in (C) stained for actin showing cell alignment at 3 days post seeding. E) Striated actin (red) and multi-nucleated (green) cells. Adapted from (Sakar et al., 2012).




Figure 7. Neuronal circuits on a chip. (a) Network of individual neurons patterned by laser, (b) 30 μm lines and 80 μm square nodes at 21 days in culture. (c) Neuropil structure separated by 500 μm with 3 μm wide lines. (d) Cross pattern of 80 μm nodes and 30 μm lines, stained for neurons (green), astroglia (red), and nuclei (blue). Adapted from ⁴².




Figure 8.

Vascular networks formed in microfluidic platforms. Left: A) Microfluidic system. B) Schematic showing four parallel channels of the device with two outside gel regions (LO, RO), two media channels (LI, LO) and a central gel region. C) – F) Different seeding conditions for forming vascular networks. Reproduced from ²¹. Right top: A) – C) Vascular networks formed in the central gel region as in Left (A). Right bottom: A) Confocal image of a perfusable vascular network grown in a microfluidic gel system. B) Slices of the network in (A) showing lumens and the 3D nature of the formed vessels. Reproduced from ⁷.




Figure 9.

Autonomous bio-hybrid muscle actuators capable of (a) swimming in fluid – bioengineering of a artificial jelly fish like structure capable of swimming in fluid autonomously or being pulsed by an external electric field (Adapted from Nawroth, et al. 2012), (b) walking in fluid - Biological Biomorph cantilever structure actuation with the beating of primary cardiac cells resulting in a net motion with maximum velocity 236 $\mu\text{m/s}$, average displacement 354 $\mu\text{m/stroke}$ and average beating frequency ~ 1.5 Hz Adapted from ⁸.




Figure 10.

Body-on-a-chip. Reproduced from ³⁴. Conceptual image of how the various existing organs-on-a-chip might be assembled to simulate the entire physiological system of a human for the purpose of drug screening. A) Lung. Reproduced from ²⁵. B) Blood brain barrier. Reproduced from ⁶. C) Heart tissue. Reproduced from ¹⁶. D) Liver. Reproduced from ¹³. E) Intestinal villi. Reproduced from ³⁵. F) Muscle. Reproduced from ⁴³. G) Blood vessels. Reproduced from ⁴⁷.