



Published in final edited form as:

*Inorg Chem.* 2013 June 17; 52(12): 6767–6769. doi:10.1021/ic400353j.

## Inelastic X-Ray Scattering (IXS) of a Transition Metal Complex (FeCl<sub>4</sub><sup>-</sup>)— Vibrational Spectroscopy for All Normal Modes

Hongxin Wang<sup>&‡</sup>, Weibing Dong<sup>&‡</sup>, Marilyn M. Olmstead<sup>&</sup>, James C. Fettinger<sup>&</sup>, Jay Nix<sup>‡</sup>, Hiroshi Uchiyama<sup>¥</sup>, Satoshi Tsutsui<sup>¥</sup>, Alfred Q. R. Baron<sup>¥,§</sup>, Eric Dowty<sup>§</sup>, and Stephen P. Cramer<sup>\*,&,‡</sup>

<sup>&</sup> Department of Chemistry, University of California, Davis, CA 95616

<sup>‡</sup> Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

<sup>†</sup> Department of Biochemistry & Molecular Biology, Pennsylvania State University, University Park, PA 16802

<sup>¥</sup> Research and Utilization Division, SPring-8/JASRI, 1-1-1 Kouto, Sayo, Hyogo 679-5198, JAPAN

<sup>§</sup> Materials Dynamics Laboratory, RIKEN SPring-8 Center, 1-1-1 Kouto, Sayo, Hyogo 679-5148, JAPAN

<sup>§</sup> Shape Software, 521 Hidden Valley Road, Kinsport, TN 37663

### Abstract

The tetraethylammonium salt of the transition metal complex (FeCl<sub>4</sub><sup>-</sup>) has been examined using inelastic x-ray scattering (IXS) with 1.5 meV resolution (12 cm<sup>-1</sup>) at 21.747 keV. This sample serves as a feasibility test for more complex transition metal complexes. The IXS spectra were compared with previously recorded infrared, Raman, and NRVS spectra, revealing the same normal modes but with less strict selection rules. Calculations with a previously derived Urey Bradley force field were used to simulate the expected Q and orientation dependence of the IXS intensities. The relative merits of IXS, as compared to other photon based vibrational spectroscopies such as NRVS, Raman, and IR are discussed.

### Keywords

inelastic x-ray scattering; IXS; synchrotron radiation; normal modes; vibrational spectroscopy

Vibrational spectroscopy is one of the most fundamental tools for inorganic chemistry. The standard techniques, IR and Raman spectroscopy, are powerful probes of structure and dynamics, and they complement each other with different selection rules<sup>1</sup>. Particle scattering methods such as inelastic neutron scattering (INS) and high resolution electron energy loss (HREELS) also have their merits, but have respective requirements for multi

\*Corresponding Author spjcramer@ucdavis.edu.

†Present Addresses Weibing Dong, Department of Biochemistry, Pennsylvania State University, University Park, PA 16802

Supporting Information. Experimental details for sample preparation; crystallographic results concerning phase transformations at different temperatures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

gram samples or an ultra high vacuum environment. One synchrotron based technique, nuclear resonant vibrational spectroscopy (NRVS), has shown great potential for characterization of materials and metalloproteins<sup>2</sup>, but it can only be applied to isotopes such as <sup>57</sup>Fe with appropriate excited state energies and lifetimes. A second synchrotron based technique, resonant inelastic x-ray scattering (RIXS), has been used to observe well spaced molecular vibrations in samples such as O<sub>2</sub><sup>3a</sup>, but at the moment is limited to several hundred cm<sup>-1</sup> resolution<sup>3b</sup>. Here we employ a third synchrotron method, inelastic x-ray scattering (IXS)<sup>4</sup>. IXS has previously been applied to pure elements such as carbon in diamond or graphite<sup>5</sup>, binary solid state compounds such as MgB<sub>2</sub>, MgO, and SiC<sup>6</sup>, complex superconductors<sup>4c</sup>, liquids as diverse as water or <sup>3</sup>He<sup>7</sup>, and even DNA and proteins<sup>8</sup>, but never to a metal coordination complex.

IXS differs fundamentally from IR, Raman, and RIXS in that it depends on an  $\mathbf{A}^2$  term in the Hamiltonian for the interaction of radiation with matter, as opposed to the  $\mathbf{p} \cdot \mathbf{A}$  terms relevant to the other methods (where  $\mathbf{A}$  is the vector potential and  $\mathbf{p}$  is momentum)<sup>9</sup>. Whereas x-ray diffraction Bragg peaks derive from coherent elastic scattering of x-rays, IXS derives from the coherent inelastic scattering of x-rays by phonons in a sample. In fact, much of the thermal diffuse scattering (TDS) seen between diffraction peaks is part of the IXS signal, but in a TDS measurement there is no attempt to energy resolve that scattering. Thanks to developments in synchrotron radiation sources and x-ray optics, IXS experiments can resolve scattered radiation with an energy resolution of ~1 meV (8 cm<sup>-1</sup>). As a test of the feasibility and relevance of this technique for inorganic coordination chemistry, we recorded the IXS for a crystal of a tetrahedral Fe complex: (NEt<sub>4</sub>)(FeCl<sub>4</sub>). Despite the small cross section and competition with photoabsorption, we were able to record excellent spectra using an established spectrometer<sup>4c,10</sup>. A representative spectrum (acquired in 24 hours) is compared with previously recorded NRVS, Raman, and IR data in **Figure 1**; experimental details are provided in **Supporting Information**.

For an isolated transition metal complex in tetrahedral T<sub>d</sub> symmetry, there are 4 types of normal modes: A<sub>1</sub> and T<sub>2</sub> stretching motions, along with T<sub>2</sub> and E bending modes. The selection rule advantage of IXS for seeing these modes, compared to other methods, is clearly visible in **Figure 1**, where all 4 of the expected bands are clearly visible. This is in sharp contrast to the IR, NRVS, and Raman spectra, which are each most sensitive to only 2 out of the 4 main features. For example, the bands at 119 and 137 cm<sup>-1</sup> can be attributed to the E and T<sub>2</sub> bend modes respectively<sup>11</sup>. The former is strong in the Raman while the latter is more visible in the IR and NRVS, but both bands are strong in the IXS. At higher energies, the bands at 330 and 380 cm<sup>-1</sup> correspond to the totally symmetric A<sub>1</sub> stretching mode and the T<sub>2</sub> asymmetric stretch. As before, the first band is strong in the Raman while the second is mostly visible by IR or NRVS, but both modes are distinct in the IXS.

A closer look reveals that the FeCl<sub>4</sub><sup>-</sup> asymmetric stretching mode near 380 cm<sup>-1</sup> is split. This is clearest in the Raman spectrum, where bands at 376 and 389 cm<sup>-1</sup> are resolved. In the room temperature crystal structure, the symmetry of the Fe-Cl<sub>4</sub><sup>-</sup> ion is better described as C<sub>3v</sub>, with 3 shorter Fe-Cl bonds at 2.177 Å and 1 longer Fe-Cl bond at 2.186 Å. Descent in symmetry from T<sub>d</sub> to C<sub>3v</sub> splits the T<sub>2</sub> modes into E and A<sub>1</sub> modes. This splitting is

suggested in the IXS, since the line width of the  $A_1$  region is 12-14  $\text{cm}^{-1}$  FWHM (essentially the beamline resolution), while for the  $T_2$  region it is  $\sim 30 \text{ cm}^{-1}$ .

The IXS data in Figure 1 come from just one crystal orientation and scattering angle  $\theta$ . One of the strengths of this technique is that particular normal modes can be enhanced by variation of both settings. To illustrate the value of Q-dependent and orientation-dependent IXS for highlighting particular normal modes, we present spectra obtained for different conditions in Figure 2. (At  $\chi=0$  the crystal  $c$ -axis was vertical and the beam was along the  $a$ -axis. At  $\chi=45^\circ$  the  $c$ -axis was rotated to make a  $45^\circ$  degree angle with the beam. The additional curves each took 12 hours.)

From the data shown in Figure 2, it is clear that there is a strong Q-dependence to the relative intensities of IXS bands. For example, for  $\theta=15^\circ$  the stretching bands are  $\sim 3$ -fold weaker than the bending modes, while for  $\theta=27^\circ$  the  $T_2$  stretch and  $T_2$  bend are of comparable intensity. There are also wide variations within the stretching region. As shown better in Figure 2 (right), for  $\theta=15^\circ$  the  $A_1$  stretch intensity is 82% of the  $T_2$  intensity, while for  $\theta=27^\circ$  it is reduced to 24%. Finally, there are also significant variations in relative intensities for a fixed scattering angle. For example, at  $\theta=27^\circ$ , rotation of the crystal by  $45^\circ$  about the  $a$ -axis yields the spectrum with the weakest  $A_1$  band.

With a known crystal structure and force field, all of the observed changes are in principle calculable from the well-developed theory behind IXS. This theory has been extensively reviewed elsewhere<sup>4a,12</sup>, so before discussing the representative calculated spectra, we merely summarize the key factors that govern IXS intensities. With initial photon energy  $E_i$  and final energy  $E_f$ , the phonon energy is given by:  $E=\hbar\omega \cong E_i - E_f$ . As defined in Figure 1, where  $\vec{k}_i, \hat{\epsilon}_i, \vec{k}_f$ , and  $\hat{\epsilon}_f$  are the initial and final photon wavevectors and polarizations respectively, and for scattering angle  $\theta$  the momentum transfer is  $\hbar\vec{Q}=2\hbar\vec{k}_i \sin(\theta/2)$ . The single phonon cross-section for scattering,  $\frac{\partial^2\sigma}{\partial\Omega\partial E}$ , is proportional to  $S(\vec{Q}, \omega)$  – the ‘dynamical structure factor’. Along with terms accounting for phonon population and polarization effects,  $S(\vec{Q}, \omega)$  is in turn proportional to the ‘inelastic structure factor’,  $F_{in}(\vec{Q})$ , which involves a summation over all the atoms in the primitive cell, and thus for a particular phonon mode<sup>4a,4c, 12</sup>:

$$\frac{\partial^2\sigma}{\partial\Omega\partial E} \propto S(\vec{Q}, E) \propto F_{in}(\vec{Q}) = \left| \sum_m M_m^{-1/2} f_m(\vec{Q}) \left[ \vec{e}_m^n(\vec{q}) \cdot \vec{Q} \right] \exp(i\vec{Q} \cdot \vec{r}_m) \exp(-w_m) \right|^2$$

In the above expression, atom  $m$  with mass  $M_m$  is located at  $\vec{r}_m$  and has x-ray form factor  $f_m(Q)$  and Debye-Waller factor  $\exp(-w_m)$ . The term  $\vec{e}_m^n(\vec{q})$  is the phonon polarization eigen-vector for the direction of motion of atom  $m$  in phonon mode  $n$  with wavevector  $\vec{q}$ <sup>4c</sup>. Since as  $Q \rightarrow 0, f_m(Q) \rightarrow Z$ , at small angles the strength of scattering from a particular atom in a sample will vary approximately as  $Z^2$ . The overall signal strength depends on the product

of three terms:  $f_m(Q)$ ,  $Q^2$ , and a polarization factor that goes as  $\cos^2\theta$ . In practice, this product maximizes at  $Q \approx 10 \text{ \AA}^{-1}$  for many elements.

The product  $\vec{e}_m^n(\vec{q}) \cdot \vec{Q}$ , the projection of the phonon polarization onto the total momentum transfer vector  $\vec{Q}$ , plays a critical role in the observed intensities. Thus the intensities of the IXS signal can be directly related to the motions of particular atoms in a given normal mode <sup>4a,12</sup> projected onto  $\vec{Q}$ . Since one can choose different  $\vec{Q}$  by controlling the scattering geometry, IXS can be made more sensitive to particular phonons in a sample. This means that IXS has the flexibility to see *all* of the normal modes for a transition metal coordination complex. The appearance of the sum inside the magnitude signs means that the intensity is sensitive to the relative phase of the motions within one primitive cell.

The strength of geometry-dependent IXS for highlighting specific normal modes is illustrated in Figure 3. As examples, we show the  $Q$ -dependence for an incident beam along the crystal  $a$ -axis (with  $c$ -axis vertical and a horizontal scattering plane) and the IXS for different crystal orientations at fixed  $Q$ .

One striking result from these simulations is the extreme variability in intensity of the totally symmetric  $A_1$  stretching mode at  $330 \text{ cm}^{-1}$ . Within the range of  $\theta$  from  $24$  to  $28^\circ$ , the strength of this mode varies from almost as strong as the  $T_2$  mode to nearly invisible. A similar variability in relative intensities is observed for different crystal orientations at a fixed scattering angle. From a practical point of view, this ability to enhance or diminish particular normal modes should assist the detailed assignment of vibrational features in complex spectra.

In summary, the current results show that high quality IXS spectra can be obtained for a transition metal complex in a few hours at a third generation synchrotron radiation source. IXS intensities are amenable to straightforward calculation from a given structure and force field, with none of the complexities involved in the calculation of IR or Raman spectra. Compared to NRVS, IXS has the advantage of applicability to any element in the periodic table, and compared to INS, IXS requires orders of magnitude less material. The  $Q$ -dependence allows one to enhance the visibility of particular normal modes. There is also a benefit from orientation dependence of single crystal spectra, but it has been shown by others that powder spectra are also informative.<sup>12b</sup> Although it has limitations as a bulk-sensitive technique, there should be many applications in co-ordination chemistry where selection rules or other constraints preclude the use of more conventional vibrational methods, and where IXS will shine as a technique that allows any de-sired normal mode to be observed.

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS


This work was funded by NIH grants GM-65440 (SPC) and the DOE Office of Biological and Environmental Research (SPC). Use of SPring 8 is supported by JASRI.

## REFERENCES

1. a Nakamoto, K. *Infrared & Raman Spectra of Inorganic & Coordination Compounds*. 5th ed.. Wiley-Interscience; New York: 1997. b Squires, GL. *Introduction to the Theory of Thermal Neutron Scattering*. Dover; New York: 1978. c Richardson NV. *Curr. Opin. Solid State Mat. Sci.* 1997; 2:517–524. d Seto M, Yoda Y, Kikuta S, Zhang XW, Ando M. *Phys. Rev. Lett.* 1995; 74:3828–3831. [PubMed: 10058307]
2. a Alp E, Sturhahn W, Toellner TS, Zhao J, Hu M, Brown DE. *Hyp. Interact.* 2002; 144/145:3–20. b Sturhahn W. *J. Phys. Cond. Matt.* 2004; 16:S497–S530.
3. a Hennies F, Pietzsch A, Berglund M, Fohlisch A, Schmitt T, Strocov V, Karlsson HO, Andersson J, Rubensson JE. *Phys. Rev. Lett.* 2010; 104:193002–193005. [PubMed: 20866962] b van Schooneveld MM, Gosselink RW, Eggenhuisen TM, Al Samarai M, Monney C, Zhou KJ, Schmitt T, de Groot FMF. *Ang. Chem.* 2013; 1170:3828–3831.
4. a Burkel E. *J. Phys. Cond. Matt.* 2001; 13:7627–7644. b Krisch, M.; Sette, F. *Light Scattering in Solids IX*. Vol. 108. Springer Verlag; Berlin: 2007. p. 317–369. c Baron AQR. *J. Spect. Soc. Japan.* 2009; 58:205–214. (Japanese) arXiv 0910.5764 (English).
5. a Kulda J, Kainzmaier H, Strauch D, Dorner B, Lorenzen M, Krisch M. *Phys. Rev. B.* 2002; 66:241202. b Maultzsch J, Reich S, Thomsen C, Requardt H, Ordejón P. *Phys. Rev. Lett.* 2004; 92:075501. [PubMed: 14995866]
6. a Baron AQR, Uchiyama H, Tanaka Y, Tsutsui S, Ishikawa D, Lee S, Heid R, Bohnen KP, Tajima S, Ishikawa T. *Phys. Rev. Lett.* 2004; 92:197004. [PubMed: 15169439] b Fukui H, Katsura T, Kuribayashi T, Matsuzaki T, Yoneda A, Ito E, Kudoh Y, Tsutsui S, Baron AQR. *J. Synchrot. Radiat.* 2008; 15:618–623. c Strauch D, Dorner B, Ivanov A, Krisch M, Serrano J, Bosak A, Choyke WJ, Stojetz B, Malorny M. *Mater. Sci. Forum.* 2006; 527-529:689–694694.
7. a Sette F, Ruocco G, Krisch M, Bergmann U. *Phys. Rev. Lett.* 1995; 75:850–853. [PubMed: 10060134] b Seyfert C, Simmons RO, Sinn H, Arms DA, Burkel E. *J. Phys. Condens. Matter.* 1999; 11:3501.
8. a Liu Y, Chen SH, Berti D, Baglioni P, Alatas A, Sinn H, Alp E, Said A. *J. Chem. Phys.* 2005; 123:2235–2245. b Liu D, Chu XQ, Lagi M, Zhang Y, Fratini E, Baglioni P, Alatas A, Said A, Alp E, Chen SH. *Phys. Rev. Lett.* 2008; 101:135501. [PubMed: 18851459]
9. Baron AQR, Uchiyama H, Tsutsui S, Tanaka Y, Ishikawa D, Sutter JP, Lee S, Tajima S, Heid R, Bohnen KP. *Physica C.* 2007; 456:83–91.
10. Baron AQR, Tanaka Y, Goto S, Takeshita K, Matsushita T, Ishikawa TJ. *Phys. Chem. Sol.* 2000; 61:461–465.
11. Smith MC, Xiao Y, Wang H, George SJ, Coucovanis D, Koutmos M, Sturhahn W, Alp EE, Zhao J, Cramer SP. *Inorg. Chem.* 2005; 44:5562–5570. [PubMed: 16060605]
12. a Burkel E. *Rep. Prog. Phys.* 2000; 63:171–232. b Sinha SK. *J. Phys. Cond. Matt.* 2001; 13:7511–7523. Fischer I, Bosak A, Krisch M. *Phys. Rev. B.* 2009; 79:134302.




**Figure 1.** Comparison of IXS for  $\text{NEt}_4(\text{FeCl}_4)$  vs. other methods. Top to bottom: IXS for crystal with beam along  $a$ -axis and  $c$ -axis perpendicular to the horizontal scattering plane;  $\theta=27^\circ$  (—),  $^{57}\text{Fe}$  NRVS PVDOS (—),  $^{11}\text{B}$ , FT-IR (—), and Raman (—). Inset: definition of scattering angle  $\theta$  and other quantities.



**Figure 2.**

IXS for various  $Q$  and crystal orientations. Left: Spectra for  $(\chi, \theta)$  at  $(0^\circ, 15^\circ)$  (—),  $(0^\circ, 27^\circ)$  (—),  $(0^\circ, 45^\circ)$  (—), and  $(45^\circ, 27^\circ)$  (—). Right: stretching mode region comparison with Raman spectrum. (All curves normalized to maximum peak height.)



**Figure 3.**

Comparison of experiment and theory for IXS. Top: data for  $\chi=0^\circ$  and  $\theta=27^\circ$ . Middle: calculated IXS for  $\chi=0^\circ$  and different momentum transfers –  $\theta=10^\circ$  (—),  $12^\circ$  (— —),  $24^\circ$  (— — —), and  $28^\circ$  (— — —). Bottom: IXS at  $\theta=27^\circ$  for beam along crystal orientations  $(1\ 1\ 0)$  (—),  $(14\ 1\ 0)$  (— — —),  $(10\ 5\ 0)$  (— — —), and  $(1\ 0\ 0)$  (— — —).