

Correction. In the article "Intramolecular general base-catalyzed ester hydrolyses by the imidazolyl group" by Makoto Komiyama, Thomas R. Roesel, and Myron L. Bender, which appeared in the January 1977 issue of *Proc. Natl. Acad. Sci. USA* 74, 23–25, the authors request that two additional authors be added—namely, Masanori Utaka and Akira Takeda, Department of Synthetics Chemistry, School of Engineering, Okayama University, Tsushima, Okayama-shi 700, Japan. In addition, two deletions should be made. On page 23, "furnished by M. Utaka" should be deleted. On page 25, in the *Acknowledgements*, "We thank Dr. Masanori Utaka for kindly providing the compounds 1–4" should be deleted.

Correction. In the article "Crystallography of new ternary borides" by J. M. Vandenberg and B. T. Matthias, which appeared in the April 1977 issue of *Proc. Natl. Acad. Sci. USA* 74, 1336–1337, a printer's error resulted in an incorrect symbol in the legend for Fig. 1. The figure and correct legend are:




FIG. 1. Proposed structure of YRh_4B_4 . The centers of the Rh_4 tetrahedra are located on the origin and center of the unit cell. Only one of the four B_2 pairs is indicated. ○, Y; ○, Rh; ●, B.

Correction. In the article "Physical-chemical approach to the transient change in Na ion conductivity of excitable membranes" by Philip K. Rawlings and Eberhard Neumann, which appeared in the December 1976 issue of *Proc. Natl. Acad. Sci. USA* 73, 4492–4496, some errors were made by the Proceedings Office. Ref. 3 in the legends of Figs. 1 and 3 should be ref. 4. On page 4495, left-hand column, line 10, the sentence should begin "In ref. 4." In the same column, line 8 up from the bottom, the sentence should read "However, as was found experimentally by Goldman and Schauf, the values of τ_c are uniformly greater than the decay constants τ_h obtained from the declining portions of the curves in Fig. 1." On p. 4496, left-hand column, line 17, the symbol AR_f should be AR_F .

Correction. For the article "Nuclear magnetic resonance studies of the interaction of general anesthetics with 1,2-dihexadecyl-*sn*-glycero-3-phosphorylcholine bilayer" by D. D. Shieh, I. Ueda, H.-C. Lin, and H. Eyring, which appeared in the November 1976 issue of *Proc. Natl. Acad. Sci. USA* 73, 3999–4002, the following should be noted. The compound used to form the bilayer had hexadecanoic acid in ester linkage at glycerol carbons 1 and 2. According to the revised (1976) recommendations of the IUPAC-IUB Commission on Biochemical Nomenclature, the compound can be named 1,2-dihexadecanoyl-*sn*-glycero(3)phosphocholine.

Correction. In the article "Synthesis and activation of mitotic Ca^{2+} -adenosinetriphosphatase during the cell cycle of mouse mastocytoma cells" by C. Petzelt and D. Auel, which appeared in the April 1977 issue of *Proc. Natl. Acad. Sci. USA* 74, 1610–1613, the authors have requested the following change. In Fig. 3 on p. 1612, the units of ATPase specific activity on the ordinate should be nmol P_i /mg of protein per 30 min.