

## Paternal *RHD* zygosity determination in Tunisians: evaluation of three molecular tests

Narjes Kacem<sup>1,2,3</sup>, Saloua Jemni-Yacoub<sup>3</sup>, Jacques Chiaroni<sup>1,2</sup>, Pascal Bailly<sup>1,2</sup>, Monique Silvy<sup>1,2</sup>

<sup>1</sup>French Blood Institute-Alpes-Méditerranée, Marseille; <sup>2</sup>UMR 7268 ADES, Aix-Marseille Université-EFS-CNRS, Marseille, France; <sup>3</sup>Regional Blood Transfusion Centre, Research Unit "UR06SP05", Sousse, Tunisia

**Background.** The choice of a molecular test for first intention determination of paternal *RHD* zygosity, before entering into invasive diagnostics, is important for the management of pregnancies at risk of haemolytic disease of the foetus and newborn related to anti-RhD.

**Materials and methods.** *RHD* zygosity was evaluated in 370 RH:1 Tunisian donors by polymerase chain reaction - sequence-specific polymorphism (PCR-SSP) analysis and polymerase chain reaction - restriction fragment length polymorphism (PCR-RFLP) amplification of hybrid *Rhesus box* and by real time quantitative polymerase chain reaction (RQ-PCR) specific for *RHD* exon 5. To evaluate the accuracy of molecular tests in the cases of discordant results, the ten exons of *RHD* and *Rhesus boxes* were amplified by PCR and sequenced.

**Results.** Molecular investigations revealed that our 370 donors comprise 193 dizygous and 145 hemizygous individuals and 32 subjects whose zygosity remains unknown. Positive predictive values were higher than 99% for all the methods, reaching 100% for RQ-PCR. Negative predictive values were 83.24%, 87.27% and 98% for PCR-SSP, PCR-RFLP and RQ-PCR respectively. This study also revealed 19 novel *Rhesus box* polymorphisms and three novel *RHD* alleles: *RHD*(Trp185Stop), *RHD*(Ala176Thr) and *RHD*(Ile342Ile).

**Discussion.** RQ-PCR is the most convenient method for first intention determination of paternal *RHD* zygosity in Tunisians. However, taking into account positive and negative predictive values, PCR-RFLP could be an alternative despite the heterogeneity of *Rhesus boxes* and the complexity of *RHD*.

**Keywords:** zygosity, *RHD* alleles, *Rhesus box* polymorphisms.

### Introduction

RhD is a complex blood group antigen and anti-D has been implicated in haemolytic disease of the foetus and newborn (HDFN). Adoption of antenatal and postpartum use of Rh immune globulin in industrialised countries has resulted in a major decrease in the frequency of this disease<sup>1</sup>. In Tunisia, HDFN due to RhD immunisation is currently prevented in the vast majority of cases by administration of anti-D immunoglobulin to D-negative women within 72 hours of delivery of a RhD-positive neonate as well as in cases of abortion<sup>2</sup>. Systematic antenatal prophylaxis could enhance the prevention for all pregnant women but this strategy could create a shortage in anti-D immunoglobulin and carry significant potential costs. In order to develop an antenatal prophylaxis limited to fetuses that are potentially at risk of HDFN related to anti-D, it has been proposed that paternal zygosity could be determined as the first step in the management of the red cell alloimmunised pregnancy<sup>3</sup>. If the father is found to have a heterozygous genotype, genetic testing could be undertaken, through amniocentesis, to determine whether the foetus is at

risk of foetal anaemia. The overall sensitivity and specificity of polymerase chain reaction (PCR) typing on amniotic fluid has been reported to be 99.5% and 98.6%, respectively, and both positive and negative predictive values were 99.1%<sup>4</sup>. However this invasive procedure is risky and might cause further sensitisation in RhD-negative women. *RHD* zygosity was once determined through serological testing using population statistics. Ethnicity played a major role in these calculations and recently molecular tests, which circumvent this issue, were shown to be more accurate<sup>5,6</sup>. A general problem concerning determination of the zygosity of the father is non-paternity. However, in a conservative society such as Tunisia, non-paternity is unlikely. The newest technology used in the determination of the foetal RhD type in the case of heterozygous paternity involves free foetal DNA in the maternal circulation. Based on the concept that cell-free tumour DNA could be found in the peripheral circulation of patients with cancer, Lo *et al.*<sup>7</sup> were the first to report the presence of the *RHD* gene in the plasma of women pregnant with a RhD-positive foetus. Detection of foetal *RHD* in maternal

plasma is used as a non-invasive method for assessing the risk of HDFN, but a remaining pitfall that hampers its use is the limited reliability of negative results since, without an internal control, true negative results cannot be distinguished from false negative results due to insufficient amounts of free foetal DNA. A recent report described the use of single nucleotide polymorphisms as internal controls, but a large study using this strategy is lacking<sup>8</sup>. Moreover, non-invasive antenatal diagnostic testing to target anti-D prophylaxis was shown to be unlikely to produce important clinical benefits and its reliability in different ethnic minority populations needs to be demonstrated rigorously<sup>9</sup>.

Since *RHD* is a very polymorphic gene attention should be paid to the specificity and sensitivity of molecular tests. In Caucasians, the most common mechanism of RH:-1 phenotype is deletion of the *RHD* gene<sup>10</sup> occurring as a result of unequal crossing-over between the two *Rhesus boxes* flanking the *RHD* gene. This leaves a single hybrid *Rhesus box* as a target for *RHD* zygosity testing and detection of this hybrid *Rhesus box* has been used to demonstrate the presence of the *RHD* deletion, making it possible to distinguish *RHD* dizygous individuals from *RHD* hemizygous ones<sup>11</sup>. Different approaches to detect the hybrid *Rhesus box* have been described, but these molecular methods may lead to false results in Africans because of their large genetic diversity<sup>12</sup>.

*RHD* deletion was shown to be the main background of the RH:-1 phenotype in the Tunisian population<sup>13</sup> suggesting that paternal *RHD* zygosity in Tunisians could be determined using molecular methods that are accurate in Caucasians. To identify the most accurate and reliable method for *RHD* zygosity assignment in Tunisians, *RHD* zygosity was determined in 405 Tunisian donors. Three molecular methods, based

on polymerase chain reaction - sequence-specific polymorphism (PCR-SSP) analysis, polymerase chain reaction - restriction fragment length polymorphism (PCR-RFLP) amplification and real-time, quantitative polymerase chain reaction (RQ-PCR) specific for the *RHD* exon 5, were compared.

## Materials and methods

### Blood sampling, serological typing and DNA extraction


A total of 405 random EDTA blood samples were collected from blood donors recruited with informed consent according to the approved protocol of the Regional Blood Transfusion Centre of Sousse (Tunisia) to determine *RHD* zygosity. RhD phenotyping was routinely done as described in our previous study<sup>6</sup> and the indirect antiglobulin test was performed systematically for apparently RH:-1 results. Genomic DNA was isolated from buffy coats by a salting-out method according to a standard protocol<sup>14</sup>.

### Polymerase chain reaction - sequence-specific polymorphism analysis for the hybrid *Rhesus box*

A 2778 bp product was amplified using forward primer u1-s<sup>15</sup>, which is specific for the hybrid and upstream *Rhesus boxes*, and reverse primer rnb31, which is specific for the hybrid and downstream *Rhesus boxes* (Figure 1). The PCR was performed as previously described<sup>6</sup>.

### Polymerase chain reaction - restriction fragment length polymorphism analysis for hybrid and downstream *Rhesus boxes*

The PCR-RFLP method was performed as described by Wagner and Flegel<sup>11</sup>. A 3,029 bp product was amplified using primers rez7 (universal primer for all *Rhesus boxes*) and rnb31 for amplification of the



**Figure 1** - Localisation of primers in the hybrid *Rhesus box*. Nucleotide 5250 represents the location of the *Pst*I site used in the PCR-RFLP. Nucleotide positions are given corresponding to the standard downstream *Rhesus box* where position 1 represents the start of the homology between upstream and downstream *Rhesus boxes* and corresponds to nucleotide number 23 in GenBank accession number AJ252312.

downstream and/or hybrid *Rhesus boxes* (Figure 1). The PCR amplicon was digested with 1.5 U/ $\mu$ L of *Pst* I (Fermentas, St. Leon-Rot, Germany) overnight at 37 °C and fragments were resolved using a 2% (w/v) agarose gel.

### Real-time quantitative polymerase chain reaction analysis for *RHD* exon 5

Primers and probes for *RHD* exon 5 and *CCR5* genes were from Finning<sup>16</sup>. The probes for *RHD* exon 5 and *CCR5* were labelled with 6-FAM and Yakima yellow, respectively. Each primers was used at a final concentration of 300 nM and probes specific for *RHD* exon 5 and *CCR5* were used at a final concentration of 200 nM and 100 nM, respectively. After 10 minutes at 95 °C, the reaction consisted of 40 cycles (95 °C for 10 sec and 60 °C for 30 sec). The MagNA Pure LC system (Roche Diagnostics, Indianapolis, IN, USA) was used to dispense master mix (TaqMan<sup>®</sup>, Applied Biosystems, Courtabœuf, France) and DNA (100 ng) in triplicate. Relative quantification ( $\Delta$ Cq) was done with *CCR5* as the control gene and compared to results of control samples with one or two copies of the *RHD* gene<sup>17</sup>.

### Genomic analysis of the *RHD* gene

To determine zygosity in discordant samples the ten exons of *RHD* were amplified by PCR and sequenced. Each exon amplification was performed using the primer sets described by Touinssi *et al.*<sup>18</sup>.

### Sequencing analyses of the *Rhesus boxes*

With the aim of explaining discordant results, *Rhesus boxes* were amplified by PCR and sequenced using two PCR sets (Figure 1): PCR1 was applied with primers CRBF06 and CRBR08<sup>12</sup> and PCR products were sequenced with the same couple of primers; PCR2 was done with Rez7 and CRBR02 primers<sup>12</sup> and PCR products were sequenced with Rez7 and CRBF04-R (5'cctctgccaggcagtgca3') primers.

### Statistical analysis

The positive predictive value (PPV) was calculated for samples with no hybrid *Rhesus Box* detected by PCR-SSP and PCR-RFLP and for samples with two expressed alleles by RQ-PCR using the following formula:

$$PPV = \frac{\text{Number of samples with two non-silent } RHD \text{ alleles}}{\text{Number of samples with two non-silent } RHD \text{ alleles} + \text{Number of samples with one non-silent } RHD \text{ alleles}}$$

The negative predictive value (NPV) was calculated for samples with one hybrid *Rhesus Box* detected by PCR-SSP and PCR-RFLP and for samples with one expressed alleles by RQ-PCR based on the same formula as that for the calculation of the PPV.

## Results

### RhD phenotype

RhD phenotyping of 405 donors randomly recruited from the centre of Tunisia revealed that 370 had the RH:1 phenotype and 35 had the RH:-1 phenotype. All RH:-1 phenotypes were confirmed through indirect antiglobulin testing.

### Analysis of the RH:-1 samples

The RH:-1 samples (n=35) were used as controls for the PCR-SSP, PCR-RFLP and RQ-PCR analyses. PCR-SSP detected the hybrid *Rhesus box* in all RH:-1 samples. PCR-RFLP showed homozygous *RHD* gene deletion in 32 RH:-1 samples and three samples containing a single copy of hybrid *Rhesus box*. Sequencing of these samples showed a *DIIIa-CE(4-7)-D*, a *weak D type 4.0* (omitted by serological techniques), and a new allele with a nucleotide change in *RHD* exon 7 at position 1026C/T (*RHD* [Ile342Ile]). According to the RQ-PCR analysis, *RHD* exon 5 was absent in 33 samples and present in *weak D type 4.0* and *RHD* (Ile342Ile).


### Analysis of the RH:1 samples

The comparison between the three methods for *RHD* zygosity assignment showed concordant results in 303 samples (81.9%). Discordant results (n=67, 18.1%) were classified into three groups (Figure 2). Group 1 (n=31) consisted of samples with results obtained by PCR-SSP similar to those obtained by PCR-RFLP but different from those obtained with RQ-PCR. Twenty-six samples had one copy of *RHD* gene by both PCR-SSP and PCR-RFLP and two copies by RQ-PCR, and five had two copies of *RHD* gene by both PCR-SSP and PCR-RFLP and one copy by RQ-PCR. Group 2 contained 12 samples with discordant PCR-SSP and RQ-PCR findings and without results in PCR-RFLP. We distinguished four samples in which the zygosity status differed in one copy of *RHD* gene by PCR-SSP and eight samples showing two copies. Group 3 consisted of 24 samples whose results obtained by PCR-RFLP were similar to those obtained by RQ-PCR but differed from those obtained by PCR-SSP.

### *RHD* sequencing results

*RHD* sequencing of the ten exons in 67 discordant samples are shown in Figure 2. All the *RHD* variants observed were heterozygous in *trans* to conventional *RHD*. We identified three new variants: (i) *RHD*(Trp185Stop): a G>A transition at position 555 leading to a stop codon; (ii) *RHD*(Ile342Ile): a 1026C>T transition in *RHD* exon 7, and (iii) *RHD*(Ala176Thr): a G>A transition at position 526 in *RHD* exon 4.

We also identified a probable *weak D type 4.0* (nucleotide changes: 667T>G in *RHD* exon 5, 819G>A



**Figure 2** - Classification and analysis of discordant results.

in *RHD* exon 6 because *RHD* exon 2 and 4 amplifications failed). We found homo/hemizygous intronic single nucleotide polymorphisms (n=10) and heterozygous intronic single nucleotide polymorphisms (n=18). We were unable to identify seven samples, probably because of the quality of the DNA.

In conclusion, zygosity remained unknown for 37 subjects: homo/hemizygous intronic single nucleotide polymorphisms (n=10), no single nucleotide polymorphisms (n=20), and genotyping failure (n=7). Therefore, 189 subjects were dizygous and 144 were hemizygous.

### **Rhesus boxes polymorphisms for RH:1 discordant samples**

Since *RHD* genotyping failed in seven samples, only the remaining 60 discordant samples were analysed for *Rhesus box* polymorphisms by PCR1 and/or PCR2 (Figure 2). No single nucleotide polymorphism was observed at the 5' part of *Rhesus boxes* in three samples (5%, with *RHD/RHD*(Ala176Thr), heterozygous intronic single nucleotide polymorphism in IVS1 and in IVS1+IVS8), whereas *Rhesus boxes* were mutated in 57 samples (95%). Table I summarises the different nucleotide positions of mutated *Rhesus boxes* in RH:1 discordant samples. In addition to single nucleotide

polymorphisms reported by Wagner *et al.*<sup>19</sup>, we identified 19 new polymorphisms in *Rhesus boxes*, listed and underlined in Table I.

Observation of polymorphisms under *rnb31* led us to assign a hemizygous status to one subject and a dizygous status to four subjects whose zygosity was previously unknown. Thus, zygosity remained unknown for 32 subjects because of homo/hemizygous intronic single nucleotide polymorphisms (n=7), genotyping failure (n=7) and no single nucleotide polymorphism in the *RHD* gene (n=18). Of the remaining 338 samples, 193 were dizygous and 145 hemizygous.

### **Positive and negative predictive values of the three methods**

The PPV and NPV were calculated for each technique using the 338 samples with known zygosity (Table II). The PPV were higher than 99% for all methods, reaching 100% for RQ-PCR. The NPV were 83.24%, 87.27% and 98% for PCR-SSP, PCR-RFLP and RQ-PCR, respectively.

### **Discussion**

Antibodies to the RhD antigen can be produced during pregnancy in a RhD-negative mother carrying

**Table 1** - Summary of mutated sequences of *Rhesus* boxes in discordant samples. Discrepant samples were compared with standard downstream and upstream *Rhesus* boxes.

SNP or allele	Nucleotide positions																											
	ul-s														rnb31													
	4929	4952	4978	5139	5265	5157	5177	7302	7315	7317	7320	7328	7331	7332	7359	7381	7394	7424	7454	7429	7555	7658	7689	7691	7695	7706	7712	
<b>Standard DRB</b>	G	A	A	T	A	G	C	A	A	G	T	A	A	G	A	G	G	C	C	A	G	del	A	C	A	C	A	C
<b>Standard URB</b>	A	C	A	C	A	G	C	A	A	G	T	A	A	G	A	G	C	C	C	A	G	AT	A	C	C	A	C	C
<i>RHD/DIIIa-CE(4-7)-D</i>	1	nd	nd	nd	nd	nd	nd	-	-	-	-	R	R	R	C	C	-	-	-	-	AT	R	M	C	M	M	M	
<i>RHD/weak D type 4.0</i>	1	-	-	C	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>RHD/probable weak D type 4.0</i>	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>RHD/RHDψ</i>	1	-	-	-	W	-	-	nd	nd	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
<i>RHD/DAU-0 and DAU-0/DFV</i>	2	A	-	C	-	-	Y	nd	nd	nd	nd	nd	nd	nd	nd	Nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
<i>RHD/DAU-3</i>	1	A	-	-	-	-	Y	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>RHD/D674</i>	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>RHD/RHD (Trp185Stop)</i>	1	-	-	C	-	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
<i>RHD/RHD (Ile342Ile)</i>	1	-	-	C	-	R	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
Hetero in IVS1	1	-	-	C	-	W	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
Homo/hemi in IVS1	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Hetero in IVS2	1	nd	nd	nd	nd	nd	nd	R	-	S	K	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Homo/hemi in IVS2	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	R	C	C	C	M	Y	-	-	-	-	-	-	-	-	
Homo/hemi in IVS5	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	R	-	-	-	-	Y	-	-	-	-	-	-	-	-	
Hetero in IVS8	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	
Homo/hemi in IVS8	1	-	-	-	-	W	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Hetero in IVS1, Homo/hemi in IVS1, Hetero in IVS2 and Homo/hemi in IVS4	4	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
Hetero in IVS0+IVS8	2	-	-	C	-	-	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
Hetero in IVS1+IVS8	1	nd	nd	nd	nd	nd	nd	-	R	-	K	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Homo/hemi in IVS1+IVS8, Hetero in IVS2 and in IVS5+IVS8	3	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Hetero in IVS2+IVS8	1	-	A	-	-	-	Y	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Hetero in IVS3+IVS8 and in IVS8	2	A	-	-	-	-	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
Hetero in IVS1+IVS8	1	A	-	-	C	-	-	-	-	-	-	-	R	R	C	C	C	-	Y	-	-	-	-	-	-	-	-	
Hetero in IVS1+IVS8	1	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	
No SNP	1	-	-	W	C	-	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
	1	-	-	C	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	2	-	-	C	-	-	-	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	2	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	R	-	-	-	-	-	-	-	-	-	-	-	-	-	
	3	nd	nd	nd	nd	nd	nd	-	-	-	-	-	R	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	9	-	-	C	-	-	-	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	

Positions of no described nucleotides changes of mutated *Rhesus* boxes are underlined; grey columns represent nucleotide positions with mutations under mb31 and/or ul-s primers. Homo/hemi: homozygous or hemizygous; Hetero: heterozygous. DRB: downstream *Rhesus* box; URB: upstream *Rhesus* box; SNP: single nucleotide polymorphism; "del": indicates a two-nucleotide deletion; "-" indicates that no mutation was noted; "nd": not determined.

**Table II** - Positive and negative predictive values and Youden index for the three molecular methods.

	<i>HDFN risk</i>	Two non-silent <i>RHD</i> alleles	One non-silent <i>RHD</i> allele	PPV	NPV	Youden index
		High (100%)	Moderate (50%)			
PCR-SSP	Need for amniocentesis	No	Yes	99.39	83.24	0.84
	No hybrid <i>Rhesus box</i> detected	164	1			
	One hybrid <i>Rhesus box</i> detected	29	144			
PCR-RFLP	No hybrid <i>Rhesus box</i> detected	167	1	99.4	87.27	0.88
	One hybrid <i>Rhesus box</i> detected	21	144			
RQ-PCR	Two expressed alleles	188	0	100	98	0.98
	One expressed alleles	3	147			

Positive predictive value (PPV), negative predictive value (NPV) and Youden index were calculated using: [http://www.aly-abbara.com/utilitaires/statistiques/sensibilite\\_specificite\\_vpp\\_vpn.html](http://www.aly-abbara.com/utilitaires/statistiques/sensibilite_specificite_vpp_vpn.html).

a RhD-positive foetus, in particular following foetal-maternal haemorrhage at birth. While the first baby is usually not harmed, these antibodies may cause HDFN in subsequent RhD-positive babies. RhD incompatibility is a major cause of HDFN. Establishing paternal *RHD* zygosity to determine the risk of HDFN related to anti-D has the advantage of predicting the risk of HDFN in future pregnancies. Indeed, if the father is homozygous for the *RHD* gene (dizygous), the chance of inheriting the *RHD* gene is 100% for each pregnancy. Conversely, in the case of a hemizygous father (*RHD/d*) the risk is 50%. The aim of the present study was to assess the most convenient molecular method for determination of *RHD* zygosity in the Tunisian population as a method of first intention for the evaluation of HDFN risk.

Analysis of 370 random RhD-positive Tunisians using three molecular tests showed 81.9% concordant results and through investigation of *RHD* alleles and *Rhesus box* polymorphisms in the 67 discordant samples we concluded that our cohort consisted of 193 dizygous subjects, 145 hemizygous individuals and 32 subjects whose zygosity remains unknown because of lack of heterozygous single nucleotide polymorphisms or technical failure. Based on the 338 samples with known zygosity, the PPV, NPV and Youden index were calculated for each molecular test. The highest Youden index was observed for RQ-PCR (0.98) showing that this method was the most reliable and convenient for the Tunisian population. However, taking into account the high PPV (99.4%) and despite the lower NPV (87.27%) of PCR-RFLP, this method could be an alternative to the use of RQ-PCR for determining *RHD* zygosity in Tunisia. If the father is found to be hemizygous, second intention invasive methods would then be used to evaluate the risk of HDFN. Thus, in cases of falsely labelled hemizygous fathers, invasive investigations

will be performed uselessly correcting the false zygosity assignment of the first intention test. Since some *RHD* variant alleles in the Tunisian population were shown to be related to some in Caucasians, we suggest that the false negative results using RQ-PCR (n=3) could be linked to presence in *trans* to a conventional *RHD* allele of a hybrid *RHD-CE-D* gene, such as *DVI type 1* or *2* which are the most frequent partial alleles encountered in Caucasians<sup>20,21</sup>. In the heterozygous state such a hybrid gene cannot be distinguished by either haemagglutination tests or DNA analysis. False positive and false negative results using PCR-SSP and PCR-RFLP were the result of: (i) mutations in *Rhesus boxes* leading to lack of amplification of hybrid *Rhesus boxes*, and (ii) the presence of non-functional *RHD* alleles from African origin leading to an incorrect estimation of the risk of HDFN.

The present study also led to the identification of three novel *RHD* alleles: *RHD*(Trp185Stop), *RHD*(Ala176Thr) and *RHD*(Ile342Ile). Because of the premature stop, *RHD*(Trp185Stop) is predicted to be a silent allele which is associated with over-estimation of HDFN risk in the context of zygosity determination. *RHD*(Ala176Thr) is predicted to encode a weak phenotype since amino-acid 176 is located in the fifth transmembrane helix<sup>22</sup>. However, since these alleles were in *trans* to conventional *RHD*, no serological investigations were performed and we have no certainty about the RhD phenotype associated with these alleles. Allele encoding *RHD*(Ile342Ile) was found in two samples. Despite the silent nature of the polymorphism one of the donors bearing this allele typed as RhD-negative. Unfortunately, no blood sample was available for verification of serological results and no conclusion can be made based only on the molecular analysis. This study also demonstrated the complexity of *Rhesus boxes*

with the description of 19 novel polymorphisms. The heterogeneity of *Rhesus boxes* highlights the limitations of *Rhesus box* approaches to the determination of *RHD* zygosity.

Altogether, the results of our analysis of 338 random samples with known zygosity demonstrated that RQ-PCR can accurately assign zygosity status of samples in Tunisians. In the case of determination of paternal zygosity, RQ-PCR enables prediction of the risk that a foetus will inherit *RHD* and will be useful in identifying HDFN cases which may require further genotyping.

### Acknowledgements

We thank all blood donors at the Regional Blood Transfusion Centre in Sousse (Tunisia) who made this work for a doctoral thesis possible. We acknowledge the team from EFS-AM (the French Blood Institute), especially Sophie Beley and Thomas Granier for their technical help.

### Authorship contribution

Narjes Kacem and Monique Silvy performed the research study and wrote the paper. Narjes Kacem, Monique Silvy and Pascal Bailly analysed the data. Monique Silvy, Pascal Bailly, Saloua Jemni-Yacoub and Jacques Chiaroni contributed essential reagents and tools.

*The Authors declare no conflict of interest.*

### References

- 1) Moise KJ Jr, Argoti PS. Management and prevention of red cell alloimmunization in pregnancy: a systematic review. *Obstet Gynecol* 2012; **120**: 1132-9.
- 2) Moussa H, Tsochandaridis M, Jemni-Yacoub S, et al. Fetal RhD genotyping by real time quantitative PCR in maternal plasma of RhD-negative pregnant women from the Sahel of Tunisia. *Ann Biol Clin* 2012; **70**: 683-8.
- 3) Moise KJ Jr. Management of rhesus alloimmunization in pregnancy. *Obstet Gynecol* 2008; **112**:164-76.
- 4) Goebel JC, Soergel P, Pruggmayer M, et al. Prenatal diagnosis of the Rhesus D fetal blood type on amniotic fluid in daily practice. *Arch Gynecol Obstet* 2008; **277**: 155-60.
- 5) Pirelli KJ, Pietz BC, Johnson ST, et al. Molecular determination of RHD zygosity: predicting risk of hemolytic disease of the fetus and newborn related to anti-D. *Prenat Diagn* 2010; **30**: 1207-12.
- 6) Kacem N, Chakroun T, Moussa H, et al. RHD zygosity assignments based on most probable genotype and hybrid Rhesus box detection in Tunisia. *Transfus Med* 2012; **22**: 362-6.
- 7) Lo YM, Corbetta N, Chamberlain PF, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997; **350**: 485-7.
- 8) Doescher A, Petershofen EK, Wagner FF, et al. Evaluation of single-nucleotide polymorphisms as internal controls in prenatal diagnosis of fetal blood groups. *Transfusion* 2013; **53**: 353-62.

- 9) Szczepura A, Osipenko L, Freeman K. A new fetal RHD genotyping test: costs and benefits of mass testing to target antenatal anti-D prophylaxis in England and Wales. *BMC Pregnancy Childbirth* 2011; **11**: 5.
- 10) Colin Y, Chérif-Zahar B, Le Van Kim C, et al. Genetic basis of the RhD-positive and RhD-negative blood group polymorphism as determined by Southern analysis. *Blood* 1991; **78**: 2747-52.
- 11) Wagner FF, Flegel WA. RHD gene deletion occurred in the Rhesus box. *Blood* 2000; **95**: 3662-8.
- 12) Grootkerk-Tax MG, Maaskant-van Wijk PA, van Drunen J, et al. The highly variable RH locus in nonwhite persons hampers RHD zygosity determination but yields more insight into RH-related evolutionary events. *Transfusion* 2005; **45**: 327-37.
- 13) Moussa H, Tsochandaridis M, Chakroun T, et al. Molecular background of D-negative phenotype in the Tunisian population. *Transfus Med* 2012; **22**: 192-8.
- 14) Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; **16**: 1215.
- 15) Perco P, Shao CP, Mayr WR, et al. Testing for the D zygosity with three different methods revealed altered Rhesus boxes and a new weak D type. *Transfusion* 2003; **43**: 335-9.
- 16) Finning KM, Martin PG, Soothill PW, et al. Prediction of fetal D status from maternal plasma: introduction of a new non-invasive fetal RHD genotyping service. *Transfusion* 2002; **42**: 1079-85.
- 17) Silvy M, Chapel-Fernandes S, Beley S, et al. Molecular characterization of a new D- - haplotype in a Comorian man. *Vox Sang* 2012; **103**: 352-5.
- 18) Touinssi M, Chapel-Fernandes S, Granier T, et al. Molecular analysis of inactive and active RHD alleles in native Congolese cohorts. *Transfusion* 2009; **49**: 1353-60.
- 19) Wagner FF, Moulds JM, Flegel WA. Genetic mechanisms of Rhesus box variation. *Transfusion* 2005; **45**: 338-44.
- 20) Muller TH, Wagner FF, Trockenbacher A, et al. PCR screening for common weak D types shows different distributions in three Central European populations. *Transfusion* 2001; **41**: 45-52.
- 21) Roubinet F, Apoil PA, Blancher A. Frequency of partial D phenotypes in the south western region of France. *Transfus Clin Biol* 1996; **3**: 247-55.
- 22) Silvy M, Barrault A, Velliquette RW et al. RHCE\*cE734C allele encodes an altered c antigen and a suppressed E antigen not detected with standard reagents. *Transfusion* 2013; **53**: 955-61.

---

Arrived: 11 November 2013 - Revision accepted: 11 February 2014

**Correspondence:** Monique Silvy  
Établissement Français du Sang-Alpes-Méditerranée  
149 Boulevard Bailly  
13392 Marseille Cedex, France  
e-mail: monique.silvy@efs.sante.fr

---