



Published in final edited form as:

*JAMA Neurol.* 2015 February 1; 72(2): 209–216. doi:10.1001/jamaneurol.2014.2157.

## Rarity of the Alzheimer Disease–Protective *APP* A673T Variant in the United States

Li-San Wang, PhD, Adam C. Naj, PhD, Robert R. Graham, PhD, Paul K. Crane, MD, MPH, Brian W. Kunkle, PhD, MPH, Carlos Cruchaga, PhD, Josue D. Gonzalez Murcia, BS, Lisa Cannon-Albright, PhD, Clinton T. Baldwin, PhD, Henrik Zetterberg, MD, PhD, Kaj Blennow, MD, Walter A. Kukull, PhD, Kelley M. Faber, MS, Nicole Schupf, PhD, DrPH, Maria C. Norton, PhD, JoAnn T. Tschanz, PhD, Ronald G. Munger, MPH, PhD, Christopher D. Corcoran, PhD, Ekaterina Rogaeva, PhD, Alzheimer's Disease Genetics Consortium, Chiao-Feng Lin, PhD, Beth A. Dombroski, PhD, Laura B. Cantwell, MPH, Amanda Partch, MS, Otto Valladares, MS, Hakon Hakonarson, MD, PhD, Peter St George-Hyslop, MD, FRCP, Robert C. Green, MD, MPH, Alison M. Goate, DPhil, Tatiana M. Foroud, PhD, Regina M. Carney, MD, Eric B. Larson, MD, MPH, Timothy W. Behrens, MD, John S. K. Kauwe, PhD, Jonathan L. Haines, PhD, Lindsay A. Farrer, PhD, Margaret A. Pericak-Vance, PhD, Richard Mayeux, MD, and Gerard D. Schellenberg, PhD for the National Institute on Aging–Late-Onset Alzheimer's Disease (NIA-LOAD) Family Study

Department of Pathology and Laboratory Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia (Wang, Lin, Dombroski, Cantwell, Partch, Valladares, Schellenberg); Department of Biostatistics and Epidemiology, University of Pennsylvania Perelman School of Medicine, Philadelphia (Naj); Department of Human Genetics, Genentech Inc, South San Francisco, California (Graham, Behrens); Department of Medicine, University of Washington, Seattle (Crane, Larson); John P. Hussman Institute for Human Genomics, University of Miami, Miami, Florida (Kunkle, Pericak-Vance); Department of Psychiatry, Washington University School of Medicine, St Louis, Missouri (Cruchaga, Goate); Hope Center Program on Protein Aggregation and Neurodegeneration, Washington University School of Medicine, St Louis, Missouri (Cruchaga, Goate); Department of Biology, Brigham Young University, Provo, Utah (Murcia, Kauwe, Farrer); Division of Genetic Epidemiology, Department of Medicine, University of Utah School of Medicine, Salt Lake City (Cannon-Albright); George E. Wahlen Department of Veterans Affairs Medical Center, Salt Lake City, Utah (Cannon-Albright); Biomedical Genetics, Department of Medicine, Boston University, Boston, Massachusetts (Baldwin); Institute of Neurology, University College London, London, England (Zetterberg); Clinical Neurochemistry Laboratory, Institute of Neuroscience and Physiology, Department of Psychiatry and Neurochemistry, Sahlgrenska Academy at University of Gothenburg, Sahlgrenska University Hospital, Mölndal, Sweden (Zetterberg, Blennow); Department of Epidemiology, University of Washington, Seattle (Kukull); Department of Medical and Molecular Genetics, Indiana University, Indianapolis (Faber, Foroud); Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York (Schupf); Taub Institute for Research on Alzheimer's Disease and the Aging

Copyright 2014 American Medical Association. All rights reserved.

Corresponding Author: Gerard D. Schellenberg, PhD, University of Pennsylvania Perelman School of Medicine, 422 Curie Blvd, Stellar Chance Laboratories, 609B, Philadelphia, PA 19104 (gerardsc@mail.med.upenn.edu).

Supplemental content at [jamaneurology.com](http://jamaneurology.com)

Brain, Columbia University, New York, New York (Schupf, Mayeux); Gertrude H. Sergievsky Center, Columbia University, New York, New York (Schupf, Mayeux); Department of Family, Consumer, and Human Development, Utah State University, Logan (Norton); Department of Psychology, Utah State University, Logan (Norton, Tschanz); Department of Nutrition, Dietetics, and Food Sciences, Utah State University, Logan (Munger); Department of Mathematics and Statistics, Utah State University, Logan (Corcoran); Tanz Centre for Research in Neurodegenerative Disease, University of Toronto, Toronto, Ontario, Canada (Rogaeva, St George-Hyslop); Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania (Hakonarson); Cambridge Institute for Medical Research, Department of Clinical Neurosciences, University of Cambridge, Cambridge, England (St George-Hyslop); Division of Genetics, Department of Medicine and Partners Center for Personalized Genetic Medicine, Brigham and Women's Hospital/ Harvard Medical School, Boston, Massachusetts (Green); Department of Psychiatry and Behavioral Sciences, Miller School of Medicine, University of Miami, Miami, Florida (Carney); Group Health Research Institute, Seattle, Washington (Larson); Center for Human Genetics and Research, Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee (Haines); Department of Biostatistics, Boston University, Boston, Massachusetts (Farrer); Department of Ophthalmology, Boston University, Boston, Massachusetts (Farrer); Department of Neurology, Boston University, Boston, Massachusetts (Farrer); Department of Epidemiology, Boston University, Boston, Massachusetts (Farrer); Dr John T. Macdonald Foundation Department of Human Genetics, University of Miami, Miami, Florida (Pericak-Vance); Department of Neurology, Columbia University, New York, New York (Mayeux)

## Abstract

**IMPORTANCE**—Recently, a rare variant in the amyloid precursor protein gene (*APP*) was described in a population from Iceland. This variant, in which alanine is replaced by threonine at position 673 (A673T), appears to protect against late-onset Alzheimer disease (AD). We evaluated the frequency of this variant in AD cases and cognitively normal controls to determine whether this variant will significantly contribute to risk assessment in individuals in the United States.

**OBJECTIVE**—To determine the frequency of the *APP* A673T variant in a large group of elderly cognitively normal controls and AD cases from the United States and in 2 case-control cohorts from Sweden.

**DESIGN, SETTING, AND PARTICIPANTS**—Case-control association analysis of variant *APP* A673T in US and Swedish white individuals comparing AD cases with cognitively intact elderly controls. Participants were ascertained at multiple university-associated medical centers and clinics across the United States and Sweden by study-specific sampling methods. They were from case-control studies, community-based prospective cohort studies, and studies that ascertained multiplex families from multiple sources.

**MAIN OUTCOMES AND MEASURES**—Genotypes for the *APP* A673T variant were determined using the Infinium HumanExome V1 Beadchip (Illumina, Inc) and by TaqMan genotyping (Life Technologies).

**RESULTS**—The A673T variant genotypes were evaluated in 8943 US AD cases, 10 480 US cognitively normal controls, 862 Swedish AD cases, and 707 Swedish cognitively normal

controls. We identified 3 US individuals heterozygous for A673T, including 1 AD case (age at onset, 89 years) and 2 controls (age at last examination, 82 and 77 years). The remaining US samples were homozygous for the alanine (A673) allele. In the Swedish samples, 3 controls were heterozygous for A673T and all AD cases were homozygous for the A673 allele. We also genotyped a US family previously reported to harbor the A673T variant and found a mother-daughter pair, both cognitively normal at ages 72 and 84 years, respectively, who were both heterozygous for A673T; however, all individuals with AD in the family were homozygous for A673.

**CONCLUSIONS AND RELEVANCE**—The A673T variant is extremely rare in US cohorts and does not play a substantial role in risk for AD in this population. This variant may be primarily restricted to Icelandic and Scandinavian populations.

The amyloid precursor protein gene (*APP*; GenBank NC\_011512) encodes a transmembrane protein of unknown normal function. In the normal processing of the APP protein, proteolytic cleavage yields a 39- to 43-amino acid peptide called  $\beta$ -amyloid ( $A\beta$ ). Release of  $A\beta$  from APP is catalyzed by the  $\beta$ -site APP cleaving enzyme 1 (BACE1) that cleaves APP at the N-terminal end of  $A\beta$  and by the  $\gamma$ -secretase complex that cleaves at the C-terminal end.<sup>1</sup> Together, these 2 proteases generate the  $A\beta$  peptide. In Alzheimer disease (AD),  $A\beta$  accumulates in extracellular amyloid plaques that are characteristic of this disease. Mutations in *APP* cluster around the N-terminal and C-terminal sequences that encode  $A\beta$  and cause early-onset AD. Likewise, early-onset AD is also caused by mutations in the presenilin 1 gene (*PSEN1*) and presenilin 2 gene (*PSEN2*), which encode protease subunits of the  $\gamma$ -secretase complex.<sup>1</sup> Genetic studies and a large body of functional evidence convincingly show that  $A\beta$  is a toxic molecule critical to the pathogenesis of AD. As a result, multiple drug trials are in progress designed to stimulate  $A\beta$  clearance using immunological approaches or to inhibit  $A\beta$  production using small-molecule inhibitors of  $\gamma$ -secretase or BACE1.<sup>2-5</sup>

Recently, Jonsson et al<sup>6</sup> reported that the *APP* coding mutation A673T, in which alanine is replaced by threonine at position 673, is protective against late-onset AD. This rare variant was enriched in Icelandic elderly controls compared with AD cases from the same population. The frequency was 0.13% in AD cases and ranged from 0.45% to 0.79% in controls, depending on age. The A673T variant was also observed in an individual with ischemic cerebrovascular disease but not AD<sup>7</sup> and in a 104-year-old patient with dementia who had hippocampal sclerosis and little  $A\beta$  accumulation.<sup>8</sup> The A673T variant is located at position 2 of  $A\beta$  and thus is immediately downstream of the BACE1 cleavage site. Ex vivo and in vitro experiments show that this variant inhibits BACE1 cleavage and results in reduced  $A\beta$  production. Another mutation at this site, A673V, enhances BACE1 cleavage activity<sup>9,10</sup> and is a recessive mutation causing early-onset AD. Likewise, the K670N/M671L mutation that affects the 2 amino acids immediately upstream of the BACE1 cleavage site also enhances BACE1 cleavage, increases  $A\beta$  production, and causes early-onset AD.<sup>11</sup> Thus, multiple mutations in close proximity to the BACE1 cleavage site influence risk for AD and  $A\beta$  production.

We genotyped a large number of AD cases and controls to determine whether the A673T mutation is an important protective variant in cognitively intact elderly US white individuals and in patients with AD from the same population. We found that this variant is extremely rare and does not have a discernible impact on AD risk in the US population.

## Methods

### Genotyping

We genotyped the samples listed in Table 1 using the Infinium HumanExome V1 Beadchip (Illumina, Inc). Genotyping was performed for 8410 individuals at the Robert S. Boas Center for Genomics and Human Genetics, Feinstein Institute for Medical Research, Manhasset, New York, 1990 individuals at the John P. Hussman Institute for Human Genomics, University of Miami, Miami, Florida, and 6166 individuals at the Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania. Genotypes were initially called using the default clustering profile from Illumina and recalled using clustering profiles generated by Genentech using data from 30 000 samples. We also genotyped the individuals listed in Table 2 for single-nucleotide polymorphism rs63750847 using the TaqMan assay C\_\_89522366\_10 (Life Technologies) and a 384-well plate format. Each plate contained samples from 2 heterozygotes from the exome array experiments. For both the exome array assay and the TaqMan assay, manual inspection of clustering indicated both were valid assays (eFigure 1 in the Supplement).

### Participants

The National Institute on Aging Alzheimer's Disease Centers case-control sample, the University of Toronto/GlaxoSmithKline (also called Gen ADA) case-control sample, the Vanderbilt/ Miami/Mount Sinai case-control sample, the National Institute on Aging-Late-Onset Alzheimer's Disease multiplex family-based sample, the National Cell Repository for Alzheimer's Disease multiplex family-based sample, the Multi-institutional Research in Alzheimer's Genetic Epidemiology family-based sample, and the Adult Changes in Thought (ACT) prospective cohort were described previously.<sup>12,13</sup> The Genetics Differences cohort is a population-based prevalent case-control study from the same population as the ACT study.<sup>14</sup> The Washington Heights-Inwood Columbia Aging Project sample is a multiethnic prospective cohort<sup>15</sup>; for this study, only white individuals were genotyped. The Washington University cohort is a white case-control cohort.<sup>13</sup> The Miami multiplex families and the National Institute of Mental Health multiplex families were as previously described.<sup>16-18</sup> The Cache County Study on Memory Health and Aging is a population-based study with 4 assessments of cognitive function since 1994.<sup>19</sup> The Swedish cohorts are case-control studies recruited from neuropsychiatric clinics in Sweden as described previously.<sup>20</sup> For the family-based sample, we genotyped a single affected individual from each kindred. All studies were approved by institutional review boards at the respective universities involved in each study, and the overall study was approved by the University of Pennsylvania institutional review board. The participants provided written informed consent.

## Analysis

The exome chip genotyping data (16 525 samples total) were first preprocessed using quality check steps adapted from Naj et al.<sup>13</sup> Briefly, we excluded 881 samples and markers with a missing call rate higher than 2%, samples with genotype-imputed or reported sex mismatch, and markers significant ( $P < 10^{-6}$ ) in either Hardy-Weinberg or informative missingness tests. We pruned 969 samples by relatedness test ( $\pi > 0.4$  using 15 086 linkage disequilibrium-pruned autosomal markers with a minor allele frequency  $>0.1$ ) and compared samples with HapMap 3 data to exclude nonwhite individuals. We checked population substructure in 15 644 samples with good call rates using 5848 single-nucleotide polymorphisms that were present on both the exome array and in the HapMap 3 data set. Forty-nine individuals were likely to be from non-white or admixed populations. Two hundred seventy-three individuals were excluded owing to incomplete diagnosis information. In all, 14 355 individuals (8221 cases and 6134 controls) passed quality check. Five hundred eighty-five individuals were younger than 60 years at onset (512 AD cases) or at last visit (34 controls); 61 samples lacked age information.

Genotypes for rs63750847 (A673T; c.G2017>A) were ascertained via the exm 1563596 marker (major/minor allele: G/A, 100% call rate after quality check). One sample with the A/A genotype was excluded owing to a very high missing call rate (20.84%). Two samples passing quality check carried the A/G genotype.

## Results

A total of 8221 AD cases and 6134 elderly cognitively normal controls were genotyped using Illumina exome arrays (Table 1). These arrays have single-nucleotide polymorphism assays for more than 240 000 exonic variants that are nonsynonymous, nonsense, or splice-site variants. The exome array has an assay for rs63750847, which is the A673T variant discussed earlier (eFigure 2 in the Supplement). We identified 2 heterozygotes, one in a sample from an AD case (minor allele frequency = 0.0061%) of Russian ancestry who had onset of AD symptoms at age 89 years and an *APOE* genotype of  $\epsilon 3/\epsilon 3$  and the second in a sample from a cognitively intact 82-year-old control (minor allele frequency = 0.0082%) born in Iceland with an *APOE* genotype of  $\epsilon 3/\epsilon 3$  (Table 3). Both were from the ACT study based in Seattle, Washington. Population principal components showed both to be of northern European ancestry when compared with HapMap 3 samples (eFigure 2 in the Supplement).

To validate the array genotyping results, we used a Taq-Man assay to re-genotype 1610 of the same samples, including the 2 heterozygotes. We also genotyped 983 additional samples from the ACT study (Table 2). All genotypes derived from the exome array genotyping were confirmed by the TaqMan assay, including the 2 heterozygotes initially observed. All additional ACT samples were homozygous for the normal A673 allele. TaqMan genotyping was also used to survey 506 AD cases, 383 individuals with other dementias, and 3838 controls from the Cache County Study on Memory Health and Aging. All dementia cases were homozygous for the normal A673 allele. One of the 3838 individuals without dementia carried the A673T allele (Table 2). This individual was aged 77 years at last assessment, had an *APOE* genotype of  $\epsilon 3/\epsilon 4$ , and had ancestors from Denmark, Ireland, Scotland, and

England (Table 3). Because the minor allele of the A673T variant was observed in a Swedish sample at a frequency of 0.42%,<sup>2</sup> we also genotyped 862 AD cases and 707 cognitively normal controls from Sweden (Table 2). Three controls were heterozygous for A673T. All AD cases were homozygous for the common A673 allele.

Previous work reported that an affected individual in a small family with late-onset AD was heterozygous for A673T.<sup>21</sup> Because an affected individual from the same family was genotyped using the exome array as one of the individuals from the National Institute on Aging–Late-Onset AD sample, we genotyped all other available family members for the A673T variant. This included 6 AD cases, 4 married-in spouses, and 7 blood relatives of affected individuals. We observed 1 spouse and the child of that spouse as heterozygous for A673T. The child is a blood relative of affected individuals in the family. Both the child and parent were unaffected individuals (ages at last visit, 84 and 72 years, respectively). The heterozygous individual originally reported as an AD case was actually the unaffected child described here. This family is from the United States with a Scandinavian background.

## Discussion

The *APP* A673T variant is overrepresented in Icelandic controls when compared with Icelandic AD cases.<sup>6</sup> Therefore, like the *APOE*  $\epsilon$ 2 allele,<sup>22</sup> the A673T variant appears to protect against late-onset AD, defined as disease onset after age 60 years. Studies on the functional consequences of the A673T substitution showed that this amino acid substitution inhibited BACE1 cleavage of APP, potentially reducing or eliminating the production of A $\beta$  peptide from APP encoded by this allele.<sup>6</sup> That this allele appears to be protective provides additional genetic evidence that A $\beta$  is a critical toxic molecule contributing to AD. Overproduction of A $\beta$  (eg, from the *APP* K670N/M671L mutation) causes early-onset AD, and a variant associated with reduced A $\beta$  production, A673T, protects against AD. Amino acid 673 appears to be critical for BACE1 cleavage since a different allele at the same amino acid (an alanine change to a valine, A673V) enhances A $\beta$  production and causes recessive early-onset dementia.<sup>9</sup> However, the A673T variant is exceedingly rare in the white individuals tested here (carrier frequency = 0.011% in US individuals with AD and 0.018% in cognitively normal controls), so it could not have a substantial role in risk for AD in the US population. The frequency in Swedish controls tested here was somewhat higher (0.42%) and is in line with the frequency in the Icelandic population. The effect size (Fisher exact test 99% CI, 3.97– $\infty$ ) is weaker than in the Icelandic population (oddsratio = 5.29), although this observation is suggestive and a larger sample size is needed.

The protective effect of A673T against late-onset AD reported in the Icelandic study<sup>6</sup> supports the hypothesis that A $\beta$  not only plays a critical role in early-onset familial AD but also is important for late-onset AD. This hypothesis is also supported by earlier work showing that the *APOE*  $\epsilon$ 4 allele not only increases risk for late-onset AD and lowers the age at onset<sup>23</sup> but similarly also modifies AD risk in carriers of *PSEN1*<sup>24</sup> and *PSEN2*<sup>25</sup> mutations. Mutations in *PSEN1* and *PSEN2* alter the C-terminal cleavage  $\gamma$ -secretase site and thus cause AD by an A $\beta$ -related mechanism. The fact that *APOE* genotype influences age at onset of this A $\beta$ -driven process suggests that late-onset AD and early-onset familial

AD share at least 1 common mechanism. The age cutoff distinguishing early- and late-onset disease is also blurred by the observation that *PSEN2* mutations occur in early- and late-onset AD cases.<sup>26,27</sup>

Jonsson et al<sup>6</sup> reported that in Icelandic individuals, the A673T allele frequencies were 0.13% in AD cases and 0.45% to 0.79% in controls. They also reported frequencies of 0.21%, 0.51%, and 0.42% in Norwegian, Finnish, and Swedish controls, respectively. In our Swedish cohort, we observed 3 controls who were heterozygous for A673T (age at sampling, 72, 55, and 59 years). Two of these controls are too young to definitively conclude they will not develop AD in the future. In fact, 1 control (aged 55 years) has an *APOE* genotype of  $\epsilon 4/\epsilon 4$ , suggesting increased risk for late-onset AD. In our US sample, we observed much lower frequencies in cases and controls, and 1 of our 3 carriers was born in Iceland. While our results are inconclusive regarding whether A673T is an AD-protective allele, they do indicate that this allele has little influence on risk in white individuals living in the United States. Our findings and the absence of the A673T allele in a large Chinese sample (n = 8721)<sup>28</sup> suggest the *APP* A673T allele may have an appreciable effect on AD risk only in Scandinavian and Icelandic populations.

## Conclusions

We show that the *APP* A673T allele is extremely rare in US white populations and thus does not play a substantial role in risk of developing AD in this group. Our study of a Swedish cohort showed a higher carrier frequency of *APP* A673T, and thus this variant appears primarily present in Icelandic and Scandinavian populations. Our results are consistent with this mutation being protective because carriers in our study were mostly controls. However, because of the rarity of this mutation in our populations, we could not independently verify that *APP* A673T is a protective allele.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Authors/Members in the Alzheimer's Disease Genetics Consortium

Marilyn S. Albert, PhD; Roger L. Albin, MD; Liana G. Apostolova, MD; Steven E. Arnold, MD; Robert Barber, PhD; M. Michael Barmada, PhD; Lisa L. Barnes, PhD; Thomas G. Beach, MD, PhD; James T. Becker, PhD; Gary W. Beecham, PhD; Duane Beekly, BS; David A. Bennett, MD; Eileen H. Bigio, MD; Thomas D. Bird, MD; Deborah Blacker, MD, ScD; Bradley F. Boeve, MD; James D. Bowen, MD; Adam Boxer, MD, PhD; James R. Burke, MD, PhD; Joseph D. Buxbaum, PhD; Nigel J. Cairns, PhD; Chuanhai Cao, PhD; Chris S. Carlson; Steven L. Carroll, MD, PhD; Helena C. Chui, MD; David G. Clark, MD; David H. Cribbs, PhD; Elizabeth A. Crocco, MD; Charles DeCarli, MD; Steven T. DeKosky, MD; F. Yesim Demirci, MD; Malcolm Dick, PhD; Dennis W. Dickson, MD; Ranjan Duara, MD; Nilufer Ertekin-Taner, MD, PhD; Kenneth B. Fallon, MD; Martin R. Farlow, MD; Steven Ferris, PhD; Matthew P. Frosch, MD, PhD; Douglas R. Galasko, MD; Mary Ganguli, MD, PhD; Marla Gearing, PhD; Daniel H. Geschwind, MD, PhD;

Bernardino Ghetti, MD; John R. Gilbert, PhD; Jonathan D. Glass, MD; Neill R. Graff-Radford, MD; John H. Growdon, MD; Ronald L. Hamilton, MD; Kara L. Hamilton-Nelson; Lindy E. Harrell, MD, PhD; Elizabeth Head, PhD; Lawrence S. Honig, MD, PhD; Christine M. Hulette, MD; Bradley T. Hyman, MD, PhD; Gail P. Jarvik, MD, PhD; Gregory A. Jicha, MD, PhD; Lee-Way Jin, MD, PhD; Gyungah Jun, PhD; M. Ilyas Kamboh, PhD; Anna Karydas, BA; Jeffrey A. Kaye, MD; Ronald Kim, MD; Edward H. Koo, MD; Neil W. Kowall, MD; Joel H. Kramer, PhD; Patricia Kramer, PhD; Frank M. LaFerla, PhD; James J. Lah, MD, PhD; James B. Leverenz, MD; Allan I. Levey, MD, PhD; Ge Li, MD, PhD; Andrew P. Lieberman, MD, PhD; Oscar L. Lopez, MD; Kathryn L. Lunetta, PhD; Constantine G. Lyketsos, MD, MHS; Wendy J. Mack, PhD; Daniel C. Marson, MD, PhD; Eden R. Martin, PhD; Frank Martiniuk, PhD; Deborah C. Mash, PhD; Eliezer Masliah, MD; Wayne C. McCormick; Susan M. McCurry, PhD; Andrew N. McDavid, BA; Ann C. McKee, MD; M. Marsel Mesulam, MD; Bruce L. Miller, MD; Carol A. Miller, MD; Joshua W. Miller, MD; Thomas J. Montine, MD, PhD; John C. Morris, MD; Jill R. Murrell, PhD; John M. Olichney, MD; Joseph E. Parisi, MD; William Perry; Elaine Peskind, MD; Ronald C. Petersen, MD, PhD; Aimee Pierce; Wayne W. Poon, PhD; Huntington Potter, PhD; Joseph F. Quinn, MD; Ashok Raj, MD; Murray Raskind, MD; Eric M. Reiman, MD; Barry Reisberg, MD; Christiane Reitz, MD, PhD; John M. Ringman, MD, MS; Erik D. Roberson, MD, PhD; Howard J. Rosen, MD; Roger N. Rosenberg, MD; Mary Sano, PhD; Andrew J. Saykin, PsyD; Julie A. Schneider, MD; Lon S. Schneider, MD; William W. Seeley, MD; Amanda G. Smith; Joshua A. Sonnen, MD; Salvatore Spina, MD; Robert A. Stern, PhD; Rudolph E. Tanzi, PhD; Tricia A. Thornton-Wells, PhD; John Q. Trojanowski, MD, PhD; Juan C. Troncoso, MD; Debby W. Tsuang, MD; Vivianna M. Van Deerlin, MD, PhD; Linda J. Van Eldik, PhD; Badri N. Vardarajan, PhD; Harry V. Vinters, MD; Jean Paul Vonsattel, MD; Sandra Weintraub, PhD; Kathleen A. Welsh-Bohmer, PhD; Jennifer Williamson, MS, MPH; Sarah Wishnek; Randall L. Woltjer, MD, PhD; Clinton B. Wright, MD, MS; Steven G. Younkin, MD, PhD; Chang-En Yu, PhD; Lei Yu, PhD.

## Affiliations of Authors/Members in the Alzheimer's Disease Genetics Consortium

Department of Neurology, Johns Hopkins University, Baltimore, Maryland (Albert); Department of Neurology, University of Michigan, Ann Arbor (Albin); Geriatric Research, Education, and Clinical Center, VA Ann Arbor Healthcare System, Ann Arbor, Michigan (Albin); Michigan Alzheimer's Disease Center, Ann Arbor (Albin); Department of Neurology, University of California, Los Angeles (Apostolova, Ringman, Vinters); Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia (Arnold); Department of Pharmacology and Neuroscience, University of North Texas Health Science Center, Fort Worth (Barber); Department of Human Genetics, University of Pittsburgh, Pittsburgh, Pennsylvania (Barmada, Demirci, Kamboh); Department of Neurological Sciences, Rush University Medical Center, Chicago, Illinois (Barnes, Bennett, J. A. Schneider, L. Yu); Department of Behavioral Sciences, Rush University Medical Center, Chicago, Illinois (Barnes); Civin Laboratory for Neuropathology, Banner Sun Health Research Institute, Phoenix, Arizona (Beach); Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh,



Pennsylvania (Becker, Ganguli); Department of Neurology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (Becker); Department of Psychology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (Becker); John P. Hussman Institute for Human Genomics, University of Miami, Miami, Florida (Beecham, Gilbert, Hamilton-Nelson, Martin, Perry, Wishnek); Dr John T. Macdonald Foundation Department of Human Genetics, University of Miami, Miami, Florida (Beecham, Gilbert, Martin); National Alzheimer's Coordinating Center, University of Washington, Seattle (Beekly); Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, Illinois (Bennett); Department of Pathology, Northwestern University Feinberg School of Medicine, Chicago, Illinois (Bigio); Cognitive Neurology and Alzheimer's Disease Center, Northwestern University Feinberg School of Medicine, Chicago, Illinois (Bigio, Mesulam, Weintraub); Geriatric Research, Education, and Clinical Center, VA Puget Sound Healthcare System, Seattle, Washington (Bird, Tsuang); Department of Neurology, University of Washington, Seattle (Bird); Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts (Blacker); Department of Psychiatry, Massachusetts General Hospital/Harvard Medical School, Boston (Blacker); Department of Neurology, Mayo Clinic, Rochester, Minnesota (Boeve, Petersen); Swedish Medical Center, Seattle, Washington (Bowen); Department of Neurology, University of California, San Francisco (Boxer, Karydas, B. L. Miller, Rosen, Seeley); Department of Medicine, Duke University, Durham, North Carolina (Burke, Welsh-Bohmer); Department of Neuroscience, Mount Sinai School of Medicine, New York, New York (Buxbaum); Department of Psychiatry, Mount Sinai School of Medicine, New York, New York (Buxbaum, Sano); Department of Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, New York (Buxbaum); Department of Pathology and Immunology, Washington University, St Louis, Missouri (Cairns, Morris); USF Health Byrd Alzheimer's Institute, University of South Florida, Tampa (Cao, Potter, Raj, Smith); Fred Hutchinson Cancer Research Center, Seattle, Washington (Carlson, McDavid); Department of Pathology, University of Alabama at Birmingham, Birmingham (Carroll, Fallon); Department of Neurology, University of Southern California, Los Angeles (Chui, L. S. Schneider); Department of Neurology, University of Alabama at Birmingham, Birmingham (Clark, Harrell, Marson, Roberson); Department of Neurology, University of California, Irvine (Cribbs, Pierce); Department of Psychiatry and Behavioral Sciences, Miller School of Medicine, University of Miami, Miami, Florida (Crocco); Department of Neurology, University of California, Davis, Sacramento (DeCarli, Olichney); University of Virginia School of Medicine, Charlottesville (DeKosky); now with University of Pittsburgh, Pittsburgh, Pennsylvania (DeKosky); Institute for Memory Impairments and Neurological Disorders, University of California, Irvine (Dick, Poon); Department of Neuroscience, Mayo Clinic, Jacksonville, Florida (Dickson, Ertekin-Taner, Graff-Radford, Younkin); Wien Center for Alzheimer's Disease and Memory Disorders, Mount Sinai Medical Center, Miami Beach, Florida (Duara); Department of Neurology, Mayo Clinic, Jacksonville, Florida (Ertekin-Taner, Graff-Radford); Department of Neurology, Indiana University, Indianapolis (Farlow); Department of Psychiatry, New York University, New York (Ferris, Reisberg); C. S. Kubik Laboratory for Neuropathology, Massachusetts General Hospital, Charlestown (Frosch); Department of Neurosciences, University of California, San Diego, La Jolla (Galasko, Koo, Masliah); Department of Pathology and Laboratory Medicine, Emory University, Atlanta, Georgia

(Gearing); Alzheimer's Disease Research Center, Emory University, Atlanta, Georgia (Gearing); Neurogenetics Program, University of California, Los Angeles (Geschwind); Department of Pathology and Laboratory Medicine, Indiana University, Indianapolis (Ghetti, Murrell, Spina); Department of Neurology, Emory University, Atlanta, Georgia (Glass, Lah, Levey); Department of Neurology, Massachusetts General Hospital/Harvard Medical School, Boston (Growdon, Hyman, Tanzi); Division of Neuropathology, Department of Pathology, University of Pittsburgh, Pittsburgh, Pennsylvania (Hamilton); Department of Molecular and Biomedical Pharmacology, Sanders-Brown Center on Aging, University of Kentucky, Lexington (Head); Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University, New York, New York (Honig, Reitz, Vardarajan, Vonsattel, Williamson); Department of Neurology, Columbia University, New York, New York (Honig, Reitz, Vardarajan); Department of Pathology, Duke University, Durham, North Carolina (Hulette); Department of Genome Sciences, University of Washington, Seattle (Jarvik); Division of Medical Genetics, Department of Medicine, University of Washington, Seattle (Jarvik); Department of Neurology, Sanders-Brown Center on Aging, University of Kentucky, Lexington (Jicha); Department of Pathology and Laboratory Medicine, University of California, Davis, Sacramento (Jin, J. W. Miller); Genetics Program, Department of Medicine, Boston University, Boston, Massachusetts (Jun); Department of Biostatistics, Boston University, Boston, Massachusetts (Jun, Lunetta); Department of Ophthalmology, Boston University, Boston, Massachusetts (Jun); Alzheimer Disease Research Center, University of Pittsburgh, Pittsburgh, Pennsylvania (Kamboh, Lopez); Department of Neurology, Oregon Health & Science University, Portland (Kaye, P. Kramer, Quinn); Department of Neurology, Portland Veterans Affairs Medical Center, Portland, Oregon (Kaye); Department of Pathology and Laboratory Medicine, University of California, Irvine (Kim); Department of Neurology, Boston University, Boston, Massachusetts (Kowall, McKee, Stern); Department of Pathology, Boston University, Boston, Massachusetts (Kowall, McKee); Department of Neuropsychology, University of California, San Francisco (J. H. Kramer); Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland (P. Kramer); Department of Neurobiology and Behavior, University of California, Irvine (LaFerla); Department of Pathology, University of Washington, Seattle (Leverenz, Montine, Sonnen); Department of Psychiatry and Behavioral Sciences, University of Washington School of Medicine, Seattle (Li, Peskind, Raskind, Tsuang); Department of Pathology, University of Michigan, Ann Arbor (Lieberman); Department of Psychiatry, Johns Hopkins University, Baltimore, Maryland (Lyketsos); Department of Preventive Medicine, University of Southern California, Los Angeles (Mack); Division of Pulmonary, Critical Care, and Sleep Medicine, Department of Medicine, New York University, New York (Martiniuk); Department of Neurology, University of Miami, Miami, Florida (Mash); Department of Pathology, University of California, San Diego, La Jolla (Masliah); Department of Medicine, University of Washington, Seattle (McCormick, C.-E. Yu); Northwest Research Group on Aging, School of Nursing, University of Washington, Seattle (McCurry); Department of Neurology, Northwestern University Feinberg School of Medicine, Chicago, Illinois (Mesulam); Department of Pathology, University of Southern California, Los Angeles (C. A. Miller); Department of Neurology, Washington University, St Louis, Missouri (Morris); Department of Medical and Molecular Genetics, Indiana University, Indianapolis (Murrell, Saykin);

Department of Anatomic Pathology, Mayo Clinic, Rochester, Minnesota (Parisi); Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, Minnesota (Parisi); Neurogenomics Division, Translational Genomics Research Institute, Phoenix, Arizona (Reiman); Arizona Alzheimer's Consortium, Phoenix (Reiman); Banner Alzheimer's Institute, Phoenix, Arizona (Reiman); Alzheimer's Disease Center, New York University, New York (Reisberg); Gertrude H. Sergievsky Center, Columbia University, New York, New York (Reitz, Vardarajan); Department of Neurology, University of Texas Southwestern Medical Center, Dallas (Rosenberg); Department of Radiology and Imaging Sciences, Indiana University, Indianapolis (Saykin); Department of Pathology (Neuropathology), Rush University Medical Center, Chicago, Illinois (J. A. Schneider); Department of Psychiatry, University of Southern California, Los Angeles (L. S. Schneider); Center for Human Genetics and Research, Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee (Thornton-Wells); Department of Pathology and Laboratory Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia (Trojanowski, Van Deerlin); Department of Pathology, Johns Hopkins University, Baltimore, Maryland (Troncoso); Department of Anatomy and Neurobiology, Sanders-Brown Center on Aging, University of Kentucky, Lexington (Van Eldik); Department of Pathology and Laboratory Medicine, University of California, Los Angeles (Vinters); Department of Pathology, Columbia University, New York, New York (Vonsattel); Department of Psychiatry, Northwestern University Feinberg School of Medicine, Chicago, Illinois (Weintraub); Department of Psychiatry and Behavioral Sciences, Duke University, Durham, North Carolina (Welsh-Bohmer); Department of Pathology, Oregon Health & Science University, Portland (Woltjer); Evelyn F. McKnight Brain Institute, Department of Neurology, Miller School of Medicine, University of Miami, Miami, Florida (Wright).

## Author Contributions

Drs Wang and Schellenberg had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Study concept and design:* Wang, Naj, Cruchaga, Zetterberg, Blennow, Kauwe, Haines, Farrer, Pericak-Vance, Mayeux, Schellenberg.

*Acquisition, analysis, or interpretation of data:* Wang, Naj, Graham, Crane, Kunkle, Cruchaga, Gonzalez Murcia, Cannon-Albright, Baldwin, Zetterberg, Blennow, Kukull, Faber, Schupf, Norton, Tschanz, Munger, Corcoran, Rogaeva, Albert, Albin, Apostolova, Arnold, Barber, Barmada, Barnes, Beach, Becker, Beecham, Beekly, Bennett, Bigio, Bird, Blacker, Boeve, Bowen, Boxer, Burke, Buxbaum, Cairns, Cao, Carlson, Carroll, Chui, Clark, Cribbs, Crocco, DeCarli, DeKosky, Demirci, Dick, Dickson, Duara, Ertekin-Taner, Fallon, Farlow, Ferris, Frosch, Galasko, Ganguli, Gearing, Geschwind, Ghetti, Gilbert, Glass, Graff-Radford, Growdon, Hamilton, Hamilton-Nelson, Harrell, Head, Honig, Hulette, Hyman, Jarvik, Jicha, Jin, Jun, Kamboh, Karydas, Kaye, Kim, Koo, Kowall, J. H. Kramer, P. Kramer, LaFerla, Lah, Leverenz, Levey, Li, Lieberman, Lopez, Lunetta, Lyketsos, Mack, Marson, Martin, Martiniuk, Mash, Masliah, McCormick, McCurry, McDavid, McKee, Mesulam, B. L. Miller, C. A. Miller, J. W. Miller, Montine, Morris, Murrell, Olichney,

Parisi, Perry, Peskind, Petersen, Pierce, Poon, Potter, Quinn, Raj, Raskind, Reiman, Reisberg, Reitz, Ringman, Roberson, Rosen, Rosenberg, Sano, Saykin, J. A. Schneider, L. S. Schneider, Seeley, Smith, Sonnen, Spina, Stern, Tanzi, Thornton-Wells, Trojanowski, Troncoso, Tsuang, Van Deerlin, Van Eldik, Vardarajan, Vinters, Vonsattel, Weintraub, Welsh-Bohmer, Williamson, Wishnek, Woltjer, Wright, Younkin, C.-E. Yu, L. Yu, Lin, Dombroski, Cantwell, Partch, Valladares, Hakonarson, St George-Hyslop, Green, Goate, Foroud, Carney, Larson, Behrens, Kauwe, Pericak-Vance, Mayeux, Schellenberg.

*Drafting of the manuscript:* Wang, Naj, Schellenberg.

*Critical revision of the manuscript for important intellectual content:* All authors.

*Statistical analysis:* Wang, Naj, Gonzalez Murcia, Corcoran, Carlson, Ghetti, Hamilton-Nelson, Jun, Van Deerlin, Foroud, Kauwe, Haines, Schellenberg.

*Obtained funding:* Munger, Barnes, Bennett, DeKosky, Galasko, Ganguli, Ghetti, Jarvik, Kamboh, Lopez, Mesulam, Montine, Morris, Reiman, Saykin, Hakonarson, St George-Hyslop, Larson, Haines, Farrer, Mayeux, Schellenberg.

*Administrative, technical, or material support:* Naj, Cruchaga, Zetterberg, Blennow, Norton, Munger, Arnold, Beach, Becker, Beekly, Bennett, Boeve, Cairns, Cao, Chui, Cribbs, DeKosky, Ertekin-Taner, Ferris, Geschwind, Ghetti, Hamilton, Harrell, Hyman, Jarvik, Jin, LaFerla, Lieberman, Mash, McCormick, Parisi, Potter, Rosen, Sano, Trojanowski, Vonsattel, Larson, Behrens, Kauwe, Schellenberg.

*Study supervision:* Munger, Leverenz, Hakonarson, Pericak-Vance, Mayeux, Schellenberg.

## Conflict of Interest Disclosures

Drs Graham and Behrens are full-time employees of Genentech, Inc. Dr DeKosky is a consultant for AstraZeneca, Eli Lilly and Co, Merck and Co, and Rivermend and is involved in clinical trials with Baxter, Elan, Janssen, Novartis, and Pfizer. Dr Farlow receives research support from Accera, Biogen, Eisai, Eli Lilly and Co, Genentech, MedAvante/AstraZeneca, and Navidea; is on the speaker's bureau at Eisai, Pfizer, Forest, Novartis, and Eli Lilly and Co; and is a consultant or advisory board member for Accera, Alltech, Avanir, Eisai, Med Res, Hilicon, Medavante, Medivation, Merck and Co, Novartis, Pfizer, Prana Biotech, QR Pharma, Roche, Sanofi-Aventis, Schering-Plough, Toyama Pharm, Eli Lilly and Co, and UCB Pharma. Dr Ghetti has consulted for Piramal Imaging. Dr Rosenberg is the *JAMA Neurology* Editor and Dr Honig is the *JAMA Neurology* Web Editor but neither was involved in the review process or the acceptance of the manuscript. Dr Leverenz is a consultant for Boehringer-Ingelheim, Navidea Biopharmaceuticals, and Piramal Healthcare. Dr Petersen is chair for the data monitoring committee for Pfizer and Janssen Alzheimer Immunotherapy and is a consultant for GE Healthcare and Roche. Dr Wright receives royalties from UpTo Date for 2 chapters; has done legal consulting for the law firms of Abali, Milne, and Faegre Baker Daniels; is a consultant for Merck and Co; and does stroke adjudication for a National Institutes of Health clinical trial. No other disclosures were reported.

## Funding/Support

The National Institute on Aging supported this work through the following grants: U01 AG032984 and RC2 AG036528 (Alzheimer's Disease Genetics Consortium); U01 AG016976 (National Alzheimer's Coordinating Center); U24 AG021886 (National Cell Repository for Alzheimer's Disease); U24 AG026395, R01 AG041797, and U24 AG026390 (National Institute on Aging–Late-Onset Alzheimer's Disease); P30 AG019610 (Banner Sun Health Research Institute); P30 AG013846, R01 HG02213, K24 AG027841, U01 AG10483, R01 CA129769, and R01 MH080295 (Boston University); P50 AG008702 (Columbia University); P30 AG028377 (Duke University); AG025688 (Emory University); P30 AG10133 (Indiana University); P50 AG005146 (Johns Hopkins University); P50 AG005134 (Massachusetts General Hospital); P50 AG016574 (Mayo Clinic); P50 AG005138 (Mount Sinai School of Medicine); P30 AG08051, U01 AG16976, MO1 RR00096, and UL1 RR029893 (New York University); P30 AG013854 (Northwestern University); P30 AG008017 (Oregon Health & Science University); P30 AG010161 (Rush University); P50 AG016582 and UL1 RR02777 (Center for Clinical and Translational Science, University of Alabama at Birmingham); P30 AG010129 (University of California, Davis); P50 AG016573, P50 AG016574, P50 AG016575, P50 AG016576, and P50 AG016577 (University of California, Irvine); P50 AG016570 (University of California, Los Angeles); P50 AG005131 (University of California, San Diego); P50 AG023501 and P01 AG019724 (University of California, San Francisco); P30 AG028383 (University of Kentucky); P50 AG008671 (University of Michigan); P30 AG010124 (University of Pennsylvania); P50 AG005133 (University of Pittsburgh); P50 AG005142 (University of Southern California); P30 AG012300 (University of Texas Southwestern Medical Center); P50 AG005136 and R01 AG007584 (University of Washington); and P50 AG005681 and P01 AG03991 (Washington University). The Alzheimer's Disease Genetics Consortium is also supported by a grant from a private foundation wishing to remain anonymous. The work completed by Boston University is also supported by grant IIRG-08-89720 from the Alzheimer's Association and by the VA New England Geriatric Research Education and Clinical Center. The work by Miami University is also supported by grant R01 AG027944 from the National Institute on Aging and by the Alzheimer's Association. This project was also made possible by the many contributions of individual study data sets, supported in part by the National Institutes of Health, including Adult Changes in Thought and Electronic Medical Records and Genetics (eMERGE) (grants U01 AG06781, U01 HG004610, and U01 HG006375 from the National Human Genome Research Institute), Columbia University Washington Heights–Inwood Columbia Aging Project (R01 AG037212 from the National Institutes of Health), and MIRAGE (R01 AG009029 from the National Institute on Aging). The University of Miami/Vanderbilt University/Mount Sinai School of Medicine work was supported by grants AG010491, AG002219, AG005138, AG027944, AG021547, AG019757, and R01 AG027944 from the National Institute on Aging and IIRG-05-14147 from the Alzheimer's Association. A subset of these participants was ascertained while Dr Pericak-Vance was a faculty member at Duke University. Clinical data and genotyping efforts for the Religious Orders Study and the Memory and Aging Project were supported by grants P30 AG10161, R01 AG15819, R01 AG17917, and R01 AG30146 from the National Institute on Aging and by the Illinois Department of Public Health and the Translational

Genomics Research Institute. Data and samples from the National Institute on Aging–Late-Onset Alzheimer’s Disease Family Study, which receives government support under cooperative agreement grant U24AG026390 from the National Institute on Aging, were used in this study. Data from the Cache County Study on Memory Health and Aging were supported by grants R01 AG042611 from the National Institute on Aging and MNIRG-11-205368 from the Alzheimer’s Association and by the Brigham Young University Gerontology Program and the Charleston Conference on Alzheimer’s Disease to Brigham Young University; they were also supported by grants R01 AG11380, R01 AG21136, and R01AG3272 from the National Institutes of Health and by the Utah Science, Technology, and Research Initiative and the Utah State University Agricultural Experiment Station to Utah State University. Samples from the National Cell Repository for Alzheimer’s Disease, which receives government support under cooperative agreement grant U24 AG21886 from the National Institute on Aging, were used in this study. Dr St George-Hyslop is supported by the Wellcome Trust, Howard Hughes Medical Institute, and Canadian Institute of Health Research. Drs Barmada, Demirci, and Kamboh are supported by grant AG030653 from the National Institute on Aging. Dr Wright is supported by grants K02 NS 059729 from the National Institute of Neurological Disorders and Stroke, R01 HL 108623 from the National Heart, Lung, and Blood Institute, and SDG 0735387N from the American Heart Association.

## Role of the Funder/Sponsor

The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

## Group Information

The National Institute on Aging–Late-Onset Alzheimer’s Disease (NIA-LOAD) Family Study investigators include the following: Boston University, Boston, Massachusetts: Robert C. Green, MD, MPH, Neil W. Kowall, MD, Lindsay A. Farrer, PhD; Columbia University, New York, New York: Jennifer Williamson, MS, MPH, Vincent Santana, MBA; Duke University, Durham, North Carolina: Donald Schmechel, MD, Perry Gaskell, BS, Kathleen A. Welsh-Bohmer, PhD; Indiana University, Indianapolis: Bernardino Ghetti, MD, Martin R. Farlow, MD, Kelly Horner; Massachusetts General Hospital, Boston: John H. Growdon, MD, Deborah Blacker, MD, ScD, Rudolph E. Tanzi, PhD, Bradley T. Hyman, MD, PhD; Mayo Clinic, Rochester, Minnesota: Bradley F. Boeve, MD, Karen Kuntz, RN, Lindsay Norgaard, BS, Nathan Larson, BS; Mayo Clinic, Jacksonville, Florida: Dana Kistler, BSH, Francine Parfitt, MS, Jenny Haddow, BS; Mount Sinai School of Medicine, New York, New York: Jeremy Silverman, PhD, Michal Schnaider Beerli, PhD, Mary Sano, PhD, Joy Wang, BA, Rachel Lally, BA; Northwestern University, Chicago, Illinois: Nancy Johnson, PhD, M. Marsel Mesulam, MD, Sandra Weintraub, PhD, Eileen H. Bigio, MD; Oregon Health & Science University, Portland: Jeffery A. Kaye, MD, Patricia Kramer, PhD, Jessica Payne-Murphy, BA; Rush University, Chicago, Illinois: David A. Bennett, MD, Holli Jacobs, BA, Jeen-Soo Chang, MD, Danielle Arends, RN; University of Alabama at Birmingham, Birmingham: Lindy E. Harrell, MD, PhD; University of California, Los Angeles: George

Bartzokis, MD, Jeffery Cummings, MD, Po H. Lu, PsyD, Usha Toland, MS; University of Kentucky, Lexington: William Markesbery, MD, Charles Smith, MD, Alise Brickhouse, BA; University of Pennsylvania, Philadelphia: Gerard D. Schellenberg, PhD, John Q. Trojanowski, MD, PhD, Viviana M. Van Deerlin, MD, PhD, Elisabeth McCarty Wood, MS; University of Pittsburgh, Pittsburgh, Pennsylvania: Steven T. DeKosky, MD, Robert Sweet, MD, Elise Weamer, MPH; University of Southern California, Los Angeles: Helena C. Chui, MD, Arousiak Varpetian, MD; University of Texas Southwestern Medical Center, Dallas: Ramon Diaz-Arrastia, MD, PhD, Roger N. Rosenberg, MD, Barbara Davis, MA; University of Washington, Seattle: Thomas D. Bird, MD, Malia Rumbaugh, MS, Murray Raskind, MD; Washington University, St Louis, Missouri: Alison M. Goate, DPhil, John C. Morris, MD, Joanne Norton, MSN, RN, Denise Levitch, RN, Betsy Grant, MSW, PhD, Mary Coats, MSN, RN.

## Additional Contributions

Annette Lee, PhD, and Peter K. Gregersen, MD, Feinstein Institute for Medical Research, Manhasset, New York, provided genotyping services. Creighton Phelps, PhD, Stephen Snyder, PhD, and Marilyn Miller, PhD, National Institute on Aging, Bethesda, Maryland, helped in acquiring samples and data and are ex officio members of the Alzheimer's Disease Genetics Consortium. Duke University acknowledges John Ervin, BA, from the Brain Bank and Kathleen Hayden, PhD, in the Clinical Core for their respective efforts in the DNA and data pulls required. These individuals received no compensation from the funders for their contributions. We thank the contributors, including the Alzheimer's Disease Centers who collected samples used in this study, as well as the patients and their families, whose help and participation made this work possible.

## References

- Schellenberg GD, Montine TJ. The genetics and neuropathology of Alzheimer's disease. *Acta Neuropathol.* 2012; 124(3):305–323. [PubMed: 22618995]
- Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell.* 2012; 148(6):1204–1222. [PubMed: 22424230]
- Ghosh AK, Brindisi M, Tang J. Developing  $\beta$ -secretase inhibitors for treatment of Alzheimer's disease. *J Neurochem.* 2012; 120(suppl 1):71–83. [PubMed: 22122681]
- De Strooper B, Vassar R, Golde T. The secretases: enzymes with therapeutic potential in Alzheimer disease. *Nat Rev Neurol.* 2010; 6(2):99–107. [PubMed: 20139999]
- Delrieu J, Ousset PJ, Caillaud C, Vellas B. "Clinical trials in Alzheimer's disease": immunotherapy approaches. *J Neurochem.* 2012; 120(suppl 1):186–193. [PubMed: 21883222]
- Jonsson T, Atwal JK, Steinberg S, et al. A mutation in *APP* protects against Alzheimer's disease and age-related cognitive decline. *Nature.* 2012; 488(7409):96–99. [PubMed: 22801501]
- Peacock ML, Warren JT Jr, Roses AD, Fink JK. Novel polymorphism in the A4 region of the amyloid precursor protein gene in a patient without Alzheimer's disease. *Neurology.* 1993; 43(6):1254–1256. [PubMed: 8170579]
- Kero M, Paetau A, Polvikoski T, et al. Amyloid precursor protein (*APP*) A673T mutation in the elderly Finnish population. *Neurobiol Aging.* 2013; 34(5):e1–e3. [PubMed: 23102935]
- Di Fede G, Catania M, Morbin M, et al. A recessive mutation in the *APP* gene with dominant-negative effect on amyloidogenesis. *Science.* 2009; 323(5920):1473–1477. [PubMed: 19286555]
- Giliberto L, Borghi R, Piccini A, et al. Mutant presenilin 1 increases the expression and activity of BACE1. *J Biol Chem.* 2009; 284(14):9027–9038. [PubMed: 19196715]

11. Mullan M, Crawford F, Axelman K, et al. A pathogenic mutation for probable Alzheimer's disease in the APP gene at the N-terminus of beta-amyloid. *Nat Genet.* 1992; 1(5):345–347. [PubMed: 1302033]
12. Jun G, Naj AC, Beecham GW, et al. Alzheimer's Disease Genetics Consortium. Meta-analysis confirms *CRI*, *CLU*, and *PICALM* as Alzheimer disease risk loci and reveals interactions with *APOE* genotypes. *Arch Neurol.* 2010; 67(12):1473–1484. [PubMed: 20697030]
13. Naj AC, Jun G, Beecham GW, et al. Common variants at *MS4A4/MS4A6E*, *CD2AP*, *CD33* and *EPHA1* are associated with late-onset Alzheimer's disease. *Nat Genet.* 2011; 43(5):436–441. [PubMed: 21460841]
14. Kukull WA, Schellenberg GD, Bowen JD, et al. Apolipoprotein E in Alzheimer's disease risk and case detection: a case-control study. *J Clin Epidemiol.* 1996; 49(10):1143–1148. [PubMed: 8826994]
15. Tang MX, Stern Y, Marder K, et al. The *APOE*-epsilon4 allele and the risk of Alzheimer disease among African Americans, whites, and Hispanics. *JAMA.* 1998; 279(10):751–755. [PubMed: 9508150]
16. Blacker D, Haines JL, Rodes L, et al. ApoE-4 and age at onset of Alzheimer's disease: the NIMH genetics initiative. *Neurology.* 1997; 48(1):139–147. [PubMed: 9008509]
17. Naj AC, Beecham GW, Martin ER, et al. Dementia revealed: novel chromosome 6 locus for late-onset Alzheimer disease provides genetic evidence for folate-pathway abnormalities. *PLoS Genet.* 2010; 6(9):e1001130. [PubMed: 20885792]
18. Beecham GW, Martin ER, Li YJ, et al. Genome-wide association study implicates a chromosome 12 risk locus for late-onset Alzheimer disease. *Am J Hum Genet.* 2009; 84(1):35–43. [PubMed: 19118814]
19. Breitner JCS, Wyse BW, Anthony JC, et al. *APOE*-epsilon4 count predicts age when prevalence of AD increases, then declines: the Cache County Study. *Neurology.* 1999; 53(2):321–331. [PubMed: 10430421]
20. Zetterberg M, Landgren S, Andersson ME, et al. Association of complement factor H Y402H gene polymorphism with Alzheimer's disease. *Am J Med Genet B Neuropsychiatr Genet.* 2008; 147B(6):720–726. [PubMed: 18163432]
21. Cruchaga C, Haller G, Chakraverty S, et al. NIA-LOAD/NCRAD Family Study Consortium. Rare variants in *APP*, *PSEN1* and *PSEN2* increase risk for AD in late-onset Alzheimer's disease families. *PLoS One.* 2012; 7(2):e31039. [PubMed: 22312439]
22. Corder EH, Saunders AM, Risch NJ, et al. Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer disease. *Nat Genet.* 1994; 7 (2):180–184. [PubMed: 7920638]
23. Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science.* 1993; 261(5123):921–923. [PubMed: 8346443]
24. Lendon CL, Martinez A, Behrens IM, et al. E280A PS-1 mutation causes Alzheimer's disease but age of onset is not modified by ApoE alleles. *Hum Mutat.* 1997; 10(3):186–195. [PubMed: 9298817]
25. Wijsman EM, Daw EW, Yu X, et al. APOE and other loci affect age-at-onset in Alzheimer's disease families with PS2 mutation. *Am J Med Genet B Neuropsychiatr Genet.* 2005; 132B(1):14–20. [PubMed: 15389756]
26. Levy-Lahad E, Wijsman EM, Nemens E, et al. A familial Alzheimer's disease locus on chromosome 1. *Science.* 1995; 269(5226):970–973. [PubMed: 7638621]
27. Levy-Lahad E, Wasco W, Poorkaj P, et al. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science.* 1995; 269 (5226):973–977. [PubMed: 7638622]
28. Ting SK, Chong MS, Kandiah N, et al. Absence of A673T amyloid- $\beta$  precursor protein variant in Alzheimer's disease and other neurological diseases. *Neurobiol Aging.* 2013; 34(10):e7–e8. [PubMed: 23652020]



**Table 1**  
**Samples Genotyped for A673T (rs63750847) With the Exome Array That Passed Quality Checking**

Characteristic	Cohort											
	ADC <sup>a</sup>	University of Toronto/ GSK <sup>b</sup>	Miami <sup>c</sup>	NIMH <sup>b</sup>	NIA-LOAD <sup>b</sup>	NCRAD <sup>b</sup>	MIRAGE <sup>b</sup>	Genetic Differences <sup>b</sup>	ACT <sup>b</sup>	WHICAP <sup>b</sup>	Washington University <sup>d</sup>	Total
Samples, No. (A673T heterozygotes, No.)												
Cases	3930 (0)	152 (0)	936 (0)	354 (0)	749 (0)	395 (0)	576 (0)	239 (0)	282 (1)	54 (0)	554 (0)	8221 (1)
Controls	2326 (0)	0	995 (0)	0	458 (0)	0	12 (0)	216 (0)	1445 (1)	322 (0)	360 (0)	6134 (1)
Total	6256 (0)	152 (0)	1931 (0)	354 (0)	1207 (0)	395 (0)	588 (0)	455 (0)	1727 (2)	376 (0)	914 (0)	14 355 (2)
Male, %	43.6	46.1	38.5	28.0	39.4	31.9	38.4	35.8	43.4	39.9	40.4	41.1
Case age at onset												
Mean (SD), y	72.4 (9.3)	77.8 (6.8)	72.6 (7.1)	71.6 (8.1)	73.8 (7.2)	71.0 (8.5)	68.6 (8.7)	76.4 (6.2)	83.7 (4.7)	84.5 (7.2)	79.8 (9.7)	73.4 (9.1)
No.	3930	152	885	354	749	393	572	239	282	54	554	6130
Control age at last examination												
Mean (SD), y	77.2 (9.3)	NA	73.5 (7.9)	NA	79.6 (8.7)	NA	76.1 (7.8)	80.8 (6.6)	81.6 (6.1)	81.0 (6.1)	74.6 (8.7)	78.0 (8.6)
No.	2326	NA	991	NA	458	NA	12	216	1445	322	360	8164
Cohort type	Case-control	Case-control	Case-control	Multiplex families	Multiplex families	Multiplex families	Multiplex families	Population-based incident cases with matched controls	Prospective cohort	Prospective cohort	Case-control	Case-control

Abbreviations: ACT, Adult Changes in Thought; ADC, Alzheimer's Disease Centers; GSK, GlaxoSmithKline; MIRAGE, Multi-institutional Research in Alzheimer's Genetic Epidemiology; NA, not applicable; NCRAD, National Cell Repository for Alzheimer's Disease; NIA-LOAD, National Institute on Aging-Late-Onset Alzheimer's Disease; NIMH, National Institute of Mental Health; WHICAP, Washington Heights-Inwood Columbia Aging Project.

<sup>a</sup>In this cohort, 3640 cases and 2223 controls were genotyped at the Robert S. Boas Center for Genomics and Human Genetics, Feinstein Institute for Medical Research, Manhasset, New York (as 2 batches) and 282 cases and 103 controls were genotyped at the Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania.

<sup>b</sup>Genotyped at the Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania.

<sup>c</sup>Genotyped at the John P. Hussman Institute for Human Genomics, University of Miami, Miami, Florida.

<sup>d</sup>Genotyped at the Robert S. Boas Center for Genomics and Human Genetics, Feinstein Institute for Medical Research.

Table 2

Samples Genotyped for A673T (rs63750847) by TaqMan Assay<sup>a</sup>

Characteristic	Cohort					
	United States			Sweden		
	ACT <sup>b</sup>	Cache County Study on Memory Health and Aging	Total White Individuals	Cohort 1	Cohort 2	Total Swedish Individuals
Samples, No. (A673T heterozygotes, No.)						
Cases	498 (1)	506 (0)	1004 (1)	298 (0)	564 (0)	862 (0)
Controls	1953 (1)	3838 (1)	5791 (2)	220 (0)	487 (3)	707 (3)
Dementia, other <sup>c</sup>	138 (0)	383 (0)	521 (0)	0	0	0
Total	2589 (2)	4727 (1)	7313 (3)	518 (0)	1051 (3)	1569 (3)
Male, No. (%)						
Cases	34.2	32.3		36.9	41.7	
Controls	43.5	41.7		25.0	72.1	
Age, mean (SD), y						
At onset for cases	84.1 (5.6)	82.4 (6.9)		75.7 (12.9)		
At sampling for cases					76.4 (10.7)	
At last examination for controls	81.2 (7.0)	79.8 (6.3)		98.0 (1.5)		
At sampling for controls					62.7 (10.7)	
Cohort type	Prospective cohort	Prospective cohort		Case-control	Case-control	

Abbreviation: ACT, Adult Changes in Thought.

<sup>a</sup>Life Technologies.

<sup>b</sup>Includes ACT samples genotyped using the exome array and an additional 364 cases, 1344 controls, and 117 other dementias. The two A673T variants are the same as in Table 1.

<sup>c</sup>Includes vascular dementia, mixed dementias, and other rare forms of dementia.

**Table 3**

## Heterozygotes for A673T

Cohort and Participant	Age, y <sup>a</sup>	APOE Genotype
ACT		
Case	89	ε3/ε3
Control	77	ε3/ε4
Cache County Study on Memory Health and Aging control		
	82	ε3/ε4
Swedish cohort 2		
Control 1	72	ε3/ε3
Control 2	55	ε4/ε4
Control 3	59	ε3/ε3
Family 1		
Control, mother	72	ε3/ε3
Control, daughter	84	ε3/ε3

Abbreviation: ACT, Adult Changes in Thought.

<sup>a</sup> Age indicates age at onset for the ACT case, age at last examination for the ACT, Cache County Study on Memory Health and Aging, and family 1 controls, and age at sampling for the 3 Swedish controls.