

# Double-stranded RNA made in *C. elegans* neurons can enter the germline and cause transgenerational gene silencing

Sindhuja Devanapally, Snusha Ravikumar, and Antony M. Jose<sup>1</sup>

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742

Edited by Gary Ruvkun, Massachusetts General Hospital, Boston, MA, and approved January 9, 2015 (received for review December 10, 2014)

An animal that can transfer gene-regulatory information from somatic cells to germ cells may be able to communicate changes in the soma from one generation to the next. In the worm *Caenorhabditis elegans*, expression of double-stranded RNA (dsRNA) in neurons can result in the export of dsRNA-derived mobile RNAs to other distant cells. Here, we show that neuronal mobile RNAs can cause transgenerational silencing of a gene of matching sequence in germ cells. Consistent with neuronal mobile RNAs being forms of dsRNA, silencing of target genes that are expressed either in somatic cells or in the germline requires the dsRNA-selective importer SID-1. In contrast to silencing in somatic cells, which requires dsRNA expression in each generation, silencing in the germline is heritable after a single generation of exposure to neuronal mobile RNAs. Although initiation of inherited silencing within the germline requires SID-1, a primary Argonaute RDE-1, a secondary Argonaute HRDE-1, and an RNase D homolog MUT-7, maintenance of inherited silencing is independent of SID-1 and RDE-1, but requires HRDE-1 and MUT-7. Inherited silencing can persist for >25 generations in the absence of the ancestral source of neuronal dsRNA. Therefore, our results suggest that sequence-specific regulatory information in the form of dsRNA can be transferred from neurons to the germline to cause transgenerational silencing.

epigenetics | mobile RNAs | soma to germline | Weismann barrier

The germline is separated from the rest of the body, or soma, during early development in most animals, consistent with the suggestion that environmental effects on soma throughout the lifetime of an animal cannot influence inheritance through the germline (1). However, some environmental changes can cause effects that last for three or more generations, even in the apparent absence of changes in the genotype (reviewed in ref. 2). These transgenerational epigenetic effects are presumably initiated either by direct changes within the ancestral germline or by the transfer of information from ancestral somatic cells to the ancestral germline. It is difficult to distinguish between these possibilities because complex ancestral changes that affect subsequent generations, such as diet (3–5) or endocrine disruption (6), perturb many genes in many tissues in ways that are as yet unclear. Manipulating the activity of a single gene in specific tissues and across generations can help distinguish between these possibilities. Such specific inactivation of a single gene can be achieved by using double-stranded RNA (dsRNA) to trigger RNA interference (RNAi) in the worm *Caenorhabditis elegans* (7).

As in most animals, the *C. elegans* germline is set aside early in development—after four cell divisions (8). Gene silencing initiated through RNAi-related mechanisms within the *C. elegans* germline can last for many generations (9–13). Such transgenerational silencing can be triggered by both injected dsRNA (14–16) and ingested dsRNA (16–19). However, both injection and ingestion can deliver dsRNA directly into the fluid-filled body cavity that surrounds the germline, without entry into the cytosol of any somatic cell (20, 21). Thus, it remains unknown

whether somatic cells in *C. elegans* can export signals for delivery into the germline to cause transgenerational gene silencing.

The transfer of gene-specific information from one somatic tissue to another somatic tissue during RNAi has been observed in *C. elegans* (22). Such intertissue transfer of gene-regulatory information appears to occur through the transport of forms of dsRNA called mobile RNAs (23). Entry of these mobile RNAs into the cytosol requires the dsRNA-selective importer SID-1 (22, 24, 25). Consequently, when dsRNA is expressed in a variety of somatic tissues such as the gut, muscles, or neurons, SID-1-dependent silencing of genes of matching sequence is observed in other somatic tissues (20). Because gene silencing by mobile RNAs from neurons appears to be stronger than that by mobile RNAs from other somatic tissues (20), we examined whether neurons export mobile RNAs that can enter the germline to cause transgenerational gene silencing.

Here, we show that neuronal mobile RNAs can enter both somatic and germ cells to trigger gene silencing. Although silencing in somatic tissues is not detectably inherited despite multigenerational exposure to neuronal mobile RNAs, silencing in the germline is inherited for many generations after a single generation of exposure to neuronal mobile RNAs.

## Results

**Neuronal Mobile RNAs Can Enter Most Somatic Tissues and the Germline.** Genetic analyses suggest that neuronal mobile RNAs are forms of dsRNA (23). Mobile RNAs generated from dsRNA expressed in neurons against the muscle gene *unc-22* can enter muscle cells through the dsRNA importer SID-1 and cause *unc-22* silencing (23). To examine silencing of a gene expressed in multiple

### Significance

The germline, which produces sperm or oocyte, is separated from other cells that generate the rest of the body, the soma, during early development in most animals. Somatic cells experience and respond to the environment in each generation, and it is unknown whether they can transmit information to the germline for inheritance into subsequent generations. We found that neurons of the worm *Caenorhabditis elegans* can transmit double-stranded RNA to the germline to initiate transgenerational silencing of a gene of matching sequence. To our knowledge, these results demonstrate for the first time that a somatic tissue of an animal can have transgenerational effects on a gene through the transport of double-stranded RNA to the germline.

Author contributions: S.D., S.R., and A.M.J. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: amjose@umd.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1423333112/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1423333112/-DCSupplemental).

tissues by a single source of dsRNA in neurons, we used animals that expressed cytosolic *gfp* (*Peft-3::gfp*) in all somatic tissues and *gfp-dsRNA* in all neurons (*Prgef-1::gfp-dsRNA*) (Fig. 1). GFP expression was detectably reduced in most somatic tissues (with the notable exception of the pharynx) in the presence of *Prgef-1::gfp-dsRNA* (Fig. 1*A*, *Top* vs. Fig. 1*A*, *Middle*), and this silencing was enhanced in the absence of the exonuclease ERI-1 (Fig. S1), consistent with ERI-1 acting to inhibit silencing by imported neuronal mobile RNAs (23). Silencing in all somatic tissues, even in the *eri-1(-)* background, was lost upon removal of the mobile RNA importer SID-1 (Fig. 1*A*, *Bottom*, and Fig. S1), suggesting that all observed silencing was due to mobile RNAs made in neurons.

To test whether the germline is susceptible to silencing by mobile RNAs, we examined silencing of GFP expression in animals that express *gfp* in the germline (*Pmex-5::gfp*) and neuronal mobile RNAs from a *Prgef-1::gfp-dsRNA* transgene. Like most somatic cells, the germline was susceptible to silencing by neuronal mobile RNAs, and the silencing was predominantly dependent on SID-1 (Fig. 1*B*). The silencing was sequence-specific and did not occur in animals with transgenic expression of a co-injection marker (Fig. S2*A*) or in animals with transgenic expression of *unc-22* dsRNA in neurons (Fig. S2*B*). Furthermore, silencing, as detected by the loss of GFP fluorescence within the germline (Fig. S3*A*), was correlated with a reduction in *gfp* mRNA levels (Fig. S3*B*). Consistent with mobile RNAs that are imported into the germline being forms of *gfp-dsRNA*, silencing was strongly dependent on the dsRNA importer SID-1 and the primary Argonaute RDE-1 that acts on short dsRNA (26), but independent of the RNA-dependent RNA polymerase RRF-1 that generates single-stranded secondary small RNAs in somatic cells (27) (Fig.

S3*C*). The residual silencing observed in *sid-1(-)* and *rde-1(-)* animals may reflect additional *sid-1-* and *rde-1-*independent gene silencing mechanisms that can act in the germline (9–12). Because silencing of a germline target due to dsRNA expression in neurons is greatly reduced in the absence of SID-1 (Fig. 1*B*, *Bottom*, and Fig. S3*C*), we conclude that SID-1-dependent neuronal mobile RNAs can enter the germline.

Together, our results suggest that neuronal mobile RNAs can enter most somatic tissues as well as the germline to silence genes of matching sequence. Because injection of in vitro-synthesized dsRNA can generate signals that are inherited in *C. elegans* (7, 14–16), our observations raise the possibility that neuronal mobile RNAs may also generate such inherited signals upon silencing a gene within the germline or upon silencing a gene in other somatic cells.


### Silencing in the Germline by Neuronal Mobile RNAs Is Inherited for Many Generations.

Injected or ingested dsRNA can cause transgenerational gene silencing of germline genes in *C. elegans* (10, 14–18). However, both forms of dsRNA delivery could result in the direct entry of dsRNA into the germline without entry into the cytoplasm of somatic cells. Ingested dsRNA is transcytosed across the gut into the body cavity that surrounds the germline (20, 21), and it is difficult to avoid spillage of injected dsRNA into the body cavity. These experimental considerations suggest that to test the possibility of somatic tissues initiating transgenerational gene silencing, it is necessary to express silencing triggers within somatic tissues and examine gene silencing within the germline. Although induction using heat shock of a transgene that encodes a viral genome in somatic tissues caused transgenerational silencing in *C. elegans* (13), such heat-shock induction also leads to expression within the germline (figure S5 in ref. 28). Therefore, because of the inherent difficulty in ensuring lack of expression within the germline from transgenes, only germline silencing that is reduced in the absence of the dsRNA importer SID-1 (Fig. 1*B*, *Bottom*, and Fig. S3*C*) can be interpreted as being caused by mobile RNAs.


To determine whether neuronal mobile RNAs that are imported into the germline can cause transgenerational silencing, we examined animals that lack the DNA for *gfp-dsRNA* but whose ancestors expressed neuronal dsRNAs. Because stable transgenic lines of extrachromosomal arrays are generated in *C. elegans* two generations after an animal [parental generation (P0)] is transformed with DNA (i.e., in the F2 generation) (29), we examined the silencing of GFP expression in wild-type animals of the F3 generation that lacked the *gfp-dsRNA* transgene and in their descendants (Fig. 2*A*, *Left*). Animals that lack the *gfp-dsRNA* transgene can be identified by the loss of a red fluorescent co-injection marker, the DNA for which is expected to be incorporated along with the DNA for *gfp-dsRNA* into a single extrachromosomal array upon cotransformation. All F3 animals without the extrachromosomal array showed silencing of GFP expression in the germline (Fig. 2*A*, *Right*). Inherited silencing due to the ancestral production of neuronal mobile RNAs persisted for >25 subsequent generations, despite unbiased passaging of worms from one generation to the next (Fig. 2*A*, *Right*, and Fig. S4). Consistent with the loss of the *gfp-dsRNA* transgene in animals that lack fluorescence from the co-injection marker, we failed to detect the *gfp-dsRNA* transgene in the DNA of worms that lacked the co-injection marker after 35 cycles of PCR amplification (Fig. 2*B*). These results suggest that neuronal mobile RNAs imported into the germline can initiate gene silencing that lasts for many generations in the absence of the ancestral source of neuronal dsRNA.

### Transgenerational Silencing by Neuronal Mobile RNAs Has Distinct Genetic Requirements for Initiation and Maintenance.

Although transgenerational silencing is reliably observed by using multiple transgenic sources of neuronal mobile RNAs (Fig. S4), the



**Fig. 1.** Neuronal mobile RNAs can cause gene silencing in most somatic tissues and in the germline. (*A*) Representative fourth larval (L4)-staged animals that express GFP (black) in somatic tissues (*Peft-3::gfp*) in a wild-type (*Top*) background and animals that in addition express dsRNA in neurons against *gfp* (*Prgef-1::gfp-dsRNA*) in wild-type (*Middle*) or *sid-1(-)* (*Bottom*) backgrounds are shown. Silenced tissues and unsilenced pharynx are indicated (*Middle*). Detectable silencing was observed in 100% of wild-type animals ( $n = 135$ ) and 0% of *sid-1(-)* animals ( $n = 115$ ). (Scale bars, 50  $\mu\text{m}$ .) Also see Fig. S1. (*B*) Representative L4-staged animals that express GFP (black) in the germline (*Pmex-5::gfp*; outlined in cyan) in a wild-type (*Top*) background and animals that in addition express *Prgef-1::gfp-dsRNA* in wild-type (*Middle*) or *sid-1(-)* (*Bottom*) backgrounds are shown. Because of the long exposure time required to acquire these images, variable and irregular autofluorescence due to gut granules was also detected. Detectable silencing was observed in 87% of wild-type animals ( $n = 54$ ) and 27% of *sid-1(-)* animals ( $n = 59$ ). (Scale bars, 50  $\mu\text{m}$ .)




**Fig. 2.** Neuronal mobile RNAs can cause transgenerational silencing of a germline gene. (A) Inherited silencing in the germline lasts for >25 generations after the source of neuronal mobile RNAs is lost. (A, Left) *Pmex-5::gfp* animals (P0) were injected with constructs to express neuronal mobile RNAs (*Prgef-1::gfp-dsRNA*) along with a co-injection marker (*Pmyo-2::DsRed*) to generate F2 transgenic lines (blue worm). (A, Right) The proportions of animals that all lack fluorescence from the co-injection marker (gray worm) but that show either strong (dark gray bars) or weak (light gray bars) silencing in the F3 generation and in successive generations (F4–F30) were determined. Error bars indicate 95% CI and  $n > 14$  L4-staged animals for each generation. Also see Fig. S4. Dark gray bars and light gray bars are as in Fig. S3C. (B) Animals that lack the co-injection marker also lack the *gfp-dsRNA* transgene. Genomic DNA from wild-type animals (no dsRNA), from wild-type animals that express *Prgef-1::gfp-dsRNA*, from *Pmex-5::gfp* animals, and from *Pmex-5::gfp* animals that either have or whose ancestors had extrachromosomal transgenes [i.e., *Pmex-5::gfp* animals that in addition express the co-injection marker alone (marker; orange worm) or along with *Prgef-1::gfp-dsRNA* (*Prgef-1::gfp-dsRNA & marker*; blue worm) or apparently lack these extrachromosomal transgenes (gray worm) but that were derived from ancestors that expressed these transgenes] were analyzed. Although the control gene was detected in all cases, a PCR product for the *gfp-dsRNA* transgene was detected only in wild-type animals with *gfp-dsRNA* and in *Pmex5::gfp* animals with *gfp-dsRNA* as evidenced by fluorescence from the co-injection marker.

number of generations that show silencing varied from one transgenic line to another, possibly due to differences in the levels of expression of dsRNA in different transgenic lines. To facilitate comparison of transgenerational silencing across multiple genetic backgrounds and to expose animals to mobile RNAs in defined generations, we chose a single extrachromosomal transgenic line that expresses neuronal mobile RNAs against *gfp* in wild-type animals and crossed it into animals that express *gfp* in the germline. This experimental scheme was then used to determine the genetic requirements for the initiation and maintenance of transgenerational gene silencing.


Using this experimental scheme, we found that exposure of a germline target gene to neuronal mobile RNAs for a single generation was sufficient to cause transgenerational silencing (Fig. 3A). Specifically, when animals with *Pmex-5::gfp* and animals with *Prgef-1::gfp-dsRNA* were mated, the F1 cross progeny that inherited the *Prgef-1::gfp-dsRNA* transgene could initiate transgenerational silencing. This silencing persisted for many

generations, despite the loss of the source of neuronal mobile RNAs in the F2 generation (Fig. 3A).

To test whether a gene is required for germline silencing by neuronal mobile RNAs, we used the same experimental scheme as above, but with animals that also had a mutation in the gene being tested (Fig. 3B). For example, to test the requirement for *sid-1*, we mated *sid-1* null mutants [*sid-1(-)*] that express *Pmex-5::gfp* with *sid-1(-)* animals that express *Prgef-1::gfp-dsRNA* and examined silencing in *sid-1(-)* animals of a later generation that express both *Pmex-5::gfp* and *Prgef-1::gfp-dsRNA*. Germline



**Fig. 3.** Neuronal mobile RNAs have distinct requirements for the initiation and maintenance of transgenerational silencing. (A) Expression of neuronal mobile RNAs for one generation is sufficient to initiate multigenerational silencing. *Pmex-5::gfp* animals were crossed with animals that express neuronal dsRNA from an extrachromosomal array (*Ex[gfp-ds]*) and the proportions of animals that lack the extrachromosomal array (gray worm) but that show either strong (dark gray bars) or weak (light gray bars) silencing in the F2 generation and in successive generations (F3–F10) were determined. The loss of *Ex[gfp-ds]* was determined by the loss of the fluorescent co-injection marker. (B) Initiation of silencing by neuronal mobile RNAs requires *sid-1*, *rde-1*, *hrde-1*, and *mut-7*. Wild-type (wt), *sid-1(-)*, *rde-1(-)*, *hrde-1(-)*, or *mut-7(-)* animals that all express *Pmex-5::gfp* were mated with animals of identical genetic backgrounds that all express neuronal dsRNA (*Ex[gfp-ds]*), and the silencing in descendants that had both *Pmex-5::gfp* and *Ex[gfp-ds]* was measured as in A. (C) Maintenance of germline gene silencing by neuronal mobile RNAs requires HRDE-1 and MUT-7, but not SID-1 or RDE-1. Wild-type (wt), *sid-1(+/-)*, *rde-1(+/-)*, *hrde-1(+/-)*, or *mut-7(+/-)* animals that all had both *Pmex-5::gfp* and *Ex[gfp-ds]* were allowed to have progeny, and the silencing in wild-type (wt), *sid-1(-)*, *rde-1(-)*, *hrde-1(-)*, or *mut-7(-)* grand progeny animals that all had *Pmex-5::gfp* but that all lacked *Ex[gfp-ds]* was measured as in A. The analyzed grand progeny were progeny of animals that also lacked *Ex[gfp-ds]*. Error bars indicate 95% CI. \* $P < 0.05$ .  $n > 19$  L4-staged animals, except for *mut-7(-)* animals in C, where  $n = 10$  L4-staged animals. Dark gray bars and light gray bars are as in Fig. S3C.



**Fig. 4.** Silencing of a somatic gene by neuronal mobile RNAs in parents is not detectably inherited by progeny. (A) Neuronal mobile RNAs can silence GFP expression in gut cells. Representative L4-staged animals that express GFP (black) in all somatic cells (*Psur-5::sur-5::gfp*) (Upper) or that in addition express dsRNA in all neurons (*Prgef-1::gfp-dsRNA*) (Lower) are shown. Brackets indicate strongly silenced gut nuclei. (Scale bars, 50  $\mu$ m.) Also see Fig. S6. (B) Double-stranded RNAs expressed in neurons against *gfp* require SID-1 to silence GFP expression in gut cells. The numbers of GFP-expressing gut nuclei were counted in wild-type animals that do not express dsRNA against *gfp* (no dsRNA; gray) and in wild-type, or *sid-1(-)* animals that express *Prgef-1::gfp-dsRNA* (blue). Gray line indicates average number of gut nuclei in L4-staged animals,  $n > 19$  L4-staged animals, and red bar in box plots indicates median.  $*P < 0.05$  (Student's *t* test). (C–E) An enhanced RNAi background [*eri-1(-)*] was used to maximize the ability to detect inherited silencing. (C) Unlike silencing by ingested dsRNA, silencing by neuronal mobile RNAs is not detectably inherited by progeny. Numbers of GFP-positive gut nuclei in genetically identical progeny of animals that were not exposed to *gfp-dsRNA* (none) or that were exposed to ingested *gfp-dsRNA* or that had one copy of an integrated transgene that expresses *Prgef-1::gfp-dsRNA* (*gfp-ds/+*) were counted. Errors indicate SEM. (D) Unbiased passaging of worms for multiple generations can lead to small differences in gene silencing. Worms that express *Prgef-1::gfp-dsRNA* (P0) were passaged for five generations (F1–F5) by picking a random worm at each generation, and the numbers of GFP-positive gut nuclei in animals of each generation were determined (see Fig. S8 for additional data). Gray line,  $n$ , red bar, and asterisks are as in B, except for F5, which had  $n = 8$  L4-staged animals. (E) SID-1 is required for silencing by neuronal mobile RNAs even after 17 generations of ancestral silencing by neuronal mobile RNAs. (E, Left) Schematic of experimental design to test the requirement for SID-1 in each generation for silencing by neuronal mobile RNAs. At each generation, the numbers of GFP-positive gut nuclei in *sid-1(-/-)* animals were counted, and heterozygous [*sid-1(+/-)*] siblings of any *sid-1(-/-)* animal (F1–F11) or heterozygous siblings of the most silenced *sid-1(-/-)* animal (F12–F18) were passaged. (E, Right) The extent of silencing in F1, F11, and F18 are shown (see Fig. S9 for additional data). Gray line, red bar, and  $n$  are as in B.

silencing using this experimental scheme also required SID-1 and RDE-1, in agreement with the results obtained for silencing by

transgenic lines that were independently generated in mutant backgrounds (Fig. S3C). Thus, germline silencing due to neuronal mobile RNAs likely relies on the import of forms of dsRNA through SID-1 and subsequent processing by the primary Argonaute RDE-1 within the germline. Further processing within the germline leads to the production of secondary single-stranded small RNAs. These secondary small RNAs eventually cause gene silencing through mechanisms that require many proteins (reviewed in ref. 30), including the secondary nuclear Argonaute HRDE-1 (18) and the RNase D homolog MUT-7 (31). We found that both HRDE-1 and MUT-7 were required for silencing by neuronal mobile RNAs, suggesting that silencing within the germline is executed by secondary small RNAs and downstream genes. For all genes tested above, the source of neuronal mobile RNAs was present in the animals that were tested. Therefore, the lack of silencing in *sid-1(-)*, *rde-1(-)*, *hrde-1(-)*, and *mut-7(-)* animals reflects a requirement for the corresponding genes in the initiation of germline silencing by neuronal mobile RNAs.

The observed genetic requirements for silencing by neuronal mobile RNAs are distinct from those observed for silencing by ingested or injected dsRNA. Whereas the requirement for SID-1 and RDE-1 is in agreement with the requirement for these genes when silencing is triggered using ingested and injected dsRNA, the requirement for HRDE-1 and MUT-7 is in contrast to the HRDE-1-independent silencing observed in response to ingested dsRNA (ref. 18 and Fig. S5) and the MUT-7-independent silencing observed in response to injected dsRNA (14). These differences might reflect differences in the dosage of dsRNA delivered into the germline using the different methods or the differential engagement of silencing machinery by the different sources of dsRNA used to trigger gene silencing.

To test whether a gene is required for the maintenance of transgenerational silencing by neuronal mobile RNAs, we examined silencing in animals that had mutations in the gene but were descendants of ancestors that had a wild-type copy of the gene during exposure to neuronal mobile RNAs (Fig. 3C). For example, to test the requirement for *sid-1* in the maintenance of transgenerational silencing, we examined silencing in *sid-1* null mutants [*sid-1(-)*] that were grand progeny of *sid-1(+/-)* heterozygous animals that were exposed to *Prgef-1::gfp-dsRNA*. Grand progeny were examined for silencing instead of progeny because maternal deposition of mRNA or protein from heterozygous parents can complicate interpretation of results in *sid-1(-)* progeny. We observed silencing in the *sid-1(-)* grand progeny of *sid-1(+/-)* heterozygous animals that were exposed to *Prgef-1::gfp-dsRNA*, which suggests that SID-1 is not required for the maintenance of transgenerational silencing. Similar experiments with null mutants of *rde-1*, *hrde-1*, and *mut-7* revealed that RDE-1, like SID-1, is dispensable for the maintenance of transgenerational silencing, but HRDE-1 and MUT-7 are required for the maintenance of transgenerational silencing.

In summary, our results suggest a model where mobile RNAs exported from neurons enter the germline through SID-1 to cause RDE-1-, MUT-7-, and HRDE-1-dependent silencing in the parent, which is subsequently maintained through a MUT-7- and HRDE-1-dependent, but SID-1- and RDE-1-independent mechanism. Because HRDE-1 has been shown to use secondary small RNAs to guide trimethylation of the histone H3 on lysine 9 (H3K9me3) at genes of matching sequence (17, 18), our results suggest that the initiation and maintenance of transgenerational silencing by neuronal mobile RNAs is associated with the deposition of H3K9me3 marks on genes of matching sequence. Although the response to ingested or injected dsRNA strongly suggests that secondary small RNAs are inherited (14, 17, 19), it is possible that in response to neuronal mobile RNAs chromatin marks are inherited across generations. Furthermore, although silencing of somatic genes has been reported to be inherited for

a few generations when the silencing is triggered by using ingested dsRNA (17, 19), it is unclear whether silencing of a somatic gene by neuronal mobile RNAs is inherited and whether transgenerational silencing by neuronal mobile RNAs within the germline can spread to somatic cells.

**Silencing in Somatic Cells by Neuronal Mobile RNAs Is Not Detectably Inherited.** To measure silencing by neuronal mobile RNAs in somatic cells, we used animals that have two different integrated transgenes—one that expresses nuclear-localized GFP (*sur-5::gfp*) under the control of a promoter that drives expression in all somatic cells (*P<sub>sur-5</sub>*) and one that expresses *gfp-dsRNA* under the control of a promoter that drives expression in all neurons (*P<sub>prgef-1</sub>*). Silencing due to neuronal mobile RNAs made from the *P<sub>prgef-1</sub>::gfp-dsRNA* transgene results in a reduction in fluorescence of nuclear-localized GFP made from the *P<sub>sur-5</sub>::sur-5::gfp* transgene (Fig. 4A). This silencing can be most easily observed in the large intestinal cell nuclei, and counting the number of GFP-positive gut nuclei provides a reliable measure of silencing that correlates with reduction in *gfp* mRNA levels (Fig. S6A and B). Wild-type animals with neuronal mobile RNAs had, on average, fewer GFP-positive gut nuclei than did animals without neuronal mobile RNAs (Fig. 4B; 24.2 vs. 29.9 GFP-positive gut nuclei;  $P < 0.05$ ). Consistent with silencing by neuronal mobile RNAs, this silencing was abolished in *sid-1(-)* animals (Fig. 4B) and not observed in wild-type animals that were merely cocultured with animals that express neuronal mobile RNAs (Fig. S6C).

Because the initiation of inherited silencing occurs more frequently in animals that lack the exonuclease ERI-1 (16), we examined the ability of neuronal mobile RNAs to trigger inherited silencing in an *eri-1(-)* background, where trace amounts of dsRNA (32) and additional mobile RNAs (20) made from the multicopy *P<sub>sur-5</sub>::sur-5::gfp* transgene could also contribute to silencing. Using this sensitive genetic background, we did not detect inherited silencing by neuronal mobile RNAs (Fig. 4C), but did detect inherited silencing by ingested dsRNAs as reported earlier (Fig. 4C and ref. 19). We noticed a correlation between an increase in silencing by neuronal mobile RNAs and an increase in parental or ancestral exposure to mobile RNAs


(Fig. S7). However, the increases in silencing were small and comparable to the small variations in silencing observed in successive generations when worms with *P<sub>prgef-1</sub>::gfp-dsRNA* of identical genotype were simply passaged (Fig. 4D). Furthermore, selection of the most silenced or most desilenced animal for four generations introduced marginal differences in silencing between the first and fifth generations (Fig. S8). Nevertheless, if marginal increases in inherited silencing accrued over many generations due to the presence of parental neuronal mobile RNAs, such inherited silencing might become independent of neuronal mobile RNAs and thus independent of SID-1 in later generations. However, we did not detect such SID-1-independent silencing, even after exposure to 17 generations of silencing by neuronal mobile RNAs (Fig. 4E and Fig. S9). The requirement for *sid-1* in every generation for silencing by neuronal mobile RNAs suggests that transport of neuronal mobile RNAs must occur in every generation to observe silencing in somatic cells.

The absence of robust inherited silencing by neuronal mobile RNAs of genes expressed in somatic cells could be either because somatic silencing does not generate signals for transmission to the next generation or because such signals require a template of matching sequence in the germline for stability. To test this latter possibility, we examined inherited somatic silencing by neuronal mobile RNAs in animals that express the target gene (*gfp*) in somatic cells as well as in the germline either from a single transgene (germline expression due to *P<sub>mex-5</sub>::gfp* and pharyngeal expression due to an additional promoter in the *P<sub>mex-5</sub>::gfp* transgene) (Fig. 5A) or from two separate transgenes (germline expression due to *P<sub>mex-5</sub>::gfp* and gut expression due to *sur-5::gfp*) (Fig. 5B). In both cases, no inherited silencing was detected in somatic cells.

Together, our results suggest that neuronal mobile RNAs generate transgenerational silencing signals that have a strong effect on gene expression in the germline and a minimal effect, if any, on gene expression in somatic tissues.

## Discussion

We found that neurons can transport forms of dsRNA into the germline to cause silencing that can last for many generations



**Fig. 5.** Inherited silencing of a germline gene by neuronal mobile RNAs in the parent does not spread to the soma of the progeny. (A) Silencing of GFP expression within the germline by neuronal mobile RNAs does not cause detectable inherited silencing of GFP expressed from the same locus in the pharynx of progeny. *Pmex-5::gfp* animals and *Pmex-5::gfp* animals that in addition express an extrachromosomal source of either the co-injection marker (*Ex[marker]*) or neuronal mobile RNAs (*Ex[gfp-dsRNA]*) were passaged and L4-staged progeny that lack the extrachromosomal arrays were imaged under identical conditions. The pharyngeal expression of GFP (black) is from an additional uncharacterized promoter (*P<sub>phar</sub>::gfp*) within the *Pmex-5::gfp* transgene and is absent in wild-type worms. Germline (outlined in cyan), GFP expression in the germline nuclei (cyan brackets), and GFP expression in pharyngeal nuclei (purple brackets) are indicated. (Scale bars, 50  $\mu$ m.) (B) Silencing of GFP expression within the germline by neuronal mobile RNAs in the parent does not cause detectable inherited silencing of GFP expressed from a different locus in gut cells of progeny. *Pmex-5::gfp* animals (P0 hermaphrodite) that in addition expressed *Ex[marker]* or *Ex[gfp-dsRNA]* were crossed with *Psur-5::sur-5::gfp* animals (P0 male), and the numbers of GFP-positive gut nuclei were counted in the resulting F1 progeny that lack extrachromosomal arrays. Errors indicate SEM, and  $n > 18$  L4-staged animals.

and that such transgenerational silencing is restricted to the germline with distinct genetic requirements for initiation and maintenance.

Mobile RNAs that enter the germline can provide an organism with the ability to transfer gene-specific regulatory information from somatic cells across generations and could be one mechanism by which the environment elicits transgenerational effects in animals. Although restricted to the germline, transgenerational silencing by mobile RNAs could underlie effects of the environment across generations in some cases. For example, expression of some genes within the germline can affect longevity (33), and transgenerational silencing of such genes might underlie the longevity that results from ancestral starvation in *C. elegans* (5). Thus, additional experiments are needed to determine the role of mobile RNAs, if any, in the transport of such experience-dependent information from somatic cells to subsequent generations in *C. elegans*.

The presence of a mammalian homolog of the dsRNA importer SID-1 that is also required for the uptake of dsRNAs into cells (34) raises the possibility that dsRNA generated from distant somatic cells—potentially in response to environmental influences—may be imported through SID-1 into the mammalian germline to trigger transgenerational epigenetic changes. Consistent with this possibility, small RNAs have been found in circulation in mammals (35); dsRNAs have been detected in mammalian germ cells (36–38); and injection of RNAs into the early mouse embryo can trigger epigenetic silencing (39). However, even if RNAs from somatic cells are transported to the germline in mammals, they may not always initiate transgenerational inherited effects because they have to escape mechanisms that reprogram epigenetic information

in each generation (40). Additional studies are required to determine whether specific mechanisms prevent environmental influences from triggering transmission of information in the form of mobile RNAs from somatic cells to the germline.

## Materials and Methods

All *C. elegans* strains were generated and maintained by using standard methods (41). Transgenic animals were generated by injecting PCR fragments or plasmids into the germline (29) of wild-type or mutant animals, and transgenes were also introduced into different genetic backgrounds through genetic crosses. Visible markers were used to balance *sid-1(-)* and to mark *Pmex-5::gfp*. Silencing of *Pmex-5::gfp* and *Peft-3::gfp* were measured by imaging under identical nonsaturating conditions using a Nikon AZ100 microscope. Silencing of *sur-5::gfp* was quantified by counting the number of gut nuclei that showed GFP expression above a fixed threshold of brightness. DNA of *Prgef-1::gfp-dsRNA* and *Pmex-5::gfp* transgenes were detected by using PCR in crosses and in inheritance experiments. Semi-quantitative RT-PCR was used to determine relative mRNA levels by carrying out reverse transcription with gene-specific primers for *gfp* and *tbb-2* followed by <31 cycles of PCR. Inherited silencing by feeding RNAi (19) and statistical analyses (20) were performed as described earlier. Detailed procedures are provided in *SI Materials and Methods*.

**ACKNOWLEDGMENTS.** We thank Tom Kocher, Geraldine Seydoux, Norma Andrews, Steve Mount, Steve Wolniak, Karen Carleton, and members of the A.M.J. laboratory for critical reading of the manuscript; the *Caenorhabditis elegans* Genetic Stock Center and the Hunter laboratory (Harvard University) for some worm strains; the Hamza laboratory (University of Maryland) for bacteria that express *gfp-dsRNA*; and Amy Beaven from the Department of Cell Biology and Molecular Genetics imaging core for microscopy advice. This work was supported by National Institutes of Health Grants R00GM085200 and R01GM111457 (to A.M.J.).

- Weismann A (1883) On heredity. *Essays upon Heredity and Kindred Biological Problems*, eds Poulton EB, Schonland S, Shipley AE (Clarendon, Oxford), pp 67–106.
- Heard E, Martienssen RA (2014) Transgenerational epigenetic inheritance: Myths and mechanisms. *Cell* 157(1):95–109.
- Carone BR, et al. (2010) Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell* 143(7):1084–1096.
- Ng SF, et al. (2010) Chronic high-fat diet in fathers programs  $\beta$ -cell dysfunction in female rat offspring. *Nature* 467(7318):963–966.
- Rechavi O, et al. (2014) Starvation-induced transgenerational inheritance of small RNAs in *C. elegans*. *Cell* 158(2):277–287.
- Anway MD, Cupp AS, Uzumcu M, Skinner MK (2005) Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* 308(5727):1466–1469.
- Fire A, et al. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391(6669):806–811.
- Strome S (2005) Specification of the germ line. *WormBook*, ed The *C. elegans* Research Community, 10.1895/wormbook.1.9.1, www.wormbook.org.
- Shirayama M, et al. (2012) piRNAs initiate an epigenetic memory of nonself RNA in the *C. elegans* germline. *Cell* 150(1):65–77.
- Ashe A, et al. (2012) piRNAs can trigger a multigenerational epigenetic memory in the germline of *C. elegans*. *Cell* 150(1):88–99.
- Lee HC, et al. (2012) *C. elegans* piRNAs mediate the genome-wide surveillance of germline transcripts. *Cell* 150(1):78–87.
- Luteijn MJ, et al. (2012) Extremely stable Piwi-induced gene silencing in *Caenorhabditis elegans*. *EMBO J* 31(16):3422–3430.
- Rechavi O, Minevich G, Hobert O (2011) Transgenerational inheritance of an acquired small RNA-based antiviral response in *C. elegans*. *Cell* 147(6):1248–1256.
- Grishok A, Tabara H, Mello CC (2000) Genetic requirements for inheritance of RNAi in *C. elegans*. *Science* 287(5462):2494–2497.
- Alcazar RM, Lin R, Fire AZ (2008) Transmission dynamics of heritable silencing induced by double-stranded RNA in *Caenorhabditis elegans*. *Genetics* 180(3):1275–1288.
- Vastenhouw NL, et al. (2006) Gene expression: Long-term gene silencing by RNAi. *Nature* 442(7105):882.
- Gu SG, et al. (2012) Amplification of siRNA in *Caenorhabditis elegans* generates a transgenerational sequence-targeted histone H3 lysine 9 methylation footprint. *Nat Genet* 44(2):157–164.
- Buckley BA, et al. (2012) A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature* 489(7416):447–451.
- Burton NO, Burkhart KB, Kennedy S (2011) Nuclear RNAi maintains heritable gene silencing in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 108(49):19683–19688.
- Jose AM, Smith JJ, Hunter CP (2009) Export of RNA silencing from *C. elegans* tissues does not require the RNA channel SID-1. *Proc Natl Acad Sci USA* 106(7):2283–2288.
- Calixto A, Chelur D, Topalidou I, Chen X, Chalfie M (2010) Enhanced neuronal RNAi in *C. elegans* using SID-1. *Nat Methods* 7(7):554–559.
- Winston WM, Molodowitch C, Hunter CP (2002) Systemic RNAi in *C. elegans* requires the putative transmembrane protein SID-1. *Science* 295(5564):2456–2459.
- Jose AM, Garcia GA, Hunter CP (2011) Two classes of silencing RNAs move between *Caenorhabditis elegans* tissues. *Nat Struct Mol Biol* 18(11):1184–1188.
- Feinberg EH, Hunter CP (2003) Transport of dsRNA into cells by the transmembrane protein SID-1. *Science* 301(5639):1545–1547.
- Shih JD, Hunter CP (2011) SID-1 is a dsRNA-selective dsRNA-gated channel. *RNA* 17(6):1057–1065.
- Tabara H, et al. (1999) The *rde-1* gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* 99(2):123–132.
- Sijen T, et al. (2001) On the role of RNA amplification in dsRNA-triggered gene silencing. *Cell* 107(4):465–476.
- Sheth U, Pitt J, Dennis S, Priess JR (2010) Perinuclear P granules are the principal sites of mRNA export in adult *C. elegans* germ cells. *Development* 137(8):1305–1314.
- Mello CC, Kramer JM, Stinchcomb D, Ambros V (1991) Efficient gene transfer in *C. elegans*: Extrachromosomal maintenance and integration of transforming sequences. *EMBO J* 10(12):3959–3970.
- Grishok A (2013) Biology and mechanisms of short RNAs in *Caenorhabditis elegans*. *Adv Genet* 83:1–69.
- Ketting RF, Haverkamp TH, van Luenen HG, Plasterk RH (1999) Mut-7 of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. *Cell* 99(2):133–141.
- Hellwig S, Bass BL (2008) A starvation-induced noncoding RNA modulates expression of Dicer-regulated genes. *Proc Natl Acad Sci USA* 105(35):12897–12902.
- Greer EL, et al. (2011) Transgenerational epigenetic inheritance of longevity in *Caenorhabditis elegans*. *Nature* 479(7373):365–371.
- Wolfrum C, et al. (2007) Mechanisms and optimization of in vivo delivery of lipophilic siRNAs. *Nat Biotechnol* 25(10):1149–1157.
- Arroyo JD, et al. (2011) Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci USA* 108(12):5003–5008.
- Song R, et al. (2011) Male germ cells express abundant endogenous siRNAs. *Proc Natl Acad Sci USA* 108(32):13159–13164.
- Tam OH, et al. (2008) Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 453(7194):534–538.
- Watanabe T, et al. (2008) Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 453(7194):539–543.
- Rassoulzadegan M, et al. (2006) RNA-mediated non-mendelian inheritance of an epigenetic change in the mouse. *Nature* 441(7092):469–474.
- Feng S, Jacobsen SE, Reik W (2010) Epigenetic reprogramming in plant and animal development. *Science* 330(6004):622–627.
- Brenner S (1974) The genetics of *Caenorhabditis elegans*. *Genetics* 77(1):71–94.