



HHS Public Access

Author manuscript

J Subst Abuse Treat. Author manuscript; available in PMC 2016 April 01.

Published in final edited form as:

J Subst Abuse Treat. 2015 April ; 51: 1–18. doi:10.1016/j.jsat.2014.09.001.

Brief Alcohol Interventions for Adolescents and Young Adults: A Systematic Review and Meta-analysis

Emily E. Tanner-Smith, Ph.D.* and Mark W. Lipsey, Ph.D.

Peabody Research Institute, Vanderbilt University

Abstract

This study reports findings from a meta-analysis summarizing the effectiveness of brief alcohol interventions for adolescents (age 11-18) and young adults (age 19-30). We identified 185 eligible study samples using a comprehensive literature search and synthesized findings using random-effects meta-analyses with robust standard errors. Overall, brief alcohol interventions led to significant reductions in alcohol consumption and alcohol-related problems among adolescents ($\bar{g} = 0.27$ and $\bar{g} = 0.19$) and young adults ($\bar{g} = 0.17$ and $\bar{g} = 0.11$). These effects persisted for up to one year after intervention and did not vary across participant demographics, intervention length, or intervention format. However, certain intervention modalities (e.g., motivational interviewing) and components (e.g., decisional balance, goal-setting exercises) were associated with larger effects. We conclude that brief alcohol interventions yield beneficial effects on alcohol-related outcomes for adolescents and young adults that are modest but potentially worthwhile given their brevity and low cost.

Keywords

adolescents; brief alcohol intervention; meta-analysis; systematic review; young adults

1. Introduction

A sizeable portion of adolescents and young adults engage in heavy episodic consumption of alcohol, and thus put themselves at risk for numerous detrimental consequences related to their physical, mental, and social well-being (Brown et al., 2009). In 2011, for example, the estimated rate of past month binge drinking (five or more drinks on the same occasion for males, four or more for females) was 15% for 16-17 year olds, 31% for 18-20 year olds, and 45% for 21-25 year olds (SAMHSA, 2012). In response, a growing body of research has sought to identify early intervention programs that are effective for preventing or delaying the initiation of alcohol use, or intervening with heavier users before they progress to more

© 2014 Elsevier Inc. All rights reserved

*Corresponding author. Vanderbilt University, Peabody Research Institute, Department of Human and Organizational Development, Box 0181 GPC, Vanderbilt University, Nashville, TN 37203-5721; Phone 615-322-6304; Fax 615-322-0293; e.tanner-smith@vanderbilt.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

problematic levels of use. One approach is a brief intervention, defined broadly here as an intervention aimed at providing motivation for behavior change in a relatively circumscribed time (one to five sessions). Brief interventions are attractive primarily because of their brevity and the varied settings in which they can be conveniently delivered. If effective, they may therefore offer a cost-effective way to address a potentially lethal public health problem (Fleming et al., 2002; Neighbors, Barnett, Rohsenow, Colby, & Monti, 2010; Wutzke, Shiell, Gomel, & Conigrave, 2001).

Prior research reviews have found that brief interventions are indeed generally effective in reducing alcohol consumption among adolescents and young adults (e.g., Carey, Scott-Sheldon, Carey, & DeMartini, 2007; Carey, Scott-Sheldon, Elliott, Bolles, & Carey, 2009; Tait & Hulse, 2003). However, these reviews have not fully explored the characteristics of the participants and interventions that are associated with the strongest intervention effects or the persistence of those effects over time. The accumulating research is ripe for a comprehensive meta-analysis that examines how much, when, for whom, and for how long such interventions are effective in this population—information that can guide future research and aid practitioners planning to implement brief alcohol interventions.

1.1 Brief Alcohol Interventions for Adolescents and Young Adults

The defining characteristic of a brief intervention is the relatively brief contact time—generally one to five sessions—with a provider such as a physician, nurse, psychologist, counselor, or other service professional. In other regards, these interventions vary considerably, e.g., in length, structure, targets, media communication, underpinning theory, and intervention philosophy (Heather, 1995). Brief interventions are typically not intended to provide a full treatment regimen for individuals with alcohol use disorders but, rather, are designed to motivate and provide resources to participants to help them moderate their alcohol consumption, or, if needed, seek more intensive treatment options. As such, they can be used as universal, selective, or indicated prevention strategies (Barry, 1999). Most brief alcohol interventions include at least one of the following components: a discussion of alcohol consumption, feedback on risk or levels of use, comparisons to local or national norms, information on potential harms, or coping strategies and goal-setting plans for dealing with drinking situations. These therapeutic components are most often based on the principles of cognitive behavioral therapy, motivational interviewing (Miller & Rollnick, 1991), the transtheoretical model of behavior change (ProChaska & DiClemente, 1984), or social norms theory (Berkowitz, 2004), all of which emphasize the stimulation of participants' abilities, capacities, and motivations to self-evaluate and self-regulate their behaviors.

Since the launch of the Substance Abuse and Mental Health Services Administration's Screening, Brief Intervention, and Referral to Treatment Initiative in 2003 (SAMHSA, 2014), there has been an expanding body of methodologically rigorous research on brief interventions targeting alcohol and illicit substance use. Prior meta-analyses have shown that brief interventions are capable of reducing alcohol use among adults (Ballesteros, Duffy, Querejeta, Arino, Gonzalez-Pinto, 2004; Beich, Thorsen, & Rollnick, 2003; Bertholet, Daepfen, Wietlisbach, Fleming, & Burnand, 2005; Bien, Miller, & Tonigan, 1993; Burke,

Arkowitz, & Menchola, 2003; Kaner et al., 2007; Poikolainen, 1999; Vasilaki, Hosier, & Cox, 2006; Wilk, Jensen, & Havighurst, 1997). Their effectiveness for younger populations, however, has been less well documented despite the prevalence of excessive alcohol use among youth. The one meta-analysis of which we are aware that focused exclusively on brief interventions for adolescents found a moderate positive benefit for alcohol use outcomes (Cohen's $d = 0.28$), but included only 11 studies (Tait & Hulse, 2003). Although several existing meta-analyses have focused on alcohol interventions for college-age students (e.g., Carey et al., 2007; Carey et al., 2009; Fachini, Aliane, Martinez, & Furtado, 2012; Moreira, Smith, & Foxcroft, 2009; Scott-Sheldon, DeMartini, Carey, & Carey, 2009), most of these reviews either included brief interventions among other types of alcohol interventions, or only focused on a specific branded intervention program (e.g., Brief Alcohol Screening and Intervention for College Students [BASICS]). The most comprehensive meta-analysis to date (Carey et al., 2007) focused on 62 studies of individually delivered alcohol interventions for college students and reported positive effects for alcohol outcomes ($d = 0.17-0.18$), although these effects were significantly attenuated over longer follow-up periods.

The large and growing body of empirical research assessing the effectiveness of brief alcohol interventions for adolescents and young adults is now sufficient to support a comprehensive systematic review and meta-analysis that can investigate the sources of variability in outcomes in more depth than has been possible in prior reviews. In particular, the current meta-analysis examines how much, when, for whom, and for how long brief alcohol interventions may be effective in youth populations.

1.2 Sources of Variability in the Effects of Brief Alcohol Interventions

The growing support for brief alcohol interventions is leading researchers and practitioners to call for research to move beyond questions of whether they work to questions of what makes them work and in which populations and conditions they work best (Cunningham et al., 2009; Nilsen, Kaner, & Babor, 2008). The diversity of brief interventions, for instance, makes it important to investigate the extent to which the effects vary with the characteristics of the interventions themselves. A central question in that regard is just how brief the interventions can be and still be effective. Some researchers have suggested that three to four intervention sessions with a few follow-up calls or brief visits are most effective (Fleming et al., 2002), whereas others have maintained that one hour or even five minute interventions can be effective, at least with college students (Kulesza, Apperson, Larimer, & Copeland, 2010).

Other intervention characteristics that may moderate treatment effects include the primary intervention modality (e.g., motivational interviewing, cognitive behavioral), specific intervention components (e.g., providing advice, personalized normative feedback, goal-setting exercises), the delivery site (e.g., emergency room, school/university), and the delivery mode (e.g., computerized, in-person) (Bewick et al., 2008; Whitlock, Polen, Green, Orleans, & Klein, 2004; Winters & Leitten, 2007).

Another question is whether brief interventions are more effective for some types of participants than others (Cunningham et al., 2009). Among young populations, a key

distinction is between adolescents who are 18 years old and under, for whom alcohol consumption is almost universally illegal, and young adults of college age, many of whom can drink legally and are in the prime years for doing so (SAMHSA, 2012). Gender, race, and baseline levels of alcohol use may also be important individual characteristics associated with intervention effectiveness (Bien et al., 1993; Kaner et al., 2007; Poikolainen, 1999; Walton et al., 2008).

Still another question is whether, given their brevity, brief interventions can produce effects that are sustained. In their meta-analysis of (mostly brief) alcohol interventions for college students, Carey and colleagues (2007) found that intervention effects were attenuated to non-significance by 27 weeks of follow-up. However, they had limited data with which to investigate this issue, so there is a question as to how generalizable these results are for other populations and across different intervention approaches.

The study reported here used meta-analytic methods to synthesize the empirical findings from brief alcohol intervention research with adolescents (age 11-18) and young adults (age 19-30) with particular emphasis on identifying variables related to differential effects. Specifically, this meta-analysis examined: 1) the overall effects of brief alcohol interventions on adolescent and young adults' alcohol consumption and alcohol-related problems, 2) the variation in effects associated with intervention and participant characteristics, and 3) the persistence of the effects of brief alcohol interventions.

2. Materials and Methods

2.1 Inclusion and Exclusion Criteria

Eligible studies were those focused on brief interventions explicitly aimed at reducing participants' alcohol use or alcohol-related problems. Interventions could target any risk level (universal, selective, or indicated) of participants, as long as they involved five or fewer hours of total contact time and four or fewer weeks between the first and last intervention session (excluding booster sessions). Specifying these criteria for the maximum amount of contact allowable for an intervention to be considered brief was necessary to make reliable decisions about which studies were eligible for inclusion in the review. The criteria used here were guided, first, by the definitions used in other discussions of brief alcohol interventions (e.g., Barry, 1999). We also examined the distributions of contact time and treatment duration values found in the studies included in our meta-analysis of outpatient treatment for adolescents with substance use disorders (Tanner-Smith, Wilson, & Lipsey, 2013). These were bimodal and five hours of contact time and four weeks duration marked points that generally separated the shorter interventions from more extensive outpatient treatment programs.

Eligible studies had to include comparison conditions of no treatment, a wait-list control, or some form of routine treatment as usual (i.e., services the participants would have received even in the absence of the brief intervention). Studies that compared two types of interventions were not eligible. Eligible participant samples included adolescents and young adults, defined as individuals age 11-25. Samples comprised entirely of undergraduate college students were also eligible even though they may have included students over the

age of 25, but no older than age 30. Eligible research designs included randomized controlled designs and controlled quasi-experimental designs that provided enough information to permit estimation of a pretest effect size that could be used (by us or the primary study authors) to adjust the posttest effect estimates for any initial group differences. Eligible studies were required to assess intervention effects on at least one outcome variable that measured alcohol use or alcohol-related problems (e.g., DUI/DWI). To be applicable to contemporary youth, only studies conducted in 1980 or later were included in the review. There were no geographic or language restrictions on eligibility.

2.2 Search Strategy

Using a comprehensive search strategy, we attempted to identify and retrieve the entire population of published and unpublished studies that met the aforementioned inclusion criteria. The following electronic bibliographic databases were searched, current through December 31, 2012: CINAHL, Clinical Trials Register, Dissertation Abstracts International, ERIC, International Bibliography of the Social Sciences, NIH RePORTER, PsycARTICLES, PsycINFO, PubMed, Social Services Abstracts, Sociological Abstracts, and WorldWideScience.org. We also searched the following sources in an attempt to locate grey literature: Australasian Medical Index, Campbell Collaboration Library, Canadian Evaluation Society's Grey Literature Database, Chestnut Health Systems website, Cochrane Collaboration CENTRAL, College on Problems of Drug Dependence conference presentations, EPPI-Centre Database of Health Promotion Research, Google Scholar, Index to Theses in Great Britain and Ireland, International Clinical Trials Registry, Joint Meeting on Adolescent Treatment Effectiveness conference presentations, KoreaMed, NIAAA website, NTIS, OpenSIGLE, SAMHSA website, Social Care Online, and SveMed+. We checked the bibliographies of all screened and eligible studies, as well as the bibliographies of prior narrative reviews and meta-analyses. We also conducted hand-searches in *Alcoholism: Clinical and Experimental Research*, *American Journal on Addictions*, and the *Journal of Substance Abuse Treatment*.

2.3 Screening and Coding Procedures

Under the supervision of the first author, a team of six master's level research assistants first screened all abstracts and titles resulting from the search to eliminate clearly irrelevant study reports. All six researchers initially screened 500 randomly selected abstracts/titles and discussed disagreements until 100% consensus was reached. The remaining abstracts/titles were screened by one team member, with the first author reviewing all their screening decisions. Any disagreements were discussed until consensus was reached. If there was any ambiguity about the potential eligibility of a report based on the abstract/title, we retrieved the full text report for further review. Full text was retrieved for all study reports that were not judged explicitly ineligible in the initial screening. The same team then used the same procedure to screen full text reports and make final eligibility decisions, with the first author again reviewing all eligibility decisions and consensus used to resolve any disagreements.

All data extraction followed a standardized coding protocol, with data entered directly into a FileMaker Pro database. The coding protocol was similar to those used in our previous meta-analyses (e.g., Wilson, Lipsey, Tanner-Smith, Huang, & Steinka-Fry, 2010), and

provided detailed instructions for extracting data related to general study characteristics, participant groups, the interventions, outcome measures, and statistical data needed for effect size calculations (coding protocol available from the authors upon request). Coding information from the eligible study reports was conducted by the same research team after several weeks of training led by the first author. During training, five studies were coded by all the coders, who then convened to resolve coding discrepancies until 100% consensus was attained on all coded variables. After the training period, coding questions were addressed in weekly meetings and decided via consensus with the group. In addition, the first author reviewed all coding and resolved any further discrepancies via consensus with the initial coder.

2.4 Statistical Methods

2.4.1 Effect size metric—The intervention effects of interest were represented with standardized mean difference effect sizes (Cohen's d), calculated as the difference between the intervention and control group means on an outcome variable after the end of the intervention divided by the pooled standard deviation. These effect sizes were adjusted with the small-sample correction factor to provide unbiased estimates (g) (Hedges, 1981). All effect sizes were coded so that positive values indicate better outcomes (e.g., lower alcohol consumption, higher abstinence). For binary outcomes (e.g., abstinence), the Cox transformation outlined by Sánchez-Meca and colleagues (2003) was used to convert log odds ratios into standardized mean difference effect sizes. Effect size and sample size outliers were Winsorized to less extreme values to prevent them from having disproportionate influence on the meta-analysis (Lipsey & Wilson, 2001).¹ When studies reported pretest-adjusted posttest means for both intervention groups (e.g., using ANCOVA or regression methods for adjustment), we used those pretest-adjusted means in the posttest effect size calculations. Otherwise, most studies provided enough information to permit estimation of a pretest effect size for baseline differences between groups, and we then used our own covariate adjustment method to control for pretest differences between groups (described in detail in section 2.4.4).

The standard errors of the effect size estimates used in the weighting function for the meta-analysis were adjusted for the nesting of participants within clusters (e.g., schools) for studies ($k = 39$) using designs in which clusters were assigned to conditions. In these cases, the standard error of the effect size was multiplied by the square root of the design effect, $SE_{adj} = SE_g * \sqrt{1 + (M - 1) * ICC}$, where SE_{adj} is the standard error adjusted for cluster assignment, SE_g is the standard error ignoring clustering, M is the average cluster size, and ICC is the intraclass correlation coefficient for the proportion of between cluster variance (Higgins, Deeks, & Altman, 2008). When cluster-assigned trials did not report the ICC , we used an estimated ICC value of .13, which was the average ICC for all alcohol consumption outcomes across the studies reporting it.²

¹Outliers were defined as values falling three times the interquartile range beyond the upper/lower fences of the distributions, and were Winsorized to the upper/lower fence values.

²Whether a study used a cluster assignment design was uncorrelated with effect size magnitude ($r = -.002$), and only modestly correlated with three moderators of interest (see Appendix C). Sensitivity analyses for all moderator analyses (not shown) that additionally controlled for cluster assignment design yielded results substantively similar to those reported here.

2.4.2 Moderator variables—The following participant characteristics were explored as potential effect size moderators in the analysis: percent of White youth in the intervention group; average participant age; percent of males in the intervention group; and whether participants were recruited based on their identification as high-risk drinkers on baseline screening assessments (1 = *high-risk screened/selective sample*; 0 = *universal sample*). The following intervention characteristics were explored as potential effect size moderators: interval between the end of the intervention and posttest measurement; focal modality (21st birthday card, cognitive behavioral/skills training, expectancy challenge, motivational enhancement, psycho-education, personalized feedback/information, cognitive behavioral + motivational enhancement, family focused therapy, or other); delivery site (school/university, emergency room, primary care/university health center, or self-administered); format (self-administered and computerized, self-administered but not computerized, individual, group, or family); treatment duration (single session less than 5 minutes, single session of 5-15 minutes, single session longer than 15 minutes, or multi-session); and the presence/absence of the following intervention components (BAC information, alcohol caloric information, decisional balance exercise, general education/information, personalized feedback, gender-specific feedback, goal-setting, providing money/cost information about drinking, or local/national drinking norm referencing).

Methodological and procedural characteristics of studies are often confounded with their substantive features. Therefore, the following methodological characteristics were coded and used in analysis as statistical controls to help disentangle those relationships: study design (randomized vs. controlled quasi-experimental); attrition between pretest and posttest; whether binary data were used to estimate the effect size (1 = *yes*; 0 = *no*)³; whether the posttest effect size was estimated using author-reported pretest-adjusted posttest means (1 = *yes*; 0 = *no*); type of control group (no treatment versus treatment as usual); and pretest differences between intervention and control groups indexed using standardized mean difference effect sizes.

2.4.3 Missing data—There was a modest amount of missing data for five of the moderators and covariates of interest (11% of studies were missing attrition data; 37% pretest effect sizes; 15% average age; 3% gender composition; 24% race composition). For these five covariates, we imputed missing data using an expectation-maximization algorithm (Allison, 2002) so that all cases could be included within any given analysis. Sensitivity analyses (not reported) using listwise deletion to handle the missing moderator and covariate values yielded substantively similar results, so we elected to present results using the imputed values. When primary studies failed to include sufficient statistical information to estimate effect sizes, we contacted the study authors for that information. Overall, we had an excellent response from authors, most of whom provided the needed information. Seventeen studies met all eligibility criteria but did not provide sufficient information to estimate effect sizes, and we were unsuccessful in obtaining that information from study authors. We did not impute missing effect sizes on any outcome variables but, rather, omitted them from any analysis involving those outcomes.

³See Sánchez-Meca et al. (2003) for a discussion on transforming effect sizes into different metrics.

2.4.4 Analytic strategies—Most studies reported multiple measures of alcohol consumption (e.g., frequency of consumption, quantity consumed, blood alcohol concentration), and/or multiple measures of alcohol-related problems (e.g., scores from the Rutgers Alcohol Problem Index, relationship problems, risky sexual behavior, DUI/DWI convictions).⁴ We therefore used robust variance estimation in all analyses (Hedges, Tipton, & Johnson, 2010), which permits inclusion of statistically dependent effect sizes (e.g., two different measures of alcohol consumption from the same participant sample), and ensures that studies contributing multiple effect sizes are not given proportionately more weight to the analysis than studies contributing fewer effect sizes.⁵ All analyses used inverse variance weighting so that the contribution of each effect size was proportionate to its statistical precision (Hedges & Olkin, 1985; Lipsey & Wilson, 2001). As suggested by Hedges and colleagues (2010), we used a conservative approach in calculating the weights by applying a weight for each effect size based on the sample size, then dividing that weight by the number of effect sizes in that study for a given outcome type. All analyses used a random-effects model given the presumed heterogeneity in effect sizes, and our desire to generalize findings beyond our analytic sample.

Meta-regression models were used to investigate the overall effects of the interventions, the influence of the intervention and participant characteristics as moderators, and the persistence of effects over time. These models were also used to generate predictions of the mean effect sizes for studies with different intervention characteristics. Because effect sizes can be influenced by the methodological characteristics of the studies, all analyses used effect sizes that were covariate-adjusted to estimate effects at the following values: randomized study design, binary data not used to estimate effect size, effect size adjusted for pretest differences, no-treatment control condition, no pretest differences, and average level of attrition. This was done by predicting the effect sizes from only these variables, retaining the residuals, and adding each residual to the constant predicted from the selected values on the covariates (see Appendix B for estimation model). This conservative technique ensures that any variance in effect sizes associated with differences in method between studies is removed from the analysis of the influence of substantive variables on those effect sizes.

We used contour enhanced funnel plots (Peters, Sutton, Jones, Abrams, & Rushton, 2008) to explore the possibility of bias resulting from omission of small sample size studies with null or negative findings due to selective publication, reporting, or other forms of dissemination biases. None of the funnel plots (not shown, but available from authors upon request) indicated asymmetry, thus providing no clear evidence of small study bias. However, many of the included studies had similar sample sizes and, overall, there was a notable lack of small sample studies regardless of whether they reported negative, null, or positive findings.

⁴Although some outcomes measures may be more commonly used with certain populations (e.g., blood alcohol concentration measures used in selective or indicated populations of youth), based on our pre-specified eligibility criteria for the systematic review we conceptualized all of these measures as representing the same underlying constructs of alcohol consumption or alcohol related problems, respectively.

⁵The robust standard error technique requires an estimate of the mean correlation (ρ) between all pairs of effect sizes within a study for calculation of the between-study sampling variance estimate, τ^2 . In all analyses, we estimated τ^2 with $\rho = .80$; sensitivity analyses (not reported) showed that the findings were robust across different plausible estimates of ρ .

All results are presented separately by the age of participant samples (adolescents ages 11-18 versus young adults ages 19-30)⁶. Although some results did not differ across age groups, we chose to present them separately given the different legal, social, and developmental norms surrounding alcohol in these groups, and researcher and practitioner interest in age-specific effects. Similarly, we present all results separately for alcohol consumption and alcohol-related problem outcomes, given their conceptual differences.

3. Results

3.1 Literature Search

We identified 7,593 candidate reports in the literature search; 2,467 were duplicates that were dropped from consideration and 2,641 were screened as ineligible at the abstract level (see Figure 1). We were unable to locate one unpublished manuscript (Leffingwell et al., 2007). Of the 2,484 articles retrieved in full text, 2,171 were deemed ineligible. Seventeen studies (in 23 reports) met all eligibility criteria but did not provide sufficient information to estimate effect sizes, and we were unsuccessful in obtaining that information from study authors.⁷ The final meta-analysis reported here was based on 185 studies consisting of independent samples that were reported in 313 documents (see Appendix A for a list of references).

3.2 Study Characteristics

Table 1 presents descriptive statistics for the studies included in the meta-analysis, shown separately for the 24 adolescent and 161 young adult samples. One-half of the adolescent studies were conducted in the United States versus 81% of the young adult studies. Most studies were randomized controlled trials, published in journal articles, had low to moderate attrition rates, and reported effects 20-24 weeks after the end of the intervention. The most common intervention modalities for adolescents were motivational enhancement/motivational interviewing (MET; 42%) and psycho-educational therapy (PET; 36%). For young adults, MET programs were common (35%) as were those focused on providing personalized feedback about drinking (28%). Most interventions were conducted in high school or university settings. The brief alcohol interventions were notably longer for adolescents than for young adults (100 versus 55 total minutes, 6 versus 3 days covered).

Table 1 also shows the weighted bivariate correlation between each study characteristic and the unadjusted alcohol consumption effect sizes for the adolescent and young adult samples respectively. Most study characteristics had small to modest bivariate correlations with the effect sizes, but several of the method-related characteristics exhibited correlations with effect size magnitude that were large enough to potentially influence results. Given this, all subsequent analyses used these method-related variables (study design, attrition, effect size estimation method, control group type, and pretest differences between groups) to adjust

⁶We split group at age 18/19 given that the majority of studies recruited participants from either high school or postsecondary educational settings, and given a specific interest in examining efficacy for adolescents.

⁷There was evidence in some of these studies that either the intervention or control group fared better at posttest but effect sizes were still not estimable (e.g., means were presented without standard deviations). We conducted sensitivity analyses (not reported) that made the conservative assumption that all missing effect sizes were equal to zero. The results were substantively similar to those reported here; thus, we are fairly confident that omitting those studies did not bias our results.

effect sizes to the standard methodological profile described earlier (see section 2.4.4 and Appendix B).

3.3 Overall Effects

3.3.1 Effectiveness for adolescents—The random-effects mean of 134 method-adjusted effect sizes estimated from 24 studies indicated that, on average, adolescents age 11-18 who received brief alcohol interventions reported significantly lower levels of alcohol consumption than those in control conditions ($\bar{g} = 0.27$, 95% CI [0.16, 0.38]). This mean effect size is significantly different from zero, and moderately large. For instance, we can translate this standardized effect size back into the metric of a commonly used measure of alcohol consumption: the number of days alcohol was consumed in the past 30 days using the Alcohol Timeline Followback (TLFB). Using the median from the control groups of studies using the TLFB outcome measure, this mean effect size of 0.27 translates into a reduction of 1.3 drinking days per month, with adolescents in the intervention groups consuming alcohol an average of 4.9 days in the past month, versus 6.2 days in the past month for those in control groups.

Only eight studies with adolescent samples also reported an alcohol-related problem outcome (e.g., scores on the Rutgers Alcohol Problem Index). Using 38 effect sizes from those 8 studies, the brief alcohol interventions were associated with significantly lower levels of alcohol-related problems among adolescents ($\bar{g} = 0.19$, 95% CI [0.06, 0.31]). Using Cohen's U3 index (1988), this mean effect size of 0.19 indicates that brief alcohol interventions produced an 8-percentile improvement on alcohol-related problem outcomes for adolescents, relative to control group participants.

3.3.2 Effectiveness for young adults—One hundred fifty-six studies contributed 1,312 method-adjusted effect sizes measuring effects on alcohol consumption among young adults age 19-30. Young adults who received brief alcohol interventions reported significantly lower levels of alcohol consumption than those in control conditions ($\bar{g} = 0.17$, 95% CI [0.13, 0.20]). This effect is modest in practical terms, equivalent to a 0.8 reduction in drinking days per month, from 6.2 to 5.4 days in the past month (using the alcohol TLFB). However, there was no evidence that this mean effect for young adults was significantly different from that observed for adolescents when tested with a meta-regression model that included a dummy variable for adolescent vs. young adult sample ($b = 0.09$, 95% CI [-0.01, 0.20]).

Ninety-six studies with young adult samples contributed 379 method-adjusted effect sizes for alcohol-related problem outcomes. These showed a significant beneficial effect on alcohol-related problems ($\bar{g} = 0.11$, 95% CI [0.08, 0.14]). Using the U3 index, this mean effect size of 0.11 indicates that brief alcohol interventions produced an 4-percentile improvement on alcohol-related problem outcomes for young adults, relative to control group participants. This mean effect size for alcohol-related problems among young adult samples was not significantly different from that observed in the adolescent samples ($b = 0.08$, 95% CI [-0.03, 0.20]).

3.4 Variability in Effects

The between-study variance estimates (τ^2) of the alcohol consumption effect sizes were 0.04 ($Q = 46.92$) for adolescents and 0.02 ($Q = 334.74$) for young adults. For effect sizes on the alcohol-related problem outcomes the τ^2 estimates were 0.00 ($Q = 7.20$) for adolescents and 0.00 ($Q = 108.25$) for young adults. Despite the homogeneity of these effect sizes evidenced by the small τ^2 and Q values, we examined whether any of the moderators of interest were related the effects of the brief alcohol interventions given our original intent to explore when, for whom, and for how long brief alcohol interventions are effective for adolescents and young adults.

3.4.1 Variability across intervention characteristics—As is evident in the descriptive summary presented in Table 1, the brief interventions represented in this sample of studies are quite varied in modality, delivery site, format, length, and intervention components. An important question, therefore, is whether any of these characteristics are associated with differential effects on alcohol consumption or alcohol-related problems. Table 2 shows the corresponding predicted mean effect sizes split by age group, outcome type, and intervention characteristic.⁸

As shown in the top section of Table 2, effects were relatively similar across different intervention modalities for both adolescents and young adults. The notable exceptions to this were 21st birthday card interventions and interventions that combined motivational enhancement and cognitive behavioral therapy components (MET/CBT), neither of which showed evidence of significant beneficial effects in the young adult samples on which they were tested. One limitation of these analyses is the low statistical power for mean effect sizes estimated from such small numbers of studies. Given this, we also estimated mean effect sizes aggregated across the adolescent and young adult samples and outcome types to increase the number of studies and effect sizes contributing to the analysis. Those results were similar, again providing no evidence that either 21st birthday card interventions ($\bar{g} = 0.07$, 95% CI [-0.03, 0.18]) or MET/CBT ($\bar{g} = 0.08$, 95% CI [-0.05, 0.20]) had significant effects.

The mean effects were also quite similar for interventions provided in most of the different delivery sites represented (school/university, primary health care clinics, or self-administered). However, interventions conducted in emergency room settings did not show significant effects on alcohol-related problem outcomes among young adults. Again, because few studies were conducted in emergency room settings (see Table 1), we aggregated results across the age groups. With that larger sample, the mean effect size for alcohol-related problems was still not statistically significant ($\bar{g} = 0.11$, 95% CI [-0.01, 0.22]).

⁸Table 2 only includes results for intervention categories with at least two studies (see Table 1). The adjusted effect sizes shown in Table 2 were estimated from meta-regression models that predicted the method-adjusted effect sizes, but also additionally controlled for participant age, gender, race, and baseline alcohol severity. Statistical tests for differences between intervention characteristic categories were estimated using dummy variables in the meta-regression models for each family of intervention characteristic, in turn (i.e., the models did not control for the other intervention characteristics shown in Table 2). There was no evidence of significant differences in effect sizes across any of the intervention characteristic categories shown in Table 2.

The results shown in Table 2 also indicate that the effects of brief alcohol interventions were similar across different formats (computerized, non-computerized, one-on-one, group). Finally, results in the bottom of Table 2 show that the effects were substantially similar for different amounts of contact time in the adolescent samples, but were more variable across contact time in the young adult samples. Single-session interventions delivered with less than 5 minutes of total contact time (e.g., providing youth with personalized feedback reports to read) showed the largest mean effect on both alcohol consumption ($\bar{g} = 0.30$, 95% CI [0.09, 0.51]) and alcohol-related problem outcomes ($\bar{g} = 0.25$, 95% CI [0.05, 0.46]). However, only a few studies contributed effect sizes to this category, so these results must be interpreted cautiously.

Table 3 shows mean covariate-adjusted effect sizes for interventions with and without specific therapeutic components (BAC information, alcohol caloric information, decisional balance exercise, general education/information, personalized feedback, gender-specific feedback, goal-setting, providing money/cost information about drinking, or local/national drinking norm referencing).⁹ Overall, the effects were relatively similar across different intervention components, but there were some notable exceptions. For adolescents, brief alcohol interventions had significantly larger effects on alcohol consumption if they included decisional balance or goal-setting exercises. For 42 effect sizes from 8 adolescent studies with interventions that included both decisional balance and goal-setting exercises, the estimated mean covariate-adjusted effect size was 0.50 (95% CI [0.18, 0.82], $\tau^2 = 0.10$, $Q = 22.83$), indicating a sizeable effect on alcohol consumption outcomes among adolescents, equivalent to a reduction from 6.2 drinking days in the past month among control participants (using the alcohol TLFB), to 3.9 drinking days among intervention participants. Interventions that included personalized feedback and norm referencing also had significantly larger effects on alcohol-related problems for adolescents, although those results should be interpreted cautiously given the small number of studies contributing effect sizes. Among young adults, however, there was no evidence that any of the intervention components shown were associated with significantly larger or smaller effects on alcohol consumption or alcohol-related problem outcomes, as shown in the right panel of Table 3.

3.4.2 Variability across participants—We estimated meta-regression models to examine whether the following characteristics of the participant samples were associated with the magnitude of effects: race composition, average age, gender composition, or whether samples were selected based on high-risk drinking at baseline screening. Results indicated that alcohol consumption outcomes were significantly better for the adolescent samples with fewer White participants, i.e., a greater proportion of minority participants ($b = -0.66$, 95% CI [-0.96, -0.36]). However, of the 24 studies in this analysis, this effect was driven by 4 studies (contributing 12 effect sizes) comprised entirely of Hispanic or Mi'kmaq Aboriginal Canadian youth who experienced notably larger benefits from the brief interventions ($\bar{g} = 1.03$, 95% CI [0.68, 1.38]). When these four studies were excluded, the

⁹The predicted effect sizes shown in Table 3 were estimated from meta-regression models that predicted the method-adjusted effect sizes, but also additionally controlled for participant age, gender, race, and baseline alcohol consumption. Statistical tests for differences between intervention component categories were estimated using dummy variables in the meta-regression models for each intervention components, in turn (i.e., the models did not control for the other intervention components shown in Table 3 due to high inter-correlations between the various intervention components).

proportion of White youth in the intervention group was no longer significantly related to the effect sizes for the alcohol consumption outcomes.

Within the adolescent and young adult samples, the age of the participants was not associated with the intervention effects for alcohol consumption outcomes. However, among adolescent samples, the effects on alcohol-related problems were significantly smaller in older samples ($b = -0.11$, 95% CI $[-0.17, -0.05]$). There was no evidence that the gender composition of the adolescent samples (percent of males) was associated with the effects on alcohol consumption ($b = -0.06$, 95% CI $[-0.79, 0.67]$) or on alcohol-related problems ($b = 0.66$, 95% CI $[-0.20, 1.53]$); nor for young adults' alcohol consumption ($b = 0.06$, 95% CI $[-1.21, 0.33]$) or alcohol-related problems ($b = -0.03$, 95% CI $[-0.21, 0.16]$). Very few studies were comprised of all male or female samples, but restricting analyses to those studies also showed little differential effectiveness across gender. For instance, the mean method-adjusted effect size for alcohol consumption for young adult female samples was 0.28 (95% CI $[0.14, 0.41]$) versus 0.34 (95% CI $[0.05, 0.63]$) for young adult male samples.

Finally, there was no consistent evidence that brief alcohol interventions were differentially effective for high-risk screened participants. The one exception to this was for alcohol-related problem outcomes among adolescent samples, whereby high-risk screened participants experienced larger beneficial intervention effects than those that were not selected for interventions based on their baseline alcohol consumption ($b = 0.49$, 95% CI $[0.27, 0.72]$). Adolescents already exhibiting heavy or hazardous alcohol consumption patterns may exhibit larger intervention effects given that they have more room for improvement after the intervention (relative to universal prevention programs where most participants may consume alcohol relatively infrequently).

3.4.3 Persistence of effects over time—For the adolescent samples, longer intervals between the brief interventions and follow-up measures of the outcomes were not associated with effects on alcohol consumption ($b = -0.005$, 95% CI $[-0.01, 0.002]$) nor with the effects on alcohol-related problems ($b = 0.0005$, 95% CI $[-0.01, 0.02]$). It is important to note, however, that none of the adolescent studies reported follow-up results longer than one-year after the end of the intervention. Several young adult samples included longer follow-up periods, up to four years post-intervention. For alcohol-related problems, the follow-up period for those young adult samples was not related to the effect sizes ($b = -0.001$, 95% CI $[-0.002, 0.001]$). However, longer follow-up intervals (in weeks) were associated with smaller effects on alcohol consumption among the young adults ($b = -0.003$, 95% CI $[-0.004, -0.001]$). For example, the mean effect size for alcohol consumption among young adults was 1.22 at 1-week follow-up ($k = 15$, $n = 66$, 95% CI $[0.17, 0.27]$) and 0.08 at 24-month follow-up ($k = 3$, $n = 24$, 95% CI $[-0.11, 0.28]$).

Figure 2 shows results from the meta-regression models that examined the persistence of effects over time, split by age group and outcome type. Each effect size is shown proportionate to its weight in the analysis (larger circles represent higher weight, larger sample size studies) and the fitted regression line with its 95% confidence interval is imposed on the graph. A dashed line is shown at the null effect size value of zero (i.e., indicating no difference between the outcomes for the brief intervention and control group).

As shown in the upper right panel of Figure 2, the effects of brief alcohol interventions were attenuated to non-significance at the two-year follow-up point for alcohol consumption among young adults. Although there was no observed attenuation of effects over time for adolescents, only relatively short follow-up intervals were used in the adolescent studies. If only follow-up intervals of less than one year post-intervention are examined for the alcohol consumption effects in the young adult samples (equivalent to the follow-up intervals in the adolescent samples), the results also indicate no significant relationship between the effect sizes and the length of follow-up intervals. The available evidence thus indicates no significant decrease in the effects of the brief interventions on alcohol consumption for up to one year after the intervention.

4. Discussion

This meta-analysis synthesized findings from 185 experimental and quasi-experimental independent study samples that examined the effects of brief alcohol interventions on alcohol-related outcomes for adolescents and young adults who were not seeking treatment. Overall, brief alcohol interventions with up to five hours of total contact time were associated with statistically significant post-intervention reductions in alcohol consumption and alcohol-related problem outcomes. These effects were modest for adolescents—equivalent to 0.27 and 0.19 standard deviation reductions in alcohol use and alcohol-related problems respectively. Although smaller in magnitude, the benefits for young adults were also positive and significant—0.17 and 0.11 standard deviation reductions in alcohol use and alcohol-related problems respectively. Overall, these results indicated that youth receiving brief alcohol interventions reduced their alcohol consumption between 1.0 – 1.3 drinking days per month (relative to control participants, who reported an average of 6.2 drinking days per month at baseline).

These effect estimates are of the same order of magnitude as those reported in previous meta-analyses examining similar interventions for youth (e.g., Jensen et al., 2011; Scott-Sheldon et al., 2009; Tait & Hulse, 2003). The primary strengths of this meta-analysis are the large number of studies included and the application of statistical techniques that permit inclusion of multiple effect sizes from each study. These combined to produce a rich data set that allowed exploration of the variability in effects rather than being restricted to estimating overall mean effects. The results yielded several findings with implications for both research and practice.

One such finding is that, despite their brevity, the effects of brief alcohol interventions persist for up to one year after the end of the interventions. Few studies have followed samples longer than that, but those that do indicate that the effects may dissipate by two years after the interventions (at least among young adults). Furthermore, although the effects were generally similar across different participant demographic profiles and various intervention formats, certain treatment modalities and therapeutic components were associated with larger or smaller effects. In particular, there were no statistically significant effects for the MET/CBT combinations and 21st birthday card interventions evaluated in these research studies. The 21st birthday card interventions typically involved mailing college students a postcard or flyer with a birthday message and safe drinking tips or

information about the dangers of binge drinking. One possible explanation for the lack of effects from this type of intervention could be poor implementation fidelity – many of the studies did not monitor the number of students who actually received the birthday card, nor how many read it. The source of the null findings for the MET/CBT interventions, on the other hand, is especially puzzling in light of the positive effects of MET and CBT by themselves. Perhaps combining these two components in a circumscribed time to provide a brief intervention dilutes the beneficial effects each has alone (see Bell, Marcus, & Goodlad, 2013, for a recent meta-analysis of component treatment studies).

Although the primary goal of this meta-analysis was to identify any features of brief interventions that are associated with differential effects, another important contribution for any meta-analysis is to guide future research. This can be accomplished not only by highlighting gaps in the evidence, but also by providing suggestions for new intervention strategies that build upon the characteristics most strongly associated with positive outcomes that can be identified in the currently available research. We therefore have used the results from this meta-analysis to create a hypothetical profile of characteristics associated with the most and least effective brief alcohol interventions for youth. Based on predicted values from the regression models developed during our analysis, the interventions for adolescents that would potentially yield the largest benefit on alcohol consumption would use MET in a single session of more than 15 minutes delivered on a high school campus. That intervention would include decisional balance, goal-setting, and norm referencing as therapeutic components and would not include BAC information, basic education/information, or personalized feedback. Our regression model predicts that this hypothetical intervention profile would produce a mean effect size for alcohol outcomes of 0.39, equivalent to a reduction from 6.2 to 4.4 drinking days in the past month (using the TLFB for alcohol consumption). This is in contrast to the hypothetical “worst” intervention profile for adolescents, which the regression model indicates would be MET/CBT conducted in multiple sessions, delivered individually in an emergency room, and with therapeutic components that include BAC information, basic education, and personalized feedback, and which do not include decisional balance, goal-setting or norm referencing. This worst-case hypothetical profile yields a predicted effect size for alcohol consumption among adolescents of 0.10, equivalent to a reduction from 6.2 to 5.7 drinking days in the past month.

Using the same regression modeling technique, we projected the hypothetical intervention profiles for young adults predicted to produce the largest and smallest effects based on the existing research. Those results suggest that the strongest program for young adults would be a self-administered computerized expectancy challenge intervention conducted on a university campus. It would include BAC information, decisional balance, goal-setting, and money/cost information as therapeutic components and would not include basic education/information or norm referencing. This profile predicts that the intervention would produce a mean effect size of 0.59, equivalent to a reduction from 6.2 to 3.5 drinking days in the past month. In contrast, the worst-case scenario profile for young adults is a self-administered, multi-session MET/CBT intervention delivered in an emergency room. That profile predicts to a negative, but non-significant mean effect size of -0.14 , equivalent to an increase from 6.2 to 6.8 drinking days in the past month. Of course, it is critical to note that these best and

worst case profiles are simulated and do not necessarily represent any actual interventions represented in the available research. Nor can we assume that the effects of brief interventions with these profiles would actually produce the predicted effects. Nonetheless, this exercise does suggest some possible brief intervention configurations that may be worth investigating in future research.

One limitation of the current meta-analysis was its necessary reliance on the information that is reported by the authors of the primary research studies that were included. The limited reporting of the details of the respective interventions inherently limits the range and depth of characteristics available for analyzing the variability in brief alcohol intervention effects. For instance, the personnel used to deliver brief alcohol interventions vary widely in terms of their background training, education level, and therapeutic expertise; e.g., undergraduate peers trained by research staff (Mayhew et al., 2010), trained professional clinicians (Borsari et al., 2012), and computerized self-administered interventions (Butler & Correia, 2009). Unfortunately, such characteristics of the personnel delivering the interventions are so inconsistently reported that we were unable to examine their relationship to the effectiveness of the brief alcohol interventions. As a practical matter, however, knowing about any influence of the characteristics of the delivery personnel on the intervention effects would be important for anyone implementing a brief alcohol intervention.

Another limitation, stemming in this case from the nature of the research that has actually been conducted rather than insufficient reporting, was our inability to examine the long-term persistence of effects, given that few primary studies reported follow-ups beyond 12-months (particularly in the adolescent samples). Although expecting large, persistent effects may be overly ambitious for brief interventions that are often delivered in a single-session, understanding the persistence of these effects is nonetheless important. Even modest reductions in alcohol consumption may have lasting benefits for youth during these critical developmental periods where progression to alcohol use disorders begins.

Despite these limitations, findings from the current meta-analysis provide compelling evidence that brief alcohol interventions can yield beneficial effects on alcohol consumption and alcohol-related problems in non-treatment seeking populations of adolescents and young adults. Although the magnitude of the effects is generally modest, the brevity and low cost of these interventions allow them to be applied on a relatively large scale where they may add incrementally to the influences that deter risky drinking among youth. To optimize the beneficial effects, providers may want to be guided by the factors this study found to be associated with the most positive effects on alcohol consumption. In particular, these research results indicate that the site, modality, and delivery characteristics of brief alcohol interventions may influence their effectiveness with adolescent and young adult populations.

Acknowledgements

This work was supported by Award Number R01AA020286 from the National Institute on Alcohol Abuse and Alcoholism. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Alcohol Abuse and Alcoholism or the National Institutes of Health. The authors wish to acknowledge Emily Fisher, Samyuktha Kashinath, Webster Kasongo, Lauren Kissinger, Yike Li, Jennifer Samson, Amber Solivan, and Katarzyna Steinka-Fry for their assistance with data collection.

Appendix A

References to Study Reports Providing Effect Sizes Included in the Meta-analysis

References

- Agostinelli G, Brown JM, Miller WR. Effects of normative feedback on consumption among heavy drinking college students. *Journal of Drug Education*. 1995; 25:31–40. doi:10.2190/XD56-D6WR-7195-EAL3. [PubMed: 7776148]
- Allison DL. Comparable effects of the use of a computer-assisted, facilitative teaching approach to a traditional lecture teaching approach on college students' learning about and subsequent use of alcohol. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 2002; 64(10):3604. (UMI No. 3110324).
- Baer JS, Kivlahan DR, Blume AW, McKnight P, Marlatt GA. Brief intervention for heavy-drinking college students: 4-year follow-up and natural history. *American Journal of Public Health*. 2001; 91:1310–1316. doi:10.2105/ajph.91.8.1310. [PubMed: 11499124]
- Bailey KA, Baker AL, Webster RA, Lewin TJ. Pilot randomized controlled trial of a brief alcohol intervention group for adolescents. *Drug and Alcohol Review*. 2004; 23:157–166. doi: 10.1080/09595230410001704136. [PubMed: 15370021]
- Barnett, NP.; Monti, PM.; Wood, MD. Motivational interviewing for alcohol-involved adolescents in the emergency room. In: Wagner, EF.; Waldron, H., editors. *Innovations in adolescent substance abuse interventions*. Bergman Press; New York, NY: 2001. p. 143-168. doi:10.1016/B978-008043577-0/50027-4
- Bendtsen P, McCambridge J, Bendtsen M, Karlsson N, Nilsen P. Effectiveness of a proactive mail-based alcohol internet intervention for university students: Dismantling the assessment and feedback components in a randomized controlled trial. *Journal of Medical Internet Research*. 2012; 14:e142. doi:10.2196/jmir.2062. [PubMed: 23113955]
- Bernstein J, Heeren T, Edward E, Dorfman D, Bliss C, Winter M, Bernstein E. A brief motivational interview in a pediatric emergency department, plus 10-day telephone follow-up, increases attempts to quit drinking among youth and young adults who screen positive for problematic drinking. *Academic Emergency Medicine*. 2010; 17:890–902. doi:10.1111/j.1553-2712.2010.00818.x. [PubMed: 20670329]
- Bersamin M, Paschall MJ, Fearnow-Kenney M, Wyrick D. Effectiveness of a web-based alcohol misuse and harm-prevention course among high- and low-risk students. *Journal of American College Health*. 2007; 55:247–254. doi:10.3200/JACH.55.4.247-254. [PubMed: 17319331]
- Bewick BM, Trusler K, Mulhern B, Barkham M, Hill AJ. The feasibility and effectiveness of a web-based personalised feedback and social norms alcohol intervention in UK university students: A randomised control trial. *Addictive Behaviors*. 2008; 33:1192–1198. doi:10.1016/j.addbeh.2008.05.002. [PubMed: 18554819]
- Bewick BM, West R, Gill J, O'May F, Mulhern B, Barkham M, Hill AJ. Providing web-based feedback and social norms information to reduce student alcohol intake: A multisite investigation. *Journal of Medical Internet Research*. 2010; 12:e59. doi:10.2196/jmir.1461. [PubMed: 21169171]
- Bingham CR, Barretto AI, Walton MA, Bryant CM, Shope JT, Raghunathan TE. Efficacy of a web-based, tailored, alcohol prevention/intervention program for college students: Initial findings. *Journal of American College Health*. 2010; 58:349–356. doi:10.1080/07448480903501178. [PubMed: 20159758]
- Bingham CR, Barretto AI, Walton MA, Bryant CM, Shope JT, Raghunathan TE. Efficacy of a web-based, tailored, alcohol prevention/intervention program for college students: 3-month follow-up. *Journal of Drug Education*. 2011; 41:405–430. doi:10.1080/07448480903501178. [PubMed: 22455103]
- Blair KE. Evaluation of the effectiveness of a physician intervention program to reduce alcohol consumption among college students. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 1999; 61(2-A):514. (UMI No. 9960550).

- Boekeloo BO, Jerry J, Lee-Ougo WI, Worrell KD, Hamburger EK, Russek-Cohen E, Snyder MH. Randomized trial of brief office-based interventions to reduce adolescent alcohol use. *Archive of Pediatrics & Adolescent Medicine*. 2004; 158:635–642. doi:10.1001/archpedi.158.7.635.
- Borsari BE. Two brief alcohol interventions for referred college students. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2003; 64(2-B):956. (UMI No. 3081625).
- Borsari B, Carey KB. Effects of a brief motivational intervention with college student drinkers. *Journal of Consulting and Clinical Psychology*. 2000; 68:728–733. doi:10.1037/0022-006x.68.4.728. [PubMed: 10965648]
- Borsari B, Hustad JT, Mastroleone NR, Tevyaw TO, Barnett NP, Kahler CW, Monti PM. Addressing alcohol use and problems in mandated college students: A randomized clinical trial using stepped care. *Journal of Consulting and Clinical Psychology*. 2012; 80:1062–1074. doi:10.1037/a0029902. [PubMed: 22924334]
- Borsari B, Tevyaw TO, Barnett NP, Kahler CW, Monti PM. Stepped care for mandated college students: A pilot study. *American Journal on Addictions*. 2007; 16:131–137. doi: 10.1080/10550490601184498. [PubMed: 17453615]
- Braitman AL. The effects of personalized boosters for a computerized intervention targeting college student drinking. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2012; 73 09, E. (UMI No. 3510998).
- Butler LH, Correia CJ. Brief alcohol intervention with college student drinkers: Face-to-face versus computerized feedback. *Psychology of Addictive Behaviors*. 2009; 23:163–167. doi:10.1037/a0014892. [PubMed: 19290702]
- Carey KB, Carey MP, Henson JM, Maisto SA, DeMartini KS. Brief alcohol interventions for mandated college students: Comparison of face-to-face counseling and computer-delivered interventions. *Addiction*. 2011; 106:528–537. doi:10.1111/j.1360-0443.2010.03193.x. [PubMed: 21059184]
- Carey KB, Carey MP, Maisto SA, Henson JM. Brief motivational interventions for heavy college drinkers: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2006; 74:943–954. doi:10.1037/0022-006x.74.5.943. [PubMed: 17032098]
- Caudill BD, Luckey B, Crosse SB, Blane HT, Ginexi EM, Campbell B. Alcohol risk-reduction skills training in a national fraternity: A randomized intervention trial with longitudinal intent-to-treat analysis. *Journal of Studies on Alcohol and Drugs*. 2007; 68:399–409. [PubMed: 17446980]
- Ceperich SD, Ingersoll KS. Motivational interviewing + feedback intervention to reduce alcohol-exposed pregnancy risk among college binge drinkers: Determinants and patterns of response. *Journal of Behavioral Medicine*. 2011; 34:381–395. doi:10.1007/s10865-010-9308-2. [PubMed: 21318412]
- Chiauzzi E, Green TC, Lord S, Thum C, Goldstein M. My Student Body: A high-risk drinking prevention web site for college students. *Journal of American College Health*. 2005; 53:263–274. doi:10.3200/jach.53.6.263-274. [PubMed: 15900990]
- Clinton-Sherrod M, Morgan-Lopez AA, Brown JM, McMillen BA, Cowell A. Incapacitated sexual violence involving alcohol among college women: The impact of a brief drinking intervention. *Violence Against Women*. 2011; 17:135–154. doi:10.1177/1077801210394272. [PubMed: 21199812]
- Collins, DL. A study of a brief alcohol education program. Francis Marion College; Florence, South Carolina: 1990. (Unpublished master's thesis)
- Collins SE. Weighing the pros and cons: Evaluating decisional balance as a brief motivational intervention for at-risk college drinkers. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2003; 64(11):5775. (UMI No. 3113234).
- Collins SE, Carey KB, Sliwinski MJ. Mailed personalized normative feedback as a brief intervention for at-risk college drinkers. *Journal of Studies on Alcohol*. 2002; 63:559–567. [PubMed: 12380852]
- Comeau MN. Framing solutions: Adolescent girls and their relationships with alcohol. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2003; 64(09):4601. (UMI No. NQ83702).

- Cox JM. A social norms approach to college alcohol use: Drinking in a low-use environment. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2010; 72(02) (UMI No. 3433393).
- Cronin C. Harm reduction for alcohol-use-related problems among college students. *Substance Use & Misuse*. 1996; 31:2029–2037. doi:10.3109/10826089609066450. [PubMed: 8969022]
- Croom K, Lewis D, Marchell T, Lesser ML, Reyna VF, Kubicki-Bedford L, Feffer M, Staiano-Coico L. Impact of an online alcohol education course on behavior and harm for incoming first-year college students: Short-term evaluation of a randomized trial. *Journal of American College Health*. 2009; 57:445–454. doi:10.3200/jach.57.4.445-454. [PubMed: 19114384]
- Cruz IY. Reducing alcohol consumption among high school students by challenging alcohol expectancies. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2006; 67(09) (UMI No. 3233876).
- Cunningham RM, Chermack ST, Zimmerman MA, Shope JT, Bingham R, Blow FC, Walton MA. Brief motivational interviewing intervention for peer violence and alcohol use in teens: One-year follow-up. *Pediatrics*. 2012; 129:1083–1090. doi:10.1542/peds.2011-3419. [PubMed: 22614776]
- Daepfen J-B, Bertholet N, Gaume J, Fortini C, Faouzi M, Gmel G. Efficacy of brief motivational intervention in reducing binge drinking in young men: A randomized controlled trial. *Drug and Alcohol Dependence*. 2011; 113:69–75. doi:10.1016/j.drugalcdep.2010.07.009. [PubMed: 20729010]
- Darkes J, Goldman MS. Expectancy challenge and drinking reduction: Process and structure in the alcohol expectancy network. *Experimental and Clinical Psychopharmacology*. 1998; 6:64–76. [PubMed: 9526147]
- Dearing, J. Reducing alcohol-related harm through utilizing a harm prevention curriculum at the University of Central Oklahoma. 2008. (Master's thesis). Available from ProQuest Dissertations & Theses Full Text. (UMI No. 1450606)
- Dempster M, Newell G, Cowan G, Marley J. Facing up to binge drinking: Reducing binge drinking in adolescent males. *British Dental Journal*. 2006; 201:587–590. doi:10.1038/sj.bdj.4814204. [PubMed: 17099670]
- DiFulvio GT, Linowski SA, Mazziotti JS, Puleo E. Effectiveness of the Brief Alcohol and Screening Intervention for College Students (BASICS) program with a mandated population. *Journal of American College Health*. 2012; 60:269–280. doi:10.1080/07448481.2011.599352. [PubMed: 22559086]
- Dimeff LA. Brief intervention for heavy and hazardous college drinkers in a student primary health care setting. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 1997; 58(12-B):6805. (UMI No. 9819231).
- D'Onofrio G, Fiellin DA, Pantalon MV, Chawarski MC, Owens PH, Degutis LC, O'Connor PG. A brief intervention reduces hazardous and harmful drinking in emergency department patients. *Annals of Emergency Medicine*. 2012; 60:181–192. doi:10.1016/j.annemergmed.2012.02.006. [PubMed: 22459448]
- Doumas DM, Andersen LL. Reducing alcohol use in first-year university students: Evaluation of a web-based personalized feedback program. *Journal of College Counseling*. 2009; 12:18–32. doi:10.1002/j.2161-1882.2009.tb00037.x.
- Doumas DM, Hannah E. Preventing high-risk drinking in youth in the workplace: A web-based normative feedback program. *Journal of Substance Abuse Treatment*. 2008; 34:263–271. doi:10.1016/j.jsat.2007.04.006. [PubMed: 17600650]
- Doumas DM, Kane CM, Navarro B, Roman J. Decreasing heavy drinking in first-year students: Evaluation of a web-based personalized feedback program administered during orientation. *Journal of College Counseling*. 2011; 14:5–20. doi:10.1002/j.2161-1882.2011.tb00060.x.
- Eggleston AM. Components analysis of a brief intervention for college drinkers. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2007; 68(08) (UMI No. 3276684).
- Ekman DS, Andersson A, Nilsen P, Ståhlbrandt H, Johansson AL, Bendtsen P. Electronic screening and brief intervention for risky drinking in Swedish university students—a randomized controlled trial. *Addictive Behaviors*. 2011; 36:654–659. doi:10.1016/j.addbeh.2011.01.015. [PubMed: 21316157]

- Ewing SWF, LaChance HA, Bryan A, Hutchison KE. Do genetic and individual risk factors moderate the efficacy of motivational enhancement therapy? Drinking outcomes with an emerging adult sample. *Addiction Biology*. 2009; 14:356–365. doi:10.1111/j.1369-1600.2009.00149.x. [PubMed: 19298319]
- Fischer Potts KA. Efficacy of the alcohol skills training program, a brief group intervention for high-risk college student drinkers. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2002; 63(06):3006. (UMI No. 3055270).
- Fleming MF, Balousek SL, Grossberg PM, Mundt MP, Brown D, Wiegel JR, Zakletskaia LI, Saewyc EM. Brief physician advice for heavy drinking college students: A randomized controlled trial in college health clinics. *Journal of Studies on Alcohol and Drugs*. 2010; 71:23–31. [PubMed: 20105410]
- Flórez Alarcón L, Gantiva Díaz CA. Terapias motivacionales breves: Diferencias entre la aplicación individual y grupal para la moderación del consumo de alcohol y de las variables psicológicas asociadas al cambio. *Acta Colombiana de Psicología*. 2009; 12:13–26. [Brief motivational therapy: Differences between individual and group application for the moderation in alcohol consumption and other psychological variables associated with change]. Retrieved from <http://regweb.ucatolica.edu.co/publicaciones/psicologia/ACTA/>.
- Foxcroft DR, Kypri K, Simonite V. Bayes' Theorem to estimate population prevalence from Alcohol Use Disorders Identification Test (AUDIT) scores. *Addiction*. 2009; 104:1132–1137. doi: 10.1111/j.1360-0443.2009.02574.x. [PubMed: 19438421]
- Fried, A. Evaluation of an expectancy challenge presentation in reducing high-risk alcohol use among Greek affiliated college students. 2010. (Unpublished master's thesis). University of Central Florida, Orlando, Florida. Retrieved from http://etd.fcla.edu/CF/CFE0003263/Fried_Abigail_B_201008_MS.pdf
- Fromme K, Corbin W. Prevention of heavy drinking and associated negative consequences among mandated and voluntary college students. *Journal of Consulting and Clinical Psychology*. 2004; 72:1038–1049. doi:10.1037/0022-006X.72.6.1038. [PubMed: 15612850]
- Gallo LA. Investigating written emotional disclosure as an intervention for college student drinking. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2007; 68(07) (UMI No. 3274107).
- Gaume J, Daeppen JB, Gmel G. Administering brief motivational interventions to young men: Results from a pilot census study among 19 year-old Francophone Swiss men. Lausanne, Switzerland: Centre hospitalier universitaire vaudois - CHUV, Département universitaire de médecine et santé communautaires. 2008
- Gaume J, Gmel G, Faouzi M, Bertholet N, Daeppen J-B. Is brief motivational intervention effective in reducing alcohol use among young men voluntarily receiving it? A randomized controlled trial. *Alcoholism: Clinical and Experimental Research*. 2011; 35:1822–1830. doi:10.1111/j.1530-0277.2011.01526.x.
- Glassman, T. 21st birthday social marketing intervention study in University of Florida. University of Florida; Gainesville, FL: (n.d.)Unpublished manuscript
- Glassman T, Dodd V, Kenzik K, Miller EM, Sheu J-J. Social norms vs. risk reduction approaches to 21st birthday celebrations. *American Journal of Health Education*. 2010; 41:38–45. doi: 10.1080/19325037.2010.10599125.
- Gmel G, Venzin V, Marmet K, Danko G, Labhart F. A quasi randomized group trial of a brief alcohol intervention on risky single occasion drinking among secondary school students. *International Journal of Public Health*. 2012; 57:935–944. doi:10.1007/s00038-012-0419-0. [PubMed: 23089675]
- Gonzalez GM. The effect of a model alcohol education module on college students' attitudes, knowledge, and behavior related to alcohol use. *Journal of Alcohol and Drug Education*. 1980; 25:1–12.
- Green PJ. Social influence and personality as moderators of the effects of psychoeducational interventions on alcohol use. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 1998; 59(05):2484. (UMI No. 9834461).
- Gregory BM. College alcohol and life skills study with student-athletes. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 2001; 62(1-A):89. (UMI No. 3001230).

- Hagger MS, Lonsdale A, Koka A, Hein V, Pasi H, Lintunen T, Chatzisarantis NL. An intervention to reduce alcohol consumption in undergraduate students using implementation intentions and mental simulations: A cross-national study. *International Journal of Behavioral Medicine*. 2012; 19:82–96. doi:10.1007/s12529-011-9163-8. [PubMed: 21562782]
- Hagger MS, Lonsdale A, Chatzisarantis NLD. A theory-based intervention to reduce alcohol drinking in excess of guideline limits among undergraduate students. *British Journal of Health Psychology*. 2012; 17:18–43. doi:10.1111/j.2044-8287.2010.02011.x. [PubMed: 22233103]
- Hendershot CS, Otto JM, Collins SE, Liang T, Wall TL. Evaluation of a brief web-based genetic feedback intervention for reducing alcohol-related health risks associated with ALDH2. *Annals of Behavioral Medicine*. 2010; 40:77–88. doi:10.1007/s12160-010-9207-3. [PubMed: 20652463]
- Henslee A. Providing personalized feedback regarding alcohol use in a group format to college freshmen. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2008; 69(10) (UMI No. 3333129).
- Henslee AM, Irons JG, Day JM, Butler L, Benson TA, Correia CJ. Using national alcohol screening day to deliver personalized feedback: A pilot study. *Journal of Drug Education*. 2006; 36:271–278. doi:10.2190/1u8h-41u2-5k03-0j24. [PubMed: 17533801]
- Journal of Drug Education*. 36:271–278. doi:10.2190/1u8h-41u2-5k03-0j24.
- Hester RK, Delaney HD, Campbell W. The College Drinker's Check-Up: Outcomes of two randomized clinical trials of a computer-delivered intervention. *Psychology of Addictive Behaviors*. 2012; 26:1–12. doi:10.1037/a0024753 *This report provides data for two unique studies. [PubMed: 21823769]
- Hogan, LM. Developing and evaluating brief, computerised interventions for excessive drinkers. 2005. (Doctoral dissertation, University of Wales, Bangor, UK). Available from ProQuest Dissertations & Theses: UK & Ireland. (UMI No. U202714)
- Horner K. Brief motivational interviewing: An intervention for alcohol abusing college students. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2010; 71(05) (UMI No. 3402364).
- Hosier SG. An evaluation of two brief interventions aimed at reducing college students' alcohol use. 2002 (Doctoral dissertation, University of Wales, Bangor, UK). Available from ProQuest Dissertations & Theses: UK & Ireland. (UMI No. U144754).
- Howat P, Robinson S, Binns C, Palmer S, Landauer A. Educational biofeedback driving simulator as a drink-driving prevention strategy. *Journal of Alcohol and Drug Education*. 1991; 37:7–14.
- Hunt WM. Effects of participant engagement on alcohol expectancies and drinking outcomes for a computerized expectancy challenge intervention. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2004; 65(12):6655. (UMI No. 3157213).
- Hustad JTP, Barnett NP, Borsari B, Jackson KM. Web-based alcohol prevention for incoming college students: A randomized controlled trial. *Addictive Behaviors*. 2010; 35:183–189. doi:10.1016/j.addbeh.2009.10.012. [PubMed: 19900763]
- Ingersoll KS, Ceperich SD, Nettleman MD, Karanda K, Brocksen S, Johnson BA. Reducing alcohol-exposed pregnancy risk in college women: Initial outcomes of a clinical trial of a motivational intervention. *Journal of Substance Abuse Treatment*. 2005; 29:173–180. doi:10.1016/j.jsat.2005.06.003. [PubMed: 16183466]
- Ivy K. Prevention of alcohol misuse and related violence for college students. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 1998; 59(08):2874. (UMI No. 9902906).
- Jewell J, Hupp SDA. Examining the effects of Fatal Vision goggles on changing attitudes and behaviors related to drinking and driving. *The Journal of Primary Prevention*. 2005; 26:553–565. doi:10.1007-s10935-005-0013-9. [PubMed: 16228116]
- Johnston BD, Rivara FP, Droesch RM, Dunn CW, Copass MK. Behavior change counseling in the emergency department to reduce injury risk: A randomized, controlled trial. *Pediatrics*. 2002; 110:267–274. doi:10.1542/peds.110.2.267. [PubMed: 12165577]
- Jones LM, Silvia LY, Richman CL. Increased awareness and self-challenge of alcohol expectancies. *Substance Abuse*. 1995; 16:77–85. doi:10.1080/08897079509444707.

- Juarez, P. A randomized trial of motivational interviewing and feedback on heavy drinking college students. 2001. (Unpublished master's thesis). University of New Mexico, Albuquerque, New Mexico
- Juarez P, Walters ST, Daugherty M, Radi C. A randomized trial of motivational interviewing and feedback with heavy drinking college students. *Journal of Drug Education*. 2006; 36:233–246. doi: 10.2190/753n-8242-727t-g63l. [PubMed: 17345916]
- Kazemi DM, Dmochowski J, Sun L, Nies MA, Walford S. Outcomes of a Targeted Capacity Expansion (TCE) brief motivational intervention for high-risk drinking freshmen: Pilot study comparison at baseline and 6 months. *Alcoholism Treatment Quarterly*. 2011; 29:219–229. doi: 10.1080/07347324.2011.586275.
- Keillor RM, Perkins WB, Horan JJ. Effects of videotaped expectancy challenges on alcohol consumption of adjudicated students. *Journal of Cognitive Psychotherapy*. 1999; 13:179–187.
- Kerr-Corrêa, F.; Simão, MO.; Martins, RA. Prevenção ao uso de álcool por estudantes universitários. Unpublished manuscript, Universidade Estadual Paulista, São Paulo, Brazil. Retrieved from. 2003. http://fmb.unesp.br/Home/Departamentos/Neurologia,PsicologiaePsiquiatria/ViverBem/Prevencao_ao_uso_de_alcool_por_estudantes_universitarios.pdfhttp://fmb.unesp.br/Home/Departamentos/Neurologia,PsicologiaePsiquiatria/ViverBem/Prevencao_ao_uso_de_alcool_por_estudantes_universitarios.pdf
- Koning IM, Vollebergh WAM, Smit F, Verdurmen JEE, van den Eijnden RJJM, ter Bogt TFM, Engels RCME. Preventing heavy alcohol use in adolescents (PAS): Cluster randomized trial of a parent and student intervention offered separately and simultaneously. *Addiction*. 2009; 104:1669–1678. doi:10.1111/j.1360-0443.2009.02677.x. [PubMed: 21265908]
- Kulesza M, Apperson M, Larimer ME, Copeland AL. Brief alcohol intervention for college drinkers: How brief is? *Addictive Behaviors*. 2010; 35:730–733. doi:10.1016/j.addbeh.2010.03.011. [PubMed: 20381972]
- Kypri K, Hallett J, Howat P, McManus A, Maycock B, Bowe S, Horton NJ. Randomized controlled trial of proactive web-based alcohol screening and brief intervention for university students. *Archives of Internal Medicine*. 2009; 169:1508–1514. doi:10.1001/archinternmed.2009.249. [PubMed: 19752409]
- Kypri K, McCambridge J, Vater T, Bowe SJ, Saunders JB, Cunningham JA, Horton NJ. Web-based alcohol intervention for Maori university students: Double-blind, multi-site randomized controlled trial. *Addiction*. 2013; 108:331–338. doi:10.1111/j.1360-0443.2012.04067.x. [PubMed: 22925046]
- Kypri K, Saunders JB, Williams SM, McGee RO, Langley JD, Cashell-Smith ML, Gallagher SJ. Web-based screening and brief intervention for hazardous drinking: A double-blind randomized controlled trial. *Addiction*. 2004; 99:1410–1417. doi:10.1111/j.1360-0443.2004.00847.x. [PubMed: 15500594]
- LaBrie JW. Weighing the pros and cons: A brief motivational intervention reduces risk associated with drinking and unsafe sex. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2002; 63(12-B):6098. (UMI No. 3074943).
- LaBrie JW, Huchting KK, Lac A, Tawalbeh S, Thompson AD, Larimer ME. Preventing risky drinking in first-year college women: Further validation of a female-specific motivational-enhancement group intervention. *Journal of Studies on Alcohol and Drugs, Suppl*. 2009; 16:77–85.
- LaBrie JW, Huchting K, Tawalbeh S, Pedersen ER, Thompson AD, Shelesky K, Neighbors C. A randomized motivational enhancement prevention group reduces drinking and alcohol consequences in first-year college women. *Psychology of Addictive Behaviors*. 2008; 22:149–155. doi:10.1037/0893-164x.22.1.149. [PubMed: 18298242]
- LaBrie JW, Hummer JF, Neighbors C, Pedersen ER. Live interactive group-specific normative feedback reduces misperceptions and drinking in college students: A randomized cluster trial. *Psychology of Addictive Behaviors*. 2008; 22:141–148. doi:10.1037/0893-164x.22.1.141. [PubMed: 18298241]
- LaBrie JW, Migliuri S, Cail J. A night to remember: A harm-reduction birthday card intervention reduces high-risk drinking during 21st birthday celebrations. *Journal of American College Health*. 2009; 57:659–663. doi:10.3200/JACH.57.6.659-663. [PubMed: 19433404]

- Larimer ME, Turner AP, Anderson BK, Fader JS, Kilmer JR, Palmer RS, Cronce JM. Evaluating a brief alcohol intervention with fraternities. *Journal of Studies on Alcohol*. 2001; 62:370–380. [PubMed: 11414347]
- Lau-Barraco C, Dunn ME. Evaluation of a single-session expectancy challenge intervention to reduce alcohol use among college students. *Psychology of Addictive Behaviors*. 2008; 22:168–175. doi: 10.1037/0893-164x.22.2.168. [PubMed: 18540714]
- Leffingwell, TR.; Leedy, MJ.; Lack, CW. Paper presented at the meeting of the Association for Behavioral and Cognitive Therapies. Washington, DC: Nov. 2005 A multimedia computer-based intervention for college student drinking: Short-term outcomes of a randomized trial.
- Lewis MA, Neighbors C. Optimizing personalized normative feedback: The use of gender-specific referents. *Journal of Studies on Alcohol and Drugs*. 2007; 68:228–237. doi:10.1037/0893-164X.22.2.176. [PubMed: 17286341]
- Lewis MA, Neighbors C, Lee CM, Oster-Aaland L. 21st birthday celebratory drinking: Evaluation of a personalized normative feedback card intervention. *Psychology of Addictive Behaviors*. 2008; 22:176–185. doi:10.1037/0893-164x.22.2.176. [PubMed: 18540715]
- Lewis MA, Neighbors C, Oster-Aaland L, Kirkeby BS, Larimer ME. Indicated prevention for incoming freshmen: Personalized normative feedback and high-risk drinking. *Addictive Behaviors*. 2007; 32:2495–2508. doi:10.1016/j.addbeh.2007.06.019. [PubMed: 17658695]
- Light LS, McCoy TP, Thompson MP, Spittler HD, Sutfin EL, Rhodes SD. Modeling the Rutgers Alcohol Problem Index (RAPI): A comparison of statistical methods. *Addiction Research & Theory*. 2011; 19:510–518. doi:10.3109/16066359.2011.569100.
- Lojewski R, Rotunda RJ, Arruda JE. Personalized normative feedback to reduce drinking among college students: A social norms intervention examining gender-based versus standard feedback. *Journal of Alcohol & Drug Education*. 2010; 54:19–40.
- Lysaught EM. A comparison of an assessment/information based group versus an assessment only group: An investigation of drinking reduction with young adults. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 1998; 60:550-A–551-A. UMI No. 9918230.
- Magill M, Kahler CW, Monti P, Barnett NP. Do research assessments make college students more reactive to alcohol events? *Psychology of Addictive Behaviors*. 2012; 26:338–344. doi:10.1037/a0025571. [PubMed: 21928871]
- Maio RF, Shope JT, Blow FC, Gregor MA, Zakrajsek JS, Weber JE, Nypaver MM. A randomized controlled trial of an emergency department-based interactive computer program to prevent alcohol misuse among injured adolescents. *Annals of Emergency Medicine*. 2005; 45:420–429. doi:10.1016/j.annemergmed.2004.10.013. [PubMed: 15795723]
- Marlatt GA, Baer JS, Kivlahan DR, Dimeff LA, Larimer ME, Quigley LA, Williams E. Screening and brief intervention for high-risk college student drinkers: Results from a 2-year follow-up assessment. *Journal of Consulting and Clinical Psychology*. 1998; 66:604–615. doi: 10.1037/0022-006x.66.4.604. [PubMed: 9735576]
- Martell, D.; Atkin, C. B.R.A.D. Card Survey, 2001-2002: Preliminary analysis of survey results. Michigan State University, Office for Survey Research, Institute for Public Policy and Social Research; East Lansing, MI: 2002.
- Martínez Martínez, KI.; Pedroza Cabrera, FJ.; Salazar Garza, ML., editors. Resultados iniciales de la aplicación de consejo breve sobre el consumo de alcohol en adolescentes rurales de México [Initial results of the application of brief advice on alcohol consumption in rural teenagers from Mexico]. TIPICA: Boletín Electrónico de Salud Escolar. 2008. 4. Retrieved from <http://www.tipica.org/>
- Martínez Martínez KI, Pedroza Cabrera FJ, Salazar Garza ML, Vacío Muro M. Evaluación experimental de dos intervenciones breves para la reducción del consumo de alcohol de adolescentes [Experimental evaluation of two brief interventions for the reduction of alcohol consumption in teenagers]. *Revista Mexicana De Análisis De La Conducta*. 2010; 36:35–53. doi: 10.5514/rmac.v36.i3.03.
- Martínez Martínez KI, Pedroza Cabrera FJ, Vacío Muro MA, Jiménez Pérez AL, Salazar Garza ML. Consejo breve para adolescentes escolares que abusan del alcohol. [School-based brief

counseling for teenage drinkers]. *Revista Mexicana de Análisis de la Conducta*. 2008; 34:245–262.

- Martins, RA.; Manzato, AJ.; Poiate, SMG.; Scarin, ACCF.; Nogueira da Cruz, LA.; Teixeira, PS.; Kawashima, RA. Alcohol usage and brief intervention in heavy-drinking teenagers; Paper presented at the meeting of the 34th Annual Alcohol Epidemiology Symposium of the Kettil Bruun Society; Victoria, BC, Canada. Jun. 2008
- Mastroleo N. Comparison of supervision training techniques in a motivational enhancement intervention on college student drinking. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 2008; 69(11) UMI No. 3336087.
- Mayhew MJ, Caldwell RJ, Hourigan AJ. The influence of curricular-based interventions within first-year "success" courses on student alcohol expectancies and engagement in high-risk drinking behaviors. *NASPA Journal*. 2008; 45:49–72.
- Mayhew, MJ.; Caldwell, R.; Hourigan, A.; Bezbatchenko, AW.; Fried, M. Intervention deliveries: The role of peer educators in reducing high-risk drinking among first-year students. New York University; New York, NY: 2010. Unpublished manuscript
- Mayhew MJ, Klein S, Behringer LB, Ulrich AS, Caldwell RJ, Hourigan A. Curricular infusion and high-risk drinking among first-year students. *Journal of the First Year Experience & Students in Transition*. 2011; 23:9–34.
- McNally AM. Motivational interventions for problematic alcohol use: In search of the mechanisms of action. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2003; 63(10-B):4914. UMI No. 3067197.
- McNally AM, Palfai TP. Brief group alcohol interventions with college students: Examining motivational components. *Journal of Drug Education*. 2003; 33:159–176. doi:10.2190/82ct-lrc5-antw-c090. [PubMed: 12929707]
- McPherson, P. Efficacy of brief alcohol interventions in an Australian tertiary education setting (Doctoral dissertation, Royal Melbourne Institute of Technology University, Melbourne, Australia). 2012. Retrieved from <http://researchbank.rmit.edu.au/view/rmit:160203>
- Michael KD, Curtin L, Kirkley DE, Jones DL, Harris R Jr. Group-based motivational interviewing for alcohol use among college students: An exploratory study. *Professional Psychology: Research and Practice*. 2006; 37:629–634. doi:10.1037/0735-7028.37.6.629.
- Mignogna J. A test of moderating factors of brief interventions for hazardous alcohol use among college students. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2010; 71(10) UMI No. 3422281.
- Miller ET. Preventing alcohol abuse and alcohol-related negative consequences among freshmen college students: Using emerging computer technology to deliver and evaluate the effectiveness of brief intervention efforts. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2000; 61(8-B):4417. UMI No. 9983525.
- Monti PM, Colby SM, Barnett NP, Spirito A, Rohsenow DJ, Myers M, Woolard R, Lewander W. Brief intervention for harm reduction with alcohol-positive older adolescents in a hospital emergency department. *Journal of Consulting and Clinical Psychology*. 1999; 67:989–994. doi:10.1037/0022-006x.67.6.989. [PubMed: 10596521]
- Moreira MT, Oskrochi R, Foxcroft DR. Personalised normative feedback for preventing alcohol misuse in university students: Solomon three-group randomised controlled trial. *PLOS One*. 2012; 7:e44120. doi:10.1371/journal.pone.0044120. [PubMed: 22984466]
- Mundt M. College alcohol abuse and alcohol-related harms. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2007; 68(12) UMI No. 3294169.
- Murgraff, V.; McDermott, M.; Abraham, C. An evaluation of an action-planning intervention to reduce the incidence of high-risk single-session alcohol consumption in high risk drinkers. Alcohol Education and Research Council; London, UK: 2006.
- Murphy JG, Dennhardt AA, Skidmore JR, Martens MP, McDevitt-Murphy ME. Computerized versus motivational interviewing alcohol interventions: Impact on discrepancy, motivation, and drinking. *Psychology of Addictive Behaviors*. 2010; 24:628–639. doi:10.1037/a0021347. [PubMed: 21198224]

- Murphy JG, Duchnick JJ, Vuchinich RE, Davison JW, Karg RS, Olson AM, Coffey TT. Relative efficacy of a brief motivational intervention for college student drinkers. *Psychology of Addictive Behaviors*. 2001; 15:373–379. doi:10.1037/0893-164x.15.4.373. [PubMed: 11767271]
- Mushquash, CJ. Personality and motives for alcohol use in Aboriginal adolescents: A culturally relevant approach to alcohol abuse early intervention (Unpublished doctoral dissertation, Dalhousie University, Nova Scotia, Canada). 2011. Retrieved from <http://hdl.handle.net/10222/14241>
- Neal DJ, Carey KB. Developing discrepancy within self-regulation theory: Use of personalized normative feedback and personal strivings with heavy-drinking college students. *Addictive Behaviors*. 2004; 29:281–297. doi:10.1016/j.addbeh.2003.08.004. [PubMed: 14732417]
- Neighbors C, Jensen M, Tidwell J, Walter T, Fossos N, Lewis MA. Social norms interventions for light and nondrinking students. *Group Processes & Intergroup Relations*. 2011; 14:651–669. doi: 10.1177/1368430210398014.
- Neighbors C, Larimer ME, Lewis MA. Targeting misperceptions of descriptive drinking norms: Efficacy of a computer-delivered personalized normative feedback intervention. *Journal of Consulting and Clinical Psychology*. 2004; 72:434–447. doi:10.1037/0022-006x.72.3.434. [PubMed: 15279527]
- Neighbors C, Lee CM, Atkins DC, Lewis MA, Kaysen D, Mittmann A, Larimer ME. A randomized controlled trial of event-specific prevention strategies for reducing problematic drinking associated with 21st birthday celebrations. *Journal of Consulting and Clinical Psychology*. 2012; 80:850–862. doi:10.1037/a0029480. [PubMed: 22823855]
- Walter T. Internet-based personalized feedback to reduce 21st-birthday drinking: A randomized controlled trial of an event-specific prevention intervention. *Journal of Consulting and Clinical Psychology*. 2009; 77:51–63. doi:10.1037/a0014386. [PubMed: 19170453]
- Neighbors C, Lewis MA, Atkins DC, Jensen MM, Walter T, Fossos N, Larimer ME. Efficacy of web-based personalized normative feedback: A two-year randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2010; 78:898–911. doi:10.1037/a0020766. [PubMed: 20873892]
- Neighbors C, Lewis MA, Bergstrom RL, Larimer ME. Being controlled by normative influences: Self-determination as a moderator of a normative feedback alcohol intervention. *Health Psychology*. 2006; 25:571–579. doi:10.1037/0278-6133.25.5.571. [PubMed: 17014274]
- Neighbors C, Spieker CJ, Oster-Aaland L, Lewis MA, Bergstrom RL. Celebration intoxication: An evaluation of 21st birthday alcohol consumption. *Journal of American College Health*. 2005; 54:76–80. doi:10.3200/JACH.54.2.76-80. [PubMed: 16255318]
- Ostafin BD. Affective learning and alcohol consumption: Correlates of risk and causes of change. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2004; 65(03):1559. UMI No. 3124866.
- Ostafin BD, Palfai TP. When wanting to change is not enough: Automatic appetitive processes moderate the effects of a brief alcohol intervention in hazardous-drinking college students. *Addiction Science & Clinical Practice*. 2012; 7:7–25. doi:10.1186/1940-0640-7-25. [PubMed: 23186245]
- Palfai TP, Zisserson R, Saitz R. Using personalized feedback to reduce alcohol use among hazardous drinking college students: The moderating effect of alcohol-related negative consequences. *Addictive Behaviors*. 2011; 36:539–542. doi:10.1016/j.addbeh.2011.01.005. [PubMed: 21295919]
- Palmer RS. Efficacy of the alcohol skills training program in mandated and nonmandated heavy drinking college students. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2004; 65(5-B):2644. UMI No. 0806362.
- Paschall MJ, Antin T, Ringwalt CL, Saltz RF. Effects of AlcoholEdu for college on alcohol-related problems among freshmen: A randomized multicampus trial. *Journal of Studies on Alcohol and Drugs*. 2011; 72:642–650. [PubMed: 21683046]
- Paschall MJ, Antin T, Ringwalt CL, Saltz RF. Evaluation of an internet-based alcohol misuse prevention course for college freshmen: Findings of a randomized multi-campus trial. *American Journal of Preventive Medicine*. 2011; 41:300–308. doi:10.1016/j.amepre.2011.03.021. [PubMed: 21855745]

- Paschall MJ, Bersamin M, Fearnow-Kenney M, Wyrick D, Currey D. Short-term evaluation of a web-based college alcohol misuse and harm prevention course (College Alc). *Journal of Alcohol and Drug Education*. 2006; 50:49–65.
- Patel AB. An experimental test of collegiate drinking norms. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2011; 73(01) UMI No. 3479973.
- Pedersen ER. Brief online interventions targeting risk and protective factors for increased and problematic alcohol use among American college students studying abroad. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2012; 74(03, E) UMI No. 3542377.
- Peeler CM, Far JM, Miller JA, Brigham TA. An analysis of the effects of a program to reduce heavy drinking among college students. *Journal of Alcohol and Drug Education*. 2000; 45:39–54.
- Pensuksan WC, Taneepanichskul S, Williams MA. A peer-drinking group motivational intervention among Thai male undergraduate students. *International Journal of Drug Policy*. 2010; 21:432–436. doi:10.1016/j.drugpo.2010.02.005. [PubMed: 20347584]
- Quigley, CF. A psychosocial model of drinking amongst young people and the effects of brief interventions (Doctoral dissertation, University of Sheffield, England). 2010. Available from ProQuest Dissertations and Theses database. (UMI No. U557325) *This report provides data for two unique studies
- Reilly DW, Wood MD. A randomized test of a small-group interactive social norms intervention. *Journal of American College Health*. 2008; 57:53–60. doi:10.3200/jach.57.1.53-60. [PubMed: 18682346]
- Reynolds KD, Coombs DW, Lowe JB, Peterson PL, Gayoso E. Evaluation of a self-help program to reduce alcohol consumption among pregnant women. *International Journal of the Addictions*. 1995; 30:427–443. [PubMed: 7607777]
- Schaus JF, Sole ML, McCoy TP, Mullett N, O'Brien MC. Alcohol screening and brief intervention in a college student health center: A randomized controlled trial. *Journal of Studies on Alcohol and Drugs, Suppl*. 2009; 16:131–141.
- Schreiner, AM. Evaluation of an expectancy challenge curriculum in reducing high risk alcohol use among college students when modified for large classes (Unpublished master's thesis). University of Central Florida; Orlando, Florida: 2010. Retrieved from http://etd.fcla.edu/CF/CFE0003114/Schreiner_Amy_M_201005_MS.pdf
- Segatto, ML. A efetividade da intervenção breve motivacional em jovens atendidos no pronto socorro após um evento relacionado ao consumo de álcool; Paper presented at the meeting of the International Network on Brief Interventions for Alcohol Problems; Ribeirão Preto, Brazil. Oct. 2008
- Sharmer L. Evaluation of alcohol education programs on attitude, knowledge, and self-reported behavior of college students. *Evaluation & the Health Professions*. 2001; 24:336–357. doi: 10.1177/01632780122034957. [PubMed: 11523322]
- Simão, MO. Avaliação da eficácia da intervenção breve para redução de danos em estudantes universitários da Unesp que fazem uso excessivo de bebidas alcoólicas [Assessment of the effectiveness of brief intervention to reduce harm in university students of UNESP who use alcohol excessively] (Doctoral dissertation, Universidade de São Paulo, Riberão Preto, Brazil). 2005. Retrieved from <http://bases.bireme.br/cgi-bin/wxislind.exe/iah/online/?IsisScript=iah/iah.xis&src=google&base=ADOLEC&lang=p&nextAction=lnk&exprSearch=468382&indexSearch=ID>
- Simão MO, Kerr-Corrêa F, Smaira SI, Trinca LA, Floripes TMF, Dalben I, Tucci AM. Prevention of "risky" drinking among students at a Brazilian university. *Alcohol and Alcoholism*. 2008; 43:470–476. doi:10.1093/alcalc/agn019. [PubMed: 18364361]
- Sivasithamparam, J. Implementation and evaluation of a classroom-based approach to expectancy challenge for reducing alcohol use among college students (Unpublished master's thesis). University of Central Florida, Orlando, Florida. 2008. Retrieved from <http://purl.fcla.edu/fcla/etd/CFE0002307>
- Smith BH. A randomized study of a peer-led, small group social norming intervention designed to reduce drinking among college students. *Journal of Alcohol and Drug Education*. 2004; 47:67–75.

- Smith BH, Bogle KE, Talbott L, Gant R, Castillo H. A randomized study of four cards designed to prevent problems during college students' 21st birthday celebrations. *Journal of Studies on Alcohol*. 2006; 67:607–615. [PubMed: 16736081]
- Spijkerman R, Roek MA, Vermulst A, Lemmers L, Huijberts A, Engels RC. Effectiveness of a web-based brief alcohol intervention and added value of normative feedback in reducing underage drinking: A randomized controlled trial. *Journal of Medical Internet Research*. 2010; 12:e65. doi:10.2196/jmir.1465. [PubMed: 21169172]
- Spirito A, Monti PM, Barnett NP, Colby SM, Sindelar H, Rohsenow DJ, Myers M. A randomized clinical trial of a brief motivational intervention for alcohol-positive adolescents treated in an emergency department. *Journal of Pediatrics*. 2004; 145:396–402. doi:10.1016/j.jpeds.2004.04.057. [PubMed: 15343198]
- Stachula JP. A social norming based proactive intervention for college student alcohol use. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2004; 65(6-B):3184. UMI No. 3135853.
- Ståhlbrandt H, Johnsson KO, Berglund M. Two-year outcome of alcohol interventions in Swedish university halls of residence: A cluster randomized trial of a brief skills training program, twelve-step-influenced intervention, and controls. *Alcoholism: Clinical and Experimental Research*. 2007; 31:458–466. doi:10.1111/j.1530-0277.2006.00327.x.
- Stamper GA. The evaluation of program fidelity in two college alcohol interventions. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2006; 67(09) UMI No. 3232540.
- Stamper GA, Smith BH, Gant R, Bogle KE. Replicated findings of an evaluation of a brief intervention designed to prevent high-risk drinking among first-year college students: Implications for social norming theory. *Journal of Alcohol and Drug Education*. 2004; 48:53–72.
- Steffian G. Correction of normative misperceptions: An alcohol abuse prevention program. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 1997; 59(1-B):429. UMI No. 9821478.
- Sugarman, DE. Web-based alcohol feedback intervention for heavy drinking college students: Does drinking control strategy use mediate intervention effects? (Doctoral dissertation). 2009. Retrieved from http://surface.syr.edu/psy_etd/8 UMI No. 3385859
- Sugarman DE, Carey KB. Drink less or drink slower: The effects of instruction on alcohol consumption and drinking control strategy use. *Psychology of Addictive Behaviors*. 2009; 23:577–585. doi:10.1037/a0016580. [PubMed: 20025364]
- Terlecki, M. The long-term effect of a brief motivational alcohol intervention for heavy drinking mandated college students (Unpublished doctoral dissertation). Louisiana State University; Baton Rouge, Louisiana: 2011.
- Terlecki MA, Buckner JD, Larimer ME, Copeland AL. The role of social anxiety in a brief alcohol intervention for heavy-drinking college students. *Journal of Cognitive Psychotherapy*. 2011; 25:7–21. doi:10.1891/0889-8391.25.1.7.
- Terry DL. Screening and brief intervention for hazardous alcohol use: A pilot study in a college counseling center. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2012; 74(02, E) UMI No. 3527869.
- Thush C, Wiers RW, Moerbeek M, Ames SL, Grenard JL, Sussman S, Stacy AW. Influence of motivational interviewing on explicit and implicit alcohol-related cognition and alcohol use in at-risk adolescents. *Psychology of Addictive Behaviors*. 2009; 23:146–151. doi:10.1037/a0013789. [PubMed: 19290699]
- Turrisi R, Larimer ME, Mallett KA, Kilmer JR, Ray AE, Mastroleo NR, Montoya H. A randomized clinical trial evaluating a combined alcohol intervention for high-risk college students. *Journal of Studies on Alcohol and Drugs*. 2009; 70:555–567. [PubMed: 19515296]
- Voogt CV, Poelen EAP, Kleinjan M, Lemmers LACJ, Engels RCME. The effectiveness of the 'What Do You Drink' web-based brief alcohol intervention in reducing heavy drinking among students: A two-arm parallel group randomized controlled trial. *Alcohol & Alcoholism*. 2013; 48:312–321. doi:10.1093/alcalc/ags133. [PubMed: 23303466]

- Author Manuscript
- Author Manuscript
- Author Manuscript
- Author Manuscript
- Author Manuscript
- Wagener TL. Genuine vs. DrAFT-CS: A randomized trial comparing computer and live personalized feedback interventions for high-risk drinking among college students. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2010; 73(07, E) UMI No. 3498642.
- Wagener TL, Leffingwell TR, Mignogna J, Mignogna MR, Weaver CC, Cooney NJ, Claborn KR. Randomized trial comparing computer-delivered and face-to-face personalized feedback interventions for high-risk drinking among college students. *Journal of Substance Abuse Treatment*. 2012; 43:260–267. doi:10.1016/j.jsat.2011.11.001. [PubMed: 22197301]
- Walters ST. In praise of feedback: An effective intervention for college students who are heavy drinkers. *Journal of American College Health*. 2000; 48:235–238. doi:10.1080/07448480009599310. [PubMed: 10778024]
- Walters ST, Bennett ME, Miller JH. Reducing alcohol use in college students: A controlled trial of two brief interventions. *Journal of Drug Education*. 2000; 30:361–372. doi:10.2190/jhml-0jpd-ye7l-14ct. [PubMed: 11092154]
- Walters ST, Vader AM, Harris TR. A controlled trial of web-based feedback for heavy drinking college students. *Prevention Science*. 2007; 8:83–88. doi:10.1007/s1121-006-0059-9. [PubMed: 17136461]
- Walters ST, Vader AM, Harris TR, Field CA, Jouriles EN. Dismantling motivational interviewing and feedback for college drinkers: A randomized clinical trial. *Journal of Consulting and Clinical Psychology*. 2009; 77:64–73. doi:10.1037/a0014472. [PubMed: 19170454]
- Walton MA, Chermack ST, Shope JT, Bingham CR, Zimmerman MA, Blow FC, Cunningham RM. Effects of a brief intervention for reducing violence and alcohol misuse among adolescents: A randomized controlled trial. *Journal of the American Medical Association*. 2010; 304:527–535. doi:10.1001/jama.2010.1066. [PubMed: 20682932]
- Weaver CC. A randomized controlled trial comparing a computer-based personalized feedback intervention with and without a moderation skills module. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2011; 73(01) UMI No. 3474706.
- Weitzel JA, Bernhardt JM, Usdan S, Mays D, Glanz K. Using wireless handheld computers and tailored text messaging to reduce negative consequences of drinking alcohol. *Journal of Studies on Alcohol and Drugs*. 2007; 68:534–537. [PubMed: 17568957]
- Werch CE, Anzalone DM, Brokiewicz LM, Felker J, Carlson JM, Castellon-Vogel EA. An intervention for preventing alcohol use among inner-city middle school students. *Archives of Family Medicine*. 1996; 5:146–152. doi:10.1001/archfami.5.3.146. [PubMed: 8620255]
- Werch CE, Carlson JM, Pappas DM, Edgemon P, DiClemente CC. A brief alcohol preventive intervention for student athletes. *Prevention Researcher*. 2002; 9:4–5.
- Werch CE, Moore MM, DiClemente CC, Owen DM, Carlson JM, Jobli E. Single vs. multiple drug prevention: Is more always better? A pilot study. *Substance Use & Misuse*. 2005; 40:1085–1101. doi:10.1081/ja-200030814.
- Werch C, Jobli E, Moore MJ, DiClemente CC, Dore H, Brown CH. A brief experimental alcohol beverage-tailored program for adolescents. *Journal of Studies on Alcohol*. 2005; 66:284–290. [PubMed: 15957680]
- White HR, Mun EY, Morgan TJ. Do brief personalized feedback interventions work for mandated students or is it just getting caught that works? *Psychology of Addictive Behaviors*. 2008; 22:107–116. doi:10.1037/0893-164x.22.1.107. [PubMed: 18298236]
- Whiteside U. A brief personalized feedback intervention integrating a motivational interviewing therapeutic style and dialectical behavioral therapy skills for depressed or anxious heavy drinking young adults. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 2010; 71(12) UMI No. 3431523.
- Wiers RW, Kummeling RHC. An experimental test of an alcohol expectancy challenge in mixed gender groups of young heavy drinkers. *Addictive Behaviors*. 2004; 29:215–220. doi:10.1016/S0306-4603(03)00081-9. [PubMed: 14667432]
- Winters KC, Fahnhorst T, Botzet A, Lee S, Lalone B. Brief intervention for drug-abusing adolescents in a school setting: Outcomes and mediating factors. *Journal of Substance Abuse Treatment*. 2012; 42:279–288. doi:10.1016/j.jsat.2011.08.005. [PubMed: 22000326]

- Wood MD, Capone C, Laforge R, Erickson DJ, Brand NH. Brief motivational intervention and alcohol expectancy challenge with heavy drinking college students: A randomized factorial study. *Addictive Behaviors*. 2007; 32:2509–2528. doi:10.1016/j.addbeh.2007.06.018. [PubMed: 17658696]
- Wood MD, Fairlie AM, Fernandez AC, Borsari B, Capone C, Laforge R, Carmona-Barros R. Brief motivational and parent interventions for college students: A randomized factorial study. *Journal of Consulting and Clinical Psychology*. 2010; 78:349–361. doi:10.1037/a0019166. [PubMed: 20515210]
- Wooten BT. Challenging alcohol expectancies: An application to adolescents. *Dissertation Abstracts International: Section B. Sciences and Engineering*. 1995; 56(12-B):706056. UMI No. 9610329.
- Zhang Y. Effects of an alcohol education program for high school students in Wuhan, China. *Dissertation Abstracts International: Section A. Humanities and Social Sciences*. 2009; 70(30) UMI No. 3350261.

References

- Allison, PD. *Missing data*. Sage; Thousand Oaks, CA: 2002.
- Ballesteros J, Duffy JC, Querejeta I, Arino J, Gonzalez-Pinto A. Efficacy of brief interventions for hazardous drinkers in primary care: Systematic review and meta-analyses. *Alcoholism: Clinical and Experimental Research*. 2004; 28:608–618. doi:10.1097/01.ALC.0000122106.84718.67.
- Barry, KL. *Brief interventions and brief therapies for substance use*. Rockville, MD: Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment: 1999. Treatment Improvement Protocol (TIP) Series 34. DHHS Publication No. SMA 99-3353
- Bell EC, Marcus DK, Goodlad JK. Are the parts as good as the whole? A meta-analysis of component treatment studies. *Journal of Consulting and Clinical Psychology*. 2013; 81:722–736. doi: 10.1037/a0033004. [PubMed: 23688145]
- Beich A, Thorsen T, Rollnick S. Screening in brief intervention trials targeting excessive drinkers in general practice: Systematic review and meta-analysis. *BMJ*. 2003; 327:1–7. doi:10.1136/bmj.327.7414.536. [PubMed: 12842922]
- Berkowitz AD. The social norms approach: Theory, research, and annotated bibliography. Retrieved from. 2004 http://www.alanberkowitz.com/articles/social_norms.pdf
- Bertholet N, Daeppen J, Wietlisbach V, Fleming M, Burnand B. Reduction of alcohol consumption by brief alcohol intervention in primary care. *Archives of Internal Medicine*. 2005; 165:986–995. doi:10.1001/archinte.165.9.986. [PubMed: 15883236]
- Bewick BM, Trusler K, Barkham M, Hill AJ, Cahill J, Mulhern B. The effectiveness of web-based interventions designed to decrease alcohol consumption – A systematic review. *Preventive Medicine*. 2008; 47:17–26. doi:10.1016/j.ypmed.2008.01.005. [PubMed: 18302970]
- Bien TH, Miller WR, Tonigan JS. Brief interventions for alcohol problems: A review. *Addiction*. 1993; 88:315–336. doi:10.1111/j.1360-0443.1993.tb00820.x. [PubMed: 8461850]
- Brown SA, McGue M, Maggs J, Schulenberg J, Hingson R, Swartzwelder S, Martin C, Chung T, Tapert SF, Sher K, Winters KC, Lowman C, Murphy S. Underage alcohol use: Summary of developmental processes and mechanisms: Ages 16-20. *Alcohol Research & Health*. 2009; 32:41–52. [PubMed: 23104446]
- Burke BL, Arkowitz H, Menchola M. The efficacy of motivational interviewing: A meta-analysis of controlled clinical trials. *Journal of Consulting and Clinical Psychology*. 2003; 71:843–861. doi: 10.1037/0022-006x.71.5.843. [PubMed: 14516234]
- Carey KB, Scott Sheldon, L AJ, Carey MP, DeMartini KS. Individual-level interventions to reduce college student drinking: A meta-analytic review. *Addictive Behaviors*. 2007; 32:2469–2494. doi:10.1016/j.addbeh.2007.05.004. [PubMed: 17590277]
- Carey KB, Scott-Sheldon LAJ, Elliott JC, Bolles JR, Carey MP. Computer-delivered interventions to reduce college student drinking: A meta-analysis. *Addiction*. 2009; 104:1807–1819. doi: 10.1111/j.1360-0443.2009.02691.x. [PubMed: 19744139]
- Cohen, J. *Statistical power for the behavioral sciences*. 2nd. Erlbaum; Hillsdale, NJ: 1988.

- Cunningham RM, Bernstein SL, Walton M, Broderick K, Vaca FE, Woolard R, Bernstein E, Blow F, D'Onofrio G. Alcohol, tobacco, and other drugs: Future directions for screening and intervention in the emergency department. *Academic Emergency Medicine*. 2009; 16:1078–1088. doi: 10.1111/j.1553-2712.2009.00552.x. [PubMed: 20053226]
- Fachini A, Aliane PP, Martinez EZ, Furtado EF. Efficacy of Brief Alcohol Screening Intervention for College Students (BASICS): A meta-analysis of randomized controlled trials. *Substance Abuse Treatment, Prevention, and Policy*. 2012; 7:1–20. doi:10.1186/1747-597X-7-40.
- Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL. Brief physician advice for problem drinkers: Long-term efficacy and benefit-cost analysis. *Alcoholism: Clinical and Experimental Research*. 2002; 26:36–43. doi:10.1111/j.1530-0277.2002.tb02429.x.
- Heather N. Interpreting the evidence on brief interventions for excessive drinkers: The need for caution. *Alcohol & Alcoholism*. 1995; 30:287–296. [PubMed: 7545982]
- Hedges LV. Distribution theory for Glass's estimator of effect size and related estimators. *Journal of Educational Statistics*. 1981; 6:107–128. doi:10.2307/1164588.
- Hedges, LV.; Olkin, I. *Statistical methods for meta-analysis*. Academic Press; San Diego, CA: 1985.
- Hedges LV, Tipton E, Johnson MC. Robust variance estimation in meta-regression with dependent effect size estimates. *Research Synthesis Methods*. 2010; 1:39–65. doi:10.1002/jrsm.5.
- Higgins, JPT.; Deeks, JJ.; Altman, DG.; Higgins, JPT.; Green, S., editors. *Cochrane handbook for systematic reviews of interventions*. John Wiley & Sons; Chichester, UK: 2008. Chapter 16: Special topics in statistics; p. 481-529.
- Jensen CD, Cushing CC, Aylward BS, Craig JT, Sorell DM, Steele RG. Effectiveness of motivational interviewing interventions for adolescent substance use behavior change: A meta-analytic review. *Journal of Consulting and Clinical Psychology*. 2011; 79:433–440. doi:10.1037/a0023992. [PubMed: 21728400]
- Kaner EFS, Dickinson HO, Beyer FR, Campbell F, Schlesinger C, Heather N, Saunders JB, Burnand B, Pienaar ED. Effectiveness of brief alcohol interventions in primary care. *Cochrane Database of Systematic Reviews*. 2007; (2) Art. No.: CD004148. doi:10.1002/14651858.CD004148.pub3.
- Kulesza M, Apperson M, Larimer ME, Copeland AL. Brief alcohol intervention for college drinkers: How brief is? *Addictive Behaviors*. 2010; 35:730–733. doi:10.1016/j.addbeh.2010.03.011. [PubMed: 20381972]
- Leffingwell TR, Horton R, Mignogna J, Jackson M, Leedy M, Lack C. A randomized trial of a computerized multimedia feedback intervention for high risk drinking among college students. 2007 Unpublished manuscript.
- Lipsey, MW.; Wilson, DB. *Practical meta-analysis*. Sage; Thousand Oaks, CA: 2001.
- Miller, WR.; Rollnick, S. *Motivational interviewing: Preparing people to change addictive behavior*. Guilford Press; New York, NY: 1991.
- Moreira MT, Smith LA, Foxcroft D. Social norms interventions to reduce alcohol misuse in university or college students. *Cochrane Database of Systematic Reviews*. 2009; (3) Art. No.: CD006748. doi:10.1002/14651858.CD006748.pub2.
- Neighbors CJ, Barnett NP, Rohsenow DJ, Colby SM, Monti PM. Cost-effectiveness of a motivational intervention for alcohol-involved youth in a hospital emergency department. *Journal of Studies on Alcohol and Drugs*. 2010; 71:384–394. [PubMed: 20409432]
- Nilsen P, Kaner E, Babor TF. Brief intervention, three decades on: An overview of research findings and strategies for more widespread implementation. *Nordic Studies on Alcohol and Drugs*. 2008; 25:453–467.
- Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry. *Journal of Clinical Epidemiology*. 2008; 61:991–996. doi:10.1016/j.jclinepi.2007.11.010. [PubMed: 18538991]
- Poikolainen K. Effectiveness of brief interventions to reduce alcohol intake in primary health care populations: A meta-analysis. *Preventive Medicine*. 1999; 28:503–509. [PubMed: 10329341]
- Prochaska, J.; DiClemente, C. *The transtheoretical approach: Crossing traditional boundaries of therapy*. Dow Jones-Irwin; Homewood, IL: 1984.

- Sánchez-Meca J, Marín-Martínez F, Chacón-Moscoso S. Effect-size indices for dichotomized outcomes in meta-analysis. *Psychological Methods*. 2003; 8:448–467. doi:10.1037/1082-989X.8.4.448. [PubMed: 14664682]
- Scott-Sheldon LAJ, DeMartini KS, Carey KB, Carey MP. Alcohol interventions for college students improves antecedents of behavioral change: Results from a meta-analysis of 34 randomized controlled trials. *Journal of Social and Clinical Psychology*. 2009; 7:799–823. doi:10.1521/jscp.2009.28.7.799.
- Substance Abuse and Mental Health Services Administration (SAMHSA). Results from the 2011 National Survey on Drug Use and Health: Summary of national findings. Substance Abuse and Mental Health Services Administration; Rockville, MD: 2012. NSDUH Series H-44, HHS Publication No. (SMA) 12-4713
- Substance Abuse and Mental Health Services Administration (SAMHSA). Screening, Brief Intervention, and Referral to Treatment (SBIRT). 2014 Retrieved from <http://beta.samhsa.gov/sbirt>.
- Tait RJ, Hulse GK. A systematic review of the effectiveness of brief interventions with substance using adolescents by type of drug. *Drug and Alcohol Review*. 2003; 22:337–346. doi: 10.1080/0959523031000154481. [PubMed: 15385228]
- Tanner-Smith EE, Wilson SJ, Lipsey MW. The comparative effectiveness of outpatient treatment for adolescent substance abuse: a meta-analysis. *Journal of Substance Abuse Treatment*. 2013; 44:145–158. doi:10.1016/j.jsat.2012.05.006. [PubMed: 22763198]
- Vasilaki EI, Hosier SG, Cox WM. The efficacy of motivational interviewing as a brief intervention for excessive drinking: A meta-analytic review. *Alcohol & Alcoholism*. 2006; 41:328–335. doi: 10.1093/alcalc/agl016. [PubMed: 16547122]
- Walton MA, Goldstein AL, Chermack ST, McCammon RJ, Cunningham RM, Barry KL, Blow FC. Brief alcohol intervention in the emergency department: Moderators of effectiveness. *Journal of Studies on Alcohol and Drugs*. 2008; 69:550–560. [PubMed: 18612571]
- Whitlock EP, Polen MR, Green CA, Orleans T, Klein J. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: A summary of the evidence for the U.S. preventive services task force. *Annals of Internal Medicine*. 2004; 140:557–568. doi: 10.7326/0003-4819-140-7-200404060-00017. [PubMed: 15068985]
- Wilk AI, Jensen NM, Havighurst TC. Meta-analysis of randomized control trials addressing brief interventions in heavy alcohol drinkers. *Journal of General Internal Medicine*. 1997; 12:274–283. doi:10.1007/s11606-006-5063-z. [PubMed: 9159696]
- Wilson SJ, Lipsey MW, Tanner-Smith EE, Huang C, Steinka-Fry K. Dropout prevention and intervention programs: Effects on school completion and dropout among school-aged children and youth (Protocol). *Campbell Systematic Reviews*. 2010 Available at: <http://www.campbellcollaboration.org/lib/project/158/>.
- Winters KC, Leitten W. Brief intervention for drug-abusing adolescents in a school setting. *Psychology of Addictive Behaviors*. 2007; 21:249–254. doi:10.1037/0893-164x.21.2.249. [PubMed: 17563146]
- Wutzke SE, Shiell A, Gomel MK, Conigrave KM. Cost effectiveness of brief interventions for reducing alcohol consumption. *Social Science and Medicine*. 2001; 52:863–870. doi:10.1016/S0277-9536(00)00189-1. [PubMed: 11234861]

Highlights

- Brief alcohol interventions for youth yield modest beneficial effects.
- Effects persist for up to one year after intervention receipt.
- Decisional balance and goal-setting exercises are beneficial for adolescents.
- Effects are consistent across diverse settings, formats, and populations.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1.
Study identification flow diagram




Figure 2. Scatter plots of method-adjusted effect sizes and posttest follow-up timing, by age group and outcome type
Notes: Effect sizes adjusted for study method characteristics and shown proportionate to random-effects inverse variance weights. Fitted values and confidence intervals from meta-regression with robust variance estimates. Dashed line shown at null value of zero.

Table 1

Characteristics of the Studies, Participant Samples, and Interventions in the Metaanalysis, by Age Group

	Adolescents <i>k</i> = 24 <i>n</i> = 172		Young Adults <i>k</i> = 161 <i>n</i> = 1,691	
	<i>M</i> (<i>SD</i>) % (<i>n</i>)	<i>r</i> _{ES}	<i>M</i> (<i>SD</i>) % (<i>n</i>)	<i>r</i> _{ES}
Study/Design Characteristics				
U.S. sample; % (<i>n</i>) ^a	50 (12)	-.03	81 (130)	-.03
Journal article publication; % (<i>n</i>) ^a	71 (17)	.09	75 (120)	.04
Randomized controlled trial; % (<i>n</i>) ^a	79 (19)	-.07	90 (145)	-.04
Attrition; <i>M</i> (<i>SD</i>) ^b	.12 (0.14)	.16	.23 (0.19)	-.09
Binary data used to estimate ES; % (<i>n</i>) ^b	37 (63)	-.21	7 (117)	-.09
Pretest adjusted ES; % (<i>n</i>) ^b	8 (14)	-.12	14 (241)	.13
Follow-up timing (weeks); <i>M</i> (<i>SD</i>) ^b	24 (14)	-.23	20 (24)	-.14
No treatment control condition; % (<i>n</i>) ^b	39 (67)	.32	59 (990)	.15
Pretest effect size; <i>M</i> (<i>SD</i>) ^b	-0.00 (0.59)	.15	-0.07 (0.37)	.26
Participant Characteristics				
Average age; <i>M</i> (<i>SD</i>) ^c	15 (1.5)	.07	20 (1.7)	.14
Percent male composition; <i>M</i> (<i>SD</i>) ^c	.53 (0.17)	.22	.47 (0.19)	-.00
Percent White composition; <i>M</i> (<i>SD</i>) ^c	.49 (0.29)	-.33	.76 (0.18)	.01
High-risk screened sample; % (<i>n</i>) ^a	29 (7)	.19	52 (83)	-.05
Intervention Modality; % (<i>n</i>) ^c				
21 st birthday card	0	-	8 (20)	-.06
Cognitive behavioral therapy (CBT)	3 (1)	-.03	5 (12)	.00
Motivational enhancement therapy (MET)	42 (14)	.25	35 (91)	.13
CBT + MET	12 (4)	-.14	4 (10)	-.05
Expectancy challenge	3 (1)	-.02	6 (16)	.11
Feedback/information only	0 (0)	-	28 (73)	-.09
Psycho-educational therapy (PET)	36 (12)	-.14	12 (31)	-.09
Intervention Site; % (<i>n</i>) ^c				
Primary care/student health center	9 (3)	-.29	5 (13)	.05
School/university	82 (27)	.19	55 (143)	.09
Self-administered	0 (0)	-	34 (87)	-.12
Emergency room	9 (3)	-.01	2 (6)	-.05
Intervention Format; % (<i>n</i>) ^c				
Subject alone – computerized	6 (2)	-.29	13 (33)	-.04
Subject alone – non-computerized	6 (2)	-.09	27 (70)	-.00

	Adolescents <i>k</i> = 24 <i>n</i> = 172		Young Adults <i>k</i> = 161 <i>n</i> = 1,691	
	<i>M</i> (<i>SD</i>) % (<i>n</i>)	<i>r</i> _{ES}	<i>M</i> (<i>SD</i>) % (<i>n</i>)	<i>r</i> _{ES}
Individual	39 (13)	.26	32 (82)	-.00
Group	42 (14)	-.05	28 (72)	.03
Family	3 (1)	-.06	0 (0)	-
Intervention Length; <i>M</i> (<i>SD</i>) ^c				
Total contact time (minutes)	99.9 (80.3)	.19	54.6 (57.2)	.13
Total number of sessions	1.8 (1.2)	.06	1.3 (1.0)	.12
Total days covered	6.2 (9.4)	.03	2.9 (5.4)	.14

Notes. Means and standard deviations shown for continuous measures; percentages and counts shown for dichotomous measures. *k* = number of studies; *n* = number of effect sizes. *r*_{ES} = bivariate correlation with alcohol consumption effect sizes.

^aEstimates calculated at study level.

^bEstimates calculated at effect size level.

^cEstimates calculated at intervention group level.

Table 2

Predicted Mean Covariate-Adjusted Effect Sizes and 95% Confidence Intervals Split by Intervention Characteristic, by Age Group and Outcome Type

	Adolescents		Young Adults	
	Alcohol Consumption	Alcohol Related Problems	Alcohol Consumption	Alcohol Related Problems
Modality				
21 st birthday card	--	--	0.07 [-0.02, 0.17] _{8, 36}	0.04 [-0.07, 0.14] _{2, 9}
CBT	--	--	0.13 [0.04, 0.22] _{8, 67}	0.10 [0.01, 0.20] _{4, 15}
MET	0.33 [0.23, 0.42] _{12, 71}	0.30 [0.19, 0.40] _{4, 26}	0.20 [0.14, 0.26] _{68, 623}	0.17 [0.11, 0.24] _{52, 182}
MET/CBT	0.16 [0.03, 0.28] _{3, 18}	0.13 [0.01, .24] _{3, 11}	0.03 [-0.10, 0.17] _{6, 52}	0.00 [-0.12, 0.13] _{6, 20}
Expectancy challenge	--	--	0.36 [0.10, 0.62] _{12, 46}	0.34 [0.08, 0.59] _{5, 9}
Feedback/information	--	--	0.20 [0.13, 0.26] _{54, 372}	0.17 [0.10, 0.24] _{30, 99}
PET	0.28 [0.18, 0.39] _{8, 39}	--	0.16 [0.09, 0.23] _{20, 104}	0.13 [0.06, 0.20] _{12, 44}
Delivery Site				
Primary care/health center	0.29 [0.11, 0.48] _{2, 19}	--	0.17 [0.05, 0.28] _{10, 97}	0.12 [0.01, 0.24] _{9, 51}
School/university	0.29 [0.19, 0.39] _{20, 94}	0.25 [0.15, 0.35] _{6, 26}	0.21 [0.14, 0.28] _{87, 751}	0.17 [0.11, 0.23] _{57, 192}
Emergency room	0.25 [0.11, 0.38] _{2, 21}	0.21 [0.07, 0.34] _{2, 12}	0.11 [0.04, 0.19] _{4, 37}	0.07 [-0.02, 0.16] _{3, 19}
Self-administered	--	--	0.21 [0.15, 0.27] _{57, 377}	0.17 [0.10, 0.23] _{31, 109}
Format				
Subject alone – comp.	--	--	0.23 [0.14, 0.33] _{18, 79}	0.19 [0.09, 0.28] _{8, 18}
Subject alone – not comp.	0.29 [0.19, 0.38] _{2, 10}	0.24 [0.14, 0.35] _{2, 7}	0.20 [0.15, 0.26] _{54, 383}	0.16 [0.10, 0.22] _{30, 108}
Individual	0.28 [0.19, 0.38] _{12, 71}	0.24 [0.13, 0.34] _{5, 29}	0.20 [0.14, 0.26] _{57, 560}	0.16 [0.09, 0.22] _{46, 192}
Group	0.29 [0.17, 0.41] _{10, 31}	0.25 [0.13, 0.36] _{2, 2}	0.21 [0.11, 0.31] _{43, 286}	0.16 [0.08, 0.25] _{22, 60}
Length				
Single session, < 5 min.	--	--	0.30 [0.09, 0.51] _{3, 14}	0.25 [0.05, 0.46] _{2, 8}
Single session, 5-15 min.	0.27 [0.15, 0.38] _{2, 19}	--	0.19 [0.13, 0.26] _{55, 330}	0.15 [0.08, 0.23] _{23, 71}
Single session, > 15 min.	0.29 [0.19, 0.39] _{16, 78}	0.25 [0.14, 0.36] _{5, 18}	0.22 [0.15, 0.28] _{82, 694}	0.17 [0.11, 0.24] _{57, 195}
Multi-session	0.25 [0.14, 0.35] _{8, 29}	0.20 [0.09, 0.31] _{3, 20}	0.17 [0.10, 0.25] _{30, 274}	0.13 [0.06, 0.20] _{23, 105}

Notes: Results of statistical significance tests provided no evidence of differences in mean effect sizes across intervention characteristic categories. Effect sizes adjusted for study method and participant characteristics. 95% confidence intervals estimated with robust standard errors. CBT – cognitive behavioral therapy. MET –motivational enhancement therapy. PET – psycho-educational therapy. MET/CBT – motivational enhancement therapy and cognitive behavioral therapy. -- results not available (fewer than two studies in cell). Subscripts indicate *k* (number of studies), *n* (number of effect sizes).

Table 3

Predicted Mean Covariate-Adjusted Effect Sizes and 95% Confidence Intervals Split by Presence/Absence of Intervention Components, by Age Group and Outcome Type

	Adolescents		Young Adults	
	Alcohol Consumption	Alcohol Related Problems	Alcohol Consumption	Alcohol Related Problems
BAC information				
Yes	0.36 [0.07, 0.65] _{4, 21}	--	0.20 [0.14, 0.26] _{63, 650}	0.14 [0.08, 0.20] _{51, 204}
No	0.43 [0.25, 0.61] _{20, 113}	--	0.19 [0.12, 0.26] _{100, 662}	0.09 [0.03, 0.15] _{51, 175}
Caloric information				
Yes	--	--	0.19 [0.11, 0.27] _{19, 177}	0.06 [-0.03, 0.15] _{13, 48}
No	--	--	0.19 [0.13, 0.26] _{141, 1135}	0.11 [0.06, 0.16] _{86, 331}
Decisional balance				
Yes	0.52 [0.33, 0.71]_{9, 45}	0.46 [0.32, 0.61] _{5, 32}	0.26 [0.14, 0.38] _{32, 284}	0.15 [0.07, 0.23] _{26, 77}
No	0.30 [0.14, 0.46]_{15, 89}	0.29 [-0.18, 0.75] _{3, 6}	0.17 [0.12, 0.22] _{129, 1028}	0.10 [0.05, 0.16] _{74, 302}
Education/information				
Yes	0.35 [0.15, 0.55] _{20, 122}	0.36 [-0.54, 1.26] _{7, 28}	0.18 [0.13, 0.24] _{104, 872}	0.11 [0.05, 0.16] _{72, 274}
No	0.50 [0.34, 0.66] _{6, 12}	0.49 [-0.10, 1.07] _{2, 10}	0.21 [0.12, 0.30] _{64, 440}	0.12 [0.05, 0.20] _{31, 105}
Feedback, personalized				
Yes	0.39 [0.21, 0.57] _{12, 76}	1.34 [1.17, 1.50]_{6, 19}	0.19 [0.13, 0.26] _{110, 1030}	0.12 [0.07, 0.18] _{76, 307}
No	0.44 [0.24, 0.64] _{13, 58}	0.03 [-0.04, 0.10]_{2, 19}	0.20 [0.11, 0.29] _{55, 282}	0.07 [-0.01, 0.15] _{27, 72}
Feedback, gender-specific				
Yes	0.41 [0.13, 0.69] _{2, 21}	0.84 [0.52, 1.16] _{2, 12}	0.19 [0.11, 0.27] _{36, 348}	0.10 [0.02, 0.17] _{23, 75}
No	0.41 [0.24, 0.59] _{22, 113}	0.38 [0.18, 0.59] _{6, 26}	0.19 [0.12, 0.27] _{128, 964}	0.12 [0.06, 0.18] _{77, 304}
Goal-setting				
Yes	0.48 [0.30, 0.66]_{10, 53}	0.45 [0.33, 0.57] _{6, 33}	0.22 [0.13, 0.31] _{38, 349}	0.15 [0.08, 0.21] _{27, 116}
No	0.31 [0.16, 0.46]_{14, 81}	0.21 [-0.21, 0.63] _{2, 5}	0.18 [0.12, 0.24] _{125, 963}	0.09 [0.03, 0.16] _{74, 263}
Money/cost information				
Yes	--	--	0.20 [0.13, 0.27] _{24, 207}	0.10 [0.03, 0.18] _{18, 61}
No	--	--	0.19 [0.13, 0.26] _{136, 1105}	0.11 [0.06, 0.17] _{82, 318}
Norm referencing				
Yes	0.43 [0.29, 0.57] _{8, 57}	0.88 [0.71, 1.06]_{3, 13}	0.18 [0.13, 0.24] _{111, 1015}	0.11 [0.06, 0.17] _{76, 306}
No	0.39 [0.13, 0.65] _{16, 77}	0.34 [0.31, 0.36]_{5, 25}	0.22 [0.11, 0.32] _{61, 297}	0.10 [0.01, 0.19] _{28, 73}

Notes: Results in bold indicate significant differences between studies with versus without intervention component (within age and outcome categories). Effect sizes adjusted for study method and participant characteristics. 95% confidence intervals estimated with robust standard errors. BAC – blood alcohol concentration. -- results not available (fewer than two studies in cell). Subscripts indicate *k* (number of studies), *n* (number of effect sizes).

Appendix B

Unstandardized Coefficients and 95% Confidence Intervals from Meta-Regression Model Used to Create Method-Adjusted Effect Sizes

	b	95% CI
Main effects		
Adolescent sample (vs. young adult)	-0.02	[-0.22, 0.19]
Alcohol consumption outcome (vs. alcohol-related problem)	0.05 *	[0.01, 0.09]
Randomized controlled trial (vs. quasi-experimental)	-0.03	[-0.16, 0.09]
Attrition	-0.16 *	[-0.29, -0.04]
Effect size estimated from binary data (vs. not)	-0.16 *	[-0.27, -0.05]
Effect size estimated with pretest adjusted data (vs. not)	0.06	[-0.02, 0.14]
No treatment control condition (vs. treatment as usual)	0.05	[-0.00, 0.11]
Pretest effect size	0.27 *	[0.10, 0.44]
Interactions with age of sample		
Adolescent sample X Attrition	0.38	[-0.57, 1.34]
Adolescent sample X Pretest adjusted effect size data	-0.19	[-0.39, 0.02]
Intercept	0.14 *	[0.01, 0.28]
Residual between-studies variance τ^2		0.02
Number of studies (<i>k</i>)		185
Number of effect sizes (<i>n</i>)		1,863

Notes: Unstandardized coefficients and 95% confidence intervals estimated with meta-regression model using robust standard errors. Multiplicative interaction terms were used for all study method moderators that had bivariate correlations with alcohol consumption effect sizes that were in opposite directions (positive/negative) for adolescent and young adult samples (see Table 1).

* $p < .05$.

Appendix C

Bivariate Correlations between Study, Participant, and Intervention Characteristics

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.	14.	15.	16.	17.	18.	19.	20.	21.	22.	23.	
1. U.S. sample	1.0																							
2. RCT	.13	1.0																						
3. Follow-up timing	-.07	.00	1.0																					
4. No treatment control	-.09	-.05	.00	1.0																				
5. Cluster assignment	.04	-.17	-.04	-.10	1.0																			
6. High-risk sample	.02	-.02	.19	.17	-.45	1.0																		
7. Average age	-.16	-.06	-.11	.12	-.13	.20	1.0																	
8. Percent male	-.15	-.16	.16	.05	-.12	.13	.03	1.0																
9. Percent White	.09	.05	.07	.25	.03	.16	.05	.06	1.0															
10. 21 st birthday card	.07	.04	-.13	-.08	-.04	-.18	.11	-.08	.02	1.0														
11. CBT	.00	-.05	.01	.09	.18	-.20	.03	.07	.04	-.03	1.0													
12. MET	.04	-.05	.10	.06	-.04	.32	-.06	.02	.05	-.15	-.21	1.0												
13. CBT + MET	.09	.01	.01	.00	-.07	-.12	-.08	-.02	-.08	-.04	-.05	-.23	1.0											
14. Expectancy challenge	.04	-.06	-.10	-.05	.07	-.10	.07	.15	.03	-.03	-.04	-.17	-.04	1.0										
15. Feedback only	-.20	.14	.03	.02	-.18	.13	.19	-.03	-.04	-.09	-.13	-.56	-.14	-.10	1.0									
16. PET	.09	-.09	-.12	-.10	.22	-.30	-.22	-.05	-.03	-.05	-.07	-.32	-.08	-.06	-.19	1.0								
17. Primary care	-.20	.01	.12	-.26	-.09	.13	.10	.03	.10	-.05	-.07	-.02	.02	-.06	.03	.06	1.0							
18. School/univ.	.19	-.19	-.03	.28	.19	.01	-.09	-.04	.15	-.18	.13	.37	.06	.15	-.45	-.12	-.36	1.0						
19. Self-admin.	-.02	.16	-.08	-.04	-.08	-.16	.13	-.10	-.07	.26	-.06	-.49	-.14	-.10	.52	.16	-.19	-.68	1.0					
20. Emergency room	.05	.06	.10	-.25	-.09	.18	-.11	.11	-.30	-.04	-.05	.14	.15	-.04	-.13	-.08	-.07	-.26	-.13	1.0				
21. Subject alone (comp.)	.02	.07	-.10	-.06	-.08	-.13	.04	-.09	.01	.59	.01	-.21	-.06	-.01	.01	.03	.05	-.21	.25	-.06	1.0			
22. Subject alone (non-comp.)	-.15	.17	-.13	.04	-.10	-.07	.10	-.02	-.03	-.08	-.10	-.51	-.06	-.02	.60	.16	.02	-.55	.62	-.09	-.16	1.0		
23. Individual	.12	-.01	.25	.08	-.35	.51	-.03	.16	.08	-.14	-.16	.68	-.09	-.16	-.35	-.24	.03	.30	-.43	.20	-.23	-.56	1.0	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.	14.	15.	16.	17.	18.	19.	20.	21.	22.	23.
24. Group	.03	-.18	-.12	-.10	.56	-.45	-.08	-.13	-.08	-.08	.31	-.15	.21	.23	-.22	.09	-.08	.35	-.30	-.11	-.13	-.31	-.46