



Published in final edited form as:

J Subst Abuse Treat. 2015 April ; 51: 38–46. doi:10.1016/j.jsat.2014.11.004.

Characteristics of students participating in Collegiate Recovery Programs: A national survey

Alexandre B. Laudet, Ph.D.¹, Kitty Harris, Ph.D.², Thomas Kimball, Ph.D.², Ken C. Winters, Ph.D.³, and D. Paul Moberg, Ph.D.⁴

¹Center for the Study of Addictions and Recovery, National Development and Research Institutes, Inc., New York City, NY 10010

²Center for the Study of Addiction and Recovery, Texas Tech University, Lubbock, TX 79409-1160

³University of Minnesota

⁴University of Wisconsin

Abstract

Relapse rates are high among individuals with substance use disorders (SUD), and for young people pursuing a college education, the high rates of substance use on campus can jeopardize recovery. Collegiate Recovery Programs (CRPs) are an innovative campus-based model of recovery support that is gaining popularity but remains under-investigated. This study reports on the first nationwide survey of CRP-enrolled students (N = 486 from 29 different CRPs). Using an online survey, we collected information on background, SUD and recovery history, and current functioning. Most students (43% females, mean age = 26) had used multiple substances, had high levels of SUD severity, high rates of treatment and 12-step participation. Fully 40% smoke. Many reported criminal justice involvement and periods of homelessness. Notably, many reported being in recovery from, and currently engaging in multiple behavioral addictions-e.g., eating disorders, and sex and love addiction. Findings highlight the high rates of co-occurring addictions in this under-examined population and underline the need for treatment, recovery support programs and college health services to provide integrated support for mental health and behavioral addictions to SUD-affected young people.

Keywords

Recovery; college students; recovery support services; substance use disorder; addiction

© 2014 Elsevier Inc. All rights reserved.

Please address all correspondence to the first author at alexandrelaudet@gmail.com T: +1-646-387-6568.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

1.1 In recovery and in college: Double jeopardy

Rates of substance use disorders (SUD) triple from 7% in adolescence to 20% in early adulthood (Substance Abuse and Mental Health Services Administration, 2011), making this developmental stage critical to young people's future. In spite of effective interventions (Becker & Curry, 2008; Chung et al., 2003; M. Dennis et al., 2004; Tanner-Smith, Wilson, & Lipsey, 2013; Winters, Stinchfield, Lee, & Latimer, 2008), relapse rates are typically high (Substance Abuse and Mental Health Services Administration, 2008). Post-treatment continuing support is effective at sustaining recovery (M. Dennis & Scott, 2007; S. H. Godley et al., 2010; McKay et al., 2009; Substance Abuse and Mental Health Services Administration Office of Communications, 2009).

The need for recovery support is especially high for SUD-affected college students: Attending college and transitioning into adulthood can both be demanding, offering new freedoms but also less structure and supervision. For youths in SUD recovery, these challenging transitions are compounded by the need to remain sober in an 'abstinence-hostile environment' (Cleveland, Harris, & Wiebe, 2010): The high rates of substance use on campuses (Hingson, Zha, & Weitzman, 2009; Wechsler & Nelson, 2008) make college attendance a severe threat to sobriety that must often be faced without one's established support network (Bell et al., 2009; Woodford, 2001). Combined, these factors can lead to isolation when 'fitting in' is critical, and/or to yielding to peer pressure to use alcohol or drugs, both enhancing relapse risks (Harris, Baker, Kimball, & Shumway, 2008; Woodford, 2001).

Experts' calls for campus-based services for recovering students (Dickard, Downs, & Cavanaugh, 2011; Doyle, 1999) have thus far been largely unheeded (Bell, et al., 2009; Botzet, Winters, & Fahnhorst, 2007; Cleveland, Harris, Baker, Herbert, & Dean, 2007). The US Department of Education noted that 'the education system's role as part of the nation's recovery and relapse prevention support system is still emerging.' (p.10 (Dickard, et al., 2011). Preventing students relapse is especially critical as SUDs are associated with college attrition (Hunt, Eisenberg, & Kilbourne, 2010). Thus, youths' developmental stage, and the unique challenges of college, both underline the need for a recovery support infrastructure on campus (Botzet, et al., 2007; Misch, 2009). This includes the need for a recovery supportive social environment that fosters social connectedness, given the influence of peers on youths' substance use (Cimini et al., 2009; Substance Abuse and Mental Health Services Administration Office of Communications, 2009; White, 2008). Federal agencies recently called for the expansion of community based recovery support models to extend the continuum of care, *including in schools and colleges* (Office of National Drug Control Policy, 2010; U.S. Dept. of Education, 2010).

1.2 Collegiate Recovery Programs

Collegiate Recovery Programs (CRPs) started at a few universities in the 1980's to meet recovering students' support needs, as part of a broader effort to address substance use on campus. CRPs generally offered onsite sober housing, self-help meetings (e.g. 12-step), and

counseling provided by a small staff (Botzet, et al., 2007; Cleveland, et al., 2010; Smock, Baker, Harris, & D'sauza, 2011; White & Finch, 2006). CRPs' strive to create a campus-based 'recovery friendly' space and a supportive social community to enhance educational opportunities while supporting students' recovery and emotional growth (Harris, et al., 2008). The model fits into the continuing care paradigm of a 'recovery management' system (M. D. Godley, Godley, Dennis, Funk, & Passetti, 2002). Site-level records from a handful of CRPs suggest encouraging outcomes (Cleveland, et al., 2007), as do data from the site survey arm of this study (A. Laudet, Harris, Winters, Moberg, & Kimball, 2013): Across the 29 CRPs nationwide, annual relapse rates range from 0 to 25% (mean = 8%), and academic achievement (GPA and graduation) surpass the host institution's overall outcomes.

Several factors lead to increased interest in CRPs about a decade ago. This includes academic institutions and federal agencies' growing recognition of youth substance use and in particular, campus-based use, as a major public health concern, and federal agencies' shift to a recovery-oriented 'chronic care' approach to SUD services (Clark, 2008). These factors fueled a rapid growth of CRPs, from 4 in 2000 to a 29 in 2012 (A. Laudet, et al., 2013) with 5 to 7 starting annually (Kimball, 2014). While CRPs vary in orientation, budget, and in the breadth of services (A. Laudet, Harris, Kimball, Winters, & Moberg, 2014; A. Laudet, et al., 2013), most are peer-driven, 12-step based, provide onsite support groups, sober events, and seminars on SUD and recovery. The need for CRPs is bolstered by many sites reporting that demand surpasses capacity (A. Laudet, et al., 2013).

1.3 Need for Research on Collegiate Students in Recovery and Study Objectives

In spite of CRPs' rapid growth, they remain largely unexamined. Noting the lack of recovery resources in academic settings, the US Department of Education has called for research about CRPs and their students to inform the higher education system's response to college students in recovery (Dickard, et al., 2011). Information about CRP students can also to inform key stakeholder groups beyond the education system, starting with CRPs themselves. Unlike treatment programs that collect patient history upon admission to guide services, CRPs do not. Many operate with limited staff and budget (A. Laudet, et al., 2013) and lack the resources to collect student information. While five CRPs have operated for ten years or longer and some serve up to eighty students, two thirds emerged in the past five years, and over half serve fewer than ten students. Students at a given CRP are unlikely to represent the breadth of experiences and issues that a large data collection effort can document. That information can guide the development of support groups and related services, and prepare CRPs to address behavioral patterns they may face as the membership grows. Details about the broader CRP membership can also inform referral sources (e.g., high school counselors, therapists, treatment, university health staff) to determine the suitability of a CRP referral to a given student's needs.

Documenting the characteristics of CRP students will also yield knowledge about young people in recovery, an unexplored population. Clinicians and researchers understandably focus on individuals who are actively using substances or in early remission. Little is known about overall recovery paths (i.e., the totality of recovery supports used to achieve and sustain recovery), about how young people in stable recovery function, or the issues they

face. Moreover, because clinical practice and research tend to be specialized, researchers and clinicians may not explore co-morbid behavioral addiction patterns although they are likely relevant to SUD recovery-and vice versa. Thus, documenting the characteristics and experiences of college students in SUD recovery can also contribute to the knowledge base about persons in recovery to inform research and improve clinical practice.

Data for our study were collected in the context of a broader project designed to answer the U.S. Department of Education's call for research on CRPs. The first phase of the project was a nationwide survey of CRPs' structures and services, described elsewhere (A. Laudet, Harris, Kimball, et al., 2014; A. Laudet, et al., 2013). This study sought to characterize CRP students nationwide in terms of their background and current functioning.

Methods

2.1 Procedures and Participants Recruitment

New CRPs start organically; there is no centralized office or updated list of programs, though this may change with the formation of the emerging Association of Recovery in Higher Education. We had worked collaboratively with current and emerging CRPs since the planning of the study. We identified 29 CRPs nationwide when the program survey launched in the Fall of 2012, and recontacted these sites to enroll their assistance in recruiting students to take the survey. The 29 CRPs represented 19 U.S. states: 44% in the South, 22% in the Midwest, 19% in the Northeast, and 15% on the West coast. Most CRPs (85%) are hosted in public (vs. private) academic institutions; 76% operate in universities, 16% in a 4-year college, and 8% in a 2-year college.

Information sheets about the study procedures and a weblink to the confidential student survey were emailed to each CRP director; programs were instructed to email participating students, to make announcements about the survey, post the link on their internal website, and to post the study information sheet on bulletin boards at their site. The study was reviewed and approved by the ethics board (IRB) of the first author's institution, and we obtained a Certificate of Confidentiality from our funding agency. At the end of the survey, students had the option of providing their academic email address to receive a \$40 e-gift certificate at Amazon. A total of 486 unduplicated surveys were completed. Based on an estimated pool of 600 participants enrolled in CRPs over the data collection period, this represents an 81% participation rate.

2.2 Data collection and Instrument

Data collection ran from February 2013 through the spring, summer and fall semesters. The confidential survey, administered online using Survey Monkey®, started with the informed consent that described the study purpose and procedures and other required consent elements. The instrument consisted of measures and inventories summarized below, all of which we have used in previous federally funded studies of persons in SUD remission (Kaskutas et al., 2014; A. Laudet, 2007; A. Laudet, Stanick, & Sands, 2007; A. B. Laudet et al., 2004; Magura et al., 2003), with the exception of the behavioral addiction inventory that

was developed as described below. In addition to sociodemographics, background, and physical health, we collected data on the following domains:

2.2.1 Mental health—Participants answered the following questions: (a) Ever treated for an emotional/mental health problem (yes/no); IF YES: (b) Age at first treatment; (c) Ever hospitalized for an emotional/mental health problem (yes/no); (d) Ever diagnosed with a mental health disorder; IF YES: Diagnoses (up to three were coded); (e) Received treatment for an ('ongoing') mental health problem in the past year (yes/no) (f) Currently taking medications prescribed for a mental health/emotional problem (yes/no); and (g) Mental health self rating on a scale ranging from 1 (poor) to 4 (excellent).

2.2.2 Substance use involvement

Drug and alcohol use history: This was collected using a list of substance categories shown in Table 3. For each substance 'ever' used once or more, participants reported any 'regular' use - i.e., once a week or more for at least a one-year period'. For each substance used regularly, students were asked age at first regular use, duration of regular use (in years) and date last used. Students were also asked which of the 'regular use' substances had caused them the most serious problem - i.e., primary problem -, and which other substance(s) 'caused you serious problems?' (i.e., secondary problems). We computed the duration of abstinence from each regular substance, and summary variables representing the shortest duration of abstinence from any *drug*, from *alcohol*, and from *any substance* (i.e., *drug and/or alcohol*). Information was also collected on cigarette smoking.

Drug and alcohol Dependence severity: As in our previous studies (Kaskutas, et al., 2014; A. Laudet, 2007; A. Laudet, et al., 2007), our questionnaire combines the Lifetime versions of the Alcohol and Non-alcohol Psychoactive Substance Use Disorders subscales of the Mini International Neuropsychiatric Interview (M.I.N.I.), a short structured diagnostic interview developed in the U.S., and Europe for DSM-IV and ICD-10 psychiatric disorders (Sheehan et al., 1998). The MINI has been validated against the much longer Structured Clinical Interview for DSM diagnoses (SCID-P) and against the Composite International Diagnostic Interview for ICD-10 (CIDI). The 14 dichotomous items yield a single severity score that can range from 0 to 14. Cronbach Alpha = .81 in this sample.

2.2.3 Behavioral addictions—Although the study focused primarily on substance use, we were also interested in learning about students' behavioral addictions. Based on a review of the extant literature and on the team's expertise in young adults' health, we collected information about the following: eating disorders (i.e., anorexia, bulimia, bingeing), sex/love addiction, self harm/injury, gaming/gambling, compulsive shopping, Internet addiction (other than for sex, gambling or shopping), and compulsive exercise. Students first indicated every behavior from which they were in recovery (substance use and behavioral addictions). Next they indicated which problem they regarded as *primary*, and all the behaviors they regarded as *secondary* problems. Finally they reported past 90 days behaviors in the above mentioned behavioral categories: 'Independently of any alcohol or drug use, please mark any of the other addictive behaviors you may have engaged in *at a problematic or harmful level* during the past 90 days'.

2.2.4 Perceived harm of substance use and benefit of recovery—We use three summary items from the *Primary Appraisal Measure* (Morgenstern, Labouvie, McCrady, Kahler, & Frey, 1997) to assess a) perceived harm from past substance use; b) Likely negative impact if substance use were to resume/continue; and c) Likely benefit/improvement from being/remaining in recovery from drug/alcohol use. Answers are provided using a 5-point Likert-type scale ranging from No (harm/benefit) to extreme (harm/benefit).

2.2.5 Utilization of services and recovery resources

Substance abuse treatment: This section consisted of the following items: (a) Ever received addiction treatment services (yes/no); if yes, (b) age received first treatment; c) History of participation in various treatment modalities (e.g., detoxification, methadone maintenance, therapeutic community, 21/28-day inpatient rehab, outpatient treatment).

Twelve-step fellowships: Participants reported (a) whether they had ever attended a 12-step meeting (e.g., Alcohol Anonymous) to deal with a substance use problem; if yes, (b) age first attended; c) Number of meetings attended past year and past month. The section also included a nine items inventory of 12-step suggested behaviors that constitute 12-step involvement (e.g., having a sponsor, doing service, working the 12-step program) in the past year.

Non 12-step addiction recovery support groups: This section consisted of the following items: a) Ever attended a substance use recovery support group that is not 12-step based (e.g., Moderation Management, SMART Recovery); if yes, (b) age first attended; c) Number of meetings attended past year and past month.

Other recovery resources: Participants answered the following questions: a) Ever visited an emergency department because of drugs or alcohol problems; b) Ever prescribed any anticraving or other medications to deal with a drug or alcohol problem (e.g., naltrexone, acamprosate, buprenorphine); c) Ever enrolled in a wilderness program to deal with drugs or alcohol problem (e.g., Outward Bound); and d) Ever attended a recovery high school -a school designed for students in SUD recovery (D. P. Moberg & Finch, 2008).

3 Results

3.1 Sociodemographics and history

Slightly more students were males, and almost all were Caucasian (Table 1). While the mean age was 26, the range was broad (17 to 58) with 50% being over 23, including a quarter who was 28 or older (75th percentile). Most were enrolled in coursework full-time and lived off campus. Half lived with one or more roommate(s); of those, nearly half lived with someone in SUD recovery and nearly a quarter, with someone actively using drugs and/or alcohol (whether in moderation or to excess). One third had experienced a period of homelessness. A history of criminal justice involvement history was relatively prevalent, with over half reporting being arrested and charged, and over a third having been

incarcerated. However, most students had no current involvement with the criminal justice system.

3.2 Physical and mental health

Although one quarter of students had been to the emergency department for a medical problem or injury in the past year, and 40% smoke, they generally reported good physical health (Table 2). Three quarter had a history of mental health treatment and had been diagnosed with a mental health disorder. The top three diagnoses were unipolar depression (74%), anxiety disorders (48% including 8% post traumatic stress disorder), and bipolar disorder (23% - not shown in table 2).^a These reports none withstanding, most students rated their mental health as 'good' or 'very good'.

3.3 Substance use and behavioral addictions

Over half reported *drug* addiction - be it an illegal substance or abuse of prescribed medication - as their primary lifetime problem (52.6%), with alcohol a distant second (38.9%, top of Table 3).. Most also reported a secondary problem behavior, the majority citing a substance-related issue but approximately ten percent also citing a behavioral addiction - e.g., sex and love addiction, and disordered eating.

3.3.1 Drug and alcohol use history—Most had used multiple substances, with regular marijuana and alcohol use as the most frequently cited, starting in mid-adolescence and lasting for up to 7 years (alcohol). Alcohol was the single most frequently cited *primary* individual problem substance (41.2%) and was among 'other problem substance(s)' for another half (See Substances section of Table 3). Reports for any single *primary drugs* were significantly lower than for alcohol, with heroin, crack or cocaine, and pain medications as the top three mentions. However several *drugs* were cited a secondary problem substance by a quarter of students or more including marijuana, pain medications, and cocaine or crack.

The average lifetime addiction severity score was high (Mean = 11.4 on a 0-14 range; Std. Dev. = 2.6). Correspondingly, students reported high levels of perceived past harm from their substance use - not shown in table 3- (40.8% 'considerable harm' and 32.4% 'extreme harm'), a high degree of potential harm were they to continue or resume regular substance use (15.2% 'considerable harm' and 76.8% 'extreme harm'), and almost all viewed 'extreme' or 'considerable' benefit to future continued sobriety (83.2% and 13.1%).

3.3.2 Substance use and behavioral addictions: Current status—Most students had not drank alcohol or used drugs in several years (Mean days since last drank = 952; mean days since last used drugs = 1,053) although the range of durations since last use were both wide (Table 3). Five percent (5.4%) drank alcohol and/or used drugs in the past month. Restricting the analyses to students considering themselves in recovery from a substance use problem (N = 433) - see next section, 4.8% had used a substance in the past month (2.3 % alcohol and 3.6% drugs, not shown). Independently of any recent substance use, a small

^aUp to three diagnoses were coded.

percentage of students had engaged in one of more behavioral addiction in the past 90 days, sex and love, and disordered eating being the two most common (Table 3).

3.4 Recovery

Consistent with their history of multiple problem behaviors, most students considered themselves in recovery from more than one substance and/or behavioral problem (Table 4). Recovery from alcohol and from drug problems were cited most, as may be expected. While much less prevalent, several behavioral addictions were also cited, the top three being disordered eating (15.6%), self harm (10.5%), and sex and/or love addiction (9.5%).

3.5 Utilization of SUD services and recovery resources: Treatment and self-help groups

Most participants had received professional substance use treatment (detailed in Table 4). Mean age at first treatment was 20.6 years of age. In addition to treatment, frequently reported forms of help included individual counseling, and being prescribed medications. Finally though not strictly fitting in the addiction services category, 43.5% had visited an emergency room department because of drug or alcohol use (e.g., overdose) at least once over their lifetime.

Turning to self-help group participation, most had attended 12-step meetings, and level of 12-step involvement was generally high (Table 4). Asked to rate the helpfulness of 12-step attendance to their recovery (not shown in Table 4), over half (56.3%) selected 'extremely' and another quarter, 'very much'. A small percentage of students reported attendance at a non 12-step addiction recovery support meeting, a practice whose helpfulness was rated as low by most (19.2% 'not at all', 46.2% 'a little/moderately', 19.2% 'very much and 15%, 'extremely').

4. Discussion

4.1 Reprise of key findings

Consistent with the only published report - one that bears on a single site (Cleveland, et al., 2007), our national CRP student sample is somewhat older than traditional college students, mostly Caucasian, and enrolled in college full-time. On average, students had been in their CRP for 7 semesters, and had remained substance-free for nearly three years.

The use of multiple substances was the norm. *Alcohol* was primary for nearly four out of ten, and secondary for another 41%; no individual *drug* emerged as primary. While mean duration of substance use was shorter than in typical addiction study samples that consist of adults, lifetime addiction severity was high. Several other data points also suggest severe SUD problems and consequences. This includes high rates of treatment utilization including residential treatment and 12-step meetings, a third of students reporting a period of homelessness, and a high rate of past criminal justice involvement.

Among notable findings is the high rate of smoking: 40% of students smoke, twice the national rate (18.1%), even when considering only the age group with the highest smoking prevalence, adults aged 25–44 years (21.6%) (Center for Disease Control and prevention, 2014). Little is known of smoking patterns in SUD recovery. One study of adults in recovery

(N = 285, mean age 43.5) reported a 69.5% smoking rate (A. Laudet, 2009). Moreover, a large minority of smokers in that study reported that smoking had become more important since they stopped using substances, emphasizing the need to address smoking cessation as part of SUD services for young adults.

CRP students reported an extensive history of mental health problems. This is the first large scale study of young people (and college students) in recovery; we were unable to locate relevant reports bearing on this population. However our findings are consistent with several converging lines of research. This includes high rates of co-occurring mental health issues among adults in SUD recovery (A. Laudet, 2012; A. Laudet, Timko, & Hill, 2014) and among students in recovery high schools (D. P. Moberg, Finch, & Lindsley, 2014). Moreover, the prevalence of mental health disorders is generally high among college age individuals, regardless of college attendance status (Blanco et al., 2008).

While most students reported a substance as their primary problem as expected given the substance use focus of CRPs, behavioral addiction (BA) reports as a secondary problem were not infrequent, with sex and love addiction, and eating disorders reported by more than one in ten. Consistent with this finding, up to one in six students reported being in recovery from a behavioral addiction. Similar percentages had also engaged in a BA in the past 90 days. While the prevalence of BAs among SUD affected persons is likely not surprising to many clinicians, this is, to our knowledge, the first systematic attempt to document comprehensively such patterns among college students (be they in SUD recovery or not) or among SUD-affected persons.

Discussing behavioral addictions in depth is beyond our scope. We sought only to obtain a broad overview of the types and prevalence of BA in this population. Studies typically examining a single behavioral addiction have documented high rates of co-occurrence with substance use. For example, our findings on eating disorders (the most prevalent BA here) are in line with reports, most obtained among females, documenting a high co-occurrence of substance use (Cohen et al., 2010; Czarlinski, Aase, & Jason, 2012; National Center on Addiction and Substance Abuse, 2003).

4.2. Study Limitations

This is the first national study of students participating in a Collegiate Recovery Program. As such, it is an important initial step toward acquiring the information necessary information to develop and refine supports and services to promote recovery and a healthy life style in this population. This exploratory study has a number of limitations. The cross-sectional design relies on retrospective information for several key domains (e.g., substance use history), and the need to use a relatively short online survey precluded in-depth diagnostic interviews. Moreover, 19% of the estimated 600 CRP membership nationwide at the time of the survey did not participate. While our 81% participation is strong, as with all studies, some members of the target population are not represented. We cannot determine whether and how non completers compare to completers. However, our sample characteristics are highly consistent with CRP directors' reports of the students they serve (A. Laudet, et al., 2013; A. Laudet, Harris, Winters, Moberg, & Kimball, 2014), and with student characteristics reported in two single site CRP studies (Bell, et al., 2009; Cleveland,

et al., 2007). This affords us strong confidence about the representativeness of our sample and the generalizability of findings to CRP students nationwide. Our results have several implications for services development, and highlight a number of critical areas that need to be investigated further.

4.3 Implications for services and research

One of the unique aspects of this study in addition to its focus on SUD recovery, is that it also examines a number of unhealthy behavioral patterns that are ongoing in the context of SUD recovery although rarely examined simultaneously. While functioning well enough to attend college and remain substance free, many students continue to be treated for mental health, a significant proportion smoke, and a smaller percentage engage in behavioral addictions. As summarized in the Study Objectives section, disseminating such findings has implications for several stakeholder groups.

For CRPs, whose focus is on supporting SUD recovery, findings emphasize the need to broaden the scope of support groups, counseling (where available) and weekly addiction education seminars, to incorporate discussions about past and current behavioral addictions. Many CRPs may not have the necessary resources to address these issues as fully as they do SUDs, but could draw on institutional and community-based resources (e.g., faculty, local treatment agencies). Most importantly, openly discussing behavioral addictions can help decrease the stigma attached to these behaviors, enhancing the likelihood that students struggling with these issues discuss, and more importantly address their unhealthy behavior patterns. The same holds for mental health problems: CRPs may not have the local resources to treat these issues, but can sensitize students to the fact that common negative feelings such as depression or loneliness may threaten SUD recovery.

Findings can also inform CRP outreach to referral sources: CRPs are a new model about which many potential referral sources may not have heard. While treatment and 12-step utilization are the two most prevalent forms of help reported here, the utilization of emergency departments is also high. ER and local hospital staff may not be aware of the availability of a local CRP. Informing students treated in an ER for substance use about this resource as part of an ER based SUD intervention can potentially connect the student with a valuable source of recovery support.

College health personnel is anecdotally aware of the multiplicity of problems some students face. However, findings can be valuable to them in at least two ways. First, they highlight a growing campus-based recovery support resource many institutions may not know is available. Disseminating our findings can encourage institutions to explore hosting a CRP. Second, health clinics typically come into contact with students experiencing a specific concern- e.g., depression, or referred/mandated because of an offense resulting from substance use. It is critical that these contacts be used as an opportunity to explore potential co-occurring problematic behavior patterns. A recent national college study reported that few institutions have adequate strategies to address alcohol problems (Lenk, Erickson, Winters, Nelson, & Toomey, 2012; Toomey et al., 2013). While their effectiveness at addressing drug use and other behavioral problems was not examined, it is likely even less adequate than are strategies to deal with alcohol abuse. In addition to substance use and

behavioral addiction, the very high prevalence of smoking in our sample emphasizes the urgent need for academic institutions in general, and health programs serving college students in recovery in particular, to increase prevention efforts and to address smoking as an integrated component of recovery support and health services. Note that while campuses are smoke free, the majority of CRP students live off campus where smoking among peers is likely highly prevalent.

Finally, our findings have implications for SUD treatment programs serving young people, and for addiction researchers. SUD clinicians may be anecdotally aware of the potential for co-morbid behavioral addictions. However, although individuals grapple with multiple life problems simultaneously, professionals (i.e., providers, funders, researchers) typically focus on a single area of functioning (A. Laudet, 2012). For instance, until two decades ago, many SUD treatments excluded clients with mental health problems and vice versa. Our findings highlighting the relatively high prevalence of multiple behavioral health issues among young persons in SUD recovery underscore the need to adopt an integrated approach to promoting recovery and healthy functioning.

The need to adopt a more integrated approach applies to research as well. Historically, we have mostly focused on studying on a single ‘problem’, treating any co-occurring issue as a confound to be controlled for in analyses. The approach has merit in many contexts. Nonetheless, we ought to recognize the need to cast a wider net, especially when describing a previously under-examined population as we do here, or seeking to elucidate patterns of dynamic processes and broader outcomes, such as behavioral health. SUD researchers can incorporate indices of potentially related domains to begin bridging the knowledge gap about the prevalence of co-occurring behavioral addictions in SUD recovery. Failure to do can yield but a partial picture of overall clinical functioning that is insufficient to inform recovery and overall health services and supports.

Acknowledgments

The project was supported by Award Number R21DA033448 from the National Institute on Drug Abuse. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse or the National Institutes of Health.

Literature Cited

- Becker SJ, Curry JF. Outpatient interventions for adolescent substance abuse: a quality of evidence review. *J Consult Clin Psychol*. 2008; 76(4):531–543. [PubMed: 18665683]
- Bell NJ, Kanitkar K, Kerksiek KA, Watson W, Das A, Kostina-Ritchey E, Harris K. “It has made college possible for me”: feedback on the impact of a university-based center for students in recovery. *J Am Coll Health*. 2009; 57(6):650–657. [PubMed: 19433403]
- Blanco C, Okuda M, Wright C, Hasin DS, Grant BF, Liu SM, Olfson M. Mental health of college students and their non-college-attending peers: results from the National Epidemiologic Study on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2008; 65(12):1429–1437. [PubMed: 19047530]
- Botzet A, Winters K, Fahnhorst T. An exploratory assessment of a college substance abuse recovery program: Augsburg College’s StepUP Program. *Journal of Groups in Addiction and Recovery*. 2007; 2(2-4):257–287.
- Center for Disease Control and prevention. Current Cigarette Smoking Among Adults—United States, 2005-2012. *Morbidity and Mortality Weekly Report*. 2014; 63(2):29–34. [PubMed: 24430098]

- Chung T, Martin CS, Grella CE, Winters KC, Abrantes AM, Brown SA. Course of alcohol problems in treated adolescents. *Alcohol Clin Exp Res.* 2003; 27(2):253–261. [PubMed: 12605074]
- Cimini MD, Martens MP, Larimer ME, Kilmer JR, Neighbors C, Monserrat JM. Assessing the effectiveness of peer-facilitated interventions addressing high-risk drinking among judicially mandated college students. *J Stud Alcohol Drugs Suppl.* 2009; (16):57–66. [PubMed: 19538913]
- Clark, W. Recovery-Oriented Systems of Care: SAMHSA/CSAT’s Public Health Approach to Substance Use Problems & Disorders. Aligning Concepts, Practice, and Contexts to Promote Long Term Recovery: An Action Plan. 2008. from www.ireta.org
- Cleveland HH, Harris KS, Baker AK, Herbert R, Dean LR. Characteristics of a collegiate recovery community: maintaining recovery in an abstinence-hostile environment. *J Subst Abuse Treat.* 2007; 33(1):13–23. [PubMed: 17588485]
- Cleveland, HH.; Harris, KS.; Wiebe, R., editors. *Substance Abuse Recovery in College: Community Supported Abstinence.* Springer; New York: 2010.
- Cohen LR, Greenfield SF, Gordon S, Killeen T, Jiang H, Zhang Y, Hien D. Survey of eating disorder symptoms among women in treatment for substance abuse. *Am J Addict.* 2010; 19(3):245–251. [PubMed: 20525031]
- Czarlinski JA, Aase DM, Jason LA. Eating disorders, normative eating self-efficacy and body image self-efficacy: women in recovery homes. *Eur Eat Disord Rev.* 2012; 20(3):190–195. doi: 10.1002/erv.1116. [PubMed: 21751298]
- Dennis M, Godley SH, Diamond G, Tims FM, Babor T, Donaldson J, Funk R. The Cannabis Youth Treatment (CYT) Study: main findings from two randomized trials. *J Subst Abuse Treat.* 2004; 27(3):197–213. [PubMed: 15501373]
- Dennis M, Scott CK. Managing addiction as a chronic condition. *NIDA Addiction Science and Clinical Practice Perspectives.* 2007; 4(1):45–55.
- Dickard, N.; Downs, T.; Cavanaugh, D. Recovery/Relapse Prevention in Educational Settings For Youth With Substance Use & Co-occurring mental health disorders: 2010 Consultative Sessions Report. US department of Education, Office of Safe and Drug-Free Schools; Washington, DC: 2011. <http://www2.ed.gov/about/offices/list/osdfs/recoveryrpt.pdf>
- Doyle KS. The recovering college student: Factors influencing accommodation and service provision. *Dissertation Abstracts International Section A: Humanities & Social Sciences.* 1999; 60(6-A)
- Godley MD, Godley SH, Dennis ML, Funk R, Passetti LL. Preliminary outcomes from the assertive continuing care experiment for adolescents discharged from residential treatment. *J Subst Abuse Treat.* 2002; 23(1):21–32. [PubMed: 12127465]
- Godley SH, Garner BR, Passetti LL, Funk RR, Dennis ML, Godley MD. Adolescent outpatient treatment and continuing care: main findings from a randomized clinical trial. *Drug Alcohol Depend.* 2010; 110(1-2):44–54. [PubMed: 20219293]
- Harris K, Baker A, Kimball T, Shumway S. Achieving systems-Based sustained recovery: A comprehensive model for collegiate recovery communities. *Journal of Groups in Addiction and Recovery.* 2008; 2(2-4):220–237.
- Hingson RW, Zha W, Weitzman ER. Magnitude of and trends in alcohol-related mortality and morbidity among U.S. college students ages 18-24, 1998-2005. *J Stud Alcohol Drugs Suppl.* 2009; (16):12–20. [PubMed: 19538908]
- Hunt J, Eisenberg D, Kilbourne AM. Consequences of receipt of a psychiatric diagnosis for completion of college. *Psychiatr Serv.* 2010; 61(4):399–404. [PubMed: 20360280]
- Kaskutas L, Borkman T, Laudet A, Ritter L, Witbrodt J, Subbaraman M, Bond J. Elements that define recovery: The experiential perspective. *Journal of Studies on Alcohol and Drugs.* 2014; 75(6):999–1010. [PubMed: 25343658]
- Kimball, T. Update on number of operating CRPs. May 3. 2014 2014
- Laudet A. What does recovery mean to you? Lessons from the recovery experience for research and practice. *J Subst Abuse Treat.* 2007; 33(3):243–256. [PubMed: 17889296]
- Laudet, A. Longitudinal patterns and correlates of smoking in formerly polysubstance dependent individuals; Paper presented at the 71th Annual Scientific Meeting of the College on Problems of Drug Dependence (CPDD); Reno, NV. June 2009; 2009. 2009

- Laudet A. Rate and predictors of employment among formerly polysubstance dependent urban individuals in recovery. *Journal of Addictive Diseases*. 2012; 31(3):288–302. [PubMed: 22873190]
- Laudet A, Harris K, Kimball T, Winters KC, Moberg DP. Collegiate Recovery Communities Programs: What do we know and what do we need to know? *Journal of Social Work Practice in the Addictions*. 2014; 14:84–100. [PubMed: 24634609]
- Laudet, A.; Harris, K.; Winters, K.; Moberg, P.; Kimball, T. Nationwide survey of collegiate recovery programs: Is there a single model?; Paper presented at the 75th Annual Meeting - College on Problems of Drug Dependence; San Diego, CA. 2013;
- Laudet, A.; Harris, K.; Winters, K.; Moberg, P.; Kimball, T. Results from the first nationwide survey of students in Collegiate Recovery Programs; Paper presented at the 76th Annual Meeting - College on Problems of Drug Dependence; San Juan, PR. 2014;
- Laudet A, Stanick V, Sands B. The effect of onsite 12-step meetings on post-treatment outcomes among polysubstance-dependent outpatient clients. *Evaluation Review*. 2007; 31(6):613–646. [PubMed: 17986710]
- Laudet A, Timko C, Hill C. Comparing life experiences in active addiction and recovery between veterans and non-veterans: A national study. *Journal of Addictive Diseases*. 2014; 33(2):148–162. [PubMed: 24783976]
- Laudet AB, Magura S, Cleland CM, Vogel HS, Knight EL, Rosenblum A. The effect of 12-step based fellowship participation on abstinence among dually diagnosed persons: a two-year longitudinal study. *J Psychoactive Drugs*. 2004; 36(2):207–216. PMID: 1797895. [PubMed: 15369202]
- Lenk KM, Erickson DJ, Winters KC, Nelson TF, Toomey TL. Screening services for alcohol misuse and abuse at four-year colleges in the U.S. *J Subst Abuse Treat*. 2012; 43(3):352–358. [PubMed: 22377390]
- Magura S, Knight EL, Vogel HS, Mahmood D, Laudet AB, Rosenblum A. Mediators of effectiveness in dual-focus self-help groups. *Am J Drug Alcohol Abuse*. 2003; 29(2):301–322. PMID: 1828912. [PubMed: 12765208]
- McKay JR, Carise D, Dennis ML, Dupont R, Humphreys K, Kemp J, Schwartzlose J. Extending the benefits of addiction treatment: practical strategies for continuing care and recovery. *J Subst Abuse Treat*. 2009; 36(2):127–130. [PubMed: 19161893]
- Misch DA. On-campus programs to support college students in recovery. *J Am Coll Health*. 2009; 58(3):279–280. [PubMed: 19959442]
- Moberg DP, Finch AJ. Recovery High Schools: A Descriptive Study of School Programs and Students. *J Groups Addict Recover*. 2008; 2:128–161. [PubMed: 19165348]
- Moberg DP, Finch AJ, Lindsley S. Recovery High Schools: Students and Responsive Academic and Therapeutic Services. *Peabody Journal of Education*. 2014; 89(2):165–182. [PubMed: 24976659]
- Morgenstern J, Labouvie E, McCrady BS, Kahler CW, Frey RM. Affiliation with Alcoholics Anonymous after treatment: a study of its therapeutic effects and mechanisms of action. *J Consult Clin Psychol*. 1997; 65(5):768–777. [PubMed: 9337496]
- National Center on Addiction and Substance Abuse. Food for thought: Substance abuse and eating disorders. NYC; NY: 2003.
- Office of National Drug Control Policy. , editor. National Drug Control Strategy. Office of National Drug Control Policy; Washington, DC: 2010.
- Sheehan D, Lecrubier Y, Harnett-Sheehan K, Amorim P, Janavs J, Weiller E. The Mini International Neuropsychiatric Interview (M.I.N.I.): The Development and Validation of a Structured Diagnostic Psychiatric Interview. *Journal of Clinical Psychiatry*. 1998; 59:22–33. [PubMed: 9881538]
- Smock S, Baker A, Harris K, D'sauza C. The Role of Social Support in Collegiate Recovery Communities: A Review of the Literature. *Alcoholism Treatment Quarterly*. 2011; 29:35–44.
- Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: Volume I. Summary of National Findings. Office of Applied Studies; Rockville, MD: 2011.
- Substance Abuse and Mental Health Services Administration. , editor. The NSDUH Report: Trends in Substance Use, Dependence or Abuse, and Treatment among Adolescents: 2002 to 2007.

- Substance Abuse and Mental Health Services Administration, Office of Applied Studies; Rockville, MD: 2008.
- Substance Abuse and Mental Health Services Administration Office of Communications. , editor. Designing a Recovery-Oriented Care Model for Adolescents and Transition Age Youth with Substance Use or Cooccurring Mental Health Disorders. U.S. Department of Health and Human Services; Rockville, MD: 2009.
- Tanner-Smith EE, Wilson SJ, Lipsey MW. The comparative effectiveness of outpatient treatment for adolescent substance abuse: a meta-analysis. *J Subst Abuse Treat.* 2013; 44(2):145–158. [PubMed: 22763198]
- Toomey TL, Nelson TF, Winters KC, Miazga MJ, Lenk KM, Erickson DJ. Characterizing college systems for addressing student alcohol use: latent class analysis of U.S. four-year colleges. *J Stud Alcohol Drugs.* 2013; 74(5):777–786. [PubMed: 23948538]
- U.S. Dept. of Education. College Academic Performance and Alcohol and Other Drug Use. 2010. Retrieved October 26, 2010, from <http://www.higheredcenter.org/services/publications/college-academic-performance-and-alcohol-and-other-drug-use>
- Wechsler H, Nelson TF. What we have learned from the Harvard School Of Public Health College Alcohol Study: focusing attention on college student alcohol consumption and the environmental conditions that promote it. *J Stud Alcohol Drugs.* 2008; 69(4):481–490. [PubMed: 18612562]
- White, W., editor. Recovery Management and Recovery-oriented Systems of Care: Scientific Rationale and Promising Practices. Northeast Addiction Technology Transfer Center, Great Lakes Addiction Technology Transfer Center, Philadelphia Department of Behavioral Health & Mental Retardation Services; Pittsburgh, PA: 2008.
- White W, Finch A. The recovery school movement: Its history and future. *Counselor Magazine.* 2006; 7(2):54–57.
- Winters KC, Stinchfield RD, Lee S, Latimer WW. Interplay of psychosocial factors and the long-term course of adolescents with a substance use disorder. *Subst Abus.* 2008; 29(2):107–119. [PubMed: 19042330]
- Woodford M. Recovering college students' perspectives: investigating the phenomena of recovery from substance abuse among undergraduate students. *Dissertation Abstracts International Section A: Humanities & Social Sciences.* 2001; 62(7-A)

HIGHLIGHTS

- This is the first nationwide survey of college students in addiction recovery
- Students (N = 486) were participating in Collegiate Recovery Programs
- They report use of numerous substances, and a severe addiction history in spite of their young age
- Behavioral addictions, past and current,, were reported by a significant minority
- The smoking rate (40%) is twice the national CDC-reported average

Table 1
CRP Students' Demographics and Background (N = 486)

		% or Mean/SD
Gender	Female	42.8
Age		26.2 (8.19)
Ethny	Latino (yes)	5
Race	African american	1.9
	Caucasian	91.3
	Native American/Alaskan native	1.0
	Asian	1.0
	Native Hawaiian or Other Pacific Islander	0.4
	Other	4.3
Employment status	Part-time	45.9
	Full-time	14.0
	Not employed	40.1
Employment type (among employed; check all that apply)	Student project, research/teaching assistant	19.2
	Some other on campus job (e.g., clerical, maintenance)	22.0
	Off-campus job	67.0
Marital status	Married (including "common law")	5.7
	Widowed	0.4
	Separated	0.6
	Divorced	5.9
	Single (never married)	87.4
Has children (yes)		14.0
Ever served in the military (active or reserve)		11.6
Academic rank	Freshman	16.9
	Sophomore	18.4
	Junior	22.9
	Senior	28.9
	Graduate	12.8
Grade Point Average (GPA)		3.22 (0.621)
Course load	Full-time	13.8
	Part-time	1.4
	Other	
Residence	On-campus	25.2
	Off-campus	74.8
Lives in sober dorm or housing (yes)		32.4

	% or Mean/SD
Who lives with	
Living alone in room or apartment/house	25.3
Living with one or more roommate(s)	50.0
Living with spouse or partner	10.6
Living with family member(s) other than spouse or partner	7.9
Temporarily "double up" with family or friends or couch surfing	0.2
Other (please specify)	6.0
Any roommate in recovery from substance use problem (yes)	43.2
Any roommate using drugs/alcohol even in moderation (yes)	22.8
CRIMINAL JUSTICE INVOLVEMENT	
Current	
No involvement with legal system	85.9
On probation	8.8
On parole	0.8
On bail awaiting trial	0.8
Released on own recognizance, case pending	1.5
Other (please specify)	2.1
Ever arrested and charged as a juvenile or as an adult	
No, neither	41.9
Yes, as a juvenile	10.3
Yes, as an adult	34.3
Both as juvenile and adult	13.5
Ever in juvenile detention	6.4
Ever in jail or prison (as adult)	37.7

Table 2
CRP Students' physical and mental health (N = 486)

	% or Mean/SD
PHYSICAL HEALTH AND RELATED BEHAVIORS	
Health self-rating	
Poor	1.0
Fair	12.8
Good	57.7
Very good	28.5
Treated for chronic medical condition past year (yes)	18.8
Emergency department visit for medical problem or physical injury past year	24.1
Hospitalized for medical problem or physical injury past year (yes)	7.8
Smokes cigarette (yes)	40.0
Tested for Hep C	67.8
Hep C status: positive (among tested)	3.4
Tested for HIV	74.1
HIV serostatus: positive (among tested)	0.8
MENTAL HEALTH	
Ever treated for chronic mental health condition	76.1
Ever hospitalized for chronic mental health condition (among ever treated)	37.8
Ever diagnosed with a mental health disorder (among ever treated)	79.6
Treated for mental health disorder past year (among ever treated)	65.7
Currently taking medications for mental health disorder (among treated past year)	76.5
Mental Health self-rating	
Poor	1.3
Fair	13.9
Good	51.5
Very good	33.3
STRESS past month (1 - 10)	6.49 (2.08)

Table 3
CRP Students' Substance Use and Behavioral Addiction History and Current Status (N = 486)

PRIMARY AND SECONDARY ADDICTIONS	% or Mean/SD					
	Primary	Secondary				
Alcoholism	38.9	40.7				
Drug addiction	52.6	32.5				
Eating disorder	5.5	10.9				
Sex/love addiction	0.6	12.8				
Self harm/injury	0.4	7.8				
Gaming/gambling addiction	0.4	4.1				
Compulsive shopping	0.0	2.9				
Internet addiction (other than for sex, gambling or shopping)	0.2	2.5				
Exercise	0.0	4.1				
Other	1.3	--				
No secondary problem		10.3				
SUBSTANCES	Ever regularly	Primary	Other problematic*	Age started regular use	Years regular use	
Alcohol including alcohol-containing energy drinks	61.1	41.2	48.6	15.9 (2.58)	6.9 (5.8)	
Sedatives (e.g., barbiturates)	29.6	0.2	16.3	18.4 (9.9)	4.2 (3.9)	
Tranquilizers or anti-anxiety drugs	30.5	3.3	24.9	18.1 (5.0)	4.3 (3.6)	
Pain relievers (e.g., Codeine)	37.7	11.1	32.5	17.4 (4.2)	4.7 (4.7)	
Stimulants (e.g., uppers)	36.2	6.9	27.8	17.4 (4.2)	5.1 (5.2)	
Marijuana, hash, THC, or grass	74.5	10.0	42.6	15.3 (2.5)	6.4 (5.2)	
Cocaine or crack	33.1	11.7	29.4	18.9 (5.7)	4.9 (4.3)	
Hallucinogens (e.g., Ecstasy)	21.0	2.0	23.0	16.9 (3.2)	4.1 (4.2)	
Inhalants or Solvents (e.g., amyl nitrate)	6.4	0.2	7.4	17.2 (6.5)	3.9 (3.6)	
Heroin	15.0	11.9	10.9	19.5 (5.2)	3.3 (3.5)	
Performance enhancing drugs (e.g., steroids, HGH)	1.9	0.0	0.8	20.2 (4.8)	2.3 (2.1)	
Any OTHER medicines, drugs, or substances	10.7	1.5	7.8	19.6 (6.3)	3.7 (3.5)	
TIME SINCE LAST DRANK/USED DRUGS						
Among ever used regularly)						
Days since last drank alcohol	952 (962)					
Days since last used any drug	1053 (1196)					
Days since last used any drug or drank alcohol	975 (1073)					
Alcohol use past month	3.4					
Any drug past month	4.5					

	<u>% or Mean/SD</u>
Alcohol or drug use past month	5.4
PROBLEMATIC/COMPULSIVE BEHAVIOR PATTERNS Past 90 days	
Eating disorder	11.3
Sex/love addiction	11.7
Self harm/injury	5.3
Gaming/gambling addiction	5.1
Compulsive shopping	8
Internet addiction (other than for sex, gambling or shopping)	3.1
Exercise	2.9

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4
CRP Students' Utilization of Recovery Resources History (N = 486)

	<u>% or Mean/SD</u>
RECOVERY	
Considers self in recovery from a substance and/or behavior addiction or compulsion	87.6
In recovery from...	
Alcoholism	71.6
Drug addiction	72.6
Eating disorder	15.6
Sex/love addiction	9.5
Self harm/injury	10.5
Gaming/gambling addiction	2.5
Compulsive shopping	3.1
Internet addiction (other than for sex, gambling or shopping)	0.6
Exercise	3.3
Other (please specify)	2.9
UTILIZATION of RECOVERY RESOURCES	
Ever received addiction treatment	82.5
(IF YES) age first treatment	20.6 (5.0)
Treatment type (among ever treated)	
Detoxification (drug or alcohol)	49.5
Drug	10.6
Maintenance (e.g., Methadone Maintenance) (drugs only)	
Current among 'ever'	4.95
Therapeutic Community or other long-term residential (> 30 days)	65.7
Short-term residential (e.g., 21/28 day Inpatient Rehab)	55.1
'Drug free' Outpatient Treatment or Day Treatment (Non-Methadone)	60
Treatment in jail or prison (alcohol or drugs)	7.8
Ever in emergency room because of drugs or alcohol	43.2
Ever prescribed medication for substance use pb (e.g., naltrexone, acamprosate.	19.5
(IF YES)	9.9
Currently taking Rx medication for substance use pb	

	<u>% or Mean/SD</u>
Ever attended wilderness program (e.g., Outward Bound) to deal w/ substance use pb	
Ever attended recovery high school	4.9
Ever in individual counseling for substance use pb	51.8
(IF YES) Currently in individual counseling for substance use pb	35.1
TWELVE STEP FELLOWSHIPS AND OTHER SUPPORT GROUPS	
Ever attended a meeting (e.g., Alcoholics Anonymous, Narcotics Ano	93.3
(IF YES) age started 12 step attendance	21.3 (5.9)
Attended 12-step meeting past year (among ever attenders)	99.3
Attended 12-step meeting past month (among ever attenders)	89.8
12-step involvement past year (0 - 9; among ever attenders)	7.53 (2.15)
Ever attended non 12-step (e.g. Life Ring)	11.3
(IF YES) age started Non 12 step attendance	27.1 (9.7)
Attended 12-step meeting past year (among ever attenders)	73.1
Attended 12-step meeting past month (among ever attenders)	34.6