



An Overview of Nd:YAG Laser Capsulotomy

Eyyup Karahan ¹, Duygu Er ², Suleyman Kaynak ²

¹ Egepol Hospital, Department of Ophthalmology, Izmir, Turkey

² Dokuz Eylul University, Department of Ophthalmology, Izmir, Turkey

ABSTRACT

It has been revealed that posterior capsule opacification (PCO) is the most common delayed complication of cataract surgery. On the other hand, Nd:YAG laser capsulotomy is accepted as standard treatment for PCO. Although, Nd:YAG laser capsulotomy is a noninvasive and safe treatment it carries risk of some complications. Using less total energy and performing smaller capsulotomies are effective choices to decrease complications after Nd:YAG capsulotomy. The purpose of this review is to look through the complications associated with Nd:YAG laser capsulotomy, and the effect of capsulotomy size and used total energy on such complications.

KEY WORDS

Nd:YAG Laser; Capsulotomy; Posterior capsule opacification (PCO)

©2014, Med Hypothesis Discov Innov Ophthalmol.

This is an open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0), which allows users to read, copy, distribute and make derivative works for non-commercial purposes from the material, as long as the author of the original work is cited properly.

Correspondence to:

Dr. Eyyup Karahan, Narlı Mah Ecem Sok 42C/7, Narlıdere, İzmir, Turkey; Tel: +902322788111; Fax: + 902322786804E-Mail: karahaneyup@yahoo.com

INTRODUCTION

Posterior capsule opacification (PCO) is the most common delayed complication of cataract surgery (1). The incidence of PCO was reported to be 20.7% at two years and 28.5% at 5 years after cataract surgery (2). In the early 1980s, the application of Neodymium:yttrium-aluminum-garnet (Nd:YAG) laser capsulotomy as treatment for PCO was presented by Aron-Rosa (3) and Fankhauser (4). Nd:YAG laser capsulotomy showed itself to be an effective alternative to surgical dissection, avoiding such complications as endophthalmitis and vitreous loss (5,6).

Improvement in visual acuity after Nd:YAG laser capsulotomy in patients with significant PCO has been

well documented (7–9). Improvements in glare and contrast sensitivity may also be important outcome measures for many patients (10–12). Although Nd:YAG laser capsulotomy is accepted as standard treatment for PCO and has been found to be safe and effective, it is not without complications, some of which can be sight-threatening such as retinal edema and detachment (13). Several studies have described damages in the intraocular lens (IOL), increased intraocular pressure (IOP), glaucoma, retinal hemorrhage, iritis, vitreous prolapse, corneal injury, vitritis, pupil blockage, hyphema, cystoid macular edema, retinal detachment (RD), IOL dislocation or exacerbation of endophthalmitis (3,6,14–16).



Some recent studies have been concentrated to the influence of capsulotomy size and total energy level on complications after Nd:YAG laser capsulotomy (17–19).

The purpose of this review is to look through the complications associated with Nd:YAG laser capsulotomy, and the effect of capsulotomy size and used total energy on such complications.

COMPLICATIONS ASSOCIATED WITH ND:YAG LASER CAPSULOTOMY

IOL Movement and Refractive Changes

There are several reports of displaced IOLs after laser treatment (21–24). Levy et al. reported two instances of hydrogel implant dislocation into vitreous following Nd:YAG laser capsulotomy (21). Using dual-beam partial coherence interferometry the procedure of Nd:YAG laser capsulotomy has been shown to induce a small but measurable backward movement of the IOL. They stated that the larger capsulotomy openings induce greater backward movement, and recommend small openings to avoid this complication. No significant refractive change was reported in this study (25). However, Thornval and Naeser (26) failed to observe this effect. In our recently reported study we found that the hyperopic shift was higher in patients with capsulotomy size larger than 3.9 mm when compared with patients with smaller capsulotomy sizes. The hyperopic shift was progressive until 4 weeks in larger capsulotomy group. We recommended to prescribe new spectacles at least 1 week or 4 weeks if the capsulotomy is large after Nd:YAG laser capsulotomy (17). Zaidi et al., were also recorded a significant hyperopic shift which was especially important 1 week after Nd:YAG laser capsulotomy (27). Also the magnitude of shift can be affected by IOL style. The hyperopic shift was found higher with plate haptic implants than with polymethyl methacrylate and three piece foldable lenses (25).

IOL Damage/Pitting

Hassan KS et al. has noted IOL pitting 19.8% in a study of 86 eyes (28) and Haris WS noted 11.7% significant marks on IOL during laser capsulotomy in 342 eyes (29). Khanzada et al. (30) reported the range of 9.4% (30 eyes)

in 320 eyes. The retro-focusing of laser aiming beam can reduce the risk of IOL damage.

Iritis/Uveitis

Keates et al. found iritis persisting in 0.4% and vitritis persisting in 0.7% after a 6-month postoperative period (13). Chambless, in a study with an average follow-up period of 7 months, found persistent anterior uveitis in 1.4% of the patients (6). Gore et al. reported that 33.5% of patients had iritis after Nd:YAG laser capsulotomy manifested as cells and flare in the anterior chamber on slit lamp examination. They were given topical steroid, and reaction had subsided leaving no delayed complication (31). In summary, transient anterior chamber flare may be seen post-laser treatment; persistent iritis or vitritis is rare.

Rise of Intraocular Pressure

The most common complication of posterior capsulotomy is increased IOP. Different explanations which have been given for the pressure rise following Nd:YAG laser treatment include the deposition of debris in the trabecular meshwork, (32,33) pupillary block, (34,35) and inflammatory swelling of the ciliary body or iris root associated with angle closure (36). Despite the prophylactic treatment, increased IOP was reported in 15% to 30% of patients in several studies (37,38). Keates et al. (13) found elevation of IOP in 0, 6% of his patients, whereas Stark et al. (8) reported that the elevation of IOP was 1.0% after Nd:YAG capsulotomy. Ge et al. (39) found that the rise in IOP was more pronounced in patients with glaucoma in those who experienced a higher rise of IOP within hour of capsulotomy. However, Shani et al. (40) could not find any elevation of IOP and postulated that healthy pseudophakic eyes do not show elevation of IOP after Nd:YAG laser capsulotomy. Ficker et al. (31) noted 13 patients to have IOP over 23 mmHg and 9 patients to have IOP between 30–48 mmHg, within 2–3 hours after laser capsulotomy. In this group of 24 patients there was a tendency for IOP to rise when higher pulse energies were used, particularly when these exceeded 1.5 mg, and the raised IOP was controlled with antiglaucoma therapy. Ari et al. (20) underlined that the severity and duration of increased IOP and macular



thickness were less when a total energy level less than 80 mg is used.

In our study, one patient (2.7%) in small size group and three patients (9.3%) in larger capsulotomy size group had mild elevation of IOP one week after Nd: YAG laser capsulotomy. Rise in IOP was higher than previous studies. Previous studies did not give any information about the capsulotomy size. So, a comparison of rises in IOP with previous studies is not appropriate. More capsule particles released with larger capsulotomies might be the reason of higher rates of elevation in larger capsulotomy group (17).

Cystoid Macular Edema

Cystoid macular edema (CME) occurs after intraocular surgical procedures, trauma, and a variety of other inflammatory conditions affecting the retina. The etiology of CME following Nd: YAG laser capsulotomy most likely involves movement of the vitreous cavity and vitreous damage, which results in the release of inflammatory mediators. Vitreoretinal traction caused by the procedure may also play a part (41).

Previous studies have investigated changes of macular thickness after Nd: YAG laser capsulotomy. Although some of the previous studies have reported CME, many of them found no significant changes in macular thickness following Nd: YAG laser capsulotomy (42–47).

Lewis et al. found a low rate of CME when capsulotomy was delayed for over 6 months from the initial IOL implant date (48). Ari et al. (18) evaluated how different energy levels of Nd: YAG laser capsulotomy affect macular thickness. They divided patients into two groups based on the energy levels used in Nd: YAG laser capsulotomy. They found that both groups had increased macular thickness compared to preoperative levels; macular thickness measurements of the patients treated with high energy levels were significantly greater compared to low energy levels. In another study a series of 897 Nd: YAG laser capsulotomies were reviewed for the complications of CME. After Nd: YAG laser capsulotomy, 11 patients developed CME. The numbers of laser pulses and energy delivered were not risk factors (19).

In our study, energy levels were similar in both small size and large size capsulotomy groups. Comparison of two

groups with respect to macular thickness did not reveal any difference preoperatively or 1 week, 4 weeks or 12 weeks postoperatively. There was a significant thickening in macular thickness at 1 week in both groups; this difference was not statistically significant between groups. The mean macular thicknesses were decreased to preoperative levels at 4 and 12 weeks measurements (17).

Retinal Tear and Detachment

The risk for RD after Nd: YAG laser capsulotomy is estimated to be 4-fold that of the risk after uneventful surgery without a capsulotomy (49,50). Raza (51) reported 11 patients (2%) of RD after Nd: YAG laser capsulotomy. Steinert et al. (19) reported that eight patients of 897 patients treated with Nd: YAG laser capsulotomy developed RD.

Retrospective analysis of data based on Medicare claims in the US suggests that Nd:YAG laser capsulotomy is associated with a significantly elevated risk for RD, stronger associations were found for a history of RD or lattice degeneration, an axial length greater than 24.0 mm, and posterior capsule rupture during surgery (49). Several other retrospective studies confirm the higher risk of RD after capsulotomy in eyes with intraoperative complications, axial myopia, and vitreoretinal pathology (52–55); however, 2 studies show no association in the absence of these risk factors (56,57).

The precise mechanisms that lead to retinal breaks and RD after Nd:YAG laser capsulotomy are not known, Sheard et al. designed a study to determine whether RD after Nd:YAG laser capsulotomy is due to a greater incidence of posterior vitreous detachment (PVD) than in controls and whether vitreous status at the time of capsulotomy is useful in predicting the risk for RD. The prevalence of PVD was significantly higher in eyes after extra-capsular cataract extraction and IOL implantation than in Phakic eyes independent of Nd: YAG laser capsulotomy. Capsulotomy was not associated with a significantly greater incidence of new PVD, they concluded that the presence or absence of PVD at the time of capsulotomy is not helpful in assessing the risk for RD in the first year after laser treatment (58).

Other Complications



Pupillary block glaucoma (8) as well as aqueous misdirection syndrome, (59) macular hole, (6) retinal hemorrhage, (8) spreading of endocapsular low-grade endophthalmitis, (60) and secondary closure of capsulotomy aperture (61) are other complications that have been reported in isolation.

DISCUSSION/CONCLUSION

Nd: YAG laser capsulotomy is fast and noninvasive procedure with immediate improvement. Although, it is noninvasive and considered safer than surgical approach it carries the risk of some complications. Some recent studies including our study observed the effects of capsulotomy size and laser energy levels on postcasulotomy complications (17–19).

Capsulotomy size is important as patients subjected to lower amounts of laser energy for perhaps a smaller capsulotomy may benefit from fewer complications of RD, IOP rise, (62,63) and perhaps to less CME (64). Risk of IOL dislocation may be significantly less, especially with plate haptic silicone IOLs. In our study, despite energy levels were similar in small and large capsulotomy groups hyperopic shift, IOP rise and increased macular thickness was found to be lesser in patients with a smaller capsulotomy size.

Another parameter that is believed to be important is laser energy level. Ari et al. reported that IOP rise and rise in macular thickness were higher with higher energy levels. However, Steiner et al. reported that the numbers of laser pulses and energy delivered, not risk factors for the development of cystoid macular edema.

In conclusion, some complications especially rise in IOP and macular thickness seems to be unavoidable after Nd: YAG laser capsulotomy. Using less total energy and performing smaller capsulotomies are practical choices to decrease complications after Nd: YAG capsulotomy.

DISCLOSURE

Conflicts of Interest: None declared.

REFERENCES

1. Wormstone IM. Posterior capsule opacification: a cell biological perspective. *Exp Eye Res* 2002;74:337-347.

2. Nakazawa M, Ohtsuki K. Apparent accommodation in pseudophakic eyes after implantation of posterior chamber intraocular lenses. *Am J Ophthalmol.* 1983 Oct;96(4):435-8. PMID: 6624824

3. Aron-Rosa D, Aron JJ, Griesemann M, Thyzel R: Use of the neodymium-YAG laser to open the posterior capsule after lens implant surgery: a preliminary report. *J Am Intraocul Implant Soc.* 1980 Oct;6(4):352-4. PMID: 7440377

4. Fankhauser F, Roussel P, Steffen J: Clinical studies on the efficiency of high power laser radiation upon some structures of the anterior segment of the eye. First experiences of the treatment of some pathological conditions of the anterior segment of the human eye by means of a Qswitched. *Int Ophthalmol.* 1981 May;3(3):129-39. PMID: 7196390

5. Shah GR, Gills JP, Durham DG, Ausmus WH. Three thousand YAG lasers in posterior capsulotomies: an analysis of complications and comparison to polishing and surgical discission. *Ophthalmic Surg.* 1986 Aug;17(8):473-7. PMID: 3748538

6. Chambless WS. Neodymium: YAG laser posterior capsulotomy results and complications. *J Am Intraocul Implant Soc.* 1985 Jan;11(1):31-2. PMID: 3838167

7. Gardner KM, Straatsma BR, Pettit TH. Neodymium:YAG laser posterior capsulotomy: the first 100 cases at UCLA. *Ophthalmic Surg.* 1985 Jan;16(1):24-8. PMID: 3838376

8. Stark WJ, Worthen D, Holladay JT, Murray G. Neodymium:YAG lasers. An FDA report. *Ophthalmology.* 1985 Feb;92(2):209-12. PMID: 3982799

9. Wasserman EL, Axt JC, Sheets JH. Neodymium:YAG laser posterior capsulotomy. *J Am Intraocul Implant Soc.* 1985 May;11(3):245-8. PMID: 4008310

10. Magno BV, Datiles MB, Lasa MS, Fajardo MR, Caruso RC, Kaiser-Kupfer MI. Evaluation of visualfunction following neodymium:YAG laser posterior capsulotomy. *Ophthalmology.* 1997 Aug;104(8):1287-93. PMID: 9261315

11. Sunderraj P, Villada JR, Joyce PW, Watson A. Glare testing in pseudophakes with posterior capsule opacification. *Eye (Lond).* 1992;6 (Pt 4):411-3. PMID: 1478316

12. Tan JC, Spalton DJ, Arden GB. The effect of neodymium: YAG capsulotomy on contrast sensitivity and the evaluation of methods for its assessment. *Ophthalmology.* 1999 Apr;106(4):703-9. PMID: 10201590

13. Keates RH, Steinert RF, Puliafito CA, Maxwell SK. Long-term follow-up of Nd:YAG laser posterior capsulotomy. *J Am Intraocul Implant Soc.* 1984 Spring;10(2):164-8. PMID: 6547424

14. Ambler JS, Constable IJ. Retinal detachment following capsulotomy. *Aust N Z J Ophthalmol.* 1988 Nov;16(4):337-41. PMID: 3248183

15. Bath PE, Fankhauser F. Long-term results of Nd:YAG laser posterior capsulotomy with the Swiss laser. *J Cataract Refract Surg.* 1986 Mar;12(2):150-3. PMID: 3754577

16. Billotte C, Berdeaux G. Adverse clinical consequences of neodymium:YAG laser treatment of posterior capsule opacification. *J Cataract Refract Surg.* 2004 Oct;30(10):2064-71. PMID: 15474815

17. Karahan E, Tuncer I, Zengin MO. The Effect of ND:YAG Laser Posterior Capsulotomy Size on Refraction, Intraocular Pressure, and



- Macular Thickness. *J Ophthalmol.* 2014;2014:846385. doi: 10.1155/2014/846385. PMID: 24724016
18. Ari S, Cingu AK, Sahin A, inar YC, and Caca I. The effects of Nd:YAG laser posterior capsulotomy on macular thickness, intraocular pressure, and visual acuity. *Ophthalmic Surg Lasers Imaging.* 2012 Sep-Oct;43(5):395-400. doi: 10.3928/15428877-20120705-03. PMID: 22785102
19. Steinert RF, Puliafito CA, Kumar SR, Dudak SD, Patel S. Cystoid macular edema, retinal detachment, and glaucoma after Nd:YAG laser posterior capsulotomy *Am J Ophthalmol.* 1991 Oct 15;112(4):373-80. PMID: 1928237
20. Framme C, Hoerauf H, Roider J, Laqua H: Delayed intraocular lens dislocation after neodymium:YAG capsulotomy. *J Cataract Refract Surg.* 1998 Nov;24(11):1541-3. PMID: 9818350
21. Levy JH, Pisacano AM, Anello RD: Displacement of bagplaced hydrogel lenses into the vitreous following neodymium: YAG laser capsulotomy. *J Cataract Refract Surg.* 1990 Sep;16(5):563-6. PMID: 2231370
22. Maguire AM, Blumenkranz MS, Ward TG, Winkelman JZ: Scleral loop fixation for posteriorly dislocated intraocular lenses. *Arch Ophthalmol.* 1991 Dec;109(12):1754-8. PMID: 1841589
23. Schneiderman TE, Johnson MW, Smiddy WE: Surgical management of posteriorly dislocated silicone plate haptic intraocular lenses. *Am J Ophthalmol.* 1997 May;123(5):629-35. PMID: 9152068
24. Shakin EP, Carty JB Jr.: Clinical management of posterior chamber intraocular lens implants dislocated in the vitreous cavity. *Ophthalmic Surg Lasers.* 1995 Nov-Dec;26(6):529-34. PMID: 8746574
25. Findl O, Drexler W, Menapace R et al. Changes in intraocular lens position after neodymium: YAG capsulotomy. *Ophthalmic Surg Lasers.* 1995 Nov-Dec;26(6):529-34. PMID: 8746574
26. Thornval P, Naeser K. Refraction and anterior chamber depth before and after neodymium:YAG laser treatment for posterior capsule opacification in pseudophakic eyes: a prospective study. *J Cataract Refract Surg.* 1995 Jul;21(4):457-60. PMID: 8523294
27. Zaidi M, Askari NS. Effect of Nd:YAG laser posterior capsulotomy on anterior chamber depth, Intraocular pressure, and refractive status. *J Cataract Refract Surg.* 2000 Aug;26(8):1183-9. PMID: 11008046
28. Hasan KS, Adhi MI, Aziz M, et al. Nd:YAG Laser Posterior Capsulotomy. *Pak J Ophthalmol.* 1996;12: 3-7.
28. Harris WS, Herman WK, Fagadau WR. Management of the posterior capsule before and after the YAG laser. *Trans Ophthalmol Soc U K.* 1985;104 (Pt 5):533-5. PMID: 3863342
29. Ficker LA, Steel AD. Complications of Nd: YAG laser posterior capsulotomy. *Trans Ophthalmol Soc U K.* 1985;104 (Pt 5):529-32. PMID: 3863341
30. Mahtab Alam Khanzada, Shafi Muhammad Jatoy, Ashok Kumar Narsani, Syed Asher Dabir, Siddiqa Gul. Is the Nd: YAG Laser a Safe Procedure for Posterior Capsulotomy?. *Pak J Ophthalmol* 2008;24:73-78.
31. Gore VS. The study of complications of Nd:YAG laser capsulotomy. *Klin Monbl Augenheilkd.* 1994 May;204(5):286-7. PMID: 8051851
32. Kraff MC, Sanders DR, Lieberman HL. Intraocular pressure and the corneal endothelium after neodymium-YAG laser posterior capsulotomy. Relative effects of aphakia and pseudophakia. *Arch Ophthalmol.* 1985 Apr;103(4):511-4. PMID: 3985828
- 33 Vine AK. Ocular hypertension following Nd-YAG Laser Capsulotomy: A potentially blinding complication. *Ophthalmic Surg.* 1984 Apr;15(4):283-4. PMID: 6547221
34. Parker MD, Clofeine GS, Stocklin RD. Marked intraocular pressure rise following Nd-YAG laser capsulotomy. *Ophthalmic Surg.* 1984 Feb;15(2):103-4. PMID: 6546789
35. Ruderman JM, Mitchell PG, Kraff M. Pupillary block following Nd-YAG laser capsulotomy. *Ophthalmic Surg.* 1983 May;14(5):418-9. PMID: 6877743
36. MacEwen CJ, Dutton GN, Holding D. Angle closure following Neodymium-YAG (Nd-YAG) laser capsulotomy in the Aphakic Eye. *Br J Ophthalmol.* 1985 Oct;69(10):795-6. PMID: 3840385
37. Minello AAP, Prata JA, de Arruda Mello PA. Efficacy of topic ocular hipotensive agents after posterior capsulotomy. *Arquivos Brasileiros de Oftalmologia.* 2008;5:706-710.
38. Lin JC, Katz LJ, Spaeth GL, Klancnik JM. Intraocular pressure control after Nd: YAG laser posterior capsulotomy in eyes with glaucoma. *Arq Bras Oftalmol.* 2008 Sep-Oct;71(5):706-10. PMID: 19039468
39. Ge J, Wand M, Chiang R, Paranhos A, Shields B. Long term effect of Nd : YAG laser posterior capsulotomy on intraocular pressure. *Arch Ophthalmol.* 2000 Oct;118(10):1334-7. PMID: 11030814
40. Shani L, David R, Tessler Z, Rosen S, Schneck M, Yassur Y. Intraocular pressure after neodymium: YAG laser treatments in the anterior segment *J Cataract Refract Surg.* 1994 Jul;20(4):455-8. PMID: 7932138
41. Murrill CA, Stanfield DL, Van Brocklin MD: Capsulotomy. *Optom Clin.* 1995;4(4):69-83. PMID: 7488799
42. Henderson BA, Kim JY, Ament CS, et al. Clinical pseudophakic cystoid macular edema: Risk factors for development and duration after treatment. *J Cataract Refract Surg.* 2007 Sep;33(9):1550-8. PMID: 17720069
43. Bukelman A, Abrahami S, Oliver M, Pollack A. Cystoid macular oedema following neodymium:YAG laser capsulotomy: A prospective study. *Eye (Lond).* 1992;6 (Pt 1):35-8. PMID: 1426396
44. Albert DW, Wade EC, Parrish 2nd RK, et al. A prospective study of angiographic cystoid macular edema one year after Nd: YAG posterior capsulotomy. *Ann Ophthalmol.* 1990 Apr;22(4):139-43. PMID: 2350122
45. Altiparmak UE, Ersoz I, Hazirolan D, et al. The impact of Nd:YAG capsulotomy on foveal thickness measurement by optical coherence tomography. *Ophthalmic Surg Lasers Imaging.* 2010 Jan-Feb;41(1):67-71. PMID: 20143510
46. Kara N, Yazici AT, Bozkurt E, et al. Which procedure has more effect on macular thickness: Primary posterior continuous capsulorhexis (PPCC) combined with phacoemulsification or Nd:YAG laser capsulotomy? *Int Ophthalmol.* 2011 Aug;31(4):303-7. doi: 10.1007/s10792-011-9461-9. PMID: 21842401
47. Giocanti-Aure'gan A, Tilleul J, Rohart C, et al. OCT measurement of the impact of Nd:YAG laser capsulotomy on foveal thickness. *J Fr*



- Ophthalmol. 2011 Nov;34(9):641-6. doi: 10.1016/j.jfo.2011.02.020. PMID: 21889816
48. Lewis H, Singer TR, Hanscom TA, Straatsma BR: A prospective study of cystoid macular edema after neodymium: YAG laser posterior capsulotomy. *Ophthalmology*. 1987 May;94(5):478-82. PMID: 3601362
49. Tielsch JM, Legro MW, Cassard SD, et al. Risk factors for retinal detachment after cataract surgery. A population-based case-control study. *Ophthalmology*. 1996 Oct;103(10):1537-45. PMID: 8874424
50. Javitt JC, Tielsch JM, Canner JK, et al. National out-comes of cataract extraction;increased risk of retinal complications associated with Nd:YAG laser capsulo- nal detachment. *Ophthalmology*. 1992 Oct;99(10):1487-97. PMID: 1454313
51. Raza A. Complications after Nd:Yag posterior capsulotomy. *Journal of Rawalpindi Medical College*. 2007;11:27–29.
52. Olsen GM, Olson RJ. Prospective study of cataract surgery, capsulotomy, and retinal detachment. *J Cataract Refract Surg*. 1995 Mar;21(2):136-9. PMID: 7791052
53. Koch DD, Liu JF, Gill EP, Parke DW II. Axial myopia increases the risk of retinal complications after neodymium:YAG posterior capsulotomy. *Arch Ophthalmol*. 1989 Jul;107(7):986-90. PMID: 2751470
54. Rickman-Barger L, Florine CW, Larson RS, Lindstrom RL. Retinal detachment after neodymium:YAG laser posterior capsulotomy. *Am J Ophthalmol*. 1989 May 15;107(5):531-6. PMID: 2712134
55. Dardenne MU, Gerten G-J, Kokkas K, Kermani O. Retrospective study of retinal detachment following neodymium:YAG laser posterior capsulotomy. *J Cataract Refract Surg*. 1989 Nov;15(6):676-80. PMID: 2614712
56. Nielsen NE, Naeser K. Epidemiology of retinal detachment following extracapsular cataract extraction: a follow-up study with an analysis of risk factors. *J Cataract Refract Surg*. 1993 Nov;19(6):675-80. PMID: 8271160
57. Powell SK, Olson RJ. Incidence of retinal detachment after cataract surgery and neodymium:YAG laser capsulotomy. *J Cataract Refract Surg*. 1995 Mar;21(2):132-5. PMID: 7791051
58. Sheard RM, Goodburn SF, Comer MB, Scott JD, Snead MP. Posterior vitreous detachment after neodymium:YAG laser posterior capsulotomy. *J Cataract Refract Surg*. 2003 May;29(5):930-4. PMID: 12781278
59. Mastropasqua L, Ciancaglini M, Carpineto P: Aqueous misdirection syndrome: a complication of neodymium: YAG posterior capsulotomy. *J Cataract Refract Surg*. 1994 Sep;20(5):563-5. PMID: 7996414
60. Carlson AN, Koch DD: Endophthalmitis following Nd:YAG laser posterior capsulotomy. *Ophthalmic Surg*. 1988 Mar;19(3):168-70. PMID: 3258419
61. Jones NP, McLeod D, Boulton ME: Massive proliferation of lens epithelial remnants after Nd-YAG laser capsulotomy. *Br J Ophthalmol*. 1995 Mar;79(3):261-3. PMID: 7703206
62. Aron-Rosa DS: Influence of picosecond and nanosecond YAG laser capsulotomy on intraocular pressure. *J Am Intraocul Implant Soc*. 1985 May;11(3):249-52. PMID: 4008311
63. Channell MM, Beckman H: Intraocular pressure changes after neodymium-YAG laser posterior capsulotomy. *Arch Ophthalmol*. 1984 Jul;102(7):1024-6. PMID: 6547596
64. Smith RT, Moscoso WE, Trokel S, Auran J: The barrier function in neodymium-YAG laser capsulotomy. *Arch Ophthalmol*. 1995 May;113(5):645-52. PMID: 7748137