

Prevalence of congenital muscular dystrophy in Italy

A population study

Alessandra Graziano, MD*
Flaviana Bianco, MD*
Adele D'Amico, MD
Isabella Moroni, MD
Sonia Messina, MD
Claudio Bruno, MD
Elena Pegoraro, MD
Marina Mora, MD
Guja Astrea, MD
Francesca Magri, MD
Giacomo P. Comi, MD
Angela Berardinelli, MD
Maurizio Moggio, MD
Lucia Morandi, MD
Antonella Pini, MD
Roberta Petillo, MD
Giorgio Tasca, MD
Mauro Monforte, MD
Carlo Minetti, MD
Tiziana Mongini, MD
Enzo Ricci, MD
Ksenija Gorni, MD
Roberta Battini, MD
Marcello Villanova, MD
Luisa Politano, MD
Francesca Gualandi, MD
Alessandra Ferlini, MD
Francesco Muntoni, MD
Filippo Maria Santorelli, MD
Enrico Bertini, MD
Marika Pane, MD
Eugenio Mercuri, MD

Correspondence to
Dr. Mercuri:
mercuri@rm.unicatt.it

ABSTRACT

Objective: We provide a nationwide population study of patients with congenital muscular dystrophy in Italy.

Methods: Cases were ascertained from the databases in all the tertiary referral centers for pediatric neuromuscular disorders and from all the genetic diagnostic centers in which diagnostic tests for these forms are performed.

Results: The study includes 336 patients with a point prevalence of 0.563 per 100,000. Mutations were identified in 220 of the 336 (65.5%). The cohort was subdivided into diagnostic categories based on the most recent classifications on congenital muscular dystrophies. The most common forms were those with α -dystroglycan glycosylation deficiency (40.18%) followed by those with laminin $\alpha 2$ deficiency (24.11%) and collagen VI deficiency (20.24%). The forms of congenital muscular dystrophy related to mutations in *SEPN1* and *LMNA* were less frequent (6.25% and 5.95%, respectively).

Conclusions: Our study provides for the first time comprehensive epidemiologic information and point prevalence figures for each of the major diagnostic categories on a large cohort of congenital muscular dystrophies. The study also reflects the diagnostic progress in this field with an accurate classification of the cases according to the most recent gene discoveries. *Neurology*® 2015;84:904-911

GLOSSARY

α DG = α -dystroglycan; CK = creatine kinase; CMD = congenital muscular dystrophy; MEB = muscle-eye-brain disease; RSMD1 = rigid spine muscular dystrophy type 1.

The term congenital muscular dystrophy (CMD) classically includes a group of genetically, clinically, and biochemically distinct entities sharing clinical and pathologic features such as early presentation of weakness and hypotonia and dystrophic features on muscle biopsy.¹

The rapidly increasing identification of several genes responsible for different forms of CMD has dramatically expanded the spectrum of the known forms, allowing a better understanding of the individual forms and different underlying pathomechanisms.²

The incidence and prevalence of CMD in populations is not sufficiently known with only a few studies reporting incidence or point prevalence in relatively small regions.³⁻⁵ Furthermore, several new genes have only been described in the last few years and the subtyping and classification in those reports is therefore not updated.

*These authors contributed equally to this work.

From the Departments of Paediatrics and Neurology (A.G., F.B., G.T., M. Monforte, E.R., M.P., E.M.), Catholic University, Rome; Unit of Neuromuscular and Neurodegenerative Disorders (A.D., G.T., E.B.), Department of Neurosciences, Bambino Gesù Children's Hospital, Rome; Pediatric Neurology and Neuromuscular Disease and Immunology Unit (I.M., M. Mora, L.M.), Istituto Neurologico Besta, Milan; Department of Neurosciences, Psychiatry and Anaesthesiology (S.M.), University of Messina; Neuromuscular Disease Unit (C.B., C.M.), G. Gaslini Institute, Genoa; Department of Neurosciences (E.P.), University of Padua, Italy; Department of Developmental Neuroscience and Molecular Medicine Neuromuscular Unit (G.A., R.B., F.M.S.), Stella Maris Institute, Pisa; Department of Neurological Sciences (F. Magri, G.P.C.), IRCCS Ospedale Maggiore Policlinico, University of Milan; Child Neurology and Psychiatry Unit (A.B.), IRCCS C. Mondino Foundation; Neuromuscular and Rare Diseases Unit (M. Moggio), IRCCS Foundation Ca' Granda Ospedale Maggiore Policlinico, Dino Ferrari Center, Milano; Child Neurology and Psychiatry Unit (A.P.), IRCCS Istituto delle Scienze Neurologiche di Bologna, Bellaria Hospital, Bologna; Cardiomiologia e Genetica Medica (R.P., L.P.), Dipartimento di Medicina Sperimentale Seconda Università di Napoli; Neuromuscular Center (T.M.), S.G. Battista Hospital, University of Turin; Centro Nemo (K.G.), Milan; Neuromuscular Unit (M.V.), Nigrisoli Hospital, Bologna; Section of Medical Genetics (F.G., A.F.), Department of Experimental and Diagnostic Medicine, Ferrara, Italy; and Dubowitz Neuromuscular Centre (F. Muntoni), UCL Institute of Child Health & Great Ormond Street Hospital for Children, London, UK.

Go to Neurology.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

In this study, we report the results of a detailed population study and the frequency of the various forms of CMD in all the patients with CMD referred to all the tertiary care centers for pediatric neuromuscular disorders in Italy.

METHODS Our inclusion criteria were all patients with a diagnosis of CMD currently seen in the participating centers. The diagnosis of CMD was based on a combination of clinical and biopsy findings according to the recently proposed diagnostic criteria for CMD.⁶ This included clinical signs suggestive of CMD and/or dystrophic or myopathic features on muscle biopsy, with exclusion of other identifiable neuromuscular disorders. Clinical signs included weakness and hypotonia or contractures at birth or in the first months, but as in some CMD variants (e.g., Ullrich CMD), the onset can be delayed.⁶⁻⁸

All patients with a potential neuromuscular disorder, with weakness and/or increased creatine kinase (CK) serum levels, are referred to a tertiary care center for further testing. Because all the Italian tertiary care centers for pediatric and adult neuromuscular disorders participated in this study, we are therefore likely to have included all the patients who have undergone a diagnostic process for CMD. Cases were ascertained from the databases in each center, only including patients who were under current follow-up. To exclude that patients may have been referred by other physicians, we also checked all the molecular genetic centers performing analysis of the CMD genes. Other sources available for this study included the Collagen VI Database held at the University of Ferrara and the CMD Dystroglycan Database, a nationwide database used for our previous study^{9,10} and that is constantly updated.

Disease groups were established using the most recent CMD classifications² (table 1):

1. Forms of CMD due to mutations in genes encoding for structural proteins of the basal lamina or extracellular matrix or receptors for extracellular matrix proteins. This category includes mutations in the collagen 6 genes, laminin $\alpha 2$ (merosin), and integrin $\alpha 7$ and $\alpha 9$.
2. Forms secondary to genes encoding for putative or demonstrated glycosyltransferases, which affect the glycosylation of α -dystroglycan (α DG). This includes Fukuyama CMD, muscle-eye-brain disease (MEB), and Walker-Warburg syndrome, and other phenotypes.
3. Forms due to mutations in genes encoding for nuclear envelope proteins (LMNA and nesprin).
4. Forms due to abnormalities of proteins with, so far, unknown function localized in the endoplasmic reticulum, which include the form of CMD with rigid spine syndrome secondary to mutations in *SEPN1*.

For each group, and for the whole cohort, point prevalence was established. The prevalence day was January 31, 2013. The 95% confidence interval was measured using the Wilson score interval with continuity correction.

Standard protocol approvals, registrations, and patient consents. The ethical committees of the participating centers approved the study.

RESULTS Three hundred thirty-six subjects fulfilled the inclusion criteria (191 male, 145 female). All were followed by one of the participating clinical centers who were also responsible for referring them for the

genetic diagnosis with the exception of 2 boys with Ullrich CMD referred to the genetic labs by a specialist in connective tissue disorders. After diagnosis, both patients were referred and followed by one of the clinical participating centers.

The combined prevalence was 0.563 per 100,000 total population. More than 90% could be classified in one of the major diagnostic groups. Mutations were identified in 220 of the 336. Table 2 shows details for each subgroup.

Major diagnostic categories. Dystroglycanopathies. This group comprised 135 patients, 40.18% of the CMD population, with a prevalence of 0.226 per 100,000. In these patients, the diagnosis of dystroglycanopathy followed the evidence of predominant weakness and wasting of the upper compared with the lower limbs, and markedly elevated serum CK levels. In a minority of cases, the diagnosis followed clinical and radiologic evidence of CNS involvement in the first months of life, associated with highly increased values of CK serum levels (more than 5-fold the normal values). In all, muscle biopsy showed absent or reduced α DG.

The form with mental retardation and microcephaly with or without cerebellar involvement and MEB were the most frequent phenotypes.

Mutations were identified in 75 of the 135 patients with α DG reduction included in the present study. All 135 were screened for the 6 genes identified before 2012 (namely, *FKRP*, *POMT1*, *POMT2*, *POMGnT1*, *Fukutin*, *LARGE*). Sixty-seven patients (49.62%) had 2 allelic mutations identified in one of these 6 genes. By subtype of mutations, *POMT1* was the most common, representing 15.5% of the total dystroglycanopathies, followed by *FKRP*, *POMGnT1*, and *POMT2* (12.59%, 10.37%, and 9.62%, respectively). Details of the occurrence of each form are shown in table 2. The other categories were individually much smaller with only isolated cases with mutations in *LARGE* or *Fukutin*.

Of the 68 subjects who did not have mutations in the first 6 genes identified, only a subset of 18 was screened for the more recently discovered genes. Mutations were found in 8 of these 18, raising the number of patients with identified mutation from 67 to 75 of 135. Three subjects had mutations in *ISPD*, 2 in *DPM2*, 2 in *GSPMD*, and 1 in *B3GALNT2*. Because genetic analysis for the remaining 50 is in progress, we cannot therefore yet fully establish the prevalence of mutations in the new genes.

Nine of the 135 patients were familial cases (5 sets of siblings; one of them had a sibling not included in the present study). All the patients in this group were of Italian origin with the exception of 16 who were resident in Italy but with parents originally from

Table 1 Classification of CMD

Locus	Gene	Disease phenotype
Extracellular matrix proteins		
6q22-23	LAMA2	Primary merosin deficiency
21q22.3	COL6A1/COL6A2	Ullrich CMD
2q37	COL6A3	
External sarcolemmal proteins		
12q13	ITGA7	Integrin α 7-related CMD
3p23-21	ITGA9	Integrin α 9-related CMD
Dystroglycan and glycosyltransferase enzymes		
9q34.1	POMT1	WWS, MEB, CMD with cerebellar involvement, CMD with mental retardation and microcephaly
1q32-34	POMGnT1	WWS, MEB, CMD with cerebellar involvement
14q24.3	POMT2	WWS, MEB, CMD with cerebellar involvement, CMD with mental retardation and microcephaly
19q13.3	FKRP	WWS, MEB, CMD with cerebellar involvement, CMD with mental retardation and microcephaly, CMD with no mental retardation
9q31	FCMD	Fukuyama CMD
22q12.3-13.1	LARGE	WWS, MEB, white matter changes
1q12-q21	DPM2/DPM3	CMD with mental retardation and severe epilepsy
7p21.2	ISPD	WWS, LGMD
3p22.1	GTDC2	WWS
11q13.2	B3GALNT2	WWS, MEB
3p21.23	GMPPB	CMD with mental retardation and severe epilepsy, LGMD
3p21	DAG1	Primary dystroglycanopathy, LGMD with early onset and mental retardation, normal brain MRI
8p11.21	SGK196	MEB
1q42	—	MDC1B
Endoplasmic reticulum protein		
1p35-36	SEPN1	CMD with spinal rigidity (RSMD1)
Nuclear envelope proteins		
6q25	SYNE1 (<i>nesprin 1</i>)	CMD with adducted thumbs
1q21.2	LMNA	Congenital laminopathy
Sarcolemmal and mitochondrial membrane protein		
22q13	CHKB	Mitochondrial CMD (CMDmt)

Abbreviations: CMD = congenital muscular dystrophy; CMDmt = CMD with mitochondrial structural abnormalities; LGMD = limb-girdle muscular dystrophy; MDC1B = muscular dystrophy, congenital, 1B; MEB = muscle-eye-brain disease; RSMD1 = rigid spine muscular dystrophy type 1; WWS = Walker-Warburg syndrome.

other countries (8 from Northern Africa, 4 from Eastern Europe, and 4 from Asia).

Merosin-deficient CMD. Eighty-one patients were classified as laminin α 2 (merosin)-deficient CMD (24.11% of the whole CMD cohort) with a prevalence of 0.136 per 100,000 for Italy.

All had a phenotype consistent with the classic course of typical phenotype reported for merosin-deficient CMD including the involvement of the white matter on brain MRI,^{8,11} and increased CK (more than 5-fold the normal values).

LAMA2 chain gene mutations were found in 51 of 81. In 7 patients, genetic analysis for *LAMA2* is in progress or consent was not obtained by the parents.

In the remaining 23 patients, mutations were not found; of these, 21 patients had absent or markedly reduced laminin α 2 and 2 partial deficiency on muscle biopsy. In the 2 cases with partial deficiency, the reduction in laminin α 2 expression was greater than the reduction observed in α DG protein expression. Both were also screened for the first 6 genes identified for dystroglycanopathies and no mutation was found.

Table 2 Analysis of cases by disease group

Disease group	No. of cases based on clinical diagnosis	Proportion (%) of clinic population affected (95% CI)	Estimated point prevalence per 100,000 (95% CI)
Dystroglycanopathies	135	40.18 (34.93-45.65)	0.226 (0.19-0.268)
Mutation found	75	22.32 (18.06-27.23)	0.126 (0.1-0.159)
FKRP	17	5.06 (3.07-8.13)	0.028 (0.017-0.045)
POMT1	21	6.25 (4-9.54)	0.035 (0.022-0.055)
POMT2	13	3.87 (2.17-6.69)	0.022 (0.012-0.039)
POMGnT1	14	4.17 (2.39-7.06)	0.023 (0.013-0.04)
LARGE	1	0.29 (0.05-1.64)	0.002 (0-0.011)
FUKUTIN	1	0.29 (0.05-1.64)	0.002 (0-0.011)
DPM2	2	0.6 (0.1-2.38)	0.003 (0-0.013)
ISPD	3	0.89 (0.23-2.8)	0.005 (0.001-0.016)
GSPPM2	2	0.6 (0.1-2.38)	0.003 (0-0.013)
B3GALNT2	1	0.29 (0.05-1.64)	0.002 (0-0.011)
In progress	11	3.27 (1.73-5.95)	0.018 (0.009-0.033)
No mutation found (6 αDG)	49	14.58 (11.08-18.92)	0.082 (0.061-0.109)
Laminin α2-deficient CMD	81	24.11 (19.71-29.12)	0.136 (0.109-0.17)
Mutation found	51	15.18 (11.61-19.57)	0.085 (0.064-0.113)
In progress	7	2.08 (0.09-4.43)	0.012 (0.005-0.026)
No mutation found	23	68.5 (4.61-10.24)	0.039 (0.025-0.059)
Ullrich CMD	68	20.24 (16.16-25.02)	0.114 (0.089-0.014)
Mutation found	55	16.37 (12.67-20.86)	0.092 (0.07-0.121)
In progress	13	3.87 (2.17-6.69)	0.022 (0.012-0.039)
Rigid spine muscular dystrophy	21	6.25 (4-9.54)	0.035 (0.022-0.055)
Mutation found	19	5.65 (3.53-8.83)	0.032 (0.02-0.051)
No mutation found	2	0.6 (0.1-2.38)	0.003 (0-0.013)
LMNA	20		
Mutation found	20	5.95 (3.77-9.19)	0.034 (0.021-0.053)
Others			
No mutation found	11	3.27 (1.73-5.95)	0.18 (0.009-0.037)

Abbreviations: α DG = α -dystroglycan; CI = confidence interval; CMD = congenital muscular dystrophy; LMNA = lamin A/C.

Sixteen cases were familial (6 sets of siblings and 2 couples of first cousins). Sixty patients in this group were of Italian origin while the other 21 were resident in Italy but had parents originally from other countries (14 from Northern Africa, 3 from Eastern Europe, 3 from Asia, and 1 from South America).

Ullrich CMD. Sixty-eight patients (20.24%) had a phenotype compatible with Ullrich CMD and reduced collagen on skin and or muscle biopsy. CK levels were normal or only mildly elevated. Muscle MRI was available in 32 of the 64 and showed the typical pattern previously reported in association with Ullrich CMD.¹² The point prevalence was 0.114 per 100,000.

Mutations were found in 55 of the 68. Regarding the phenotypes, 11 cases were familial (5 sets of siblings) and none had other familial cases not included

in the present study. All the patients in this group were of Italian origin with the exception of 9 who were resident in Italy but with parents originally from other countries (2 from South America, 2 from Northern Africa, 2 from Eastern Europe, 1 from Asia, and 2 from Turkey).

Fifty-five cases with *COL6* mutations in whom ambulation was achieved by 24 months and the onset of weakness or contractures occurred after that, were not classified as CMD and therefore not included in the present study.

Rigid spine muscular dystrophy type 1. Twenty-one patients (6.25%) had delayed ambulation, developed rigid spine and early respiratory impairment, with a clinical and MRI phenotype compatible with rigid spine muscular dystrophy type 1 (RSM1). Serum CK levels were normal or only mildly elevated. Muscle MRI

was available in 18 and showed the typical pattern previously reported in association with *SEPN1* mutations.¹² Mutations in *SEPN1* were found in 19 of the 21. The point prevalence was 0.035 per 100,000.

They were all sporadic cases, with the exception of 2 siblings. All the patients in this group were of Italian origin with the exception of 2 whose parents were originally from South Asia.

Laminopathies. Twenty patients (5.95%) had congenital presentation with dropped head or other clinical signs suggestive of the form of CMD associated with laminopathy. All but 4 achieved independent ambulation. Mutations in *LAMA A/C* were found in all 20. The point prevalence was 0.034 per 100,000.

They were all sporadic cases with the exception of a girl whose father was also affected and had died of cardiac arrhythmia. All were of Italian origin with the exception of one child whose parents were originally from South Asia.

Other cases. Eleven cases fulfilled the inclusion criteria for CMD but did not fit into any of the previous major categories. All had increased CK, ranging from 2- to 8-fold the normal values. Three had signs of CNS involvement but normal laminin $\alpha 2$ and αDG on muscle biopsy.

DISCUSSION The population prevalence figure for CMD in our study was 0.563 per 100,000. Because all the Italian tertiary care centers for neuromuscular disorders participated in this study, this figure is likely to include all cases with a clinical presentation evocative of a CMD phenotype or at least all those for whom a possible diagnosis of CMD was considered. We cannot exclude that in some cases with milder phenotypes or later onset, the diagnosis of CMD may not have been considered and these patients may have never been referred to a tertiary care center or had any diagnostic test for CMD performed.

In this study, we subdivided our cohort according to the most recent classification for CMD. Because of this, the results are not easily comparable with previous population studies as most of them only included small cohorts and/or were performed before the identification of new genes.

In 1996, the incidence of CMD was estimated at between 4.7×10^5 live births in the north of Italy,⁵ but only 20 patients with CMD were included. Another study reports a point prevalence of 6.3×10^5 in 20 patients identified in Western Sweden as part of a larger epidemiologic study in neuromuscular disorders in childhood.³ In all these cases, no subtyping of the different forms of CMD was available. A more recent study⁴ exploring the prevalence of all the genetic muscle disease in Northern England reports a prevalence of CMD of 0.76×10^5 with 26 CMD cases subdivided into 4 major subtypes.

Other recent studies have investigated larger cohorts of CMDs reporting the relative frequencies of CMD subtypes,^{13,14} but these mainly reflected the activity of diagnostic centers and were not population studies.

More than 85% of our patients are of Italian origin, and when we analyzed their postcodes, we were unable to identify any obvious correlation between individual forms and local regional areas.

Dystroglycanopathies were the most common group of CMD in our database (40.18%), followed by merosin-deficient CMD (24.11%) and Ullrich CMD (20.24%). No data on the prevalence of the individual forms are available for comparison in the population studies^{3,5} published before the last decade. Our data appear to be different from the large diagnostic UK study reporting that dystroglycanopathies were present in 12% of the whole CMD cohort, although that study only considered patients in whom a final genetic diagnosis was reached between 2001 and 2008.¹⁴ It is possible that regional differences may also have a role in the variability observed in different countries.

Among our cohort of dystroglycanopathies, the most frequent phenotypes observed were microcephaly and mental retardation, with or without cerebellar abnormalities, previously labeled as Italian MEB, and the typical MEB phenotype.¹⁵ These phenotypes were also found to be among the most frequent in the large UK study.^{16–18}

All the cases identified as dystroglycanopathies were further subdivided according to the gene mutations. *POMT1* mutations were the most frequent, followed by *FKRP*, *POMGnT1*, and *POMT2*. In 14.58%, a mutation could not be identified, but most of them have not yet been systematically screened for all the recently discovered genes. The detection rate of mutations in the subset of cases that had already been screened for all genes suggests that the full screening is likely to markedly reduce the number of cases in whom mutations cannot be identified.

The next major group was laminin $\alpha 2$ -deficient CMD (24.11% of the whole CMD cohort) with an estimated point prevalence of 0.136 per 100,000. This form has often been reported to be the most frequent form of CMD, with values up to 40% of the CMD cases.^{11,19} More recent studies^{13,14,20} report values similar to those found in our cohort.

Twenty-three of the 81 cases in our cohort (28%) had a typical clinical phenotype consistent with merosin-deficient CMD, associated with absent or marked reduction of laminin $\alpha 2$ on muscle biopsy and classic white matter changes on brain MRI, but no mutation could be identified. With the exception of 7 previous cases in whom DHPLC (denaturing high-performance liquid chromatography) was used,

Sanger sequencing of the entire coding region was performed in all. It is possible that these patients have deep intronic variants, which affected splicing or exon deletions or duplications, not easily detectable by the standard mutation detection methods used.

Forms with congenital presentation and collagen VI involvement were found in 20.24% of our cohort with similar values to that observed in the large UK series¹⁴ (21%). Mutations were found in all but one of those in whom the analysis has been completed. The high rate of detection may be explained by the fact that all these cases had the typical phenotype characterized by distal laxity, skin changes, rigid spine, and early respiratory impairment. The combination of these clinical findings with reduction of collagen 6 on skin cultures and muscle biopsy and, when performed, of the typical muscle MRI changes, appear to increase the possibility of finding mutations in the *COL6* genes. Because of our strict inclusion criteria, we may have missed some of the milder Ullrich CMD or of the recently described intermediate phenotype²¹ that had normal early milestones, including age of ambulation and onset of clinical signs after the age of 2 years. All the patients with Bethlem myopathy had onset of clinical signs after ambulation was achieved and were therefore not included.

Mutations in *SEPN1* were found in 19 of the 21 patients in whom the clinical phenotype was consistent with that reported in RSMD^{22,23} even though it was not always easy to establish retrospectively whether these should be classified as minicore myopathies or RSMD1 because of the marked clinical and pathologic overlap between these forms.²⁴

Because of its retrospective nature, our study has some limitations as the patients included had biopsies performed in different laboratories and from the reports it was not always easy to compare the severity of changes on muscle biopsies. Similarly, data on the antibodies used for assessing laminin $\alpha 2$ were not always consistent because the samples were processed at different times. This should be systematically reassessed, reevaluating the biopsies using similar classification criteria.

Mutations were identified in 220 of the 336 (65%), but in a proportion of cases, analysis of newly identified genes is still in progress. It is of interest that, even though genetic confirmation was not achieved in all, approximately 90% of our cases could be ascribed to one of the major clinical categories. This is probably related to the fact that the diagnosis was based on an integrated approach including clinical, pathologic, genetic, and often MRI features rather than on the analysis of samples sent by other centers with reported information. This approach has also allowed identification of a number of patients, formerly labeled as CMD in the past, with mutations in other genes

responsible for other disorders such as congenital myopathies or congenital myasthenia, that have a clinical and pathologic overlapping with CMD.

Another strength is that the genetic testing for the major categories such as dystroglycanopathies or collagenopathies is centralized and this guarantees that all the cases in whom the diagnosis of these categories was suspected have been included. Work is in progress to improve the detection rate of merosin-deficient CMD, using RNA studies or other genomic techniques such as MLPA (multiplex ligation-dependent probe amplification) or DNA arrays. Should this more comprehensive approach not detect any mutation, these patients would allow exploring possible genetic heterogeneity in this subgroup.

Further work has also been planned to perform further sequencing including next-generation sequencing to detect the more rare forms of CMD such as nesprin that have not been systematically investigated in our unresolved cases.

Our results clearly show how the detection rate of mutations has increased following the increased number of new genes identified. We foresee that even screening for all the known genes, a proportion will still remain without a diagnosis and that further CMD subtypes and genes are yet to be discovered.

AUTHOR CONTRIBUTIONS

Alessandra Graziano: drafting/revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Flavia Bianca Bianco: drafting/revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data, study supervision. Adele D'Amico: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Isabella Moroni: study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Sonia Messina: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Claudio Bruno: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Elena Pegoraro: drafting/revising the manuscript, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Marina Mora: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Guja Astrea: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Francesca Magri: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Giacomo Pietro Comi: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Angela Lucia Berardinelli: drafting/revising the manuscript, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Maurizio Moggio: study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Lucia Morandi: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Antonella Pini: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval. Roberta Petillo: drafting/revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Giorgio Tasca: analysis or

interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Mauro Monforte: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Carlo Minetti: drafting/ revising the manuscript, study concept or design, accepts responsibility for conduct of research and will give final approval, acquisition of data. Tiziana Mongini: study concept or design, accepts responsibility for conduct of research and will give final approval, acquisition of data. Enzo Ricci: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Ksenija Gorni: drafting/ revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Roberta Battini: drafting/ revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Marcello Villanova: drafting/ revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data. Luisa Politano: study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data, study supervision, obtaining funding. Francesca Gualandi: drafting/ revising the manuscript, accepts responsibility for conduct of research and will give final approval, acquisition of data. Alessandra Ferlini: drafting/ revising the manuscript, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, contribution of vital reagents/tools/patients. Francesco Muntoni: drafting/ revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, study supervision, obtaining funding. Filippo Santorelli: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, contribution of vital reagents/tools/patients, acquisition of data. Enrico Silvio Bertini: study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data, obtaining funding. Marika Pane: analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, contribution of vital reagents/ tools/patients, acquisition of data. Eugenio Mercuri: drafting/ revising the manuscript, study concept or design, analysis or interpretation of data, accepts responsibility for conduct of research and will give final approval, acquisition of data, obtaining funding.

STUDY FUNDING

The study is supported by a Telethon UILDM grant (GUP11001). The authors are grateful to TREAT NMD and Care CMD for their support in developing CMD registries in Italy. F. Muntoni is supported by the Great Ormond Street Children's charity and the Great Ormond Street Hospital Biomedical Research Center.

DISCLOSURE

A. Graziano, F. Bianco, A. D'Amico, I. Moroni, S. Messina, and C. Bruno report no disclosures relevant to the manuscript. E. Pegoraro has served on a scientific advisory board for BioMarin Pharmaceutical Inc.; has received funding for travel from Genzyme Corporation; has received speaker honoraria from MedaPharma; and receives research support from Wellstone and Telethon Italy. M. Mora, G. Astrea, and F. Magri report no disclosures relevant to the manuscript. G. Comi is site principal investigator for the PTC extension study of Ataluren in DMD, for the GSK study on exon skipping; receives research support from Telethon Italy and SMA Europe. A. Berardinelli, M. Moggio, L. Morandi, A. Pini, R. Petillo, G. Tasca, M. Monforte, and C. Minetti report no disclosures relevant to the manuscript. T. Mongini has served on a scientific advisory board for Telethon Italy; has received funding for travel from Genzyme Corporation; and has received research support from AIFA (Italian Government Drug Agency) and Telethon Italy. E. Ricci, K. Gorni, R. Battina, M. Villanova, L. Politano, F. Gualandi, A. Ferlini, F. Muntoni, and F. Santorelli report no disclosures relevant to the manuscript. E. Bertini is site principal investigator for the PTC extension study of Ataluren in DMD and for the GSK study on exon skipping. He also receives funds from the Italian Telethon, the Italian Ministry of Health, and SMA Europe for observational studies on outcome measures. M. Pane reports no disclosures relevant to the manuscript. E. Mercuri is site principal investigator for the PTC extension study of Ataluren in DMD and for the GSK study on exon

skipping. He also receives funds from the Italian Telethon and SMA Europe. He has acted on the advisory board for Acceleron Pharma, Prosenza, Ely Lilly, and PTC Therapeutics, Inc. Go to Neurology.org for full disclosures.

Received May 2, 2014. Accepted in final form October 15, 2014.

REFERENCES

- Dubowitz V. 22nd ENMC sponsored workshop on congenital muscular dystrophy held in Baarn, The Netherlands, 14–16 May 1993. *Neuromuscul Disord* 1994;4:75–81.
- Mercuri E, Muntoni F. The ever-expanding spectrum of congenital muscular dystrophies. *Ann Neurol* 2012;72:9–17.
- Darin N, Tulinius M. Neuromuscular disorders in childhood: a descriptive epidemiological study from western Sweden. *Neuromuscul Disord* 2000;10:1–9.
- Norwood FL, Harling C, Chinnery PF, Eagle M, Bushby K, Straub V. Prevalence of genetic muscle disease in Northern England: in-depth analysis of a muscle clinic population. *Brain* 2009;132:3175–3186.
- Mostacciolo ML, Miorin M, Martinello F, Angelini C, Perini P, Trevisan CP. Genetic epidemiology of congenital muscular dystrophy in a sample from north-east Italy. *Hum Genet* 1996;97:277–279.
- Bonnemann CG, Wang CH, Quijano-Roy S, et al. Diagnostic approach to the congenital muscular dystrophies. *Neuromuscul Disord* 2014;24:289–311.
- Nadeau A, Muntoni F. Skin changes in Ullrich congenital muscular dystrophy. *Neuromuscul Disord* 2008;18:982.
- Geranmayeh F, Clement E, Feng LH, et al. Genotype-phenotype correlation in a large population of muscular dystrophy patients with LAMA2 mutations. *Neuromuscul Disord* 2010;20:241–250.
- Messina S, Bruno C, Moroni I, et al. Congenital muscular dystrophies with cognitive impairment: a population study. *Neurology* 2011;75:898–903.
- Mercuri E, Messina S, Bruno C, et al. Congenital muscular dystrophies with defective glycosylation of dystroglycan: a population study. *Neurology* 2009;72:1802–1809.
- Tome FM, Evangelista T, Leclerc A, et al. Congenital muscular dystrophy with merosin deficiency. *C R Acad Sci III* 1994;317:351–357.
- Mercuri E, Clements E, Offiah A, et al. Muscle magnetic resonance imaging involvement in muscular dystrophies with rigidity of the spine. *Ann Neurol* 2010;67:201–208.
- Peat RA, Smith JM, Compton AG, et al. Diagnosis and etiology of congenital muscular dystrophy. *Neurology* 2008;71:312–321.
- Clement EM, Feng L, Mein R, et al. Relative frequency of congenital muscular dystrophy subtypes: analysis of the UK diagnostic service 2001–2008. *Neuromuscul Disord* 2012;22:522–527.
- Villanova M, Mercuri E, Bertini E, et al. Congenital muscular dystrophy associated with calf hypertrophy, microcephaly and severe mental retardation in three Italian families: evidence for a novel CMD syndrome. *Neuromuscul Disord* 2000;10:541–547.
- Clement E, Mercuri E, Godfrey C, et al. Brain involvement in muscular dystrophies with defective dystroglycan glycosylation. *Ann Neurol* 2008;64:573–582.
- Godfrey C, Foley AR, Clement E, Muntoni F. Dystroglycanopathies: coming into focus. *Curr Opin Genet Dev* 2011;21:278–285.
- Godfrey C, Clement E, Mein R, et al. Refining genotype phenotype correlations in muscular dystrophies with

- defective glycosylation of dystroglycan. *Brain* 2007;130:2725–2735.
19. Philpot J, Sewry C, Pennock J, Dubowitz V. Clinical phenotype in congenital muscular dystrophy: correlation with expression of merosin in skeletal muscle. *Neuromuscul Disord* 1995;5:301–305.
 20. Bönemann CG. Congenital muscular dystrophy. In: Squire LR, editor. *Encyclopedia of Neuroscience*. Oxford, UK: Academic Press; 2009:67–74.
 21. Foley AR, Quijano-Roy S, Collins J, et al. Natural history of pulmonary function in collagen VI-related myopathies. *Brain* 2013;136:3625–3633.
 22. Flanigan KM, Kerr L, Bromberg MB, et al. Congenital muscular dystrophy with rigid spine syndrome: a clinical, pathological, radiological, and genetic study. *Ann Neurol* 2000;47:152–161.
 23. Scoto M, Cirak S, Mein R, et al. SEPN1-related myopathies: clinical course in a large cohort of patients. *Neurology* 2011;76:2073–2078.
 24. Ferreiro A, Quijano-Roy S, Pichereau C, et al. Mutations of the selenoprotein N gene, which is implicated in rigid spine muscular dystrophy, cause the classical phenotype of multimincore disease: reassessing the nosology of early-onset myopathies. *Am J Hum Genet* 2002;71:739–749.


20 Minutes Pack a Punch

Neurology® Podcasts

- Interviews with top experts on new clinical research in neurology
- Editorial comments on selected articles
- Convenient—listen during your commute, at your desk, or even at the gym
- On demand—it's there when you want it
- Fun and engaging
- New topic each week
- FREE

Listen now at www.aan.com/podcast

Neurology® Genetics Call For Papers



Neurology: Genetics is an open access journal that will provide neurologists with outstanding original contributions that elucidate the role of genetic and epigenetic variation in diseases and biological traits of the central and peripheral nervous system. We welcome all submissions. For more information on how to submit, visit <http://www.neurology.org/site/gen/gen2.xhtml>.