



Published in final edited form as:

*J Dairy Sci.* 2014 December ; 97(12): 7633–7643. doi:10.3168/jds.2014-8588.

## Effect of preweaned dairy calf housing system on antimicrobial resistance in commensal *Escherichia coli*

R. V. Pereira<sup>\*,1</sup>, J. D. Siler<sup>\*</sup>, J. C. Ng<sup>\*</sup>, M. A. Davis<sup>†</sup>, and L. D. Warnick<sup>\*</sup>

<sup>\*</sup>Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY 14853

<sup>†</sup>Veterinary Microbiology and Pathology Department, Washington State University, Pullman 99164-7040

### Abstract

Group housing of preweaned dairy calves is a growing practice in the United States. The objective of this practice is to increase the average daily gain of calves in a healthy and humane environment while reducing labor requirements. However, feeding protocols, commingling of calves, and occurrence of disease in different calf-housing systems may affect the prevalence of antimicrobial drug-resistant bacteria. This study evaluated the effect of a group pen-housing system and individual pen-housing system on antimicrobial resistance trends in fecal *Escherichia coli* of preweaned dairy calves and on the prevalence of environmental *Salmonella*. Twelve farms from central New York participated in the study: 6 farms using an individual pen-housing system (IP), and 6 farms using a group pen-housing system (GP). A maximum of 3 fecal *E. coli* isolates per calf was tested for susceptibility to 12 antimicrobial drugs using a Kirby-Bauer disk diffusion assay. Calves in GP had a significantly higher proportion of *E. coli* resistant to ciprofloxacin and nalidixic acid, whereas calves in IP had a significantly higher proportion of *E. coli* resistant to ampicillin, ceftiofur, gentamycin, streptomycin, and tetracycline. Calf-housing system had an effect on resistance to individual antimicrobial drugs in *E. coli*, but no clear-cut advantage to either system was noted with regard to overall resistance frequency. No outstanding difference in the richness and diversity of resistant phenotypes was observed between the 2 calf-housing systems.

### Keywords

antimicrobial resistance; calf-housing system; group housing; individual pens

### INTRODUCTION

According to the last inventory report released by the USDA in January 2011, there were more than 14 million calves in the United States, of which 650,000 are estimated to be preweaned dairy calves (USDA, 2011). A growing practice on US dairies, especially in the

northeast, is housing calves in group pens ranging from 12 to 25 calves, and feeding free-choice (ad libitum) acidified milk. Milk is acidified to a pH of 4.0 to 4.5 to preserve it by preventing bacteria and mold growth. This practice differs from the current conventional system where calves are housed in individual pens and fed predetermined amounts of milk or milk replacer for specific number of times during the day. The objective of group housing is to increase the ADG of calves in a healthy and humane environment while reducing labor requirements (Warnick et al., 1977; O'Driscoll et al., 2006; Bernal-Rigoli et al., 2012). One calf-housing study reported that the time needed to manage a calf in an individual pen is approximately 10 min/calf per day, but the time committed to management of a calf raised in a group pen is less than 1 min/calf per day (Kung et al., 1997). A potential disadvantage of group housing is that direct contact between calves in group pens could increase the risk of transmission of pathogens between calves and, potentially, the exchange of antimicrobial resistance.

The effect of group penning on calf health is not clear and the practice remains controversial (Wells et al., 1997). Housing calves in group pens was observed by Linton et al. (1974) to result in a higher incidence of salmonellosis, with a 10.5% average incidence in calves housed in group pens compared with a 1.1% average incidence in calves housed in individual pens. Waltner-Toews et al. (1986), in a study of dairy calf management, morbidity, and mortality in Ontario Holstein herds, observed that the direction of group penning effects on calf morbidity relative to individual pens was not consistent. They observed that the odds ratio of scours or pneumonia for calves housed either in group pens or individual pens was not significantly different, and that any effects appeared to be minimal.

Commingling of calves, feeding protocols, and incidence of disease are a few differences between calves housed in group pens and calves housed in individual pens that could result in a distinct prevalence of antimicrobial drug resistance. The objective of our study was to evaluate the effects of these 2 calf-housing systems—calves in individual pens fed milk or milk replacer 2 to 3 times a day and calves in group pens fed free-choice acidified milk—on antimicrobial resistance trends in fecal *Escherichia coli* and on the prevalence of environmental *Salmonella*.

## MATERIALS AND METHODS

### Inclusion Criteria for Farms

Farms were eligible for inclusion in the present study if their calf-housing system consisted of either (1) housing calves in individual pens while feeding them milk or milk replacer (**IP**) or (2) housing calves in group pens while feeding them acidified milk ad libitum (**GP**). In addition, farms participating in the current study could not add any antimicrobial drugs to the milk fed to pre-weaned calves. A total of 658 preweaned dairy calves were enrolled from 12 farms in central New York (Table 1). Herds were selected from a convenience sample of commercial dairy farms within a 3-h radius of Cornell University (Ithaca, NY). Dairy farms were located in the following counties in central New York State: Cayuga, Cortland, Onondaga, Ontario, Tompkins, and Wayne. Farmers were requested to answer a short questionnaire on calf management practices on the dairy.

## Study Design and Sample Collection

Our study used a cross-sectional design in which rectal swabs were collected from preweaned dairy calves during a single farm visit. At each farm, preweaned calves were divided into 3 age groups and approximately the same numbers of total samples were collected from each age group using a randomization spreadsheet. The randomization spreadsheet was created using the random number function of Excel (Microsoft, Redmond, WA). The age groups were (1) 3 to 14, (2) 15 to 35, and (3) 36 to 65 d of age. All calves enrolled in the present study were female. Environmental samples were collected from the pen floors and feeder nipples of group pens in GP farms, and the pen floor and hutches of calf pens in IP farms. Farm sampling was conducted from July to October 2012.

## Bacterial Isolation, Culture, and Identification

Transport swabs with Amies medium were used to collect individual rectal fecal samples from calves (Becton Dickinson and Company, Franklin Lakes, NJ). Individual fecal swabs were streaked onto MacConkey agar plates on the day of collection and incubated overnight at 37°C. Up to 3 *E. coli* isolates were confirmed as *E. coli* and stored in Luria-Bertani broth containing 20% glycerol at -80°C, as previously described (Pereira et al., 2011). A total of 1,960 *E. coli* isolates were obtained from fecal swab cultures (Table 1).

Standard bacteriologic culture methods were used to isolate *Salmonella* from environmental samples. Individual swab specimens from each sample were enriched in 10 mL of Tetrathionate broth (Difco, Detroit, MI) containing 0.2 mL of iodine solution; the mixture was incubated at 42°C for 18 to 24 h. After incubation, the sample-broth mixture was streaked onto Brilliant Green agar with novobiocin (Northeast Laboratory, Waterville, Maine) and xylose lysine tergitol 4 (XLT-4) selective media, and both plates were incubated at 37°C for 18 to 24 h. Red colonies (lactose-nonfermenting bacteria) on Brilliant Green agar with novobiocin and black colonies (hydrogen sulfide-producing bacteria) on XLT-4 were inoculated into Kligler Iron Agar slants and incubated at 37°C for 18 to 24 h. The XLT-4 plates without suspected colonies were reincubated at 37°C for an additional 18 to 24 h before checking again for characteristic black colonies. If a Kligler Iron Agar slant exhibited the biochemical properties of *Salmonella*, the isolate was confirmed by slide agglutination using *Salmonella* O Antiserum Poly A-I & Vi (Becton Dickinson and Company). Confirmed *Salmonella* isolates were stored in Luria-Bertani broth containing 20% glycerol at -80°C. Additionally, 1 confirmed *Salmonella* isolate from each sample was sent to the National Veterinary Services Laboratories in Ames, Iowa, for serotyping.

Antimicrobial susceptibility of *E. coli* and *Salmonella* isolates was tested for a modified National Antimicrobial Resistance Monitoring system panel of 12 antimicrobial drugs. National Antimicrobial Resistance Monitoring was established in 1996 by the Food and Drug Administration's Center for Veterinary Medicine in collaboration with the Centers for Disease Control and Prevention and the USDA in an effort to prospectively monitor the occurrence of antimicrobial resistance of zoonotic pathogens from human diagnostic specimens, retail meats, and food animals. The susceptibility testing test was done using a Kirby-Bauer disk diffusion agar assay in accordance with the guidelines published by the Clinical and Laboratory Standards Institute (CLSI) and methodology previously described

(CLSI, 2008b; Hoelzer et al., 2011; Pereira et al., 2011). Internal quality control was performed by inclusion of *E. coli* ATCC 25922, previously determined to be pansusceptible, as well as a previously characterized in-house *E. coli* isolate known to have a *bla*<sub>CMY-2</sub> gene and to be resistant to 9 of the antimicrobials tested. All susceptibility results of quality control strains complied with the quality control ranges. Antimicrobial susceptibility for all isolates was assessed using the following panel: ampicillin, 10 µg; cefoxitin, 30 µg; ceftiofur, 30 µg; ceftriaxone, 30 µg; chloramphenicol, 30 µg; ciprofloxacin, 5 µg; gentamicin, 10 µg; nalidixic acid, 30 µg; neomycin, 30 µg; streptomycin, 10 µg; tetracycline, 30 µg; and trimethoprim-sulfamethoxazole, 23.75/1.25 µg. Susceptibility of the isolates to most antimicrobial drugs was categorized (susceptible, intermediate, or resistant) by measuring the inhibition zone according to interpretive criteria adhering to the CLSI guidelines (CLSI, 2008a).

### Statistical Analyses

*Escherichia coli* isolates were categorized as susceptible, intermediate, or resistant in accordance with CLSI guidelines as described previously. The herd-level effect from the group-housing system on the proportion of calves resistant to antimicrobial drugs was estimated using generalized linear models (GLM). For our analysis, a calf was considered positive for resistance if at least 1 of the *E. coli* was classified as resistant. The herd-level proportion of calves resistant to individual antimicrobial drugs was determined using PROC MEAN in SAS (SAS Institute Inc., Cary, NC). The GLM for each antimicrobial drug tested were created using PROC GLM in SAS. For these herd-level analyses, the dependent variable was the proportion of resistant calves per farm sampled and the independent variable was calf-housing system (IP and GP). The assumption of normal distribution of residuals was analyzed using PROC UNIVARIATE in SAS and by visually evaluating the distribution plot of the residuals. Assumption of normal distribution of residuals was satisfied.

To evaluate the effect of calf-housing system on the odds of *E. coli* resistance, multivariable mixed logistic regression models were fitted to the data using the GLIMMIX procedure of SAS. The variables calf-housing system (IP or GP) and age group (3 to 14, 15 to 35, and 36 to 65 d of age) were included in all models as independent variables and the variable for individual identification of each calf (to account for clustering within calf) was nested within the herd variable and included in all models as a random effects.

Antimicrobial drug susceptibility phenotypes (ADSP) consisted of patterns of resistance to the antimicrobial drugs, including a pansusceptible phenotype. Diversity and richness of ADSP at the pen and farm levels were estimated using the 9.1 version of the software *EstimateS* (Colwell, 2013). Pens represented by less than 6 *E. coli* isolates were considered inadequate for our analysis and a total of 4 pens (total of 17 *E. coli* isolates) from GP farms were excluded from this analysis. Richness was calculated using the Chao richness index and by using rarefaction curves, which are produced by repeatedly sampling all of the collected species at random and computing the expected number of species as a function number of accumulated samples (MacGregor-Fors and Payton, 2013). Diversity was estimated using the Shannon diversity index. Rarefaction curves, Chao index, and Shannon

index were estimated by calf-housing system type at the pen and herd level. To compare the richness, diversity, and biotic similarity of antimicrobial susceptibility phenotypes at the pen level for IP farms, mock group pens were created. This was done by (1) matching an IP farm to a GP farm by total number of lactating cows; (2) ordering GP pens at the farm level according to the average calf age in days in each pen and determining the proportion of total sampled calves present in each pen; and (3) ordering all calves sampled for each IP farm at the farm level by age in days and distributing them in the same number of pens present in the matched GP farm following the same proportion of total sampled calves present in each pen using the age in days range as a guide.

*Escherichia coli* ADSP similarities between pairs of calves were assessed using the Sorensen and Jaccard incidence-based similarity indices. The Jaccard index compares the number of shared species to the total number of species in the combined assemblages, whereas the Sorensen index compares the number of shared species to the mean number of species in a single assemblage (Gotelli and Chao, 2013). Additionally, the Chao-Sorensen abundance-based similarity index was used to measure *E. coli* ADSP similarity. Similarity indices were calculated at the pen and farm level using *EstimateS* (Colwell, 2013). Both the incidence- and abundance-based similarity indices range from 0 to 1, with 0 indicating that no susceptibility phenotypes are shared between a pair of calves and 1 indicating complete identity (all the susceptibility phenotypes present in calf 1 were also present in calf 2 and vice versa). The similarity indices are shown as a percentage in the results by multiplying the index by 100.

Descriptive analysis of *E. coli* resistance phenotypes and environmental samples cultured for *Salmonella* was conducted using PROC FREQ in SAS. In our study, multidrug resistance was defined as having resistance to 3 antimicrobial agents. For all statistical models and tests, variables were considered significant when a  $P < 0.05$  was observed.

## RESULTS

### Descriptive Data

A total of 12 farms participated in our study, with 6 farms using a GP housing system for calves and 6 farms using an IP housing system for calves. The number of calves in a group pen at GP farms ranged from 6 to 10 calves for 2 farms, 15 to 20 calves for 2 farms, and 20 to 26 calves for 2 farms. For all GP farms, calves were added to a group pen according to birth order until the desired number of calves for the pen was reached. For both GP and IP farms, 1 farm had less than 200 lactating cows, 2 farms had 500 to 1,000 lactating cows, and 3 farms had more than 1,000 lactating dairy cows. The total number of calves and *E. coli* isolates and the average age of calves in the study by calf-housing system are shown in Table 1. The 2 most commonly reported antimicrobial drugs used for calves in the group housing system were enrofloxacin (Bayer Animal Health, Shawnee Mission, KS) and tulathromycin (Draxxin Injectable Solution, Pfizer Animal Health, New York, NY). Information on the 2 most common antimicrobial drugs used on farms with individual-housing systems, ceftiofur (Excenel RTU, Pfizer Animal Health) and penicillin G, was available for 3 of 6 farms and were.

### Antimicrobial Resistance in *E. coli* Isolates

At the herd level, calf-housing systems did not have a significant effect on the proportion of antimicrobial resistance (Table 2). At the calf level, IP farms had a significantly greater number of resistant isolates per calf compared with GP farms for 1 or more age groups for ampicillin, ceftiofur, gentamicin, streptomycin, and tetracycline (Table 3). The GP farms had a significantly greater number of resistant isolates per calf compared with IP farms for ciprofloxacin ( $P < 0.0001$ ) and nalidixic acid ( $P = 0.001$ ; Table 3). No significant difference between the numbers of resistant isolates per calf for calf-housing systems by age group was observed for any of the antimicrobials tested.

### Distribution of *E. coli* Antimicrobial Susceptibility Phenotypes

Of the 903 *E. coli* isolates from IP calves, 11.7% were pansusceptible and 75.5% were multidrug resistant. Of the 1,057 isolates from GP calves, 18.2% were pansusceptible and 69.1% were multidrug resistant. The most common multidrug-resistant phenotype observed in IP isolates was ampicillin-cefoxitin-ceftiofur-streptomycin-tetracycline (5.4%), whereas for GP isolates chloramphenicol-streptomycin-tetracycline (7.3%) was the most common (Table 4).

Figure 1 shows the rarefaction curves where isolates were clustered by pen and by farm for IP and GP. The mean Chao richness index for *E. coli* ADSP was 11.5 and 29.3 for IP isolates ( $n = 903$ ) and 10.1 and 29.7 for GP isolates ( $n = 1,040$ ) at the pen and farm level, respectively (Table 5). The mean Shannon diversity index for *E. coli* ADSP at the pen and farm level were 1.6 and 2.4 for IP isolates ( $n = 903$ ) and 1.5 and 2.3 for GP isolates, respectively ( $n = 1,040$ ; Table 5).

Within each calf-housing system, both incidence-(Jaccard and Sorensen) and abundance-based (Chao-Sorensen) similarity indices for *E. coli* ADSP were higher between pairs of calves at the pen level compared with the farm level (Table 5). When similarity indexes were compared between calf-housing systems, both incidence- and abundance-based similarity indexes calculated either at the pen or farm level were slightly higher for GP (Table 5).

### Distribution of Environmental *Salmonella* Serovars

Among the 122 environmental samples, 36% were positive for *Salmonella*. Of these, 25 *Salmonella* isolates were from IP farms and 19 *Salmonella* isolates were from GP farms. The *Salmonella* serovars identified in IP isolates were Cerro ( $n = 23$ ) and Liverpool ( $n = 2$ ), whereas for GP isolates Cerro ( $n = 11$ ) and Newport ( $n = 8$ ) were observed (Table 6). All *Salmonella* isolates from IP farms and all Cerro isolates were pansusceptible. Eight Newport isolates had the same multidrug-resistant phenotype ampicillin-cefoxitin-ceftiofur-chloramphenicol-neomycin-streptomycin-tetracycline and belonged to a single GP farm.



## DISCUSSION

### Antimicrobial Resistance in *E. coli* Isolates

The GP calves had a higher prevalence of resistance to antimicrobials belonging to the quinolone class than did IP calves, with 0.2 and 0.3 odds for resistance in IP compared with GP for ciprofloxacin ( $P < 0.0001$ ) and nalidixic acid ( $P = 0.001$ ; Table 3). *Escherichia coli* from calf feces typically are not resistant to ciprofloxacin and nalidixic acid, and this resistance is an uncommon finding, as reported in other calf studies (Gow et al., 2008; Pereira et al., 2011). However, a recent study that inclusively screened calf *E. coli* isolates received by the Cornell University Animal Health Diagnostic Center from 2004 to 2011 observed that resistance to enrofloxacin has increased from 1.0% (6/574) in 2004 to 8.1% (15/185) in 2011 (Cummings et al., 2014). The observed higher prevalence of resistance to quinolone drugs is a major area of concern, because they are regarded as critically important antimicrobial agents for human medicine according to the World Health Organization (Collignon et al., 2009). Resistance to quinolones can limit therapy options in humans with invasive diseases due to *Salmonella* spp., *Campylobacter* spp., and multidrug-resistant *Shigella* spp. (Collignon et al., 2009).

The health effects of housing calves in group pens under field conditions are not clear and the practice remains controversial (Wells et al., 1997; Lorenz et al., 2011). Many study reports agree that, overall, calves housed in group pens tend to have a higher occurrence of respiratory disease compared with calves housed in individual pens (Hanekamp et al., 1994; Gulliksen et al., 2009). A study by Svensson et al. (2003) observed that calves younger than 90 d old housed in group pens with 6 to 30 calves had a 2.2 odds ratio for incidence of respiratory disease compared with calves housed in single pens ( $P = 0.019$ ). However, in this same study, calves housed in group pens with 3 to 8 calves had a 0.93 odds ratio of respiratory disease compared with calves housed in individual pens ( $P = 0.019$ ). A later study focusing on the effect of group pen size on health reconfirmed that the risk of respiratory disease can be reduced by reducing the number of calves in a group pen (Svensson and Liberg, 2006). The most common range of calves per group pen in our study was 15 to 20 calves, which may have resulted in a higher medication rate of GP calves with antimicrobial drugs used for treatment of bovine respiratory disease (**BRD**). Furthermore, Baytril (Bayer Animal Health) is the only quinolone drug currently approved for use in commercial dairy cattle in the United States. This is an enrofloxacin drug approved for use in nonlactating dairy cattle less than 20 mo of age for the treatment of BRD and for the control of BRD in animals at high risk of developing BRD associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni*, and *Mycoplasma bovis*. A higher occurrence of pneumonia in GP calves followed or preceded by the use of enrofloxacin could potentially be the cause for the observed higher prevalence of resistance to quinolone drugs in GP calves when compared with IP calves. This is in accordance with the descriptive data, where 1 of the 2 most common drugs used on the GP farms was enrofloxacin (Baytril). However, more information on individual animal treatment would be necessary to confirm this hypothesis.

Calves from IP presented a significantly higher proportion of *E. coli* resistant to ampicillin, ceftiofur, streptomycin, and tetracycline compared with the GP group. Reports have showed a tendency for higher incidences of diarrhea in calves housed in individual pens compared with those housed in group pens being fed milk manually or using an ad libitum automatic milk-feeding (Hänninen et al., 2003; Svensson et al., 2003). Svensson et al. (2003) suggested that this could be occurring because calves in group pens better meet their social and emotional needs and drink their milk in a more natural way (Svensson et al., 2003). This may be relevant to the selection of resistance in IP calves, because diarrhea in calves has previously been associated with identification of antimicrobial drug resistance (Orden et al., 2000; Kaneene et al., 2009). A study by de Verdier et al. (2012) evaluated the relationship between occurrence of diarrhea and prevalence of antimicrobial resistance in fecal *E. coli*. In their study, *E. coli* was isolated from fecal samples of pairs of calves where 1 calf was diagnosed with diarrhea and 1 calf was diagnosed as healthy. Their study only sampled calves not treated with antimicrobial drugs within 2 wk before sampling. They observed a higher prevalence of isolates resistant to ampicillin, tetracycline, streptomycin, and sulfonamide ( $P = 0.05$ ) in calves with diarrhea when compared with healthy calves. The selection of resistance found in diarrheic calves could be due to linkage between virulence genes and resistance genes in *E. coli* causing diarrhea in calves (Martínez and Baquero, 2002). In the de Verdier et al. (2012) study, phenotypic resistance to one or more antimicrobial drugs was associated with the presence of 3 *E. coli* virulence genes. The administration of broad-spectrum  $\beta$ -lactams, such as ampicillin (e.g., Polyflex, Boehringer Ingelheim Vetmedica, St. Joseph, MO) and ceftiofur (e.g., Excenel RTU, Pfizer Animal Health, Exton, Pennsylvania), to treat diarrhea in calves could also be a factor in the selection of resistance to antimicrobial drugs in IP calves (Constable, 2004). The fact that the 2 most commonly used antimicrobials on IP farms were  $\beta$ -lactams (ceftiofur and penicillin G) corroborates the hypothesis that more animals may have been treated for diarrhea on IP farms, and that selection for resistance to  $\beta$ -lactams could have also occurred because of the therapeutic use of these drugs.

To our knowledge, the higher proportion of resistance to gentamicin observed in IP isolates has not been previously related to diarrhea in calves. However, a study by Ahmed et al. (2009) identified class 1 integrons in up to 10.4% of *E. coli* isolates from calves with diarrhea. Integrons are mobile DNA elements capable of encoding multiple resistance genes at once (Gillings et al., 2008). Furthermore, diarrhea could have resulted in the selection of *E. coli*-carrying class 1 integrons that resulted in co-selection of resistance to gentamicin. Moreover, characterization of class 1 integron-mediated antibiotic resistance among *E. coli* from calves has revealed the capability to integrate antimicrobial resistance genes to antimicrobial classes, such as  $\beta$ -lactams and amino-glycosides (Du et al., 2005). Resistance to these antimicrobial classes was observed in higher prevalences in IP isolates compared with GP isolates, and could have resulted in co-selection of gentamicin.

Dietary differences between IP and GP calves could also play a role in the selection advantage of resistant bacteria on IP farms. A study by Khachatryan et al. (2006) observed that feeding preweaned calves milk with an added nonmedicated dietary supplement composed of nonfat dry milk and vitamins contributed to a significantly higher prevalence



of fecal *E. coli* resistant to streptomycin, sulfadiazine, and tetracycline when compared with calves fed milk without the addition of the supplement. The researchers' hypothesis for this finding was that a genetic linkage of the streptomycin, sulfadiazine, and tetracycline determinants to other genes may exist and confer a selective advantage in the presence of the dietary supplement. The difference in the proportion of antimicrobial drug resistance between IP and GP calves may be related both to dissimilarities in the nutrients in acidified and nonacidified milk or milk replacer as well as the volume and frequency of feed consumed. Acidification of milk has been shown to have little to no effect on milk nutrients commonly considered of major importance, such as fat and CP (Woodford et al., 1987; Hill et al., 2013). However, it has been claimed that the lower pH of milk and frequency of meals of GP calves could diminish diurnal pH fluctuations in the abomasum, resulting in a lower pH in the digestive tract and a reduction in the amount of pathogens (Hepola, 2003; Guler et al., 2006). Preweaned calves provided milk ad libitum have been shown to have more frequent meals (up to 7 meals a day) and more evenly distributed diurnal patterns of feeding activity compared with the restrictive feeding of milk twice a day (Miller-Cushon et al., 2013).

Independent of the calf-housing system, a shift in the proportion of resistant isolates by age group was observed in our study (Table 3). The variance in antimicrobial drug resistance observed between calf-housing systems at different age groups could be a consequence of changes in the microbiota due to different feeding protocols in IP and GP. A study by Berge et al. (2005) observed that calves 2 wk of age were more likely to have multidrug-resistant *E. coli* compared with calves 4 and 6 wk old (OR = 29.8 and 16.4, respectively). Lack of a developed intestinal microflora in young calves could be a factor resulting in higher colonization of younger calves by resistant bacteria. A degree of protection against colonization by bacteria with a higher fitness cost, such as antimicrobial-resistant bacteria and pathogenic enteric bacteria, may follow as the calves' indigenous microflora matures and the enteric microbiota diversity increases, resulting in a decreased prevalence of resistant bacteria (Cummings et al., 2009; Oikonomou et al., 2013). Studies evaluating the effects of feeding preweaned calves acidified and nonacidified milk on the composition of the gastro-intestinal microbiota are needed to test this assumption.

### Distribution of *E. coli* Antimicrobial Susceptibility Phenotypes

Calves in GP presented a higher prevalence of multi-drug resistant phenotypes than IP calves did, as seen in Table 4. To observe if the accumulation of *E. coli* ADSP between calf-housing system types clustered differently at the pen and farm levels, a rarefaction curve was used to visualize the richness of ADSP. Similar results for most rarefaction curves within GP pens and farms were observed, with signs of reaching an asymptote earlier than in IP pens and farms (Figure 1). Rarefaction curves reach an asymptote faster when accumulation occurs with fewer numbers of isolates. Therefore, when no new ADSP is added, an asymptote should be reached (Gotelli and Colwell, 2001). The rarefaction curve indicates that GP isolates are distributed in fewer ADSP, suggesting less richness of isolates. Because not all rarefaction curves reached an asymptote, ADSP richness was estimated using Chao's asymptotic mathematical function (Magurran and McGill, 2011). A slightly lower and narrower range for the mean Chao richness index and Shannon diversity index

calculated at the pen and farm level were observed for GP isolates compared with IP isolates.

Moreover, mean similarity indices of *E. coli* ADSP between pairs of calves were higher between calves in GP than calves in IP (Table 5). A potential cause for these findings is that commingling of calves in group pens increases the exchange of bacteria and resistance genes between calves, resulting in a more homogenous distribution of *E. coli* ADSP between calves in GP. Although commingling may play a role in the distribution of *E. coli* ADSP, we did not observe a significant difference in richness, diversity, and similarity indices between GP and IP isolates, indicating that it probably is not the main factor in the spread of antimicrobial resistant commensal *E. coli*.

### Distribution of Environmental Salmonella Serovars

*Salmonella enterica* serovar Cerro was the most common serovar observed in farm environmental samples. This is in accordance with a recent study that reported Cerro as the most common serotype among 1,174 dairy cattle isolates from dairy farms in the northeastern United States between 2004 and 2011 (Cummings et al., 2013). More IP farms (83%) had at least 1 *Salmonella* colony isolated from environmental samples compared with GP farms (33%; Table 6). An interesting finding among GP farms was the isolation of *S. enterica* serovar Cerro from the feeding nipples shared by calves. Although the role of *S. enterica* serovar Cerro in causing clinical illness among dairy cattle is still unclear, the isolation of Cerro from the feeding nipples represents a potential source for the dissemination of pathogenic bacteria in a group pen, reinforcing the importance of frequent sanitation and replacement of feeding nipples on GP farms (Cummings et al., 2010).

## CONCLUSIONS

The calf-housing system has an effect on resistance to individual antimicrobial drugs in *E. coli*, but no clear-cut advantage to either system with regard to overall resistance frequency was observed. Calves in GP had a significantly higher proportion of *E. coli* resistant to ciprofloxacin and nalidixic acid, whereas calves in IP had a significantly higher proportion of *E. coli* resistant to ampicillin, ceftiofur, gentamicin, streptomycin, and tetracycline. No outstanding difference in the richness and diversity of resistant phenotypes was noted between the 2 calf-housing systems. *Salmonella* Cerro was the most commonly isolated serotype in environmental samples from all farms. *Salmonella* was isolated from feeder nipples on GP farms, raising the importance of frequent sanitation and replacement of shared feeding nipples in group pens.

## Acknowledgments


We thank Pepi Leids from the New York State Cattle Health Assurance Program (Albany, NY) for assisting with the enrollment of farms in this project. Research reported in this publication was supported by the Agriculture and Food Research Initiative Competitive Grant no. 2010-51110-21131 from the USDA National Institute of Food and Agriculture (Washington, DC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the USDA. Research reported in this publication was supported by the Office of the Director, National Institutes of Health (Bethesda, MD), under Award Number T32OD0011000. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health.

## References

- Ahmed AM, Younis EE, Osman SA, Ishida Y, El-Khodery SA, Shimamoto T. Genetic analysis of antimicrobial resistance in *Escherichia coli* isolated from diarrheic neonatal calves. *Vet Microbiol.* 2009; 136:397–402. [PubMed: 19128900]
- Berge AC, Atwill ER, Sisco WM. Animal and farm influences on the dynamics of antibiotic resistance in faecal *Escherichia coli* in young dairy calves. *Prev Vet Med.* 2005; 69:25–38. [PubMed: 15899294]
- Bernal-Rigoli JC, Allen JD, Marchello JA, Cuneo SP, Garcia SR, Xie G, Hall LW, Burrows CD, Duff GC. Effects of housing and feeding systems on performance of neonatal Holstein bull calves. *J Anim Sci.* 2012; 90:2818–2825. [PubMed: 22408085]
- CLSI. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Collected from Animals; Approved Standards. 3. Clinical and Laboratory Standards Institute; Wayne, PA: 2008a.
- CLSI. Clinical and Laboratory Standards Institute. Clinical and Laboratory Standards Institute; Wayne, PA: 2008b. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals: Approved Standard.
- Collignon P, Powers JH, Chiller TM, Aidara-Kane A, Aarestrup FM. World health organization ranking of antimicrobials according to their importance in human medicine: A critical step for developing risk management strategies for the use of antimicrobials in food production animals. *Clin Infect Dis.* 2009; 49:132–141. [PubMed: 19489713]
- Colwell, RK. EstimateS: Statistical estimation of species richness and shared species from samples, version 9.1.0. R. K. Cowell; Storrs, CT: 2013.
- Constable PD. Antimicrobial use in the treatment of calf diarrhea. *J Vet Intern Med.* 2004; 18:8–17. [PubMed: 14765726]
- Cummings KJ, Aprea VA, Altier C. Antimicrobial resistance trends among *Escherichia coli* isolates obtained from dairy cattle in the northeastern United States, 2004–2011. *Foodborne Pathog Dis.* 2014; 11:61–67. [PubMed: 24134668]
- Cummings KJ, Perkins GA, Khatibzadeh SM, Warnick LD, Altier C. Antimicrobial resistance trends among *Salmonella* isolates obtained from dairy cattle in the northeastern United States, 2004–2011. *Foodborne Pathog Dis.* 2013; 10:353–361. [PubMed: 23458026]
- Cummings KJ, Warnick LD, Alexander KA, Cripps CJ, Gröhn YT, McDonough PL, Nydam DV, Reed KE. The incidence of salmonellosis among dairy herds in the northeastern United States. *J Dairy Sci.* 2009; 92:3766–3774. [PubMed: 19620658]
- Cummings KJ, Warnick LD, Elton M, Rodriguez-Rivera LD, Siler JD, Wright EM, Grohn YT, Wiedmann M. *Salmonella enterica* serotype Cerro among dairy cattle in New York: An emerging pathogen? *Foodborne Pathog Dis.* 2010; 7:659–665. [PubMed: 20187753]
- de Verdier K, Nyman A, Greko C, Bengtsson B. Antimicrobial resistance and virulence factors in *Escherichia coli* from Swedish dairy calves. *Acta Vet Scand.* 2012; 54:2. [PubMed: 22280887]
- Du X, Shen Z, Wu B, Xia S, Shen J. Characterization of class 1 integrons-mediated antibiotic resistance among calf pathogenic *Escherichia coli*. *FEMS Microbiol Lett.* 2005; 245:295–298. [PubMed: 15837385]
- Gillings M, Boucher Y, Labbate M, Holmes A, Krishnan S, Holley M, Stokes HW. The evolution of class 1 integrons and the rise of antibiotic resistance. *J Bacteriol.* 2008; 190:5095–5100. [PubMed: 18487337]
- Gotelli, NJ.; Chao, A. *Encyclopedia of Biodiversity.* 2. Vol. 5. Academic Press; Waltham, MA: 2013. Measuring and estimating species richness, species diversity, and biotic similarity from sampling data; p. 195-211.
- Gotelli NJ, Colwell RK. Quantifying biodiversity: Procedures and pitfalls in the measurement and comparison of species richness. *Ecol Lett.* 2001; 4:379–391.
- Gow SP, Waldner CL, Harel J, Boerlin P. Associations between antimicrobial resistance genes in fecal generic *Escherichia coli* isolates from cow-calf herds in western Canada. *Appl Environ Microbiol.* 2008; 74:3658–3666. [PubMed: 18424533]

- Guler O, Yanar M, Bayram B, Metin J. Performance and health of dairy calves fed limited amounts of acidified milk replacer. *S Afr J Anim Sci.* 2006; 36:149–154.
- Gulliksen SM, Jor E, Lie KI, Loken T, Akerstedt J, Osteras O. Respiratory infections in Norwegian dairy calves. *J Dairy Sci.* 2009; 92:5139–5146. [PubMed: 19762832]
- Hanekamp WJA, Smits AC, Wierenga HK. Open versus closed barn and individual versus group-housing for bull calves destined for beef production. *Livest Prod Sci.* 1994; 37:261–270.
- Hepola H. Milk feeding systems for dairy calves in groups: Effects on feed intake, growth and health. *Appl Anim Behav Sci.* 2003; 80:233–243.
- Hill TM, Bateman HG 2nd, Aldrich JM, Quigley JD, Schlotterbeck RL. Evaluation of ad libitum acidified milk replacer programs for dairy calves. *J Dairy Sci.* 2013; 96:3153–3162. [PubMed: 23466327]
- Hoelzer K, Cummings KJ, Warnick LD, Schukken YH, Siler JD, Grohn YT, Davis MA, Besser TE, Wiedmann M. Agar disk diffusion and automated microbroth dilution produce similar antimicrobial susceptibility testing results for *Salmonella* serotypes Newport, Typhimurium, and 4,5,12:I-, but differ in economic cost. *Foodborne Pathog Dis.* 2011; 8:1281–1288. [PubMed: 21877930]
- Hänninen L, Hepola H, Rushen J, de Passillé AM, Pursiainen P, Tuure V, Syrjälä-qvist L, Pyykkönen M, Saloniemi H. Resting behaviour, growth and diarrhoea incidence rate of young dairy calves housed individually or in groups in warm or cold buildings. *Acta Agric Scand A Anim Sci.* 2003; 53:21–28.
- Kaneene JB, Warnick LD, Bolin CA, Erskine RJ, May K, Miller R. Changes in multidrug resistance of enteric bacteria following an intervention to reduce antimicrobial resistance in dairy calves. *J Clin Microbiol.* 2009; 47:4109–4112. [PubMed: 19846639]
- Khachatryan AR, Besser TE, Hancock DD, Call DR. Use of a nonmedicated dietary supplement correlates with increased prevalence of streptomycin-sulfa-tetracycline-resistant *Escherichia coli* on a dairy farm. *Appl Environ Microbiol.* 2006; 72:4583–4588. [PubMed: 16820447]
- Kung L Jr, Demarco S, Siebenson LN, Joyner E, Haenlein GF, Morris RM. An evaluation of two management systems for rearing calves fed milk replacer. *J Dairy Sci.* 1997; 80:2529–2533. [PubMed: 9361225]
- Linton AH, Howe K, Pethiyagoda S, Osborne AD. Epidemiology of *Salmonella* infection in calves (1): Its relation to their husbandry and management. *Vet Rec.* 1974; 94:581–585. [PubMed: 4432360]
- Lorenz I, Earley B, Gilmore J, Hogan I, Kennedy E, More SJ. Calf health from birth to weaning. III. Housing and management of calf pneumonia. *Ir Vet J.* 2011; 64:14. [PubMed: 22018053]
- MacGregor-Fors I, Payton ME. Contrasting diversity values: Statistical inferences based on overlapping confidence intervals. *PLoS ONE.* 2013; 8:e56794. [PubMed: 23437239]
- Magurran, AE.; McGill, BJ. *Biological Diversity: Frontiers in Measurement and Assessment.* Oxford University Press; New York, NY: 2011.
- Martínez JL, Baquero F. Interactions among strategies associated with bacterial infection: Pathogenicity, epidemicity, and antibiotic resistance. *Clin Microbiol Rev.* 2002; 15:647–679. [PubMed: 12364374]
- Miller-Cushon EK, Bergeron R, Leslie KE, DeVries TJ. Effect of milk feeding level on development of feeding behavior in dairy calves. *J Dairy Sci.* 2013; 96:551–564. [PubMed: 23164223]
- O'Driscoll K, von Keyserlingk MA, Weary DM. Effects of mixing on drinking and competitive behavior of dairy calves. *J Dairy Sci.* 2006; 89:229–233. [PubMed: 16357286]
- Oikonomou G, Teixeira AG, Foditsch C, Bicalho ML, Machado VS, Bicalho RC. Fecal microbial diversity in pre-weaned dairy calves as described by pyrosequencing of metagenomic 16S rDNA. Associations of *Faecalibacterium* species with health and growth. *PLoS ONE.* 2013; 8:e63157. [PubMed: 23646192]
- Orden JA, Ruiz-Santa-Quiteria JA, Garcia S, Cid D, De La Fuente R. In vitro susceptibility of *Escherichia coli* strains isolated from diarrhoeic dairy calves to 15 antimicrobial agents. *J Vet Med B Infect Dis Vet Public Health.* 2000; 47:329–335. [PubMed: 10900823]
- Pereira RV, Santos TM, Bicalho ML, Caixeta LS, Machado VS, Bicalho RC. Antimicrobial resistance and prevalence of virulence factor genes in fecal *Escherichia coli* of Holstein calves fed milk with and without antimicrobials. *J Dairy Sci.* 2011; 94:4556–4565. [PubMed: 21854928]

- Svensson C, Liberg P. The effect of group size on health and growth rate of Swedish dairy calves housed in pens with automatic milk-feeders. *Prev Vet Med.* 2006; 73:43–53. [PubMed: 16191449]
- Svensson C, Lundborg K, Emanuelson U, Olsson SO. Morbidity in Swedish dairy calves from birth to 90 days of age and individual calf-level risk factors for infectious diseases. *Prev Vet Med.* 2003; 58:179–197. [PubMed: 12706057]
- USDA. United States Cattle and Calf Inventory: January 2011. United States Department of Agriculture; Washington, DC: 2011.
- Waltner-Toews D, Martin SW, Meek AH. Dairy calf management, morbidity and mortality in Ontario Holstein herds. III. Association of management with morbidity. *Prev Vet Med.* 1986; 4:137–158.
- Warnick VD, Arave CW, Mickelsen CH. Effects of group, individual, and isolated rearing of calves on weight gain and behavior. *J Dairy Sci.* 1977; 60:947–953.
- Wells SJ, Dargatz DA, Ott SL. Factors associated with mortality to 21 days of life in dairy heifers in the United States. *Prev Vet Med.* 1997; 29:9.
- Woodford ST, Whetstone HD, Murphy MR, Davis CL. Abomasal pH, nutrient digestibility, and growth of Holstein bull calves fed acidified milk replacer. *J Dairy Sci.* 1987; 70:888–891. [PubMed: 3584623]



**Figure 1.** Rarefaction curves for the number of *Escherichia coli* antimicrobial drug susceptibility phenotypes. Shaded area corresponds to a 95% CI. Calves housed in group pens (GP) had 3 to 25 calves fed free-choice acidified milk. Calves housed in individual pens (IP) had no direct contact with neighboring calves and were fed milk or milk replacer 2 to 3 times a day.



**Table 1**

Total number of calves, *Escherichia coli* isolates, and average age of calves in the study by calf-housing system

Description	No. of calves	No. of <i>E. coli</i> isolates	Mean age in days (minimum; maximum)
IP <sup>1</sup>			
AG-1 <sup>2</sup>	92	275	8 (0;14)
AG-2 <sup>3</sup>	128	379	26 (15; 35)
AG-3 <sup>4</sup>	83	249	48 (36; 65)
Total	303	903	
GP <sup>5</sup>			
AG-1 <sup>2</sup>	94	278	9 (1; 14)
AG-2 <sup>3</sup>	129	383	28 (15; 35)
AG-3 <sup>4</sup>	132	396	42 (37; 62)
Total	355	1,057	
Overall Total	658	1,960	

<sup>1</sup> Calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day.

<sup>2</sup> Calves with ages between 0 to 14 d.

<sup>3</sup> Calves with ages between 15 to 35 d.

<sup>4</sup> Calves with ages between 35 to 65 d.

<sup>5</sup> Calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.

**Table 2**Herd-level effect of calf-housing system on the proportion of antimicrobial-resistant *Escherichia coli*

Antimicrobial agent	PR <sup>1</sup>		P-value
	IP (n = 6)	GP (n = 6)	
Ampicillin	80.0	72.6	0.43
Cefoxitin	48.9	59.1	0.73
Ceftiofur	29.6	25.1	0.85
Ceftriaxone	8.0	4.0	0.43
Chloramphenicol	42.6	45.7	0.75
Ciprofloxacin	1.8	9.0	0.16
Gentamicin	21.2	5.8	0.73
Nalidixic acid	2.5	10.3	0.12
Neomycin	53.3	46.9	0.69
Streptomycin	83.1	72.9	0.18
Tetracycline	87.2	79.1	0.24
TMS <sup>2</sup>	45.4	54.3	0.49

<sup>1</sup> PR = proportion of calves with resistant *E. coli* isolates by calf-housing system (IP or GP); IP = calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day; GP = calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.

<sup>2</sup> Trimethoprim sulfamethoxazole.

Effect of calf-housing management (IP or GP) and calf age group (AG-1, AG-2 or AG-3) on *Escherichia coli* resistance while controlling for the random effect of herd and calf

Table 3

Antimicrobial agent	Resistance, %		Odds ratio, 95% CI		Resistance, %			Odds ratio, 95% CI		P-value <sup>10</sup>
	IP <sup>1</sup>	GP <sup>2</sup>	IP vs. GP <sup>3</sup>	P-value <sup>4</sup>	AG-1 <sup>5</sup>	AG-2 <sup>6</sup>	AG-3 <sup>7</sup>	AG-1 vs. AG-3 <sup>8</sup>	AG-2 vs. AG-3 <sup>9</sup>	
Ampicillin	72	59	7.0 (3; 18)	<0.0001	68	65	61	2 (0.8; 7)	1.0 (1; 10)	0.06
Cefoxitin	46	47	0.7 (0.3; 1)	0.4	52	43	45	2.0 (0.9; 6)	1.0 (0.5; 3)	0.14
Ceftiofur	33	27	2.0 (1; 5)	0.02	29	35	25	2.0 (0.8; 5)	4.0 (2; 10)	0.004
Ceftriaxone	10	5	2.0 (0.8; 7)	0.1	5	10	6	0.9 (0.4; 2)	0.8 (0.3; 2)	0.7
Chloramphenicol	36	35	1.0 (0.6; 2)	0.8	34	35	37	0.6 (0.3; 1)	0.7 (0.3; 2)	0.8
Ciprofloxacin	1	6	0.2 (0.1; 0.4)	<0.0001	0.3	4	7	0.1 (0.02; 0.2)	0.6 (0.3; 1)	<0.0001
Gentamicin	16	4	5.0 (1; 19)	0.03	7	8	12	0.5 (0.1; 2)	0.7 (0.1; 3)	0.7
Nalidixic acid	1	7	0.3 (0.1; 0.6)	0.001	0.5	4	9	0.1 (0.03; 0.3)	0.5 (0.2; 1)	0.006
Neomycin	32	32	1.0 (0.8; 1)	0.6	42	29	26	2.0 (2; 3)	1.0 (0.9; 1)	<0.0001
Streptomycin	70	58	4.0 (2; 7)	<0.0001	67	59	65	1.0 (0.5; 2)	0.5 (0.2; 1)	0.14
Tetracycline	77	70	2.0 (1; 5)	0.04	79	70	72	2.0 (0.8; 6)	0.6 (0.2; 1)	0.02
TMS <sup>11</sup>	33	37	0.6 (0.3; 1)	0.1	49	30	30	8.0 (3; 19)	1.0 (0.5; 2)	<0.0001

<sup>1</sup> IP = calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day.

<sup>2</sup> GP = calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.

<sup>3</sup> Odds ratio for resistance in IP when compared with GP.

<sup>4</sup> P-value for calf-housing management (IP or GP).

<sup>5</sup> Age group with a range between 0 to 14 d old (AG-1).

<sup>6</sup> Age group with a range between 15 to 35 d old (AG-2).

<sup>7</sup> Age group with a range between 36 to 65 d old (AG-3).

<sup>8</sup> Odds ratio for resistance in AG-1 when compared with AG-3.

<sup>9</sup> Odds ratio for resistance in AG-2 when compared with AG-3.

<sup>10</sup> P-value for age group (AG-1, AG-2, and AG-3).

Trimethoprim sulfamethoxazole.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 4**

Most common multidrug-resistant phenotypes among 1,960 *Escherichia coli* from calves by calf-housing system

Resistant phenotype <sup>1</sup>	No. of isolates	% of isolates
IP <sup>2</sup> (n = 903)		
AMP-FOX-TIO-STR-TET	49	5.4
AMP-FOX-CHL-STR -TET	45	4.9
AMP-FOX-CHL-GEN-STR-TET	31	3.4
Pansusceptible	106	11.7
GP <sup>3</sup> (n = 1057)		
CHL-STR-TET	78	7.3
AMP-FOX-TIO-NEO-STR-TET-COT	59	5.6
AMP-FOX-CHL-NEO-STR-TET-COT	42	4.0
Pansusceptible	193	18.2

<sup>1</sup> AMP = ampicillin; FOX= cefoxitin; TIO = ceftiofur; CHL = chloramphenicol; GEN = gentamicin; NEO = neomycin; STR = streptomycin; COT = trimethoprim sulfamethoxazole; TET = tetracycline.

<sup>2</sup> IP = calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day.

<sup>3</sup> GP = calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.

**Table 5**Mean richness, diversity, and biotic similarity of *Escherichia coli* antimicrobial drug susceptibility phenotypes

Description	IP <sup>1</sup> (n = 903 isolates)		GP <sup>2</sup> (n = 1,040 isolates)	
	Pen <sup>3</sup> (n = 29 pens)	Farm <sup>4</sup> (n = 6 farms)	Pen <sup>3</sup> (n = 29 pens)	Farm <sup>4</sup> (n = 6 farms)
Richness				
Chao index	11.5 (3.7; 19.3)	29.3 (8.5; 55.1)	10.1 (5.5; 13.4)	29.7 (16.8; 43.0)
Diversity				
Shannon index	1.6 (1.0; 2.0)	2.45 (1.5; 3.1)	1.5 (1.1; 1.8)	2.36 (1.9; 2.8)
Similarity, <sup>5</sup> %				
Shared <sup>6</sup>	33 (3; 80)	29 (11; 65)	37 (6; 73)	31 (13; 54)
Jaccard index	13. (2; 53)	11 (4; 27)	15 (2; 30)	12 (5; 20)
Sorensen index	17 (2; 57)	16 (6; 36)	21 (3; 39)	17 (6; 27)
Chao-Sorensen index	19 (3; 53)	16 (6; 34)	22 (4; 44)	18 (7; 30)

<sup>1</sup> IP = calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day.

<sup>2</sup> GP = calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.

<sup>3</sup> Mean richness, diversity, and similarity index estimated between calves within pens. In parenthesis are the minimum and maximum mean values at the farm level.

<sup>4</sup> Mean richness, diversity, and similarity index estimated between calves within farms. In parenthesis are the minimum and maximum mean values at the farm level.

<sup>5</sup> Similarity was calculated by comparing *E. coli* antimicrobial drug susceptibility phenotypes between pairs of calves. Indices are shown as a similarity percentage.

<sup>6</sup> Mean percent of resistance phenotypes shared by a pair of calves.



**Table 6***Salmonella* serotypes by calf-housing system, source, and farm for 44 environmental *Salmonella* isolates

Serovar	No. of <i>Salmonella</i> from environment		% <i>Salmonella</i> by sample <sup>1</sup>	% <i>Salmonella</i> by farm <sup>2</sup>
	Pen floor <sup>3</sup>	Nipple <sup>4</sup>		
IP <sup>5</sup> (n = 25)	25	—	41 (25/61)	83 (5/6)
Cerro	23	—	38 (23/61)	66 (4/6)
Liverpool	2	—	3 (2/61)	16 (1/6)
GP <sup>6</sup> (n = 19)	15	4	31 (19/61)	33 (2/6)
Cerro	7	4	18 (11/61)	33 (2/6)
Newport	8	0	13 (8/61)	16 (1/6)

<sup>1</sup> Percent of environmental samples culture positive for *Salmonella*.

<sup>2</sup> Percent of farms with at least one environmental sample culture positive for the specified *Salmonella* serovar.

<sup>3</sup> *Salmonella* isolates cultured from environmental samples of calf housing pen floors. A maximum of 1 isolate was cultured per sample.

<sup>4</sup> *Salmonella* isolates cultured from environmental samples of nipples used for ad libitum feeding of GP calves.

<sup>5</sup> IP = calves housed in individual pens without direct contact with neighboring calves and fed milk or milk replacer 2 to 3 times a day.

<sup>6</sup> GP = calves housed in group pens with 3 to 25 calves being fed free-choice acidified milk.