



HHS Public Access

Author manuscript

Nutr Metab Cardiovasc Dis. Author manuscript; available in PMC 2015 March 06.

Published in final edited form as:

Nutr Metab Cardiovasc Dis. 2014 April ; 24(4): 337–343. doi:10.1016/j.numecd.2013.12.008.

The low-carbohydrate diet and cardiovascular risk factors: Evidence from epidemiologic studies

T. Hu and L. A. Bazzano*

Department of Epidemiology, School of Public Health and Tropical Medicine, Tulane University,
New Orleans, LA 70112, USA

Abstract

Aims—Obesity is an important public health issue because of its high prevalence and concomitant increase in risk of cardiovascular diseases. Low carbohydrate diets are popular for weight loss and weight management but are not recommended in leading guidelines due to the perception that increases in dietary fat intake may lead to an adverse cardiovascular risk profile. To clarify the effects of a low-carbohydrate diet for weight loss on cardiovascular disease risk factors as compared to a low fat diet for weight loss, we systematically reviewed data from randomized controlled clinical trials and large observational studies.

Data synthesis—We searched the MEDLINE database (Jan 1966–Nov 2013) to identify studies that examined a low-carbohydrate diet as compared to a low-fat diet for weight loss or the improvement of cardiovascular disease risk factors.

Conclusions—Recent randomized controlled trials document that low-carbohydrate diets not only decrease body weight but also improve cardiovascular risk factors. In light of this evidence from randomized controlled trials, dietary guidelines should be re-visited advocating a healthy low carbohydrate dietary pattern as an alternative dietary strategy for the prevention of obesity and cardiovascular disease risk factors.

Keywords

Low carbohydrate diet; Cardiovascular disease; Risk factors; Weight loss

Introduction

Obesity is an important public health issue. In 2005, there were roughly 937 million overweight and 396 million obese people in the world, and the high prevalence is still projected to increase [1]. Further, obesity is a well-known risk factor for cardiovascular disease, type-2 diabetes and metabolic syndrome, and recently recognized as a disease by

© 2014 Published by Elsevier B.V.

*Corresponding author: Department of Epidemiology, School of Public Health and Tropical Medicine, Tulane University, 1440 Canal Street, SL-18, Suite 2000, New Orleans, LA 70112, USA. Tel.: +1 504 988 7323; fax: +1 504 988 1568. lbazzano@tulane.edu (L.A. Bazzano).

Disclosures

No interest conflict.

the American Medical Association. Therefore, the spread of obesity requires an intensive and effective intervention.

There has been increasing interest in the low-carbohydrate diet for weight loss and weight management. Nevertheless, the low-carbohydrate diet has never been recommended in leading guidelines because there is the perception that a low-carbohydrate diet will have adverse effects on cardiovascular disease risk factors, mostly due to the idea that saturated fats will be increased on a low-carbohydrate diet as compared to other weight-loss regimens [2–4]. Several professional organizations, such as the American Dietetic Association, have even cautioned against the use of a low-carbohydrate diet [5]. However, recent randomized controlled trials document that low-carbohydrate diets not only decrease body weight but also improve cardiovascular risk factors [6–9]. Since lowering the quantity of carbohydrate intake decreases total energy intake which is related to obesity [10], it remains unknown whether the cardiovascular effects of a low-carbohydrate diet are results of the decrease in dietary carbohydrates or total energy intake. To clarify this issue, we systematically reviewed data from randomized controlled clinical trials that examined the effects of low-carbohydrate diets as compared to an isocaloric low-fat diet for weight loss. We also reviewed data from large observational cohort studies to assess long-term associations of low carbohydrate diets with cardiovascular events, including specific food choices which may be included as a part of a low-carbohydrate dietary intervention and their potential influence on cardiovascular health.

Low-carbohydrate diets and cardiovascular risk factors: evidence from randomized controlled trials

We used the MEDLINE online database (Jan 1966–Nov 2013) to identify studies that examined the low-carbohydrate diet as compared to the low-fat diet. The following key words or medical subject headings in MEDLINE were used: (“low-carbohydrate diet”, “carbohydrate restriction”, “carbohydrate”, “fiber”, “body mass index”, “waist circumference”, “fat mass”, “obesity”, “diabetes”, “insulin resistance”, “blood glucose”, “hypertension”, “HDL”, “LDL”, “triglycerides”, “cholesterol”, “lipids”, “dyslipidemias”, “blood pressure”, “adipocytokines”, “inflammatory cytokines”, “C-reactive protein”, “heart diseases”, “cardiovascular diseases”, and “seizure”).

In this systematic literature review, we included randomized controlled trials conducted in adults, which compared a low-carbohydrate diet (45% of energy from carbohydrates) with a low-fat diet (30% of energy from fat) over an intervention duration of at least 6 months and reported cardiovascular risk factors as outcomes [11–13]. Thus we include both ketogenic and non-ketogenic low-carbohydrate diets in this review.

Body weight, body composition, and waist circumference

Randomized controlled trials have consistently shown that low-carbohydrate diets reduce body weight, with mean reductions ranging from 2.1 to 14.3 kg over at least 6 months of intervention [6,8,9,14–21]. The body weight reduction may not differ by level of carbohydrate restriction. A study comparing ketogenic diets with non-ketogenic low-

carbohydrate diets demonstrated that their efficacy at reducing body weight was equivalent over a 4-week intervention period [22], but long-term comparison has not been conducted. Moreover, both fat mass and lean mass were decreased but fat mass comprises the majority of weight lost on low-carbohydrate diets [14,15,17,18,20].

Compared to a low-fat diet, an isocaloric low-carbohydrate diet is at least as effective at reducing body weight. A 2006 meta-analysis including 5 randomized controlled trials showed that low-carbohydrate diets resulted in significantly more reduction in body weight than isocaloric low-fat diets at 6 months (net changes -3.3 kg, 95%CI: -5.3 to -1.4), but non-significantly more reduction at 12 months (net changes: -1.0 kg, 95%CI: -3.5 to 1.5) [23]. Those on low-carbohydrate diets experienced a similar reduction in fat mass and non-fat mass as those on low-fat diets, so that the changes in body composition were not significantly different between the two diets [9,14,15,17,18,20].

Low-carbohydrate diets may have the added benefit of decreasing the risk of central obesity. Brinkworth and colleagues reported that a low-carbohydrate diet decreased abdominal fat mass by approximately 30% as assessed using dual-energy X-ray absorptiometry scanning over a one-year intervention period compared to an isocaloric low fat diet [15]. While dual-energy X-ray absorptiometry scanning provides highly specific information on the anatomic location of fat, waist circumference is a much more commonly used measure of central obesity. Decreases in waist circumference on low-carbohydrate diets have ranged from 2.2 to 9.5 cm across studies and are comparable or greater than those seen on low fat diets [8,16,18].

Glucose, insulin, and HOMA

Most published randomized controlled trials also show that low-carbohydrate diets decrease fasting levels of blood glucose [9,14–16,18,21], HbA1c [18,19,24,25], serum insulin and/or HOMA [9,14–16,21] to a similar extent as seen on isocaloric low-fat diets. Moreover, reductions in insulin resistance do not appear to be different between ketogenic and non-ketogenic low-carbohydrate diets [22].

Lipids

Despite widely held beliefs, with the exception of a few studies, most randomized controlled trials have shown that low-carbohydrate diets increase HDL cholesterol and decrease total and LDL cholesterol [6,8,9,14–21]. Results from a 2006 meta-analysis did, however, show a greater increase in HDL (net change 3.1 mg/dL, 95%CI: -0.8 to 7.0) and a lesser reduction in total cholesterol (net change 10.1 mg/dL, 95%CI: 3.5 to 10.2) and LDL cholesterol (net changes 7.7 mg/dL, 95%CI: 1.9 to 13.9) on low-carbohydrate diets as compared to low-fat diets [23]. In addition, the ratio of total to HDL cholesterol decreased among those on low-carbohydrate diets, and this decrease was not significantly different from those seen on isocaloric low-fat diets in most trials [16,17,20,21] except for one study showing a greater decrease on the low-carbohydrate diet [8]. Moreover, studies consistently reported that the low-carbohydrate diet reduced triglycerides, and to a greater extent than the isocaloric low-fat diet (net change -31.0 mg/dL, 95%CI: -59.3 to -2.7) [6,8,9,14–21].

Despite fears of more atherogenic lipid profiles on low carbohydrate diets as compared to low-fat diets for weight loss, in general, the low-carbohydrate diet itself improved lipids profiles including total cholesterol, LDL cholesterol, HDL cholesterol, total/HDL ratio, and triglycerides. In fact, both low carbohydrate and low fat diets appeared to improve the lipids profiles, without strong evidence that either one was better.

Blood pressure

Current randomized controlled trials demonstrated that low-carbohydrate diets decrease systolic and diastolic blood pressure to a similar extent as isocaloric low fat diets [6,8,9,14–21]. This is logical as presumably, weight loss would be the driving factor in decreasing levels of blood pressure, and overall weight loss on both diets appears to be similar [23].

Emerging risk factors

The low-carbohydrate diets may also improve carotid endothelial function [26–29], inflammatory cytokines [30], C-reactive protein [27,30], homocysteine [27], and adipocytokines [28]. To date, however, most of the existing information comes from small clinical trials, and few studies have explored these relationships in larger samples.

In summary, low carbohydrate diets had beneficial effects on weight loss and cardiovascular risk factors, and these effects were comparable to those seen on low-fat diets in general. In addition, ketogenic low carbohydrate diets did not show a greater reduction in body weight or more improvement in cardiovascular risk factors than non-ketogenic low carbohydrate diets. Given that ketogenic low carbohydrate diets are more difficult to adhere to long-term, non ketogenic low carbohydrate diets may be more practical for inducing weight loss and improving cardiovascular risk factors. Due to the limited number of available trials, additional studies are needed to investigate the potential influence of age, gender, and race/ethnicity on the association.

Special populations

Diabetic populations

A literature review published in 2003 indicated that there was insufficient evidence to evaluate the use of low-carbohydrate diets among diabetic patients in part due to lack of data on long-term intervention [31]. Since then, several long-term trials of low-carbohydrate interventions have been conducted among diabetic patients [24,25,32]. All of these trials demonstrated that low-carbohydrate diets were as effective as isocaloric low-fat diets at reducing body weight and inducing favorable changes in lipids, blood pressure, blood glucose and plasma insulin among diabetic patients over at least 1 year of follow-up [24,32]. During the study intervention periods, no severe hypoglycemic episodes were reported. Based on this emerging evidence, low-carbohydrate diets could also be recommended among diabetic patients for the purpose of weight loss.

Children with seizure disorder

The ketogenic low-carbohydrate diet has been broadly used for seizure control in children with epilepsy [33–35]. At least one study indicated that children with seizure disorders may

prefer foods with a high fat content, compatible with a ketogenic low-carbohydrate diet [36]. However, the ketogenic low-carbohydrate diet may have adverse cardiovascular effects in epileptic children. Studies have shown that a ketogenic diet may be associated with excess risk of cardiac arrhythmias [37] and potentially unfavorable changes in total, LDL and HDL cholesterol over a two-year intervention [38]. Given these potential adverse effects, a ketogenic may not be advisable in children with seizure disorders.

Should we restrict all carbohydrates?

The source and quality of carbohydrates are additional factors in the decision of which and to what extent carbohydrates should be restricted. Carbohydrates derived from the intake of nutrient-rich fruits, vegetables, legumes and some whole grains are often accompanied by significant amounts of dietary fiber and tend to have a lower glycemic index than carbohydrates derived from the intake of refined grain products. Dietary fiber, which is abundant in many fruits, vegetables, whole grains, and legumes, cannot be broken down and absorbed by the body as are most carbohydrate fractions, and has been shown to delay the absorption of carbohydrates after a meal and thereby decrease the insulinemic response to dietary carbohydrates [39].

Many epidemiologic studies have documented that intake of dietary fiber, especially soluble fiber in cereal grains and some fruits, is inversely associated with risk of type 2 diabetes, coronary heart disease, myocardial infarction, congestive heart failure, and cardiovascular mortality [40–50]. For example, Schulze and colleagues conducted a meta-analysis of observational studies including 9 large cohorts and reported a reduced diabetes risk with higher dietary fiber intake from cereal grains (RR for the highest quartile [weighted median intakes: 11.6 g/day] vs. the lowest quartile [weighted median intakes: 3.7 g/day]: 0.67; 95%CI: 0.62 to 0.72) [51]. Intake of whole grains in general has been shown to be beneficial for the prevention of type 2 diabetes, ischemic stroke, cardiovascular disease, and all-cause mortality [52–67]. A meta-analysis of observational studies including 7 large cohorts suggested that, after adjustment for cardiovascular risk factors, greater whole grain intake (pooled average 2.5 servings versus 0.2 servings per day) was associated with 21% (OR: 0.79, 95%CI: 0.73 to 0.85), 24% (OR: 0.76, 95%CI: 0.69 to 0.83), and 17% (OR: 0.83, 95%CI: 0.68 to 1.02) lower risks of cardiovascular events, incident coronary heart disease, and incident stroke, respectively [68]. The exact underlying mechanisms that may account for the protective effects of whole grain intake are not clear but evidence suggests that whole grain intake may assist in regulation of in body weight, waist circumference, blood pressure and blood glucose, and improvement in lipid profile [69,70].

Unlike whole grain, refined grain products do not appear to have protective effects for cardiovascular and metabolic diseases. Investigators have examined the association of refined grain intake with cardiovascular and all-cause mortality using data from the Iowa Women's Health Study, Health Professional Follow-Up Study, and Nurses' Health Study. These studies did not detect a protective effect of refined grain intake [61,66,71]. A meta-analysis of observational studies including 3 large cohort studies showed no evidence for a protective effect on cardiovascular events when comparing high with low intakes of refined grain products (RR: 1.07; 95%CI: 0.94 to 1.22) [68]. In contrast observational studies have

shown that limiting white rice, as a main source of refined grain, by substituting brown rice or beans may lower the risk of metabolic syndrome or type 2 diabetes mellitus [72,73].

Thus, a healthy low-carbohydrate diet should include substantial amounts of dietary fiber; while refined grain products should not be recommended and should be restricted. In available clinical trials, a wide variety of low-carbohydrate diets have been tested, with different levels of dietary fiber and sources of carbohydrate within the same overall level of carbohydrate content. Future studies are needed to determine the effects of quality and source of carbohydrates within a carbohydrate restricted diet.

What food choices are best in a low-carbohydrate diet?

The choices of fat and protein sources in a low-carbohydrate diet may affect the risk of subsequent cardiovascular disease. Indeed, a prospective cohort study of 82,802 U.S. nurses reported that a low-carbohydrate dietary pattern which incorporated a high intake of vegetable protein and unsaturated fat was associated with a lower risk of coronary heart disease over 19 years of follow-up [74]. In contrast, a low-carbohydrate dietary pattern accompanied by a high intake of animal-based protein and fat was not associated with a lower risk of cardiovascular disease, but associated with higher all-cause mortality in both men and women [75].

Further, among sources of animal fat and protein red meat (especially meat from ruminants), a common source of both saturated fat and animal protein, and other sources such as fish and poultry may have different cardiovascular effects. Red meat, whether processed or unprocessed, has consistently been associated with elevated risks of cardiovascular events [54,76–85]. Rather than red meat, fish and poultry could be chosen because they are not related to excess risk of cardiovascular diseases and disorders, such as coronary heart disease, myocardial infarction, congestive heart failure, stroke, type 2 diabetes, atrial fibrillation, or hypertension [76,78,80,86–91].

Low-carbohydrate, fiber-rich and non-starchy vegetables are ideal substitutions for red meat products. In particular, several studies have reported that green leafy vegetables contributed to the apparent protective effect of total vegetables on ischemic stroke (RR comparing highest to lowest quintiles: 0.79; 95%CI: 0.62 to 0.99) [92]. Other researchers also reported that an increase of 1 serving/day in green leafy vegetable consumption was associated with significantly lower risk of type 2 diabetes (HR: 0.91; 95%CI: 0.84 to 0.98) and coronary heart disease (RR: 0.77; 95%CI: 0.64 to 0.93) [93,94]. Cruciferous vegetables have also been associated with a protective effect on ischemic stroke [92]. Olive, other vegetable oils, nuts and avocado are excellent sources of dietary unsaturated fats in a low-carbohydrate dietary pattern and may improve cardiovascular risk factors. For example a six-week crossover trial showed that consuming walnuts and flax oil decreased blood pressure and improved endothelial function [95]. Other studies have demonstrated that long-term, a high intake of vegetable oil is associated with a lower risk of stroke and type 2 diabetes mellitus [96,97]. In addition, intake of soy and soy products should also be encouraged because soy protein supplement can decrease blood pressure and lower lipids [98,99]. Moreover, high soy intake is associated with a lower risk of type 2 diabetes and cardiovascular diseases

[100]. For instance, Kokubo and colleagues reported that soy intake at least 5 times per week versus 0–2 times per week was associated with 45% and 69% lower risk of myocardial infarction and cardiovascular mortality among Japanese women [101]. In contrast, there is no evidence of favorable cardiovascular effects of starchy vegetables.

In summary, a healthy low-carbohydrate dietary pattern should emphasize dietary fiber intake derived from whole grains, fiber-rich fruit, low-carbohydrate vegetables (such as green leafy vegetables, legumes, and cruciferous vegetables), avocado, olive and vegetable oils, soy, fish and chicken, and restrict or eliminate consumption of processed and unprocessed red meat as well as starchy vegetables and refined grains.

Conclusion

A low-carbohydrate diet decreases body weight and improves cardiovascular risk factors. Reducing the quantity and improving the quality of carbohydrate intake by choosing healthy carbohydrate substitutes, such as olive oil, chicken, fish, avocado, green leafy vegetables, and soy products, may decrease the risk of cardiovascular disease in addition to producing weight loss. In light of recent evidence from randomized controlled trials, dietary guidelines should be re-visited advocating a healthy low-carbohydrate dietary pattern as an alternative dietary strategy for the prevention of obesity and cardiovascular disease.

References

1. Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)*. 2008; 32:1431–7. [PubMed: 18607383]
2. Hooper L, Summerbell CD, Higgins JP, Thompson RL, Capps NE, Smith GD, et al. Dietary fat intake and prevention of cardiovascular disease: systematic review. *BMJ*. 2001; 322:757–63. [PubMed: 11282859]
3. Hu FB, Stampfer MJ, Manson JE, Rimm E, Colditz GA, Rosner BA, et al. Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med*. 1997; 337:1491–9. [PubMed: 9366580]
4. Eckel RH, Jakicic JM, Ard JD, Hubbard VS, de Jesus JM, Lee IM, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines. *Circulation*. 2013 Nov 12. [Epub ahead of print].
5. Stein K. High-protein, low-carbohydrate diets: do they work? *J Am Diet Assoc*. 2000; 100:760–1. [PubMed: 10916510]
6. Foster GD, Wyatt HR, Hill JO, McGuckin BG, Brill C, Mohammed BS, et al. A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med*. 2003; 348:2082–90. [PubMed: 12761365]
7. Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med*. 2009; 360:859–73. [PubMed: 19246357]
8. Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med*. 2008; 359:229–41. [PubMed: 18635428]
9. Gardner CD, Kiazand A, Alhassan S, Kim S, Stafford RS, Balise RR, et al. Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight and related risk factors among overweight premenopausal women: the A to Z weight loss study: a randomized trial. *J Am Med Assoc*. 2007; 297:969–77.
10. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the

- American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation*. 2013 Nov 12. [Epub ahead of print].
11. American Heart Association. AHA dietary guidelines revision 2000: a statement for healthcare professionals from the Nutrition Committee of the American Heart Association. 2000; 102:2284–99.
 12. Institute of Medicine of the National Academies. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids (macronutrients). 2005.
 13. NHLBI Obesity Education Initiative Expert Panel on the identification, evaluation, and treatment of overweight and obesity in adults: clinical guidelines of the identification, evaluation, and treatment of overweight and obesity in adults. 1998:98–4083.
 14. Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA. A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Clin Endocrinol Metab*. 2003; 88:1617–23. [PubMed: 12679447]
 15. Brinkworth GD, Noakes M, Buckley JD, Keogh JB, Clifton PM. Long-term effects of a very-low-carbohydrate weight loss diet compared with an isocaloric low-fat diet after 12 mo. *Am J Clin Nutr*. 2009; 90:23–32. [PubMed: 19439458]
 16. Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, weight watchers, and zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA*. 2005; 293:43–53. [PubMed: 15632335]
 17. Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, et al. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet: a randomized trial. *Ann Intern Med*. 2010; 153:147–57. [PubMed: 20679559]
 18. Frisch S, Zittermann A, Berthold HK, Gotting C, Kuhn J, Kleesiek K, et al. A randomized controlled trial on the efficacy of carbohydrate-reduced or fat-reduced diets in patients attending a telemedically guided weight loss program. *Cardiovasc Diabetol*. 2009; 8:36. [PubMed: 19615091]
 19. Stern L, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, et al. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med*. 2004; 140:778–85. [PubMed: 15148064]
 20. Yancy WS Jr, Olsen MK, Guyton JR, Bakst RP, Westman EC. A low-carbohydrate, ketogenic diet versus a low-fat diet to treat obesity and hyperlipidemia: a randomized, controlled trial. *Ann Intern Med*. 2004; 140:769–77. [PubMed: 15148063]
 21. Lim SS, Noakes M, Keogh JB, Clifton PM. Long-term effects of a low carbohydrate, low fat or high unsaturated fat diet compared to a no-intervention control. *Nutr Metab Cardiovasc Dis*. 2010; 20:599–607. [PubMed: 19692216]
 22. Johnstone AM, Horgan GW, Murison SD, Bremner DM, Lobley GE. Effects of a high-protein ketogenic diet on hunger, appetite, and weight loss in obese men feeding ad libitum. *Am J Clin Nutr*. 2008; 87:44–55. [PubMed: 18175736]
 23. Nordmann AJ, Nordmann A, Briel M, Keller U, Yancy WS Jr, Brehm BJ, et al. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Arch Intern Med*. 2006; 166:285–93. [PubMed: 16476868]
 24. Elhayany A, Lustman A, Abel R, Attal-Singer J, Vinker S. A low carbohydrate Mediterranean diet improves cardiovascular risk factors and diabetes control among overweight patients with type 2 diabetes mellitus: a 1-year prospective randomized intervention study. *Diabetes Obes Metab*. 2010; 12:204–9. [PubMed: 20151996]
 25. Davis NJ, Tomuta N, Schechter C, Isasi CR, Segal-Isaacson C, Stein D, et al. Comparative study of the effects of a 1-year dietary intervention of a low-carbohydrate diet versus a low-fat diet on weight and glycemic control in type 2 diabetes. *Diabetes Care*. 2009; 32:1147–52. [PubMed: 19366978]
 26. Shai I, Spence JD, Schwarzfuchs D, Henkin Y, Parraga G, Rudich A, et al. DIRECT Group: dietary intervention to reverse carotid atherosclerosis. *Circulation*. 2010; 121:1200–8. [PubMed: 20194883]
 27. Keogh JB, Brinkworth GD, Noakes M, Belobrajdic DP, Buckley JD, Clifton PM. Effects of weight loss from a very-low-carbohydrate diet on endothelial function and markers of cardiovascular

- disease risk in subjects with abdominal obesity. *Am J Clin Nutr.* 2008; 87:567–76. [PubMed: 18326593]
28. Wycherley TP, Brinkworth GD, Keogh JB, Noakes M, Buckley JD, Clifton PM. Long-term effects of weight loss with a very low carbohydrate and low fat diet on vascular function in overweight and obese patients. *J Intern Med.* 2010; 267:452–61. [PubMed: 20141567]
 29. Buscemi S, Verga S, Tranchina MR, Cottone S, Cerasola G. Effects of hypocaloric very-low-carbohydrate diet vs. Mediterranean diet on endothelial function in obese women*. *Eur J Clin Invest.* 2009; 39:339–47. [PubMed: 19302563]
 30. Bladbjerg EM, Larsen TM, Due A, Stender S, Astrup A, Jespersen J. Effects on markers of inflammation and endothelial cell function of three ad libitum diets differing in type and amount of fat and carbohydrate: a 6-month randomised study in obese individuals. *Br J Nutr.* 2011; 106:123–9. [PubMed: 21320366]
 31. Bravata DM, Sanders L, Huang J, Krumholz HM, Olkin I, Gardner CD, et al. Efficacy and safety of low-carbohydrate diets: a systematic review. *JAMA.* 2003; 289:1837–50. [PubMed: 12684364]
 32. Iqbal N, Vetter ML, Moore RH, Chittams JL, Dalton-Bakes CV, Dowd M, et al. Effects of a low-intensity intervention that prescribed a low-carbohydrate vs. a low-fat diet in obese, diabetic participants. *Obesity (Silver Spring).* 2010; 18:1733–8. [PubMed: 20019677]
 33. Hartman AL, Gasior M, Vining EP, Rogawski MA. The neuropharmacology of the ketogenic diet. *Pediatr Neurol.* 2007; 36:281–92. [PubMed: 17509459]
 34. Klepper J, Scheffer H, Leiendecker B, Gertsen E, Binder S, Leferink M, et al. Seizure control and acceptance of the ketogenic diet in GLUT1 deficiency syndrome: a 2- to 5-year follow-up of 15 children enrolled prospectively. *Neuropediatrics.* 2005; 36:302–8. [PubMed: 16217704]
 35. Kang HC, Chung DE, Kim DW, Kim HD. Early- and late-onset complications of the ketogenic diet for intractable epilepsy. *Epilepsia.* 2004; 45:1116–23. [PubMed: 15329077]
 36. Amari A, Dahlquist L, Kossoff EH, Vining EP, Trescher WH, Slifer KJ. Children with seizures exhibit preferences for foods compatible with the ketogenic diet. *Epilepsy Behav.* 2007; 11:98–104. [PubMed: 17572157]
 37. Best TH, Franz DN, Gilbert DL, Nelson DP, Epstein MR. Cardiac complications in pediatric patients on the ketogenic diet. *Neurology.* 2000; 54:2328–30. [PubMed: 10881264]
 38. Kwiterovich PO Jr, Vining EP, Pyzik P, Skolasky R Jr, Freeman JM. Effect of a high-fat ketogenic diet on plasma levels of lipids, lipoproteins, and apolipoproteins in children. *JAMA.* 2003; 290:912–20. [PubMed: 12928468]
 39. Anderson JW, O'Neal DS, Riddell-Mason S, Floore TL, Dillon DW, Oeltgen PR. Postprandial serum glucose, insulin, and lipoprotein responses to high- and low-fiber diets. *Metabolism.* 1995; 44:848–54. [PubMed: 7616842]
 40. Khaw KT, Barrett-Connor E. Dietary fiber and reduced ischemic heart disease mortality rates in men and women: a 12-year prospective study. *Am J Epidemiol.* 1987; 126:1093–102. [PubMed: 2825519]
 41. Kokubo Y, Iso H, Saito I, Yamagishi K, Ishihara J, Inoue M, et al. JPHC Study Group. Dietary fiber intake and risk of cardiovascular disease in the Japanese population: the Japan Public Health Center-based study cohort. *Eur J Clin Nutr.* 2011; 65:1233–41. [PubMed: 21654702]
 42. Park Y, Subar AF, Hollenbeck A, Schatzkin A. Dietary fiber intake and mortality in the NIH-AARP diet and health study. *Arch Intern Med.* 2011; 171:1061–8. [PubMed: 21321288]
 43. Eshak ES, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, et al. JACC Study Group. Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. *J Nutr.* 2010; 140:1445–53. [PubMed: 20573945]
 44. Bazzano LA, He J, Ogden LG, Loria CM, Whelton PK. National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. Dietary fiber intake and reduced risk of coronary heart disease in US men and women: the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Arch Intern Med.* 2003; 163:1897–904. [PubMed: 12963562]
 45. Djousse L, Gaziano JM. Breakfast cereals and risk of heart failure in the physicians' health study I. *Arch Intern Med.* 2007; 167:2080–5. [PubMed: 17954802]

46. Mozaffarian D, Kumanyika SK, Lemaitre RN, Olson JL, Burke GL, Siscovick DS. Cereal, fruit, and vegetable fiber intake and the risk of cardiovascular disease in elderly individuals. *JAMA*. 2003; 289:1659–66. [PubMed: 12672734]
47. Liu S, Buring JE, Sesso HD, Rimm EB, Willett WC, Manson JE. A prospective study of dietary fiber intake and risk of cardiovascular disease among women. *J Am Coll Cardiol*. 2002; 39:49–56. [PubMed: 11755286]
48. Wolk A, Manson JE, Stampfer MJ, Colditz GA, Hu FB, Speizer FE, et al. Long-term intake of dietary fiber and decreased risk of coronary heart disease among women. *JAMA*. 1999; 281:1998–2004. [PubMed: 10359388]
49. Rimm EB, Ascherio A, Giovannucci E, Spiegelman D, Stampfer MJ, Willett WC. Vegetable, fruit, and cereal fiber intake and risk of coronary heart disease among men. *JAMA*. 1996; 275:447–51. [PubMed: 8627965]
50. Pietinen P, Rimm EB, Korhonen P, Hartman AM, Willett WC, Albanes D, et al. Intake of dietary fiber and risk of coronary heart disease in a cohort of Finnish men. The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study. *Circulation*. 1996; 94:2720–7. [PubMed: 8941095]
51. Schulze MB, Schulz M, Heidemann C, Schienkiewitz A, Hoffmann K, Boeing H. Fiber and magnesium intake and incidence of type 2 diabetes: a prospective study and meta-analysis. *Arch Intern Med*. 2007; 167:956–65. [PubMed: 17502538]
52. He M, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation*. 2010; 121:2162–8. [PubMed: 20458012]
53. Flint AJ, Hu FB, Glynn RJ, Jensen MK, Franz M, Sampson L, et al. Whole grains and incident hypertension in men. *Am J Clin Nutr*. 2009; 90:493–8. [PubMed: 19571218]
54. Nettleton JA, Steffen LM, Loehr LR, Rosamond WD, Folsom AR. Incident heart failure is associated with lower whole-grain intake and greater high-fat dairy and egg intake in the Atherosclerosis Risk in Communities (ARIC) study. *J Am Diet Assoc*. 2008; 108:1881–7. [PubMed: 18954578]
55. Kochar J, Djousse L, Gaziano JM. Breakfast cereals and risk of type 2 diabetes in the Physicians' Health Study I. Obesity (Silver Spring). 2007; 15:3039–44. [PubMed: 18198313]
56. de Munter JS, Hu FB, Spiegelman D, Franz M, van Dam RM. Whole grain, bran, and germ intake and risk of type 2 diabetes: a prospective cohort study and systematic review. *PLoS Med*. 2007; 4:e261. [PubMed: 17760498]
57. Wang L, Gaziano JM, Liu S, Manson JE, Buring JE, Sesso HD. Whole- and refined-grain intakes and the risk of hypertension in women. *Am J Clin Nutr*. 2007; 86:472–9. [PubMed: 17684221]
58. Bazzano LA, Song Y, Bubes V, Good CK, Manson JE, Liu S. Dietary intake of whole and refined grain breakfast cereals and weight gain in men. *Obes Res*. 2005; 13:1952–60. [PubMed: 16339127]
59. Jensen MK, Koh-Banerjee P, Hu FB, Franz M, Sampson L, Gronbaek M, et al. Intakes of whole grains, bran, and germ and the risk of coronary heart disease in men. *Am J Clin Nutr*. 2004; 80:1492–9. [PubMed: 15585760]
60. Montonen J, Knekt P, Jarvinen R, Aromaa A, Reunanen A. Whole-grain and fiber intake and the incidence of type 2 diabetes. *Am J Clin Nutr*. 2003; 77:622–9. [PubMed: 12600852]
61. Liu S, Manson JE, Stampfer MJ, Hu FB, Giovannucci E, Colditz GA, et al. A prospective study of whole-grain intake and risk of type 2 diabetes mellitus in US women. *Am J Public Health*. 2000; 90:1409–15. [PubMed: 10983198]
62. Liu S, Manson JE, Stampfer MJ, Rexrode KM, Hu FB, Rimm EB, et al. Whole grain consumption and risk of ischemic stroke in women: a prospective study. *JAMA*. 2000; 284:1534–40. [PubMed: 11000647]
63. Jacobs DR Jr, Meyer KA, Kushi LH, Folsom AR. Whole-grain intake may reduce the risk of ischemic heart disease death in post-menopausal women: the Iowa Women's Health Study. *Am J Clin Nutr*. 1998; 68:248–57. [PubMed: 9701180]
64. Meyer KA, Kushi LH, Jacobs DR Jr, Slavin J, Sellers TA, Folsom AR. Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr*. 2000; 71:921–30. [PubMed: 10731498]

65. Jacobs DR Jr, Meyer HE, Solvoll K. Reduced mortality among whole grain bread eaters in men and women in the Norwegian County Study. *Eur J Clin Nutr.* 2001; 55:137–43. [PubMed: 11305627]
66. Liu S, Sesso HD, Manson JE, Willett WC, Buring JE. Is intake of breakfast cereals related to total and cause-specific mortality in men? *Am J Clin Nutr.* 2003; 77:594–9. [PubMed: 12600848]
67. Fung TT, Hu FB, Pereira MA, Liu S, Stampfer MJ, Colditz GA, et al. Whole-grain intake and the risk of type 2 diabetes: a prospective study in men. *Am J Clin Nutr.* 2002; 76:535–40. [PubMed: 12197996]
68. Mellen PB, Walsh TF, Herrington DM. Whole grain intake and cardiovascular disease: a meta-analysis. *Nutr Metab Cardiovasc Dis.* 2008; 18:283–90. [PubMed: 17449231]
69. Anderson JW, Smith BM, Gustafson NJ. Health benefits and practical aspects of high-fiber diets. *Am J Clin Nutr.* 1994; 59:1242S–7S. [PubMed: 8172129]
70. Rave K, Roggen K, Dellweg S, Heise T, tom Dieck H. Improvement of insulin resistance after diet with a whole-grain based dietary product: results of a randomized, controlled cross-over study in obese subjects with elevated fasting blood glucose. *Br J Nutr.* 2007; 98:929–36. [PubMed: 17562226]
71. Jacobs DR, Pereira MA, Meyer KA, Kushi LH. Fiber from whole grains, but not refined grains, is inversely associated with all-cause mortality in older women: the Iowa women's health study. *J Am Coll Nutr.* 2000; 19:326S–30S. [PubMed: 10875605]
72. Mattei J, Hu FB, Campos H. A higher ratio of beans to white rice is associated with lower cardiometabolic risk factors in Costa Rican adults. *Am J Clin Nutr.* 2011; 94:869–76. [PubMed: 21813808]
73. Sun Q, Spiegelman D, van Dam RM, Holmes MD, Malik VS, Willett WC, et al. White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch Intern Med.* 2010; 170:961–9. [PubMed: 20548009]
74. Halton TL, Willett WC, Liu S, Manson JE, Albert CM, Rexrode K, et al. Low-carbohydrate-diet score and the risk of coronary heart disease in women. *N Engl J Med.* 2006; 355:1991–2002. [PubMed: 17093250]
75. Fung TT, van Dam RM, Hankinson SE, Stampfer M, Willett WC, Hu FB. Low-carbohydrate diets and all-cause and cause-specific mortality: two cohort studies. *Ann Intern Med.* 2010; 153:289–98. [PubMed: 20820038]
76. Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Willett WC, et al. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr.* 2011; 94:1088–96. [PubMed: 21831992]
77. Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Stampfer MJ, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch Intern Med.* 2012; 172:555–63. [PubMed: 22412075]
78. Ashaye A, Gaziano J, Djousse L. Red meat consumption and risk of heart failure in male physicians. *Nutr Metab Cardiovasc Dis.* 2011; 21:941–6. [PubMed: 20675107]
79. Aune D, Ursin G, Veierod MB. Meat consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Diabetologia.* 2009; 52:2277–87. [PubMed: 19662376]
80. Wang L, Manson JE, Buring JE, Sesso HD. Meat intake and the risk of hypertension in middle-aged and older women. *J Hypertens.* 2008; 26:215–22. [PubMed: 18192834]
81. Steffen LM, Kroenke CH, Yu X, Pereira MA, Slattery ML, Van Horn L, et al. Associations of plant food, dairy product, and meat intakes with 15-y incidence of elevated blood pressure in young black and white adults: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr.* 2005; 82:1169–77. quiz 1363–4. [PubMed: 16332648]
82. Steffen LM, Folsom AR, Cushman M, Jacobs DR Jr, Rosamond WD. Greater fish, fruit, and vegetable intakes are related to lower incidence of venous thromboembolism: the longitudinal investigation of thromboembolism etiology. *Circulation.* 2007; 115:188–95. [PubMed: 17179018]
83. Schulze MB, Manson JE, Willett WC, Hu FB. Processed meat intake and incidence of type 2 diabetes in younger and middle-aged women. *Diabetologia.* 2003; 46:1465–73. [PubMed: 14576980]

84. Wagemakers JJ, Prynne CJ, Stephen AM, Wadsworth ME. Consumption of red or processed meat does not predict risk factors for coronary heart disease; results from a cohort of British adults in 1989 and 1999. *Eur J Clin Nutr.* 2009; 63:303–11. [PubMed: 18000518]
85. Barr SI, McCarron DA, Heaney RP, Dawson-Hughes B, Berga SL, Stern JS, et al. Effects of increased consumption of fluid milk on energy and nutrient intake, body weight, and cardiovascular risk factors in healthy older adults. *J Am Diet Assoc.* 2000; 100:810–7. [PubMed: 10916520]
86. Nanri A, Mizoue T, Noda M, Takahashi Y, Matsushita Y, Poudel-Tandukar K, et al. Japan Public Health Center-based Prospective Study Group. Fish intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. *Am J Clin Nutr.* 2011; 94:884–91. [PubMed: 21775559]
87. van Woudenberg GJ, van Ballegooijen AJ, Kuijsten A, Sijbrands EJ, van Rooij FJ, Geleijnse JM, et al. Eating fish and risk of type 2 diabetes: a population-based, prospective follow-up study. *Diabetes Care.* 2009; 32:2021–6. [PubMed: 19675200]
88. Mozaffarian D, Longstreth WT Jr, Lemaitre RN, Manolio TA, Kuller LH, Burke GL, et al. Fish consumption and stroke risk in elderly individuals: the cardiovascular health study. *Arch Intern Med.* 2005; 165:200–6. [PubMed: 15668367]
89. Brouwer IA, Heeringa J, Geleijnse JM, Zock PL, Witteman JC. Intake of very long-chain n-3 fatty acids from fish and incidence of atrial fibrillation. The Rotterdam Study. *Am Heart J.* 2006; 151:857–62. [PubMed: 16569549]
90. Jarvinen R, Knekt P, Rissanen H, Reunanen A. Intake of fish and long-chain n-3 fatty acids and the risk of coronary heart mortality in men and women. *Br J Nutr.* 2006; 95:824–9. [PubMed: 16571163]
91. de Goede J, Geleijnse JM, Boer JM, Kromhout D, Verschuren WM. Marine (n-3) fatty acids, fish consumption, and the 10-year risk of fatal and nonfatal coronary heart disease in a large population of Dutch adults with low fish intake. *J Nutr.* 2010; 140:1023–8. [PubMed: 20335635]
92. Joshipura KJ, Ascherio A, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, et al. Fruit and vegetable intake in relation to risk of ischemic stroke. *JAMA.* 1999; 282:1233–9. [PubMed: 10517425]
93. Joshipura KJ, Hu FB, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med.* 2001; 134:1106–14. [PubMed: 11412050]
94. Bazzano LA, Li TY, Joshipura KJ, Hu FB. Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care.* 2008; 31:1311–7. [PubMed: 18390796]
95. West SG, Krick AL, Klein LC, Zhao G, Wojtowicz TF, McGuinness M, et al. Effects of diets high in walnuts and flax oil on hemodynamic responses to stress and vascular endothelial function. *J Am Coll Nutr.* 2010; 29:595–603. [PubMed: 21677123]
96. Brostow DP, Odegaard AO, Koh WP, Duval S, Gross MD, Yuan JM, et al. Omega-3 fatty acids and incident type 2 diabetes: the Singapore Chinese Health Study. *Am J Clin Nutr.* 2011; 94:520–6. [PubMed: 21593505]
97. Samieri C, Feart C, Proust-Lima C, Peuchant E, Tzourio C, Stapf C, et al. Olive oil consumption, plasma oleic acid, and stroke incidence: the three-city study. *Neurology.* 2011; 77:418–25. [PubMed: 21676914]
98. He J, Wofford MR, Reynolds K, Chen J, Chen CS, Myers L, et al. Effect of dietary protein supplementation on blood pressure: a randomized, controlled trial. *Circulation.* 2011; 124:589–95. [PubMed: 21768541]
99. Reynolds K, Chin A, Lees KA, Nguyen A, Bujnowski D, He J. A meta-analysis of the effect of soy protein supplementation on serum lipids. *Am J Cardiol.* 2006; 98:633–40. [PubMed: 16923451]
100. Villegas R, Gao YT, Yang G, Li HL, Elasy TA, Zheng W, et al. Legume and soy food intake and the incidence of type 2 diabetes in the Shanghai Women's Health Study. *Am J Clin Nutr.* 2008; 87:162–7. [PubMed: 18175751]
101. Kokubo Y, Iso H, Ishihara J, Okada K, Inoue M, Tsugane S. JPHC Study Group. Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in

Japanese populations: the Japan Public Health Center-based (JPHC) study cohort I. *Circulation*. 2007; 116:2553–62. [PubMed: 18025534]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript