

RESEARCH

Open Access

Diesel exhaust modulates ozone-induced lung function decrements in healthy human volunteers

Michael C Madden^{1,4*}, Tina Stevens^{1,3}, Martin Case¹, Michael Schmitt¹, David Diaz-Sanchez¹, Maryann Bassett¹, Tracey S Montilla¹, Jon Berntsen² and Robert B Devlin¹

Abstract

The potential effects of combinations of dilute whole diesel exhaust (DE) and ozone (O₃), each a common component of ambient airborne pollutant mixtures, on lung function were examined. Healthy young human volunteers were exposed for 2 hr to pollutants while exercising (~50 L/min) intermittently on two consecutive days. Day 1 exposures were either to filtered air, DE (300 µg/m³), O₃ (0.300 ppm), or the combination of both pollutants. On Day 2 all exposures were to O₃ (0.300 ppm), and Day 3 served as a followup observation day. Lung function was assessed by spirometry just prior to, immediately after, and up to 4 hr post-exposure on each exposure day. Functional pulmonary responses to the pollutants were also characterized based on stratification by glutathione S-transferase mu 1 (GSTM1) genotype. On Day 1, exposure to air or DE did not change FEV1 or FVC in the subject population (n = 15). The co-exposure to O₃ and DE decreased FEV1 (17.6%) to a greater extent than O₃ alone (9.9%). To test for synergistic exposure effects, i.e., in a greater than additive fashion, FEV1 changes post individual O₃ and DE exposures were summed together and compared to the combined DE and O₃ exposure; the p value was 0.057. On Day 2, subjects who received DE exposure on Day 1 had a larger FEV1 decrement (14.7%) immediately after the O₃ exposure than the individuals' matched response following a Day 1 air exposure (10.9%). GSTM1 genotype did not affect the magnitude of lung function changes in a significant fashion. These data suggest that altered respiratory responses to the combination of O₃ and DE exposure can be observed showing a greater than additive manner. In addition, O₃-induced lung function decrements are greater with a prior exposure to DE compared to a prior exposure to filtered air. Based on the joint occurrence of these pollutants in the ambient environment, the potential exists for interactions in more than an additive fashion affecting lung physiological processes.

Keywords: Diesel exhaust, Ozone, Co-exposure, Lung function, Greater than additive effects

Introduction

Numerous epidemiological studies have demonstrated an association between short-term exposure to ambient airborne particulate matter (PM) and adverse cardiopulmonary effects including premature mortality, increased hospitalizations for lung problems including infections, exacerbation of asthma symptoms, chronic bronchitis, and hospitalization for clinical cardiac events including arrhythmias, myocardial infarctions, and congestive heart

failure [1,2]. The health effects are more strongly associated with PM that is smaller than 2.5 µm, i.e. PM_{2.5}, which typically is derived from human based activities such as vehicular emissions. PM_{2.5} is a complex mixture of organic and inorganic compounds absorbed onto carbonaceous material with the composition varying across space and time. In this complex mixture of ambient air substances, the ubiquitous pollutants ozone (O₃) and diesel exhaust (DE) can be major and important components. DE can have "hotspots" such as bus terminals and major streets [3]. Levels of DE PM_{2.5} reached transient concentrations of several hundred µg/m³ during drive-by studies [4]. O₃ levels have generally been decreasing in the US, but can reach over 0.1 ppm on a regular basis.

* Correspondence: madden.michael@epa.gov

¹EPHD, NHEERL, US EPA, Research Triangle Park, Chapel Hill, NC 27711, USA

⁴U.S EPA Human Studies Facility, 104 Mason Farm Road, Chapel Hill, NC 27599-7315, USA

Full list of author information is available at the end of the article

Ozone is one of the best studied gas phase pollutants in terms of lung biological effects. With dozens of controlled human exposures, biological responses have been well documented particularly for induction of decrements of lung function and inflammation typically expressed as decreased FEV1 and FVC [5] and lung neutrophilia [6,7], in a concentration-response manner (typically 0.08-0.4 ppm). Repeated exposure studies with healthy young adults inhaling 0.4 ppm O₃ for 2 hrs have shown a decrement in lung function after the first exposure and an even greater decrement after a second day exposure, but then attenuated lung function decrements for the next 3 exposure days [8,9]. Some reports have linked certain genetic polymorphisms to possible sensitivities to O₃ exposure. For example, individuals exercising on a bike outdoors for 2 hr with ambient O₃ levels > 0.08 ppm had greater lung function decrements as a group when they possessed a glutathione-S-transferase mu 1 (GSTM1) null genotype coupled with wild type NQO1 compared to other combinations of those 2 gene polymorphisms [10]. BMI has also been proportionally associated with O₃-induced lung function decrements [11].

Lung function decrements are typically not induced by exposure to DE or PM alone in controlled study designs [12]. Controlled ambient PM exposure studies in humans have been used to examine biological responses as surrogates for understanding the health effects and mechanisms involved in the responses. A dose-dependent increase in total number of cells, neutrophils, and monocytes was observed in the lung lavage fluid in contrast to lung function. For instance, healthy human volunteers exposed to 300 µg/m³ DE for one hour with intermittent exercise resulted in marked systemic and pulmonary inflammatory responses [13,14].

There is interest in examining whether ambient pollutant gases can modify the biological effect induced by PM [15]. A handful of controlled exposure studies have shown an interactive effect of PM and O₃ exposure especially with blood and cardiac endpoints. For instance blood pressure increases have been noted with co-exposure of healthy and also asthmatic subjects to O₃ and concentrated ambient particles (CAPS) [16]. Increased vasoconstriction in humans occurred upon exposure to O₃ with CAPs [17]. Rodent models of multipollutant exposures have also been utilized to examine biological responses. In an allergic mouse model, both DE and O₃ were needed to increase lung resistance, which was not observed with individual pollutant exposures [18]. Exposures of rats to resuspended DE particles and O₃ induced changes in some aortic mRNA [19] suggesting changes induced by pollutant components traversing the lung.

There is the possibility that exposure to one pollutant may sensitize an individual in a manner that a biological response from an exposure to a second pollutant may be

enhanced. Sequential O₃ and DE exposures in healthy young adult participants in a controlled exposure setting has demonstrated that an initial exposure to 300 µg/m³ DE for 1 hr followed 5 hr later by a 2 hr exposure to 0.2 ppm O₃ induced more lung inflammation (e.g. neutrophils) than DE alone with no O₃ exposure [20]. The same research group, using a similar exposure regimen, showed that a prior exposure to DE increased O₃-induced lung inflammation relative to O₃ exposure alone [21]. However, these studies did not report on lung function changes. Combinations of exposure to sulfuric acid aerosols and O₃ have demonstrated slightly additive lung function changes in asthmatic subjects [22] but not in healthy individuals [23]. This study primarily examined whether combinations of exposures to O₃ and DE can induce additive or synergistic effects related to lung function changes in healthy, normal adults, whether exposure to DE alters the response to a subsequent exposure to O₃ approximately 20 hrs later, and whether GSTM1 genotype status has an effect on the lung function responses.

Materials and Methods

Population characteristics

15 healthy, non-smoking volunteers (4 female, 11 male) completed all four Day 1 exposure regimens (a-d; described below) out of 25 subjects enrolled. Volunteers ranged in age from 23 to 36 (mean 27), had no history of asthma, chronic respiratory or acute illnesses within 4 weeks, and normal lung function (greater than 75% of predicted values for FVC, FEV1, and FEV1/FVC ratio). The volunteers refrained from taking non-steroidal anti-inflammatory medications 48 hr prior to each exposure. The use of vitamin C and E were not permitted for 2 weeks prior to exposure. [See Additional file 1 (Subject Selection and Participation Criteria) for full subject inclusion and exclusion criteria and subject participation requirements as well as the study purpose]. Participants were informed of the procedures and likely potential risks and each signed a statement of informed consent. Subject participation was strictly voluntary and they could withdraw at any time. The protocol and consent form were approved by the University of North Carolina School of Medicine Committee on the Protection of the Rights of Human Subjects and the US EPA [ClinicalTrials.gov # NCT01874834]. Once consented, volunteers were trained to correctly perform pulmonary function tests, BMI calculated, and participants exercised on a recumbent bicycle to determine a load that would elicit minute ventilation of 25 L/min/m² body surface area (BSA) (~50 L/min for most subjects). BSA and BMI (in kg/m²) was calculated by the spirometry software from a subject's height and weight. GSTM1 status was determined from isolated peripheral white blood cell DNA (QIAamp DNA Mini Kit (Qiagen Inc.,

Valencia, CA)) using RT PCR [24]; blood was drawn via standard venipuncture during the training day before any exposures occurred. One subject did not consent to genotyping and was not utilized in the analyses by genotype group. Subject characteristics are presented in Table 1.

Study design

This was a randomized crossover single blind study with 4 arms (designated as a-d) separated by at least 13 days between arms. Each arm consisted of 2 consecutive exposure days and a follow-up visit (Figure 1). On Day 1 participants were exposed for 2 hr with intermittent exercise to arm a) filtered air (FA) n = 3 (where n is the initial exposure); arm b) DE (300 ug/m³ target) n = 4; arm c) O₃ (0.300 ppm target) n = 4; or arm d) DE + O₃, n = 4. On Day 2 (~22 hr post Day 1 exposure) all research subjects were exposed to O₃ (0.300 ppm target) for 2 hrs. A follow-up visit occurred on Day 3. No two subjects received the same sequence of arms (as there were 24 possible sequences). Prior to the first exposure (Day 1, pre), lung function was measured on a 10.2-L dry seal digital spirometer interfaced to a computer (SensorMedics Model 1022; SensorMedics; Palm Springs, CA) with subjects standing. Subjects then entered a stainless steel exposure chamber (approximately 1.83 m × 1.83 m × 2.44 m, ~8.17 m³; ~650 ft³/min flow; air change time ~ 27 sec), and were exposed to filtered air, DE, O₃, or DE + O₃ for 2 h. During the exposure, subjects exercised on a recumbent bicycle ergometer for 15 min intervals

beginning after the first 15 min of exposure and resting for 15 min after each exercise time for a total of 1 hr of exercise per exposure. Ventilation rate was measured approximately 6 min into each exercise session and work load adjusted if needed to a target of 25 L/min/m² BSA. Heart rate, ECG, and SpO₂ (using pulse oximetry) were monitored continuously during the exposure. Immediately following the exposure (Day 1, post), subjects underwent pulmonary function testing. Lung function measurements were obtained once an hour for the next 4 hrs. On Day 2, the procedures described for Day 1 were repeated except that participants entered a different chamber (~2.39 m × 2.16 m × 3.20 m, ~ 16.52 m³; 450 ft³/min) for an O₃ exposure. The following morning (Day 3, follow-up), there was a final round of pulmonary function testing. Subjects who had a large Day 1 decrement in lung function were not exposed on Day 2, but did return for follow-up lung function testing.

Exposure conditions

All exposures were carried out at the EPA Human Studies Facility in Chapel Hill, NC. The targeted exposure atmosphere was approximately 40 ± 10% RH and 22 ± 2°C. Outdoor air was passed through an air purification system consisting bed of Purifil (potassium permanganate on alumina), a bed of Purikol (activated charcoal), dehumidified, and passed over an unheated bed of Hopcalite and then through a HEPA filter to ensure no detectable presence of pollutant gases (O₃, NO, NO₂, SO₂, CO, and hydrocarbons with the exception of methane) in the FA. The air was then heated, humidified, and passed through a second HEPA filter before O₃ and DE were added. The concentrations of the pollutant gases and PM were monitored continuously. The DE was generated from a Cummins diesel engine/generator QSB7 series (model DSGAA) running at 50 percent load that was located outside the Facility, and subsequently introduced into the exposure chamber after dilutions with humidified FA to give a chamber concentration target of approximately 300 µg/m³. The DE particle concentration was controlled by modulating the flow of medical grade compressed air into a Vortec Transvector Air Flow Amplifier (Model 901XSS; Cincinnati OH) with real time TEOM monitoring of PM mass concentration. Diesel fuel used for the study was a commercially obtained ultra low sulfur (<15 ppm) fuel. Particle mass, numbers, and size characteristics were measured in real time by a DataRAM™, a Tapered Element Oscillating Microbalance (TEOM®), and a Condensation Particle Counter (CPC). The CPC TSI 3022A model (detection size limit 7 nm) was used for the first 11 subjects' exposures, and a TSI 3775 model (detection size limit 4 nm) was used for the last 4 subjects' exposures for detection of particle number concentration. A Scanning Mobility Particle Sizer (SMPS™) (Shorewood MN) as also used to measure particulate

Table 1 Subject characteristics of participants in the study

Gender	Race	Age	BMI	GSTM1 status
f	c	24	26.84	Pos
f	bl	27	32.16	Pos
m	c	37	24.19	Pos
m	c	27	24.37	Null
m	bl	36	28.47	Null
m	c	27	27.08	Pos
m	hisp	24	29.94	Pos
m	hisp	24	27.15	Null
m	c	23	25.22	Nos
m	bl	23	29.37	Pos
f	c	25	25.66	Null
f	bl	29	38.06	Pos
m	c	30	21.22	Pos
m	c	28	28.94	Null
m	c	25	25.72	nd
11 m/4f	9c/4bl/2hisp	27 ± 4	27.63 ± 3.95	9+/5-

nd = not determined.

Arm	Day 1	Day 2	Day 3
A	Air	O ₃	FU
B	DE	O ₃	FU
C	O ₃	O ₃	FU
D	DE + O ₃	O ₃	FU

Figure 1 Design of Study. Subjects were exposed to target levels of either filtered air, 0.300 ppm O₃, 300 µg/m³ DE, or combined DE (300 µg/m³) and O₃ (0.300 ppm) on Day 1 for 2 hr. On Day 2 subjects were exposed to 0.300 ppm O₃ for 2 hr. During exposures on Days 1 and 2 subjects performed four 30 min cycles of rest/exercise at ~50 L/min. Day 3 was a follow-up (F/U) day for measurements and no pollutant exposures occurred.

characteristics during subject exposures. The SMPS data were used to determine particle size distribution as a number median diameter in all size bins. The number median diameter is the particle size (µm) bin where the median of the total count number occurs. Particle size distribution was determined at regular intervals. A Versatile Air Pollution Sampler (VAPS) (URG, Chapel Hill NC) configured with one quartz and two teflon filters with flow controlled by a mass flow controller sampled inlet air to the chamber. After exposure and filter weight determination, filter samples were analyzed for chemical composition of particles. Elemental (EC) and Organic (OC) Carbon content were determined using a thermo-optical method based on sequential pyrolytic vaporization and detection by transmittance using a carbon analyzer (Model 107A, Sunset Laboratory Inc., Hillsborough, NC) [25]. A bacterial endotoxin characterization test on particles was conducted by Cape Cod Inc. (East Falmouth, Massachusetts) using a Limulus polyphemus gel clot method to test for endotoxin contamination following their standard operating procedures.

Ozone was generated by passing USP grade oxygen through a silent arc O₃ generator (Innovatee Geratechnik GmbH, Rheinbach Germany) (Model CMG 10-5 for the DE chamber (Day 1) and Model 500 for the O₃ chamber on Day 2). For Day 1 exposures O₃ entered the filter air stream before DE. Concentrations were controlled by a computer feedback to the O₃ generator with a 0-5 V signal allowing internal modulation control of the O₃ generator. Both DE and O₃ were delivered into the exposure chamber within 8 sec to minimize DE loss. Increased production of O₃ was needed when DE was present as NO reacted with O₃.

Statistics

The primary hypothesis of this study was to examine whether combinations of exposures to O₃ and DE induce synergistic effects on lung function and whether exposures to DE sensitize individuals for a larger response to O₃ than does the exposure by air. To evaluate changes between pollutant and filtered air exposures we used a full two-factor (arm b plus arm c (O₃ plus DE),

and arm d (O₃ + DE)) mixed effects model with a subject-specific random intercept to account for repeated measures. To examine the first hypothesis with respect to synergism of effects, changes in lung function were examined following Day 1 exposures. To examine whether exposures to DE sensitize individuals for a larger response to O₃, lung function was examined following O₃ exposure on Day 2. In both analyses changes in lung function were contrasted to null hypothesis of no change relative to pre exposure baseline and relative to air exposure as control. Prior to the analysis, lung function endpoints FEV₁ and FEV were normalized and expressed as percent change from the pre-exposure values (Post/Pre-1). All 15 subject subjects completed four exposures on Day 1. Following O₃ exposure on day 1 some subjects responded with large drop in lung function and were not asked to undergo a Day 2 O₃ exposure. On day 2, 15 subjects completed O₃ exposures following air and DE (arms a and b, respectively), 14 of these completed O₃ exposure following O₃ on Day 1 (arm c), and 11 completed Day 2 O₃ exposure following of the combined O₃ & DE exposure (arm d). We used all available data in the analysis and did not account for possibly non-random missing values in subjects who responded strongly to O₃ or O₃ & DE on day 1. Estimated mean changes in lung function within each exposed group at each time point are expressed as mean change ± standard error, and a p-value where applicable. All statistical analyses were performed using R statistical software, with nlme and gmodels packages to estimate the effect (version 3.0.1; R Development Core Team; <http://www.r-project.org/>). A *p* value of less than 0.05 was considered statistically significant.

Results

Exposure parameters

The subjects were exposed to 290 µg/m³ DE and 0.298 ppm O₃ exposure on average during individual pollutant exposure days (Table 2). On co-exposure days, the values were very similar; 291 µg/m³ and 0.296 ppm for DE and O₃ respectively. The values for other measured parameters (Table 2; CO, and TH) were also similar for the DE and DE + O₃ exposures. When both O₃ and DE were present together, there was a decrease in NO and a

Table 2 Mean Day 1 pollutant physicochemical parameters during the four exposure scenarios

Parameter	Air	O ₃	DE	DE + O ₃
PM (µg/m ³) ^a	1.0 ± 1.5	1.0 ± 1.1	297.1 ± 20.8	294.3 ± 20.1
PM # (x10 ³ /cc)	<1	<1	731 ± 90 ^b (11) 977 ± 55 ^c (4)	734 ± 50 ^b (11) 903 ± 306 ^c (4)
Median PM size				
Volume (µm)	----	----	0.200 ± 0.007	0.200 ± 0.007
Number (µm) ^d	----	----	0.064 ± 0.006	0.066 ± 0.003
O ₃ (ppm)	0.00 ± 0.00	0.30 ± 0.00	0.01 ± 0.00	0.30 ± 0.00
CO (ppm)	0.09 ± 0.10	0.09 ± 0.10	2.58 ± 0.25	2.51 ± 0.32
NO (ppm)	0.00 ± 0.01	0.01 ± 0.03	1.58 ± 0.26	0.03 ± 0.04
NO ₂ (ppm)	0.00 ± 0.00	0.00 ± 0.00	0.16 ± 0.05	1.72 ± 0.22
THC (ppm)	2.05 ± 0.10	1.98 ± 0.10	2.35 ± 0.14	2.24 ± 0.19
SO ₂ (ppm)	<0.010	<0.010	0.013	<0.010

mean ± Std Dev except for Median PM size; n = 15, except for PM# where n is indicated.

^aGravimetric weight; average of two teflon filters.

^bCPC Model TSI 3022A.

^cCPC Model TSI 3775.

^dparticle size (µm) bin where the median of the particle counts occurred.

concomitant increase in NO₂. On Day 2, when subjects were exposed to O₃ alone, the concentration was maintained within 0.010 ppm of the targeted concentration. The median DE number and volume particle size (64 nm and 200 nm, respectively) did not change with O₃ present (number size 66 nm and volume size 200 nm, respectively). There were 731 and 734 (x10³) particles/cc for DE and DE + O₃ exposures, respectively when one CPC model was used (n = 11); the numbers increased when the second model was used (n = 4) due to a different detectable size limit. Particle EC and OC concentrations on sample filters were 39.2 ± 4.6 µg/cm² and 14.2 ± 3.5 µg/cm², respectively, i.e., ~ 3:1 ratio. PM from both DE and DE + O₃ exposures had nondetectable levels of endotoxin.

Lung function after exposures to pollutants on Day 1

Fifteen subjects completed all four of the Day 1 pollution exposure scenarios. In these subjects, decreases (relative to their pre-exposures values) in FEV1 were not observed immediately after exposure to either FA or whole DE (Figure 2A). Measurement of lung function revealed no decrement in DE or air exposed subjects for up to 4 hrs post-exposure. Exposure to O₃ induced an average decrement of 9.9 ± 2.5% measured immediately post-exposure (p < 0.05 vs air exposure). Exposure to the combination of O₃ and DE induced a decrement of 17.6 ± 4.6% and was statistically different from both air and O₃ alone exposure (p < 0.01). A greater than additive effect of the DE and O₃ exposure (arm d) on FEV1 was examined by statistical comparison to the FEV1 changes following O₃ (arm c) added to the change following DE exposure (arm b). The analysis had a p value of 0.057, close to statistical significant. FEV1 values returned to


pre-exposure values by 4 hr post-exposure in the O₃ and DE + O₃ groups at similar rates.

Among the Day 1 exposures statistically significant (p < 0.01) FVC decrements were observed in O₃-exposed subjects (8.7 ± 2.3%) and DE + O₃ exposed subjects (11.8 ± 3.0%) immediately post-exposure compared to FA exposures (Figure 2B). The FVC decrements immediately post O₃ compared to post DE + O₃ exposure were not statistically different (p = 0.10), nor was greater than additive effects observed when FVC changes in arm d were compared to the sum of FVC changes post O₃ and DE. Exposure to either air or DE alone did not alter FVC from pre-exposure values. FVC values returned to pre-exposure levels by 4 hrs post-exposure for all exposure groups with no difference in the recovery rate (data not shown).


Analysis stratified by GSTM1 genotype showed relatively small and non-significant decrease in FEV1 in GSTM1null individuals (n = 5) (0.8 ± 0.7%) immediately post DE exposure compared to individuals with a GSTM1+ genotype who did not have a decrease (FEV1 increased 1.3 ± 0.5%; n = 9) (Figure 3). Immediately after the O₃ exposure, the group of GSTM1null genotype individuals had a smaller, and statistically non-significant, decrease in FEV1, 3.70 ± 2.0% compared to a 13.7 ± 4.2% decrement in the GSTM1 + group. Using linear regression, an individual's BMI did not significantly correlate with their observed O₃-induced or DE and O₃-induced FEV1 or FVC change.

Lung function after exposure to O₃ on Day 2

After Day 1 exposures, subjects returned to the facility and most were exposed to about 0.300 ppm O₃. For lung function measured immediately post-exposure, subjects in arm a (exposed to air on day 1) had a statistically




significant O₃-induced decrease in FEV₁ ($10.9 \pm 2.6\%$) and FVC ($7.8 \pm 1.8\%$) immediately post-exposure when compared to pre-exposure values (Figure 4A and 4B). Exposure to O₃ after the previous day exposure to DE (arm b) induced statistically significant decrements of $14.7 \pm 3.3\%$ for FEV₁ and $9.2 \pm 1.9\%$ for FVC (both $p < 0.05$). The Day 2 O₃-induced FEV₁ and FVC decrements for



each subject after the Day 1 exposure to DE compared to air showed that prior exposure to DE on Day 1 induced a statistically significant greater FEV₁ decrement on Day 2 ($p < 0.05$) (Figure 5).

Some subjects were not sequentially exposed to Day 2 O₃ either due to their request or due to their large decrement in FEV₁ immediately after either O₃ ($n = 1$) or DE + O₃ ($n = 4$) on Day 1 because a second sequential O₃ exposure within 24 hr is expected to typically induce an even larger FEV₁ decrement than the first day exposure in most subjects. Consequently, for those five subjects Day 2 procedures were follow-up measurements and observations, similar to Day 3 measurements for the other 10 subjects. Subjects in this study had a greater O₃-induced lung function decrements if they received two consecutive days of O₃ exposure (arm c); consecutive O₃ exposures induced a statistically significant greater decrement in FEV₁ ($18.2 \pm 4.5\%$) than the decrement associated immediately after the first day exposure, (i.e., $10.3 \pm 3.3\%$), ($p < 0.05$) (Figure 4A) or compared to the Day 1 air exposure, then O₃ on Day 2 (i.e., $10.9 \pm 2.6\%$). The FVC decrement ($7.8 \pm 1.8\%$) on Day 2 following air (arm a) (Figure 4B) was similar to the Day 1 FVC decrement (8.7%) after O₃ (Figure 2B). For subjects exposed to DE + O₃ (arm d) the FEV₁ decrement immediately after exposure to O₃ on Day 2 was $13.8 \pm 2.2\%$, which was similar to the decrement in these subjects the previous day immediately after exposure (i.e., $17.6 \pm 4.6\%$). The FVC decrement on Day 2 (Figure 4B) was similar to the Day 1 FVC decrement as well (Figure 2B). Both FEV₁ and FVC values returned to pre-exposure values by 4 hr post exposure (data not shown).


When FEV₁ responses to the Day 2 O₃ exposure were segregated by GSTM1 genotype, it was observed that



those subjects with the null genotype had a smaller decrement ($3.8 \pm 3.8\%$) than positive genotypes ($14.9 \pm 3.6\%$) after the FA exposure day; however this difference was not significant. This is a consistent effect with the observed Day 1 responses to O₃ exposure between the two groups. No other Day 2 FEV₁ differences were observed between genotypes for the other 3 exposure regimens from Day 1.

Discussion

This study examined alterations in lung function after co-exposures of DE and O₃, as well as sequential exposures to DE and O₃ separated by one day. For the former exposure regimen, DE alone did not induce lung function



as measured by FEV₁ and FVC decrements, but O₃ did induce decrements after 2 hr reaching statistical significance for FEV₁. Exposure of the subjects to the combination of the two pollutants (arm d) induced a greater FEV₁ decrement than either pollutant alone. If the pollutant-induced FEV₁ changes are added together, i.e., the sum of the changes in FEV₁ post O₃ and post DE exposures, and compared to the combined exposure (arm d), the p value of 0.057 is obtained and is close to statistical significance (Figure 2A). A possible explanation for the synergistic effect induced by DE + O₃ co-exposure may be that the lung function change is related to the chemical composition in the atmosphere containing O₃ and DE being different from DE or O₃ alone. DE-derived particle size did not change with the presence of O₃, and chamber temperature and humidity were kept similar between the two exposure regimens, suggesting that particle deposition pattern in the lung was not likely altered due to a size change. However, the possibility that other PM physicochemically related characteristics were altered, such as a change in charge or hydrophobicity that could alter the deposition pattern cannot be excluded. For all exposures regimens, lung function returned to pre-exposure values by 4 hrs after the end of the exposure, similar to the well characterized and transient nature of O₃-induced lung function changes.

With respect to component changes in the gas phase relating to the greater than additive effect of O₃ and DE on FEV₁ decrement, the NO₂ level increased while the NO level decreased. However, NO₂ alone generally does not induce lung function changes even up to 4 ppm in healthy individuals [26] though we cannot totally exclude the possibility that it contributed to changes in lung function seen during co-exposure to DE and O₃. O₃ concentrations were kept at ~0.300 ppm with or without DE

present, so the only other components with potentially altered concentrations were derived from DE. If lung function responses to O₃ exposure are mediated at least in part via C-fiber activation [27] then chemical components affecting C-fibers activity directly (but below a threshold for discharge), or indirectly by, e.g., lowering the fiber threshold for electrical discharge, might be expected to induce the synergistic change in FEV1 seen upon concurrent O₃ and DE exposure. Some carbonyls, (e.g., acrolein) and carboxylic acids (e.g., acetic acid) have been implicated in affecting pulmonary C-fiber activity [28-30]. There are several aldehydes identified in DE including acrolein, and there is increased formation of carbonyls and carboxylic acids observed upon O₃ reacting with DE particles surface components [31]. There is evidence for a possible increased carbonyls and/or carboxylic acids formation and induction of C-fiber sensitization in the DE and O₃ atmosphere, but other classes of compounds cannot be eliminate either.

Subjects in this study receiving DE exposure the day before the Day 2 O₃ exposure had greater FEV1 decrements than when they had a Day 1 clean air exposure (Figure 5). It is uncertain whether only the DE gas phase or particle phase, or both phases, are needed to induced the augmented response to a subsequent O₃ exposure. Most previous studies examining sequential exposure effects of air pollutants on O₃-induced lung effects changes were typically performed within 6 hr of the initial exposure [20,21,32]. A report demonstrated that prior exposure for 2 hr to 0.6 ppm NO₂ with intermittent exercise (~40 L/min) did not significantly alter lung function in the young, healthy subjects; however compared to air as the initial exposure, the pre-exposure to NO₂ induced a 1.8% greater FEV1 decrement upon a 2 hr exposure to 0.300 ppm O₃ 3 hr post NO₂ exposure [32]. Another study reported that a same day follow up to approximately 0.200 ppm O₃ added to the lung inflammatory response (i.e., increased neutrophilia and myeloperoxidase) induced by an initial 300 ug/m³ DE exposure compared to an initial air exposure; but lung function was not examined in that study [21]. A possible mechanism involved with the augmented FEV1 change to O₃ on Day 2 in this study may involve, at least in part, altered production of prostaglandins E₂ (PGE₂) and F_{2α} (PGF_{2α}) which have been associated with O₃-induced lung function decrements [33]. Both prostaglandins can alter lung C-fiber activity directly inducing discharge and/or altering the threshold for discharge to an agonist [34] The possible altered AA metabolism in response to sequential DE and O₃ exposures (i.e., increased AA release, decreased re-uptake, and COX-2 protein for increased conversion of AA to PGE₂ and F_{2α}), based upon *in vivo* and *in vitro* findings [35-39], may explain, at least in part, the observed greater FEV1 decrement. It

should be noted that an adequate duration of time is needed for COX-2 protein to be upregulated [40]. Additionally the literature suggests that the DE particle phase is involved in this possible biochemical mechanism (i.e., increased COX-2), but does not necessarily negate the importance of DE gas phase components.

Factors that affect an individual's sensitivity to O₃-induced lung function decrements are not clearly understood. Increased BMI has been linked to a greater O₃-induced lung function decrement though predominantly driven by females with BMI > 25 [11]. In our study no statistically significant changes were observed in the whole population, and with only four female study participants the analyses was likely underpowered. Certain genotypes, including GST isozymes, have been examined as possible biomarkers of sensitivity. Specifically for this study, GSTM1 genotype status was examined as the null genotype was not associated with a greater FEV1 decrease upon exposure of young, normal, healthy adults to a relatively low 0.06 ppm O₃ concentration for 6.6 hr with exercise (24), though lung neutrophilia was associated with the allele deletion. In our study, the GSTM1- subjects had an approximately 10% smaller decrement to 0.300 ppm O₃ exposure than GSTM1+ subjects in arm c Day 1 and arm a Day 2, though the difference did not reach statistical significance. However, no increased resistance to development of an FEV1 decrease was observed in the GSTM1-group upon the co-exposure to DE and O₃, possibly due to a similar sensitivity to DE components or DE components that were influenced by O₃.

The modulation of O₃-induced lung function changes by DE exposure raise the possibility that such interactions may occur from exposures to ambient levels of these pollutants. Additionally the components in DE that altered the magnitude of the O₃-induced FEV1 changes may be also present in our environment and potentially derived from natural or anthropomorphic sources and not exclusively from DE. It is unclear at what ambient concentrations DE may affect responsiveness to O₃ exposure, and also the threshold of O₃ level that induces lung function decrements; recent literature suggest it may be as low as the 0.06-0.08 ppm range [24,41]. Additionally populations sensitive to the DE-induced lung function effects have not been fully identified, and remains an ongoing issue.

Additional file

Additional file 1: Subject selection and participation criteria.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MCM was a principal investigator for part of the study duration, contributed substantially to the study design and protocols, prepared the initial draft of this manuscript, assisted with subject supervision, analyzed data, and performed internal clearances. TS was also a principal investigator for part of the study duration and protocols, assisted with the writing of this manuscript and data analyses, assisted with subject supervision; MC contributed substantially to the study design and protocols, assisted with subject supervision, and analyzed data; MS was the study coordinator, contributed substantially to the protocol, and performed genotyping; MB and TSM had the primary subject supervision and monitoring responsibilities and contributed to the study protocol; JB contributed substantially to the study design and protocol, contributed substantially to the initial draft of this manuscript, and was responsible for the pollutant exposure engineering oversight and quality assurance; DDS and RBD contributed substantially to the study design and protocols, contributed substantially to the review of drafts of this manuscript. All authors read and approved the final manuscript.

Acknowledgements

We gratefully acknowledge the help and support of Heidi Hiers of the US EPA Medical Staff, Ana Rappold (US EPA) for biostatistical support, the Westat recruitment staff, and the TRC engineering staff in completing the study.

Disclaimer

This report has been reviewed by the National Health and Environmental Effects Research Laboratory, United States Environmental Protection Agency and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the Agency nor does mention of trade names or commercial products constitute endorsement or recommendation for use. None of the authors has any actual or potential competing financial interests.

Author details

¹EPHD, NHEERL, US EPA, Research Triangle Park, Chapel Hill, NC 27711, USA. ²TRC Environmental, Chapel Hill, NC 27514, USA. ³Currently ORISE, Research Triangle Park, Chapel Hill, NC 27711, USA. ⁴U.S. EPA Human Studies Facility, 104 Mason Farm Road, Chapel Hill, NC 27599-7315, USA.

Received: 14 January 2014 Accepted: 5 August 2014

Published online: 02 September 2014

References

- Sydbom A, Blomberg A, Parnia S, Stenfors N, Sandström T, Dahlgren SE: **Health effects of diesel exhaust emissions.** *Eur Respir J* 2001, **17**:733–746.
- U.S. EPA: **Integrated Science Assessment for Particulate Matter (Final Report).** In *U.S. Environmental Protection Agency*. Washington, DC: EPA/600/R-08/139F; 2009.
- U.S. EPA: **Health Assessment Document for Diesel Engine Exhaust (Final 2002).** In *U.S. Environmental Protection Agency*. Washington Office, Washington, DC: EPA/600/8-90/057F; 2002.
- Buzzard NA, Clark NN, Guffey SE: **Investigation into pedestrian exposure to near-vehicle exhaust emissions.** *Environ Health* 2009, **8**:13.
- McDonnell WF, Stewart PW, Andreoni S, Seal E Jr, Kehrl HR, Horstman DH, Follinsbee LJ, Smith MV: **Prediction of ozone-induced FEV1 changes. Effects of concentration, duration, and ventilation.** *Am J Respir Crit Care Med* 1997, **156**:715.
- Schelegle ES, Siefkin AD, McDonald RJ: **Time course of ozone-induced neutrophilia in normal humans.** *Am Rev Respir Dis* 1991, **143**(6):1353–1358.
- Koren HS, Devlin RB, Graham DE, Mann R, McGee MP, Horstman DH, Kozumbo WJ, Becker S, House DE, McDonnell WF, Bromberg PA: **Ozone-induced inflammation in the lower airways of human subjects.** *Am Rev Respir Dis* 1989, **139**(2):407–415.
- Linn WS, Medway DA, Anzar UT, Valencia LM, Spier CE, Tsao FS, Fischer DA, Hackney JD: **Persistence of adaptation to ozone in volunteers exposed repeatedly for six weeks.** *Am Rev Respir Dis* 1982, **125**(5):491–495.
- Horvath SM, Gliner JA, Follinsbee LJ: **Adaptation to ozone: duration of effect.** *Am Rev Respir Dis* 1981, **123**(5):496–499.
- Corradi M, Alinovi R, Goldoni M, Vettori M, Folesani G, Mozzoni P, Cavazzini S, Bergamaschi E, Rossi L, Mutti A: **Biomarkers of oxidative stress after controlled human exposure to ozone.** *Toxicol Lett* 2002, **134**(1–3):219–225.
- Bennett WD, Hazucha MJ, Follinsbee LJ, Bromberg PA, Kissling GE, London SJ: **Acute pulmonary function response to ozone in young adults as a function of body mass index.** *Inhal Toxicol* 2007, **19**(14):1147–1154.
- Samet JM, Graff D, Berntsen J, Ghio AJ, Huang YC, Devlin RB: **A comparison of studies on the effects of controlled exposure to fine, coarse and ultrafine ambient particulate matter from a single location.** *Inhal Toxicol* 2007, **19**(Suppl 1):29–32.
- Salvi S, Blomberg A, Rudell B, Kelly F, Sandström T, Holgate ST, Frew A: **Acute inflammatory responses in the airways and peripheral blood after short-term exposure to diesel exhaust in healthy human volunteers.** *Am J Respir Crit Care Med* 1999, **159**(3):702–709.
- Behndig AF, Mudway IS, Brown JL, Stenfors N, Helleday R, Duggan ST, Wilson SJ, Boman C, Cassee FR, Frew AJ, Kelly FJ, Sandström T, Blomberg A: **Airway antioxidant and inflammatory responses to diesel exhaust exposure in healthy humans.** *Eur Respir J* 2006, **27**(2):359–365.
- National Research Council: **Research Priorities for Airborne Particulate Matter III.** Washington DC: National Academy Press; 2001:99–104.
- Fakhri AA, Ilic LM, Wellenius GA, Urch B, Silverman F, Gold DR, Mittleman MA: **Autonomic effects of controlled fine particulate exposure in young healthy adults: effect modification by ozone.** *Environ Health Perspect* 2009, **117**(8):1287–1292.
- Brook RD, Brook JR, Urch B, Vincent R, Rajagopalan S, Silverman F: **Inhalation of fine particulate air pollution and ozone causes acute arterial vasoconstriction in healthy adults.** *Circulation* 2002, **105**(13):1534–1536.
- Farrar AK, Boykin E, Ledbetter A, Andrews D, Gavett SH: **Increased lung resistance after diesel particulate and ozone co-exposure not associated with enhanced lung inflammation in allergic mice.** *Inhal Toxicol* 2010, **22**(1):33–41.
- Kodavanti UP, Thomas R, Ledbetter AD, Schladweiler MC, Shannahan JH, Wallenborn JG, Lund AK, Campen MJ, Butler EO, Gottipolu RR, Nyska A, Richards JE, Andrews D, Jaskot RH, McKee J, Kotha SR, Patel RB, Parinandi NL: **Vascular and cardiac impairments in rats inhaling ozone and diesel exhaust particles.** *Environ Health Perspect* 2011, **119**(3):312–318.
- Bosson J, Pourazar J, Forsberg B, Adelroth E, Sandström T, Blomberg A: **Ozone enhances the airway inflammation initiated by diesel exhaust.** *Respir Med* 2007, **101**(6):1140–1146.
- Bosson J, Barath S, Pourazar J, Behndig AF, Sandström T, Blomberg A, Adelroth E: **Diesel exhaust exposure enhances the ozone-induced airway inflammation in healthy humans.** *Eur Respir J* 2008, **31**(6):1234–1240.
- Frampton MW, Morrow PE, Cox C, Levy PC, Condemi JJ, Speers D, Gibb FR, Utecht MJ: **Sulfuric acid aerosol followed by ozone exposure in healthy and asthmatic subjects.** *Environ Research* 1995, **69**:1–14.
- Horvath SM, Follinsbee LJ, Bedi JF: **Combined effect of ozone and sulfuric acid on pulmonary function in man.** *Am Ind Hyg Assoc J* 1987, **48**:94–98.
- Kim CS, Alexis NE, Rappold AG, Kehrl H, Hazucha MJ, Lay JC, Schmitt MT, Case M, Devlin RB, Peden DB, Diaz-Sanchez D: **Lung function and inflammatory responses in healthy young adults exposed to 0.06 ppm ozone for 6.6 hours.** *Am J Respir Crit Care Med* 2011, **183**(9):1215–1221.
- Birch ME, Cary RA: **Elemental carbon-based method for monitoring occupational exposures to particulate diesel exhaust.** *Aerosol Sci Technol* 1996, **25**:221–241.
- U.S. EPA: **Integrated Science Assessment for Oxides of Nitrogen – Health Criteria (Final Report).** In *U.S. Environmental Protection Agency*. Washington, DC: EPA/600/R-08/071; 2008.
- Passannante AN, Hazucha MJ, Bromberg PA, Seal E, Follinsbee L, Koch G: **Noiceptive mechanisms modulate ozone-induced human lung function decrements.** *J Appl Physiol* 1998, **85**(5):1863–1870.
- Hazari MS, Rowan WH, Winsett DW, Ledbetter AD, Haykal-Coates N, Watkinson WP, Costa DL: **Potential of pulmonary reflex response to capsaicin 24 h following whole-body acrolein exposure is mediated by TRPV1.** *Respir Physiol Neurobiol* 2008, **160**(2):160–171.
- Taylor-Clarke TE, Udem BJ: **Ozone activates airway nerves via the selective stimulation of TRPA1 ion channels.** *J Physiol* 2010, **588**(Pt 3):423–433.
- Wang YY, Chang RB, Allgood SD, Silver WL, Liman ER: **A TRPA1-dependent mechanism for the pungent sensation of weak acids.** *J Gen Physiol* 2011, **137**(6):493–505.
- Madden MC, Richards JH, Dailey LA, Hatch GE, Ghio AJ: **Effect of ozone on diesel exhaust particle toxicity in Rat lung.** *Toxicol Appl Pharmacol* 2000, **168**:140–148.

32. Hazucha MJ, Hazucha MJ, Folinsbee LJ, Seal E, Bromberg PA: **Lung function response of healthy women after sequential exposures to NO₂ and O₃.** *Am J Respir Crit Care Med* 1994, **150**(3):642–647.
33. Hazucha MJ, Madden M, Pape G, Becker S, Devlin R, Koren HS, Kehrl H, Bromberg PA: **Effects of cyclo-oxygenase inhibition on ozone-induced respiratory inflammation and lung function changes.** *Eur J Appl Physiol Occup Physiol* 1996, **73**(1–2):17–27.
34. Bergren DR: **Prostaglandin involvement in lung C-fiber activation by substance P in guinea pigs.** *J Appl Physiol* 2006, **100**(6):1918–1927.
35. Ahn EK, Yoon HK, Jee BK, Ko HJ, Lee KH, Kim HJ, Lim Y: **COX-2 expression and inflammatory effects by diesel exhaust particles in vitro and in vivo.** *Toxicol Lett* 2008, **176**(3):178–187.
36. Becker S, Mundandhara S, Devlin RB, Madden M: **Regulation of cytokine production in human alveolar macrophages and airway epithelial cells in response to ambient air pollution particles: Further mechanistic studies.** *Toxicol Appl Pharmacol* 2005, **207**:S269–S275.
37. Salvemini D, Misko TP, Masferrer JL, Seibert K, Currie MG, Needleman P: **Nitric oxide activates cyclooxygenase enzymes.** *Proc Natl Acad Sci U S A* 1993, **90**(15):7240–7244.
38. Wright DT, Adler KB, Akley NJ, Dailey LA, Friedman M: **Ozone stimulates release of platelet activating factor and activates phospholipases in guinea pig tracheal epithelial cells in primary culture.** *Toxicol Appl Pharmacol* 1994, **127**(1):27–36.
39. Madden MC, Friedman, Dailey LA, Samet JM: **Inhibition of arachidonic acid esterification in human airway epithelial cells exposed to ozone in vitro.** *Inhalation Toxicology* 1998, **10**:795–811.
40. Pueringer RJ, Hunninghake GW: **Lipopolysaccharide stimulates de novo synthesis of PGH synthase in human alveolar macrophages.** *Am J Physiol* 1992, **262**(1 Pt 1):L78–L85.
41. Horstman DH, Folinsbee LJ, Ives PJ, Abdul-Salaam S, McDonnell WF: **Ozone concentration and pulmonary response relationships for 6.6-hour exposures with five hours of moderate exercise to 0.08, 0.10, and 0.12 ppm.** *Am Rev Respir Dis* 1990, **142**(5):1158–1163.

doi:10.1186/s12989-014-0037-5

Cite this article as: Madden *et al.*: Diesel exhaust modulates ozone-induced lung function decrements in healthy human volunteers. *Particle and Fibre Toxicology* 2014 **11**:37.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

