



OPEN

Identification of domestication-related loci associated with flowering time and seed size in soybean with the RAD-seq genotyping method

SUBJECT AREAS:

QUANTITATIVE TRAIT

PLANT DOMESTICATION

GENOME-WIDE ASSOCIATION
STUDIESReceived
9 October 2014Accepted
2 March 2015Published
23 March 2015Correspondence and
requests for materials
should be addressed to
Y.-M.Z. (soy Zhang@
njau.edu.cn;
soy Zhang@mail.hzau.
edu.cn)* These authors
contributed equally to
this work.Ling Zhou^{1*}, Shi-Bo Wang^{1*}, Jianbo Jian^{2*}, Qing-Chun Geng¹, Jia Wen¹, Qijian Song³, Zhenzhen Wu², Guang-Jun Li⁴, Yu-Qin Liu⁵, Jim M. Dunwell⁶, Jin Zhang¹, Jian-Ying Feng¹, Yuan Niu¹, Li Zhang¹, Wen-Long Ren¹ & Yuan-Ming Zhang^{1,7}

¹State Key Laboratory of Crop Genetics and Germplasm Enhancement, National Center for Soybean Improvement, Jiangsu Collaborative Innovation Center for Modern Crop Production, Nanjing Agricultural University, Nanjing 210095, China, ²BGI-Shenzhen, Shenzhen 518083, China, ³Soybean Genomics and Improvement Laboratory, Agricultural Research Service, United States Department of Agriculture, Beltsville, Maryland 20705, USA, ⁴College of Life Science, Linyi University, Linyi 276005, China, ⁵Crop Research Institute, Linyi Academy of Agricultural Sciences, Linyi 276012, China, ⁶School of Agriculture, Policy and Development, University of Reading, Reading RG6 6AR, United Kingdom, ⁷Statistical Genomics Lab, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan 430070, China.

Flowering time and seed size are traits related to domestication. However, identification of domestication-related loci/genes of controlling the traits in soybean is rarely reported. In this study, we identified a total of 48 domestication-related loci based on RAD-seq genotyping of a natural population comprising 286 accessions. Among these, four on chromosome 12 and additional two on chromosomes 11 and 15 were associated with flowering time, and four on chromosomes 11 and 16 were associated with seed size. Of the five genes associated with flowering time and the three genes associated with seed size, three genes *Glyma11g18720*, *Glyma11g15480* and *Glyma15g35080* were homologous to Arabidopsis genes, additional five genes were found for the first time to be associated with these two traits. *Glyma11g18720* and *Glyma05g28130* were co-expressed with five genes homologous to flowering time genes in Arabidopsis, and *Glyma11g15480* was co-expressed with 24 genes homologous to seed development genes in Arabidopsis. This study indicates that integration of population divergence analysis, genome-wide association study and expression analysis is an efficient approach to identify candidate domestication-related genes.

The transitions from vegetative to reproductive growth (days to flowering) and from generation to generation through seeds are important stages of the plant life cycle. Flowering time reflects the adaptation of a plant to its environment, and the time required to mature varies widely among cultivars¹. Seeds are important in the reproduction and spread of flowering plants, and seed size partly reflects the efficiency of plant production². Both flowering time and seed size are important traits involved in domestication, a process accompanied by reduction in genetic diversity and loss of important traits preserved in wild relatives.

Morphological, physiological and molecular markers have been widely used to measure the genetic diversity of wild and cultivated plants in rice³, soybean⁴, and wheat⁵, to deduce the geographic regions of domestication, and to screen for breeding material. Linkage and association studies have identified quantitative trait loci (QTL) associated with domestication-related (DR) traits in various plants^{6,7} and animals⁸, such as the lateral branching locus *teosinte branched1* (*tb1*) in maize⁹, the seed shattering locus *qSH1* in rice¹⁰, and the milk-production locus *DGAT1* in dairy cattle¹¹. Numerous DR genes have also been identified, including *GW8* in rice¹², *Q* in wheat¹³, *vrs1* in barley¹⁴, and *RYR1* in pig¹⁵. As whole genome sequences from almost all major plants have become available, substantial progress has been achieved, including 1) Hundreds of DR genes have been identified by comparative analysis of genomes in plants¹⁶ and animals¹⁷; 2) Candidate genes resulting from selection were detected by comparative analysis and functional tests⁷, for example, the genes *LOC_Os05g20290* and *LOC_Os08g40710* in rice¹⁸ and the genes *GRMZM2G448355* and *abph1* (*GRMZM2G035688*) in maize¹⁹; and 3) The molecular mechanisms underlying DR traits have been reported^{20,21}. In addition, the ancient DNA and genomes of fossils




Figure 1 | Detection of domestication-related loci in the soybean genome using χ^2 test of independence (a), U-test (b) and genetic diversity analysis (c). Significances of all the SNPs in the above three analyses were marked by red (significant) and blue (not significant) dots. The genomic positions of SNPs on chromosomes 1 to 20 were 7166 ~ 55874947, 2395 ~ 51643854, 45230 ~ 47773436, 15992 ~ 49228296, 9016 ~ 41933701, 1142 ~ 50641309, 37534 ~ 44659030, 3583 ~ 46944564, 25976 ~ 46841908, 24539 ~ 50962464, 30293 ~ 39163227, 13701 ~ 40093893, 10047 ~ 44402574, 13165 ~ 49710404, 8319 ~ 50879005, 14132 ~ 37370388, 47058 ~ 41905331, 426 ~ 62303776, 9282 ~ 50584403 and 18252 ~ 46703751 (bp), respectively.




Figure 2 | Genome-wide association of first (a) and full (b) flowering times, seed length (c), seed width (d) and 100-seed weight (e) in soybean during 2008–2012 with domestication-related SNPs on chromosomes 1, 6, 7, 9, 11, 12, 13, 15, 16 and 17. Dots with green, purple, black, red, and blue colors denote 2008, 2009, 2010, 2011, and 2012, respectively.

have contributed to the histories of domestication and evolution in horses²², cattle²³, and *Arabidopsis thaliana*²⁴. However, little is known about the loci/genes underlying DR traits in soybean.

The soybean plant originated in China and was first domesticated by Chinese farmers between 6,000 and 9,000 years ago²⁵. The modern

domesticated soybean (*Glycine max*) is an economically important crop because it has high protein and oil contents, can fix nitrogen through microorganisms in the soil^{26,27}, and is a model plant for legume research. The cultivated soybean (*G. max*) differs from the wild soybean (*G. soja*) in many traits, for example, the cultivated



Table 1 | Genome-wide association of flowering time (FT) and seed size (SS) traits with domestication-related SNPs (bold) in soybean

SNP	Domestication analysis					Genome-wide association study					Homologous gene in Arabidopsis			
	Chr.	Position (bp)	χ^2 (-log ₁₀ P)	U (-log ₁₀ P)	F _{ST}	Trait	Allele	Frequency	Effect	-log ₁₀ P	r ² (%)	Candidate gene	Gene	Reference
11 10957940	11	30,601,664	30.60(6.64)	-2.36(1.74)	0.4890	SS	T	0.45	-9.12 ~ -5.84	6.32 ~ 7.43	6.24 ~ 11.80	Glyma1g15300	NOT2A (AT1G07705)	49
11 11100801	11	38,401,834	38.40(8.34)	-2.53(1.95)	0.5069	SS	G	0.58	-2.10 ~ -0.58	6.52 ~ 14.49	8.14 ~ 16.26	Glyma1g15480	NOT2A (AT1G07705)	49
11 11111516	11	111,20981	111.21(25.6)			SS	C	0.38	0.90 ~ 3.90	0.60 ~ 2.40	0.91 ~ 3.20	Glyma1g15480	NOT2A (AT1G07705)	49
11 11120981	11	111,21256	111.21(25.6)			SS	C	0.53	-23.59 ~ -18.40	6.19 ~ 10.59	6.50 ~ 14.42	Glyma1g15480	NOT2A (AT1G07705)	49
11 11121256	11	111,21307	111.21(25.6)			SS	G	0.51	4.40 ~ 18.40	6.11 ~ 9.91	5.96 ~ 16.85	Glyma1g15480	NOT2A (AT1G07705)	49
11 11121307	11	111,21307	111.21(25.6)			SS	T	0.56	-0.07 ~ 4.87	6.40 ~ 11.97	6.56 ~ 16.26	Glyma1g15480	NOT2A (AT1G07705)	49
11 15362036	11	45,451,987	45.45(9.87)	-2.43(1.82)	0.5656	FT	A	0.42	-10.11 ~ -4.87	1.51 ~ 2.61	1.32 ~ 2.17	Glyma1g18720	OTLD1 (AT2G27350)	46–48
12 5895633	12	37,231,088	37.23(8.08)	-3.73(3.71)	0.7344	FT	C	0.53	-12.96 ~ -5.21	7.63 ~ 8.65	6.16 ~ 7.00	Glyma1g2g08150	AREB3 (AT3G56850)	38
12 5895655	12	38,041,826	38.04(8.26)	-4.30(4.78)	0.7830	FT	A	0.52	-13.05 ~ -5.20	6.53 ~ 7.61	5.32 ~ 6.34	Glyma1g2g08160		
12 5895785	12	38,741,841	38.74(8.41)	-4.19(4.55)	0.7938	FT	C	0.53	-4.54 ~ -1.30	8.14 ~ 9.33	6.44 ~ 7.32	Glyma1g2g08160		
12 5957099	12	30,471,662	30.47(6.62)	-2.78(2.27)	0.6017	FT	A	0.55	-12.93 ~ -9.17	6.54 ~ 7.11	5.61 ~ 5.92	Glyma1g2g08210		
15 39607166	15	32,101,697	32.10(6.97)	-3.89(3.40)	0.6496	FT	A	0.48	-14.97 ~ -10.93	1.11 ~ 1.74	1.06 ~ 1.54	Glyma1g5g35080		
16 30207175	16	30,671,666	30.67(6.66)	-3.47(3.28)	0.5348	SS	G	0.35	-7.48 ~ -3.97	6.99	8.49	Glyma1g6g26030		
16 30207191	16	30,671,911	30.67(6.66)			SS	T	0.44	-10.29 ~ -8.41	6.47 ~ 8.05	6.02 ~ 9.97	Glyma1g6g26050		
16 30210113	16	30,901,671	30.90(6.71)	-3.74(3.74)	0.6148	SS	C	0.37	-2.13 ~ 0.84	7.25 ~ 10.22	7.24 ~ 12.01	Glyma1g6g26050		

soybean forms flowers earlier and has larger seeds than the wild soybean²⁸. Previous research has led to the discovery of many QTL correlated to DR traits such as seed size^{29–33} and other traits (<http://www.SoyBase.org>). Some genes controlling DR traits in soybean have also been discovered, e.g., *E1-E4*^{34–37}, *FLC*, *VRNA*, *ELF8*, *PHYE*, *PHYA*¹⁶, *CDF3*, *VRN1*, *SVP*, *AP3* and *PIF3* for flowering time³⁸, and *Dt1* for determinate growth habit³⁹. Although the genes and their molecular mechanisms for some DR traits in soybean have been investigated^{38,39}, the genes/loci underlying many other DR traits remain to be addressed.

In this study, restriction-site-associated DNA (RAD) tags from 14 wild, 153 landrace, and 119 bred accessions were sequenced, and the sequence variants were analyzed to detect DR loci by testing the independence between the SNPs and soybean evolutionary classes (wild, landrace, and bred) and comparing the genetic diversity between the wild and cultivated soybeans. Genome-wide association of the detected DR loci with DR traits (flowering time and seed size) were also studied. Candidate genes predicted to be involved in these two traits were pinpointed using comparative genomics technology. Co-expression analysis for individual candidate genes was also conducted.

Results and Discussion

Phenotypic characteristics of flowering time and seed size. Flowering time was measured by the days from the date of emergence to the date of first and full flowering in this study. The average plus standard error were 51.83 ± 3.73 , 46.22 ± 0.83 and 35.60 ± 0.64 (days) for first flower and 55.95 ± 4.00 , 49.96 ± 0.86 and 38.57 ± 0.65 (days) for full flower in wild, landrace and bred soybeans in 2010–2012, respectively. *G. soja* flowered later than *G. max*.

Seed size was characterized by seed length (SL), seed width (SW) and 100-seed weight (100 SW) in this study. The averages plus standard errors in wild, landrace and bred soybeans in 2008–2012 were 4.76 ± 0.18 , 7.61 ± 0.08 and 7.85 ± 0.06 (mm) for SL, 3.56 ± 0.17 , 6.54 ± 0.07 and 6.63 ± 0.05 (mm) for SW, and 2.99 ± 0.45 , 15.43 ± 0.48 and 16.85 ± 0.39 (g) for 100 SW, respectively, indicating much smaller seeds in *G. soja* than in *G. max*.

Distinct difference of these traits between *G. soja* and *G. max* suggested that these traits are domestication-related. Flowering time and seed size were also considered as DR traits in other reports^{25,28,40–42}, although domestication traits in soybean include more than these two traits, such as indeterminate habit and pod dehiscence.

Detection and distribution of domestication-associated loci.

Based on the sequence obtained from 286 accessions through the RAD-seq genotyping approach, a total of 106,013 SNPs were identified. χ^2 tests of independence between the SNPs and evolutionary classes (wild, landrace, and bred) of soybean showed that 198 SNPs were significant at P-value $\leq 4.72 \times 10^{-7}$ (Fig. 1a). A U-test determined that the allelic frequency for 72 of the 198 SNPs were significantly different between the wild and cultivated classes (Fig. 1b). The fixation index (F_{ST}) analysis demonstrated that 48 of the 72 SNPs had an F_{ST} value greater than 0.45 (Fig. 1c; Table S1).

Of the 48 DR loci, 3, 4, 1, 2, 14, 5, 2, 11, 3 and 3 loci were located on chromosomes 1, 6, 7, 9, 11, 12, 13, 15, 16 and 17, respectively (Fig. 1c). However, 52% of the 48 loci were located on chromosomes 11 or 15, and these two chromosomes harbour more DR loci than other chromosomes. Similar results were also observed by Kim *et al.*⁴³, Chung *et al.*⁴⁴ and Li *et al.*⁴⁵. On the two chromosomes, Kim *et al.*⁴³ identified four DR genes (*Glyma15g14980*, *Glyma15g23400*, *Glyma11g05720* and *Glyma15g04930*) that were associated with flowering time; Chung *et al.*⁴⁴ detected 204 soybean candidate DR genes; and Li *et al.*⁴⁵ found three DR genes (*Glyma11g03340*, *Glyma11g03330* and *Glyma11g03320*) near the reported flower colour QTL.

A total of 12 DR loci were identified in 11 genes with 4 loci in the coding regions of the genes. Of the 11 genes whose functions were




Figure 3 | Expression levels of candidate genes for soybean flowering time and seed size in various developmental stages or tissues. The expression data were extracted from the soybean transcriptome data deposited at NCBI (<http://www.ncbi.nlm.nih.gov/geo/>).

previously annotated in the Pfam and NCBI non-redundant databases, three and two genes were located on chromosomes 11 and 15, respectively. The common allele for each of the 12 loci was different in the bred and wild soybean, and common alleles for each of the 3 DR loci on chromosome 11 were the same in both the landrace and wild soybeans.

Genome-wide association studies (GWAS) for flowering time and seed size. GWAS for flowering time (first and full) and seed size (SL, SW and 100 SW) was performed based on a dataset containing 286 accessions genotyped with 55,052 SNPs (Fig. S1). Because our purpose was to investigate the association of the DR loci with the two DR traits, we only analyzed the association of the DR loci listed in Fig. 2 and Table 1.

Four QTL were associated with flowering time (P-values of 4.68E-10 to 2.95E-7), and each QTL explained 5.32–7.32% of the total phenotypic variance. These QTL were positioned at 5895633, 5895655, 5895785, and 5957099 (bp) on chromosome 12 (Fig. 2a, b). The first three loci were associated with the first and full flowering times in 2010 to 2012, while the last locus was associated with the first flowering time in 2010 and 2012 and with the full flowering time in 2012. In addition, two loci at genomic positions 15362036 bp (P-values of 0.0064 to 0.0234) on chromosome 11 and 39607166 bp (P-values of 0.0183 to 0.0434) on chromosome 15 were associated with the flowering times at a 0.05 probability level in 2010 to 2012, except for the fact that the later locus was almost associated with the first flower at a 0.05 probability level in 2012 (P-value = 0.0640) (Fig. 2a, b).

A total of 8 QTL, each explaining 0.91–16.85% of the total phenotypic variance of seed size, were identified (P-values of 3.24E-15 to 7.76E-7). These QTL were mapped to genomic positions 10957940, 11100801, 11120981, 11121256, and 11121307 bp on chromosome 11 and 30207191, 30207175, and 30210113 bp on chromosome 16 (Fig. 2c–e). Among the five SNP loci on chromosome 11, the last four loci, which were close to the DR locus at 11111516 bp (~10 kb), were associated with the three seed size traits. The DR locus at 10957940 bp was associated with SW in 2008 to 2010 and with SL in 2009. Among the three loci on chromosome 16, one locus at genomic position 30207191 bp, which was close to the DR locus 30207175 bp, was correlated with SL and 100 SW in 2011 and 2012 and with SW in 2011. The DR locus at 30207175 bp was associated with 100 SW in 2012, and the DR locus at genomic position 30210113 bp was associated with SL and 100 SW in 2012. In summary, all four DR loci were associated with seed size traits.

Candidate DR genes for flowering time. Candidate DR genes were selected if: 1) genes containing DR loci or genes in the adjacent regions to which DR loci were significantly associated with the flowering time or seed size, and 2) genes with high level of tran-

scriptome expression. The locus at 15362036 bp on chromosome 11 was in the gene *Glyma11g18720*, which was homologous to the gene *OTLD1* regulating *FLC* for flowering time and seed germination in Arabidopsis^{46–48} (Table 1, Fig. S2). The locus at 39607166 bp on chromosome 15 was in the gene *Glyma15g35080* which regulated flowering time in soybean³⁸. We also observed a relatively high expression of the *Glyma11g18720* and *Glyma15g35080* at specific stages of flowering time and seed development (Fig. 3).

A multiple linear regression analysis of each selected soybean gene (y) on *Glyma11g18720* (x_1) and *Glyma05g28130* (x_2) (homologous to *FLC*, Fig. S3) based on expression data, with the model $y = b_0 + b_1x_1 + b_2x_2 + b_{12}x_1x_2 + \varepsilon$, identified five significant flowering-time-related genes *Glyma06g02470*, *Glyma08g02490*, *Glyma11g02110*, *Glyma15g17480*, and *Glyma02g11060* with P-values of 1.28E-5 to 7.81E-2 (Table 2). Of the five genes, *Glyma06g02470* was homologous to the genes *EEL* and *AREB3*³⁸ related to flowering time in Arabidopsis (Fig. S4); *Glyma08g02490* to the genes *SPA1–SPA4*³⁸ (Fig. S5); *Glyma15g17480* to the genes *FKF1*, *LKP2*, and *ZTL*³⁸ (Fig. S6); and *Glyma02g11060* to the genes *TEM2*, *RAV1*, *TEM1*, and *EDF3*³⁸ (Fig. S7).

The genes *Glyma12g08150* and *Glyma12g08160* were adjacent to the domestication regions of 5895633 to 5895785 bp on chromosome 12, and the genes *Glyma12g08210* and *Glyma12g08230* were in close proximity to the DR locus at 5957099 bp on chromosome 12. The likely association of these genes with domestication was also reported by Chung *et al.*⁴⁴. In addition, based on the soybean transcriptome data deposited at NCBI (<http://www.ncbi.nlm.nih.gov/geo/>), all the genes had relatively high expression levels in floral bud tissue (Fig. 3). Therefore, the above 10 genes may be associated with flowering time.

Candidate DR genes for seed size. The genes *Glyma11g15300* and *Glyma11g15310* were close to the DR locus at 10957940 bp on chromosome 11. *Glyma11g15480* contained a DR locus at 11111516 bp on chromosome 11, and *Glyma16g26030* and *Glyma16g26050* were close to the domestication regions of 30207175 to 30210113 bp on chromosome 16. The three genes *Glyma11g15300*, *Glyma11g15480* and *Glyma16g26030* had relatively higher expression levels during seed development than did *Glyma11g15310* and *Glyma16g26050* (Fig. 3). The soybean gene *Glyma11g15480* is homologous to the Arabidopsis gene *NOT2A*⁴⁹ (Fig. S8) and was found to be associated with seed size in this study. A correlation analysis of *Glyma11g15480* with each selected soybean gene based on the gene expression dataset showed that 24 genes were significantly correlated with *Glyma11g15480* (P-values of 1.00E-4 to 4.90E-2) (Table 2). Among the 24 genes, 11 were homologous to five epigenetic regulation endosperm genes in Arabidopsis (*FIS2*, *MSII*, *FIE/FIS3*, *DDMI*, and *MET1*)^{50–52} (Figures S9–S13); six were homologous to four endosperm development genes (*AP2*, *AGL61*, *SHB1*, and



Table 2 | Co-expressional genes of *Glyma11g18720* and *Glyma05g28130* for flowering time (FT) and *Glyma11g15480* for seed size (SS) and homologous genes in *Arabidopsis*

Trait	Co-expressional analysis				Homologous gene in <i>Arabidopsis</i>				Evolutionary relationship	
	Gene	F(P-value)	R ² (%)	Gene	Similarity (%)	Notation	Functional description	Reference		
FT	<i>Glyma06g02470</i>	3.77[7.81E-2]	86.91	AT3G56850; AT2G41070	23.2; 12.3	AREB3; EEL	DNA binding/transcription factor	38	Figure S4	
	<i>Glyma08g02490</i> ; <i>Glyma11g02110</i>	6.42[2.66E-2] 14.33[3.83E-3]	76.24;87.75	AT2G46340; AT4G11110 AT3G15354; AT1G53090	35.4–61.7	SPA1 ~ SPA4	suppress photomorphogenesis, light signaling pathway	38	Figure S5	
	<i>Glyma15g17480</i>	108.86[1.28E-5]	98.20	AT1G68050; AT2G18915 AT5G57360	68.2–85.3	FKF1, LKP2, ZTL	control flowering time by influencing the circadian clock period, light signaling pathway	38	Figure S6	
	<i>Glyma02g11060</i>	13.29[4.66E-3]	86.91	AT1G68840; AT1G13260 AT1G25560; AT3G25730	53.9–61.6	TEM2, RAV1 TEM1, EDF3	repress flowering, light signaling pathway	38	Figure S7	
	<i>Glyma01g41460</i>	43.85[2.00E-4]	84.57	AT2G35670	29.4	FIS2	female gametophyte, endosperm	50,51	Figure S9	
	<i>Glyma05g26150</i> ; <i>Glyma08g09090</i>	12.09–82.46	66.17–91.16	AT5G58230	63.0–92.5	MSI1	female gametophyte, embryo	50,51	Figure S10	
	<i>Glyma11g09700</i> ; <i>Glyma12g03700</i>	(<1.00E-4–8.40E-3)								
	<i>Glyma10g02690</i> ; <i>Glyma02g17110</i>	6.40–16.87	44.43–67.83	AT3G20740	83.7–86.7	FIE/FIS3	female gametophyte, endosperm, embryo	50,51	Figure S11	
	<i>Glyma13g36310</i> ; <i>Glyma01g38150</i> ; <i>Glyma11g07220</i>	(3.40E-3–3.53E-2) 7.40[2.62E-2] 6.70[2.95E-2]	48.07;46.66	AT5G66750	76.2; 75.5	DDM1	chromatin remodeling factor; embryo, endosperm	50,52	Figure S12	
	<i>Glyma06g18790</i> ; <i>Glyma11g05720</i> ; <i>Glyma17g18640</i>	6.42[3.51E-2] 9.85[1.38E-2] 7.34[2.67E-2]	44.51 55.18;47.85	AT5G49160 AT4G36920	73.5 46.3; 49.5	MET1 AP2	methyl transferase; embryo, endosperm AP2 domain transcription factor; Integument, endosperm, embryo	50,52 53	Figure S13 Figure S14	
<i>Glyma20g27320</i>	5.37[4.90E-2]	40.16	AT2G24840	42.0	AGL61	MADS-box transcription factor; central cell, endosperm	50	Figure S15		
<i>Glyma07g35520</i> ; <i>Glyma09g37000</i>	35.83[3.00E-4] 7.47 [2.57E-2]	81.75;48.29	AT4G25350	68.9; 64.8	SHB1	Transcription co-activator; embryo, endosperm	54	Figure S16		
<i>Glyma20g29600</i>	9.26[1.60E-2]	53.66	AT5G07280	73.9	EXS/EMS1	Leucine-rich repeat (LRR) receptor kinase; embryo, endosperm	55	Figure S17		
<i>Glyma04g37760</i> ; <i>Glyma05g38540</i>	6.88–10.96	46.24–57.80	AT5G62000	70.1–72.1	ARF2	auxin-responsive element binding transcription	56	Figure S18		
<i>Glyma06g17320</i> ; <i>Glyma08g01100</i>	(1.07E-2–3.05E-2)					factor; integument before pollination, embryo after pollination				
<i>Glyma19g36100</i> ; <i>Glyma05g22970</i> ; <i>Glyma17g17010</i>	8.42[1.98E-2] 6.33[3.61E-2] 8.46 [1.96E-2]	51.28 44.16;51.39	AT2G37260 AT4G37750	61.8 60.5; 61.6	TTG2 ANT	WRKY transcription factor; Integument AP2-like transcription factor; Integument before pollination, embryo after pollination	57 58	Figure S19 Figure S20		



EMS1)^{50,53–55} (Figures S14–S17); five were homologous to two integument development genes (*ARF2/MNT*, and *TTG2*)^{56–57} (Figures S18–S19); and two genes were homologous to the embryo development gene (*ANT*)⁵⁸ (Fig. S20). Therefore, the above 27 genes may be associated with seed size.

In addition, significant difference of allele frequencies for the SNPs in coding region of the above eight candidate DR genes in GWAS was observed between wild and cultivated soybeans (Tables S2, S3). Although the number of wild accessions was smaller than the number of cultivated soybeans, it is close to that reported in Chung *et al.*⁴⁴ and Lam *et al.*⁵⁹, and we adopted a stringent threshold of statistical significance in determining domestication loci ($\alpha = 4.72E-7$ and $F_{ST} > 0.45$) and GWAS ($\alpha = 9.08E-7$).

Some QTL (or genes) associated with DTs in this study were consistent with those reported previously (Table S4), for example, *Glyma12g08150*, *Glyma12g08210* and *Glyma12g08230* were reported as candidate domestication genes by Chung *et al.*⁴⁴.

Conclusion

A total of 48 DR loci in the soybean genome were identified. Most of these loci were on chromosomes 11 and 15. Among these DR loci, 10 loci were associated with flowering time or seed size. Eight genes near the 10 loci were associated with the two traits. Among the eight genes, three genes *Glyma11g18720*, *Glyma11g15480* and *Glyma15g35080* were homologous to Arabidopsis genes, three known DR genes *Glyma12g08150*, *Glyma12g08210* and *Glyma12g08230* were linked to flowering time; and the other two DR genes *Glyma11g15300* and *Glyma16g26030* were reported for the first time. *Glyma11g18720* and *Glyma05g28130* were co-expressed with five genes homologous to flowering time genes in Arabidopsis, and *Glyma11g15480* was co-expressed with 24 genes homologous to seed development genes in Arabidopsis. The method that integrates domestication analysis, GWAS and gene expression analysis is an efficient approach for identifying potentially useful DR genes.

Methods

Germplasm for genome-wide re-sequencing. The 286 soybean accessions used here included 14 wild, 153 landrace, and 119 bred accessions that were randomly selected from 6 geographic regions in China using a stratified random sampling method (Table S5). The seeds of the accessions were obtained from the National Centre for Soybean Improvement and Linyi Academy of Agricultural Sciences, and planted in three-row plots in a completely randomised design at the Jiangpu Experimental Station of Nanjing Agricultural University during 2008–2012. The plots were 1.5 m wide and 2 m long. A total of 20 seeds from five plants of the middle row in each plot were randomly selected for the measurement of the seed size traits, including SL and SW, using digital vernier calipers. The SL and SW for each accession were averaged based on 20 seeds and 100 SW for each replicate. The first and full flowering dates for each accession were recorded in the field during 2010 and 2012.

Genome-wide re-sequencing and sequence alignment. Approximately 0.3 g of fresh leaves obtained from each accession in 2012 was used to extract genomic DNA using the cetyltrimethylammonium bromide method. The DNA was digested with the *EcoRI* (G^{\wedge} AATTC) enzyme. The RAD-seq libraries for sequencing were prepared according to the protocols described by Baird *et al.*⁶⁰. We sequenced 50 bp at each end and used SolexaPipeline 1.0 for base call of 50-bp reads from the raw fluorescent images. To ensure quality, the raw sequence data were processed in two steps. In the first step, reads with incorrect adapter sequences were deleted, and any reads containing more than 50% low quality bases (quality value ≤ 5) were removed. In the second step, the remaining reads were demultiplexed according to the index of each sample. The sequences were subsequently aligned on the soybean reference genome Glyma1.1 (<http://www.jgi.doe.gov>) using the Burrows-Wheeler Alignment Tool (BWA)⁶¹. The base recalibration and determination of the SNP allele were performed using GATK⁶².

Detection of DR loci. The independence between each SNP and the evolutionary class (wild, landrace, and bred) was determined using the χ^2 test⁶³. The significant threshold α for each test was adjusted using the Bonferroni correction, that is, $\alpha = 0.05/106,013 = 4.72 \times 10^{-7}$ (106,013 was the total number of SNPs identified). The SNPs significantly associated with evolutionary classes were used for the subsequent analysis.

A U-test at the 0.05 significance level was used to test the significance of the allelic differences between the wild (14) and cultivated (272) accessions for each candidate SNP.

For all the SNPs with significant differences in the U-test, the fixation index (F_{ST}) was calculated for the purpose of measuring population differentiation between the wild (14) and bred (119) classes using the software Genepop v4.2⁶⁴, and a F_{ST} threshold value of 0.45 from Lam *et al.*⁵⁹ were adopted for the identification of DR loci.

Population structure analysis. The STRUCTURE 2.2 software was used to investigate the population structures based on all available accessions⁶⁵. The number of subpopulations (K) was set from 2 to 7³². In this study, the ΔK method of Evanno *et al.*⁶⁶ was used to determine the optimal value of K . The Q matrix was calculated based on inferred K .

Genome-wide association study. A GWAS was performed using the general linear model in TASSEL V4.3 (www.maizegenetics.net/tassel) with the total average and population structure as covariates. The Manhattan plot of $-\log_{10}P$ was generated using SAS software. As P -values from the software were not corrected for multiple test, the significance level α for each test was determined after Bonferroni correction, $\alpha = 0.05/55,052 = 9.08E-7$ (55,052 was the number of SNPs with the concordance rate $< 99\%$ by the TASSEL software). Elimination of the SNPs with concordance rate $> 99\%$ was conducted by the TASSEL software.

Identification of soybean DR genes and homologous Arabidopsis genes. Many genes have been annotated with high confidence in the soybean genome²⁶. The most likely functions of the genes have also been determined based on the annotation of the Arabidopsis homologs. The protein sequences of the annotated genes in soybean (Glyma1.1) and Arabidopsis (TAIR9 release) were downloaded from the Phytozome (www.phytozome.net) and The Arabidopsis Information Resource (TAIR) (www.arabidopsis.org) websites, respectively.

The Pfam gene annotations were downloaded from the Phytozome database version 9.0 (<http://www.phytozome.net/>). The annotations of the DR genes using BLASTX (e-value $1E-4$ or better) against NCBI's non-redundant database have been described at <http://www.ncbi.nlm.nih.gov/>. The top best hits from the species of interest were extracted for each gene.

Transcriptional activity and gene expression data analysis. The transcriptome sequences were obtained from the Gene Expression Omnibus database at <http://www.ncbi.nlm.nih.gov/geo/ss> (accession number GSE29163), and were used to identify the expression level of soybean gene during seed development and throughout the life cycle.

1. Wahl, V. *et al.* Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science* **339**, 704–707 (2013).
2. Adebisi, M. A. *et al.* Influence of different seed size fractions on seed germination, seedling emergence and seed yield characters in tropical soybean (*Glycine max* L. Merrill). *J Int Pharm Res* **8**, 26–33 (2013).
3. Sun, C. Q., Wang, X. K., Li, Z. C., Yoshimura, A. & Iwata, N. Comparison of the genetic diversity of common wild rice (*Oryza rufipogon* Griff.) and cultivated rice (*O. sativa* L.) using RFLP markers. *Theor Appl Genet* **102**, 157–162 (2001).
4. Li, Y. H. *et al.* Genetic diversity in domesticated soybean (*Glycine max*) and its wild progenitor (*Glycine soja*) for simple sequence repeat and single-nucleotide polymorphism loci. *New Phytol* **188**, 242–253 (2010).
5. Laidò, G. *et al.* Genetic diversity and population structure of tetraploid wheats (*Triticum turgidum* L.) estimated by SSR, DArT and pedigree data. *PLoS ONE* **8**, e67280 (2013).
6. Gupta, P. K., Kulwal, P. L. & Mir, R. R. *QTL mapping: methodology and applications in cereal breeding*. In Gupta, P. K. & Varshney, R. K. (eds.), *Cereal Genomics II* pp319–340 (2013).
7. Olsen, K. M. & Wendel, J. F. A bountiful harvest: genomic insights into crop domestication phenotypes. *Annu Rev Plant Biol* **64**, 47–70 (2013).
8. Andersson, L. & Georges, M. Domestic-animal genomics: deciphering the genetics of complex traits. *Nat Rev Genet* **5**, 202–212 (2004).
9. Doebley, J., Stec, A. & Gustus, C. *Teosinte branched1* and the origin of maize: evidence for epistasis and the evolution of dominance. *Genetics* **141**, 333–346 (1995).
10. Konishi, S. *et al.* An SNP caused loss of seed shattering during rice domestication. *Science* **312**, 1392–1396 (2006).
11. Grisart, B. *et al.* Positional candidate cloning of a QTL in dairy cattle: identification of a missense mutation in the bovine *DGAT1* gene with major effect on milk yield and composition. *Genome Res* **12**, 222–231 (2002).
12. Wang, S. *et al.* Control of grain size, shape and quality by *OsSPL16* in rice. *Nat Genet* **44**, 950–954 (2012).
13. Simons, K. J. *et al.* Molecular characterization of the major wheat domestication gene *Q*. *Genetics* **172**, 547–555 (2006).
14. Komatsuda, T. *et al.* Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class homeobox gene. *Proc Natl Acad Sci U S A* **104**, 1424–1429 (2007).
15. Fujii, J. *et al.* Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science* **253**, 448–451 (1991).
16. Kim, M. Y., Van, K., Kang, Y. J., Kim, K. H. & Lee, S. H. Tracing soybean domestication history: From nucleotide to genome. *Breeding Sci* **61**, 445–452 (2012).



17. Bactrian Camels Genome Sequencing and Analysis Consortium, Jirimutu *et al.* Genome sequences of wild and domestic bactrian camels. *Nat Commun* **3**, 1202 (2012).
18. He, Z. *et al.* Two evolutionary histories in the genome of rice: the roles of domestication genes. *PLoS Genet* **7**, e1002100 (2011).
19. Hufford, M. B. *et al.* Comparative population genomics of maize domestication and improvement. *Nat Genet* **44**, 808–811 (2012).
20. Andersson, L. How selective sweeps in domestic animals provide new insight into biological mechanisms. *J Intern Med* **271**, 1–14 (2012).
21. Lenser, T. & Theißen, G. Molecular mechanisms involved in convergent crop domestication. *Trends Plant Sci* **18**, 704–714 (2013).
22. Millar, C. D. & Lambert, D. M. Towards a million-year-old genome. *Nature* **499**, 34–35 (2013).
23. Zhang, H. *et al.* Morphological and genetic evidence for early Holocene cattle management in northeastern China. *Nat Commun* **4**, 2755 (2013).
24. Moles, A. T. *et al.* A brief history of seed size. *Science* **576**, 307 (2005).
25. Carter, T. E., Nelson, J. R., Sneller, C. H. & Cui, Z. Genetic diversity in soybean. In Boerma, H. R. & Specht, J. E. (eds.) *Soybeans: Improvement, Production and Uses*, *Am. Soc. of Agro, Madison, Wisconsin* pp303–416 (2004).
26. Schmutz, J. *et al.* Genome sequence of the paleopolyploid soybean. *Nature* **463**, 178–183 (2010).
27. Li, Q.-G., Zhang, L., Li, C., Dunwell, J. M. & Zhang, Y.-M. Comparative genomics suggests that an ancestral polyploidy event leads to enhanced root nodule symbiosis in the *Papilionoideae*. *Mol Biol Evol* **30**, 2602–2611 (2013).
28. Liu, B. *et al.* QTL mapping of domestication-related traits in soybean (*Glycine max*). *Ann Bot* **100**, 1027–1038 (2007).
29. Liang, H. Z. *et al.* Mapping quantitative trait loci for six seed shape traits in soybean. *Henan Agriculture Science* **45**, 54–60 (2008).
30. Xu, Y. *et al.* Mapping quantitative trait loci for seed size traits in soybean (*Glycine max* L. Merr.). *Theor Appl Genet* **122**, 581–594 (2011).
31. Hu, Z. *et al.* Determination of the genetic architecture of seed size and shape via linkage and association analysis in soybean (*Glycine max* L. Merr.). *Genetica* **141**, 247–254 (2013).
32. Xie, F.-T. *et al.* Fine mapping of quantitative trait loci for seed size traits in soybean. *Mol Breeding* online, DOI 10.1007/s11032-014-0171-7 (2014).
33. Niu, Y. *et al.* Association mapping for seed size and shape traits in soybean cultivars. *Mol Breeding* **31**, 785–794 (2013).
34. Liu, B. *et al.* Genetic redundancy in soybean photoresponses associated with duplication of the phytochrome A gene. *Genetics* **180**, 995–1007 (2008).
35. Watanabe, S. *et al.* Map-based cloning of the gene associated with the soybean maturity locus E3. *Genetics* **182**, 1251–1262 (2009).
36. Watanabe, S. *et al.* A map-based cloning strategy employing a residual heterozygous line reveals that the GIGANTEA gene is involved in soybean maturity and flowering. *Genetics* **188**, 395–407 (2011).
37. Xia, Z. *et al.* Positional cloning and characterization reveal the molecular basis for soybean maturity locus E1 that regulates photoperiodic flowering. *Proc Natl Acad Sci U S A* **109**, 2155–2164 (2012).
38. Jung, C. H., Wong, C. E. & Singh, M. B. Comparative genomic analysis of soybean flowering genes. *PLoS ONE* **7**, e38250 (2012).
39. Tian, Z. *et al.* Artificial selection for determinate growth habit in soybean. *Proc Natl Acad Sci U S A* **107**, 8563–8568 (2010).
40. Chen, Y. W. & Nelson, R. L. Genetic variation and relationships among cultivated, wild, and semiwild soybean. *Crop Sci* **44**, 316–325 (2004).
41. Kim, M. Y. *et al.* Whole-genome sequencing and intensive analysis of the undomesticated soybean (*Glycine soja* Sieb. and Zucc.) genome. *Proc Natl Acad Sci U S A* **107**, 22032–22037 (2010).
42. Stupar, R. M. Into the wild: The soybean genome meets its undomesticated relative. *Proc Natl Acad Sci U S A* **107**, 21947–21948 (2010).
43. Kim, M. Y., Shin, J. H., Kang, Y. J., Shim, S. R. & Lee, S. H. Divergence of flowering genes in soybean. *J Biosciences* **37**, 857–870 (2012).
44. Chung, W. H. *et al.* Population structure and domestication revealed by high-depth resequencing of Korean cultivated and wild soybean genomes. *DNA Res* **21**, 153–167 (2014).
45. Li, Y. H. *et al.* Molecular footprints of domestication and improvement in soybean revealed by whole genome re-sequencing. *BMC Genomics* **14**, 579 (2013).
46. Krichevsky, A., Zaltsman, A., Lacroix, B. & Citovsky, V. Involvement of KDM1C histone demethylase-OTL1 otubain-like histone deubiquitinase complexes in plant gene repression. *Proc Natl Acad Sci U S A* **108**, 11157–11162 (2011).
47. Zhang, X. Y. Chromatin modification in plants. In Wendel, J. F. *et al.* (eds), *Plant Genome Diversity Volume I* **15**, 237–255 (2012).
48. Stekhoven, D. J. *et al.* Causal stability ranking. *Bioinformatics* **28**, 2819–2823 (2012).
49. Wang, L. *et al.* NOT2 proteins promote polymerase II-dependent transcription and interact with multiple micro RNA biogenesis factors in Arabidopsis. *Plant Cell* **25**, 715–727 (2013).
50. Sun, X., Shantharaj, D., Kang, X. & Ni, M. Transcriptional and hormonal signaling control of Arabidopsis seed development. *Curr Opin Plant Biol* **13**, 611–620 (2010).
51. Gao, M. J., Gropp, G., Wei, S., Hegedus, D. D. & Lydiate, D. J. Combinatorial networks regulating seed development and seed filling. In: Sato, K. I. ed, *Embryogenesis*. Rijeka, Croatia: InTech; ISBN 979-953-307-439-8 (2012).
52. Xiao, W. *et al.* Regulation of seed size by hypomethylation of maternal and paternal genomes. *Plant Physiol* **142**, 1160–1168 (2006).
53. Ohto, M. A., Floyd, S. K., Fischer, R. L., Goldberg, R. B. & Harada, J. J. Effects of APETALA2 on embryo, endosperm, and seed coat development determine seed size in Arabidopsis. *Sex Plant Reprod* **22**, 277–289 (2009).
54. Zhou, Y. *et al.* SHORT HYPOCOTYL UNDER BLUE1 associates with MINISEED3 and HAIKU2 promoters in vivo to regulate Arabidopsis seed development. *Plant Cell* **21**, 106–117 (2009).
55. Canales, C., Bhatt, A. M., Scott, R. & Dickinson, H. EXS, a putative LRR receptor kinase, regulates male germline cell number and tapetal identity and promotes seed development in Arabidopsis. *Curr Biol* **12**, 1718–1727 (2002).
56. Schruff, M. C. *et al.* The AUXIN RESPONSE FACTOR 2 gene of Arabidopsis links auxin signalling, cell division, and the size of seeds and other organs. *Development* **133**, 251–261 (2006).
57. Garcia, D., Fitz Gerald, J. N. & Berger, F. Maternal control of integument cell elongation and zygotic control of endosperm growth are coordinated to determine seed size in Arabidopsis. *Plant Cell* **17**, 52–60 (2005).
58. Mizukami, Y. & Fischer, R. L. Plant organ size control: AINTEGUMENTA regulates growth and cell numbers during organogenesis. *Proc Natl Acad Sci U S A* **97**, 942–947 (2000).
59. Lam, H. M. *et al.* Resequencing of 31 wild and cultivated soybean genomes identifies patterns of genetic diversity and selection. *Nat Genet* **42**, 1053–1059 (2010).
60. Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS ONE* **3**, e3376 (2008).
61. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).
62. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297–1303 (2010).
63. Xia, Q. *et al.* Complete resequencing of 40 genomes reveals domestication events and genes in silkworm (*Bombyx*). *Science* **326**, 433–436 (2009).
64. Rousset, F. Genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol Ecol Resource* **8**, 103–106 (2008).
65. Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Mol Ecol Notes* **7**, 574–578 (2007).
66. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* **14**, 2611–2620 (2005).

Acknowledgments

This work was supported by the National Key Basic Research Program of China (grant 2011CB109300), Fundamental Research Funds for the Central Universities (grants KJQN201414, KJQN201422, KYT201002 and KYZ201202-9), the National Natural Science Foundation of China (grants 31301229 and 31301004), a Specialised Research Fund for the Doctoral Program of Higher Education (grant 20120097110023), the Priority Academic Program Development of Jiangsu Higher Education Institutions, and Huazhong Agricultural University Scientific & Technological Self-innovation Foundation (Program No. 2014RC020). The authors declare that they have no conflict of interest.

Author contributions

Y.-M.Z. designed the project. L.Z., S.-B.W., Q.-C.G., J.W., G.-J.L., Y.-Q.L., J.Z., J.-Y.F., Y.N., L.Z., W.-L.R. and Y.-M.Z. performed the experiments and analyzed the data. J.J. and Z.W. conducted DNA sequencing of all the accessions and analyzed the genetic diversity. Y.-Q.L. provided partly soybean materials. L.Z. prepared figures and tables. Y.-M.Z., L.Z., Q.S. and J.M.D. wrote the manuscript text. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhou, L. *et al.* Identification of domestication-related loci associated with flowering time and seed size in soybean with the RAD-seq genotyping method. *Sci. Rep.* **5**, 9350; DOI:10.1038/srep09350 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>