

RESEARCH ARTICLE

Open Access

Major depressive disorder seven years after the conflict in northern Uganda: burden, risk factors and impact on outcomes (The Wayo-Nero Study)

James Mugisha^{1,2,3*}, Herbert Muyinda¹, Samuel Malamba⁴ and Eugene Kinyanda^{5,6,7}

Abstract

Background: Major depressive disorder (MDD) is a major public health burden in conflict areas. However, it is not known for how long and by how much the observed high rates of MDD seen in conflict settings persist into the post-conflict period.

Methods: A cross sectional survey was employed seven years after the conflict in northern Uganda had ended in the three districts of Amuru, Gulu and Nwoya.

Results: The prevalence of major depressive disorder (MDD) was 24.7% (95% CI: 22.9%-26.4%). The distribution by gender was females 29.2% (95% CI: 14.6%-19.5%) and males 17.0% (95% CI: 26.9%-31.5%). The risk factors for MDD fell under the broad domains of socio-demographic factors (female gender, increasing age, being widowed and being separated/divorced); distal psychosocial vulnerability factors (being HIV positive, low social support, increasing war trauma events previously experienced, war trauma stress scores previously experienced, past psychiatric history, family history of mental illness, negative coping style, increasing childhood trauma scores, life-time attempted suicide, PTSD, generalized anxiety disorder and alcohol dependency disorder) and the psychosocial stressors (food insufficiency, increasing negative life event scores, increasing stress scores). 'Not receiving anti-retroviral therapy' for those who were HIV positive was the only negative clinical and behavioral outcome associated with MDD.

Conclusions: These findings indicate that post-conflict northern Uganda still has high rates for MDD. The risk factors are quite many (including psychiatric, psychological and social factors) hence the need for effective multi-sectoral programs to address the high rates of MDD in the region. These programs should be long term in order to address the long term effects of war. Longitudinal studies are recommended to continuously assess the trends of MDD in the region and remedial action taken.

Keywords: Major depressive disorder, Post-conflict, Northern Uganda, Risk factors, Outcomes

Background

Globally, non-communicable diseases such as mental illness, neurological and substance abuse disorders are recognized as a major public health burden amidst weak public health systems in post-conflict areas [1]. It has been observed that exposure to chronic civil conflict that is characterized by widespread human suffering and massive displacement is associated with high rates of major

depressive disorder (MDD) of between 39% and 97% [2-8]. It is however not known for how long and by how much these rates of MDD persists in the post-conflict situation- when there is cessation of conflict and the communities have gone back to a semblance of ordinary life. Bolton and colleagues [9] reported a rate of MDD of 15.5% five years after the genocide in Rwanda while Wong and colleagues [10] among Cambodian refugees assessed two decades after resettlement in the USA reported a rate of MDD 51%. In the latter case the situation of war trauma may have been confounded by acculturative stress which has been associated with depression [11].

* Correspondence: jmmugi77@hotmail.com

¹Makerere University, Child Health and Development Center, School of Health Sciences, P. Box 6717, Makerere Hill, Kampala, Uganda

²Butabika National Psychiatric Referral Hospital, P.o.Box 7017, Off Old Port Bell, Kampala, Uganda

Full list of author information is available at the end of the article

Several factors have been reported to be associated with MDD in conflict settings and these include: female gender [2-4,8,12-14]; indices of socio-economic disadvantage-widowhood, disability, being married for women, unemployment, no formal education, abject poverty, broken families, long periods of displacement, experiencing ill health without medical care [2,3,8,13]; and degree of exposure to traumatic events [2-5,8,13,15]. Apart from exposure to traumatic events reported by Wong and Colleagues [10], it is not known which other factors continue to contribute to the risk of MDD in the post-conflict situation after war has ceased and communities are re-settled.

MDD in conflict settings has been associated with impaired functioning [7,9] but it is not known for how long this persists in the post-conflict situation. Yet, this information is important in planning effective post conflict recovery programs, including public health interventions. It is also vital that the planning process for interventions in post areas is based on accurate data to improve program effectiveness.

In this study we undertook a community survey in post-conflict northern Uganda seven years after the 20 year civil conflict between the Lord's Resistance Army and the central government army (Uganda Peoples Defense Forces; UPDF) had ended, the population had gone back to their villages and the government is implementing a post-conflict re-construction program for the region. This conflict had led to tremendous suffering of the population including the forced displacement of over 2 million people (80% of the population in the region) into squalid living conditions in internally displaced persons camps; abduction of more than 20,000 children; and mass sexual traumatization of the population [16-21]. Studies of psychiatric disorder undertaken in this region during the conflict reported rates of MDD depression of between 31% and 67% with no known studies to date undertaken since the conflict ended [5,8]. Additionally, previous studies examined a limited range of risk factors for MDD, used non diagnostic tools to make psychiatric diagnoses and never examined the association between MDD and important negative clinical and behavior outcomes.

To address some of these methodological issues, we assessed for psychiatric disorder using a DSM IV based diagnostic tool, the Mini-International Neuropsychiatric Interview- M.I.N.I. [22] administered by trained psychiatric nurses. In this study we explored for the association between psychiatric disorder (including MDD) and a number of socio-demographic, social, clinical and psychological constructs. Our interest in exploring this wide range of risk factors is based on the notion that several studies conducted in the region after the war indicate that the population continues to be exposed to a wide range of psychosocial vulnerability factors [13,23]. We also explored

the relationship between psychiatric disorder with key psychiatric, clinical and behavior outcomes. In this paper we present our findings on major depressive disorder.

Methods

Study design and setting

This community survey was nested into a bigger project that is delivering a novel community based intervention that is using kinship relationships (Wayos [aunties] and Neros [uncles]) to increase the uptake of mental health services in post-conflict northern Uganda (Wayo-Nero) [24]. In this study we employed a cross sectional survey design where a representative sample was drawn from three mid- northern Uganda districts of Gulu, Amuru and Nwoya. These three districts were some of the most affected by the 20 year civil war in northern Uganda. Through a consultative process with community leaders at the district level of each of the selected study district, all the sub-counties in a given district were rated on how badly (intensity) affected they were by the civil conflict from the most affected to the least affected (the purpose of this grading process was to select the sub-counties which were to be involved in the main interventional study, and the same sub-counties were used in this survey). In each district, two sub counties rated as most affected by the civil war and having a health center III (a health centre at the level of the sub-county) or IV (a health centre at the level of a country) in their midst (a requirement for the main Wayo-Nero intervention) were selected. The sub-counties selected for this study were Lalogi and Koro (in Gulu district); Amuru and Atiak (in Amuru district); and Kochgoma and Alero (in Nwoya district).

Sampling

A multistage sampling method was used to select study respondents. Study respondents had to be residents in the selected sub-counties and 18 years and above in age. In each of the selected sub-counties a parish was randomly selected for inclusion in this study. A list of villages in each of the selected parishes was developed. All villages in the selected parishes were included in this study. Since there was no reliable source of information on the number of households in each village and on the number of adult individuals in each household, we could not develop a sampling frame based on households or individuals. Using the health centre III or IV as the central point in each parish, teams of research assistants fanned out in the 4 directions of the compass to interview study respondents. All adult members (18 years and above) in homesteads lying along a village road in one direction of the compass were visited and interviewed for this study by one of the teams of research assistants. All households within a particular direction were exhausted first before taking

another direction. During this movement, both near and far households from the road side were selected. However, we excluded from interviews all community members who had not stayed in the area for more than six months and, those who could not fluently speak Luo (the local dialect and could not respond to the translated tools). We also excluded those from the survey, community members that were drunk or too sick. In each household, we interview all the adult members who met the inclusion criteria. In case someone was missing at the time of the interviews, call back visits were made. However, due to resource limitations we could not make more than two call back visits to the same household. We did not find any refusals during data collection possibly because we were working alongside community gatekeepers-community leaders. Interviews in each parish were conducted until a total number of 400 study respondents had been attained (in 2 sub-counties the selected parish could not provide all the required 400 respondents so data collection was extended into the neighboring parish of the same sub-county). Study interviewers were all psychiatric nurses who had been trained on the objectives of the study, consenting procedures, administration of the study questionnaire and the referral process for those respondents found to have significant psychiatric problems.

Sample size calculation

A multistage sampling procedure was used to draw up a representative sample from each of the six study sub-counties. A sample size of 400 respondents per sub-county was calculated using the formula: $n = \left(\frac{Z}{m}\right)^2 \hat{p}(1-\hat{p}) \times \frac{deff}{R}$ (Kirkwood and Sterne, [25]). To take care of possible non-response (R) and sampling design (*deff*), assuming a design effect (*deff*) of 2 and a non-response rate (R) of 0.987%, '*m*' level of precision desired. A sample size (n) of 2,400 was calculated to produce a two-sided 95% confidence interval ranging from 26.7% to 31.9% when the sample percentage (P) is 29.3%.

Data collection tools

A structured questionnaire consisting of various standardized tools previously employed in the Ugandan context by one of the investigators (EK) was used in this study [19,26,27]. The different assessment tools in the study questionnaire we translated into the local Luo language by a process of forward and backward translation by two teams of mental health professionals working independently of each other. A consensus meeting was then held to resolve those items where there was a wide disparity in the two translations.

The sections of the study questionnaire were:

- 1) Ecological factor: *The sub-county*
- 2) Socio-demographic factors: *sex, age, marital status, highest educational attainment, religion, and employment status.*
- 3) Social factors i). '*Food insecurity*' (assessed by the question, 'in the last month, did you or your family have enough food?'); ii) '*Negative life events index*', constructed from items of the adverse life events module of the European Parasuicide Interview Schedule [28] that has previously been modified for the Ugandan situation by Kinyanda and colleagues [26]. A total score was generated to reflect the total number of life events reported; iii) '*War trauma events index*': These items were derived from the commonly reported forms of war trauma in Uganda including loss of loved ones events, sexual trauma events, physical trauma events and psychological trauma events [19,29]; iv) '*Stress Score indices*', were constructed both for 'negative life events' and for 'war trauma events'. Each of the items of the modules on 'negative life events' and for 'war trauma experiences' was scored on a 3 point Likert scale where respondents were asked the question, 'how stressful did you find the event?' with possible responses being: 0 = (not stressing/minimal stressing), 1 = (moderately stressing), 2 = (severely stressing). A total score was generated for each module where high scores reflected more stress; v) '*Social support index*' was assessed using the Multidimensional Scale of Perceived Social Support (MSPSS) [30]. The MSPSS is a 12-item instrument that was designed to assess perceptions about support from family, friends and a significant other. Originally designed to have each item on this scale graded according to a 7 point Likert Scale, this was shortened to 5 choices namely, 1 = 'strongly disagree'; 2 = 'mildly disagree'; 3 = 'mildly agree'; 4 = 'strongly agree'; 5 = 'if you very strongly agree' as some choices did not make sense in the local cultural context. The α -Cronbach of this scale in this study was 0.92.
- 4) Psychological and clinical factors: i) *Psychiatric disorders* were assessed using the M.I.N.I. neuropsychiatric interview (MINI Plus) which is a modular DSM IV based structured interview [22]. The psychiatric disorder modules used in this study were: major depressive disorder, suicidality, alcohol dependency/ abuse disorders, generalized anxiety disorder and post-traumatic stress disorder; ii) *Family history of psychiatric disorder* was assessed by the question, 'has any member of your family (immediate family- parent, sibling, child) ever been diagnosed with a psychiatric disorder?'; iii) *Past psychiatric history* by asking the question, 'have you ever suffered from any nervous or psychiatric

- condition?'; iv) *Adverse life events in childhood* was assessed using the Childhood Trauma Questionnaire-Short Form (CTQ-SF) [31]. This is a 28 item questionnaire that asks about negative life events experienced as a child and as an adolescent. The respondent is required to respond based on a 5 point Likert format where 1 = 'never true'; 2 = 'rarely true'; 3 = 'sometimes true'; 4 = 'often true'; 5 = 'very often true'. A total score was generated. To ensure that all the items of the CTQ-SF are scored in the same direction, the following items were reverse scored [2,5,7,10,13,16,19,22,26], and 28. The α -Cronbach of this scale in this study was 0.89; v) *Positive Coping style index*. This was assessed using the Mental Adjustment to Cancer Scale (MAC) [32] whose items have been previously adapted to the Uganda situation [33]. Each of this scale's 17 questionnaire items is scored on a 4-item Likert scale 1 = (definitely does not apply to me), 2 = (does not apply to me), 3 = (applies to me), 4 = (definitely applies to me). To score all the questionnaire items so that they are all in the same direction i.e. higher scores reflecting less negative coping style, questionnaires items that were cast negatively were reverse scored at analysis. A total score was generated so that higher scores reflected a more negative coping style, the α -Cronbach of this scale in this study was 0.93; vi) *Self report on HIV status*. This was assessed by asking the question, 'what is your HIV status'.
- 5) Clinical and behavioral outcomes. The following clinical and behavioral outcomes were assessed in this study: i) 'Whether currently receiving treatment for psychiatric illness' by asking those who reported a past history of ever suffering from any nervous or psychiatric condition the question, 'are you currently receiving treatment for the above condition'; ii) for those who are HIV positive, 'whether they were receiving HIV treatment' by asking the question, 'are you receiving treatment for HIV/AIDS?'; iii) A *High risk sexual behavior index* was constructed by asking respondents whether in the past month they had engaged in high risk sexual practices that have been associated with HIV transmission in the Ugandan cultural context as reported by various HIV sero-behavioral surveys [33-35], the sexual behavioral practices assessed were: 'have you had sex with anyone other than your regular partner?', 'have you engaged in sex in exchange for gifts/money?', 'have you engaged in forced sex including rape?', 'have you engaged in sex with someone much older than you who is not your partner?', 'have you engaged in sex with someone you have just met?'. The high risk sexual behavior index was positive if a respondent reported at least one of these items.

Statistical analysis

Data was analyzed using STATA software (StataCorp LP, 4905 Lakeway Drive, College Station, TX, USA). Logistic regression models were used to assess univariate associations between the dependent variable (major depressive disorder) and the independent variables including socio-demographic, social, psychological and clinical factors. Unadjusted Odds Ratio and those adjusted for sex and age are reported. Analyses accounted for the multi-sampling design of the study including stratification, unequal selection probabilities, and clustering. Survey analyses using the 'svyset' command and the 'svy' prefix/procedures in STATA were used to account for the study design. The primary sampling unit (PSU) was the parish and the stratification variables include the 9 strata (6 certainty primary frame sub-county unit strata plus three district strata). The sampling weights were based on the selection probabilities at each level of selection. All factors associated with MDD at a level of significance of ≤ 0.1 in the bivariate models were entered into a multivariable model to determine their independent effect on MDD. A backward elimination method that dropped, at each step, all variables that did not reach the <0.05 level of significance was used to develop the final multi-variable model. Only those factors which remained associated with MDD at $p < 0.05$ were retained in the final model.

Ethical considerations

Approvals were obtained from Makerere University College of Health Sciences Science and Ethical Committee. We also secured ethical approval from the Uganda National Council for Science and Technology. Study approvals from all participating district authorities were also secured. Informed consent was obtained from all study respondents after explaining the objectives and procedures of the study. Informed consent was secured from the respondents before they could be subjected to the study tools. Written consent was secured from those (respondents) who could read and write. For respondents who could not read and write, we used verbal consent. In addition to the verbal consent, they could put their thumb print on the consent paper. Study respondents found to have significant psychiatric problems were referred to the health centre III or IV [16]. The parent interventional study had helped establish a psychiatric service in all health centre III and IV's in the survey sub-counties. Respondents were assured of confidentiality before the start of each interview. All data collected was kept under key and lock and only accessible to the research team. Personal identifiers were not used at data entry.

Results

Study population

Fieldwork was conducted between 2nd January 2013 and 2nd June 2013 where a total of 2,406 respondents were

enrolled into this study (819 from Amuru district, 770 from Gulu district and 817 from Nwoya district). At analysis, a total of 45 records were excluded because of missing data on key variables, the excluded records did not differ significantly from those retained on district, age and sex.

Socio-demographic characteristics of the respondents

Information in this study is adduced from a total sample of 2361 population of 1475 (62.5%) females and 886 (37.5%) males. Table 1, the study participants were distributed in nearly equal proportions among the age

groups of 18-24 years (552, 23.8%), 25-34 years (628,27.1%), 35-44 years (479, 20.7%) and those 45 years and above (659, 28.4%). Majority of respondents 658 (69.1%) were married with only 172 (7.2%) single or separated. Majority of respondents 1423 (59.9%) had only attained a primary level education with only 393 (16.5%) having attained a secondary level education and above. On religion, majority were Catholics 1872 (78.2%) with Protestants 304 (12.7%) and Moslems 12 (0.5%). In terms of employment, majority were peasant farmers at 1917 (80%) with only 32 (1.3%) employed as salary earners.

Table 1 Socio-demographic correlates of Major Depressive Disorder in the Wayo-Nero study population, Northern Uganda

	Number in study (N, %)	Current major depressive disorder (n) n, (n/N%)	Unadjusted OR (95% CI)	Adjusted OR ^S (95% CI)	P-value
Sex					
Male	886 (37.5)	151 (17.0)	1	1	
Female	1475 (62.5)	431 (29.2)	2.0 (1.63-2.48)	2.12 (1.71-2.63)	<0.001
Age (Years)					
18-24	552 (23.8)	95 (17.2)	1	1	
25-34	628 (27.1)	147 (23.4)	1.47 (1.10-1.96)	1.43 (1.07-1.92)	0.017
35-44	479 (20.7)	125 (26.1)	1.70 (1.26-2.30)	1.71 (1.26-2.33)	0.001
45+	659 (28.4)	215 (32.6)	2.33 (1.76-3.08)	2.33 (1.76-3.09)	<0.001
Marital Status					
Currently married/cohabiting	1649 (69.1)	385 (23.4)	1	1	
Widowed	294 (12.3)	112 (38.1)	2.02 (1.55-2.63)	1.55 (1.15-2.09)	0.004
Separated/Divorced	172 (7.2)	64 (37.2)	1.95 (1.40-2.71)	1.95 (1.38-2.74)	<0.001
Single	273 (11.4)	32 (11.7)	0.44 (0.30-0.64)	0.62 (0.39-0.98)	0.039
Education					
No education	569 (23.9)	184 (32.3)	1	1	
Primary only	1423 (59.6)	368 (25.9)	0.73 (0.59-0.90)	1.02 (0.80-1.29)	0.874
Secondary/above	393 (16.5)	45 (11.5)	0.27 (0.19-0.39)	0.55 (0.36-0.82)	0.004
Religion					
Catholics	1872 (78.2)	472 (25.2)	1	1	
Protestants	304 (12.7)	68 (22.4)	0.85 (0.64-1.14)	0.85 (0.69-1.38)	0.291
Muslims	12 (0.5)	2 (16.7)	0.59 (0.13-2.72)	-	
SDA/Pentecostal/Other	207 (8.6)	56 (27.1)	1.10 (0.80-1.52)	0.98 (0.69-1.38)	0.888
Employment					
Peasant farmer/Fisherman	1917 (80.0)	508 (26.5)	1	1	
housewife	129 (5.4)	32 (24.8)	0.92 (0.61-1.38)	0.80 (0.52-1.22)	0.292
Professional/clerical	32 (1.3)	3 (9.4)	0.29 (0.09-0.95)	0.41 (0.12-1.38)	0.151
Tradesperson/artisan/transport	56 (2.3)	11 (19.6)	0.68 (0.35-1.32)	0.96 (0.47-1.96)	0.909
Student/ unemployed but able	137 (5.7)	10 (7.3)	0.22 (0.11-0.42)	0.36 (0.17-0.77)	0.008
Unemployed and disabled	80 (3.3)	22 (27.50)	1.05 (0.64-1.74)	0.76 (0.44-1.30)	0.312
Others	44 (1.8)	9 (20.5)	0.71 (0.34-1.49)	1.09 (0.51-2.34)	0.825

Key: ^SAdjusted for sex and age.

Prevalence of major depressive disorder (MDD)

The prevalence of major depressive disorder (MDD) in this study population was 24.7% (95% CI: 22.9%-26.4%). The distribution by gender was 17.0% (95% CI: 14.6%-19.5%) for males and 29.2% (95% CI: 26.9%-31.5%) for females.

Socio-demographic correlates of MDD

Table 1, shows the socio-demographic correlates for MDD. The following socio-demographic factors were significantly associated with MDD: female gender 2.1 (95% CI: 1.71-2.63); increasing age when compared with those in the age group of 18-24 years: 25-34 years age group 1.43 (95% CI: 1.07-1.92), 35-44 years age group 1.71 (95% CI: 1.26-2.33) and 45+ years age group 2.33

(95% CI: 1.76-3.09); being widowed 1.55 (95% CI: 1.15-2.09) and separated/divorced 1.95 (95% CI: 1.38-2.74) when compared with those who were married; and being a student/ 'unemployed but able bodied' was protective against MDD 0.36 (95% CI: 0.17-0.77) when compared against being a peasant farmer.

Social correlates for major depressive disorder (MDD) at multi-variable analysis

Table 2, shows the social factors that were correlated with MDD. The social factors that were significantly associated with MDD were: food insecurity 1.99 (95% CI: 1.61-2.45); self-report of being HIV positive 2.63 (95% CI: 1.87-3.70); low social support 0.41(95% CI: 0.34-0.51), increasing negative life events scores, with those reporting

Table 2 Social correlates of Major Depressive Disorder in the Wayo-Nero study population, Northern Uganda

	Number in study (N, %)	Major depressive disorder (n, %)	Unadjusted OR (95% CI)	Adjusted OR ^s (95% CI)	P-value
Food Security					
Enough	1002 (43.3)	173 (17.3)	1	1	
Not enough	1310 (56.7)	407 (31.1)	2.16 (1.76-2.65)	1.99 (1.61-2.45)	<0.001
Reported HIV Status					
Negative	1798 (75.9)	403 (22.4)	1	1	
Positive	164 (6.9)	74 (45.1)	2.85 (2.04-3.96)	2.63 (1.87-3.70)	<0.001
Don't know	407 (17.2)	114 (28.0)	1.35 (1.06-1.72)	1.17 (0.89-1.52)	0.261
Social support index					
Low (< median)	1176 (49.0)	398 (33.8)	1	1	
High (Median plus)	1223 (51.0)	200 (16.4)	0.38 (0.31-0.46)	0.41 (0.34-0.51)	<0.001
Negative life events index					
Low (0-5 events)	611 (25.4)	66 (10.8)	1	1	
Medium (6-9 events)	1347 (56.0)	334 (24.8)	2.72 (2.05-3.62)	2.97 (2.19-4.04)	<0.001
High (10-18 events)	448 (18.6)	199 (44.4)	6.60 (4.81-9.05)	7.26 (5.17-10.20)	<0.001
Negative life events stress index					
0	260 (10.8)	13 (5.0)	1	1	
1-5	517 (21.5)	59 (11.4)	2.45 (1.32-4.55)	2.20 (1.14 -4.22)	0.018
6-10	710 (29.5)	149 (30.0)	5.05 (2.81-9.07)	4.99 (2.71-9.19)	<0.001
11-15	546 (22.7)	188 (34.4)	9.98 (5.56-17.91)	9.16 (4.97-16.88)	<0.001
16+	373 (15.5)	190 (50.9)	19.73 (10.90-35.71)	20.37 (10.95-37.87)	<0.001
War trauma events index					
None	410 (17.4)	47 (11.5)	1	1	
1-3 events	763 (32.3)	155 (20.3)	1.97 (1.386-2.798)	1.92 (1.340-2.739)	<0.001
4-7 events	576 (24.4)	167 (29.0)	3.15 (2.215-4.489)	2.83 (1.971-4.078)	<0.001
8+ events	612 (25.9)	213 (34.8)	4.12 (2.916-5.830)	4.68 (3.255-6.727)	<0.001
War trauma stress index					
Low	429 (18.2)	49 (11.4)	1	1	
Medium	1128 (47.8)	251 (22.3)	2.22 (1.597-3.084)	2.07 (1.480-2.893)	<0.001
High	804 (34.1)	282 (35.1)	4.19 (3.009-5.833)	4.42 (3.130-6.245)	<0.001

Key: ^sAdjusted for sex and age.

6-9 events having an aOR of 2.97 (95% CI: 2.19-4.04), those reporting 10-18 events having an aOR of 7.26 (95% CI: 5.17-10.20) when all were compared with those reporting 0-5 events; increasing negative life events stress scores, those with scores of 6-10 had an aOR of 4.99 (95% CI: 2.71-9.19), those with scores of 11-15 had an aOR of 9.16 (95% CI: 4.97-16.88) and those with scores of 16+ had an aOR of 20.37 (95% CI: 10.95-27.87) when all were compared with those with zero scores; increasing war trauma event scores, those reporting 1-3 events had an aOR of 1.92 (1.34-2.74), those reporting 4-7 events had an aOR of 2.83 (1.97-4.08) and those reporting 8+ events had an aOR of 4.68 (3.26-6.73) when all were compared with those reporting no war trauma event; and increasing war trauma stress scores, with those reporting medium scores having an aOR of 2.07 (95% CI: 1.48-2.89), those reporting high scores an aOR of 4.42

(95% CI: 3.13-6.25) when all were compared with those reporting low scores.

Clinical and psychological factors associated with major depressive disorder

Table 3, shows the psychological and clinical factors associated with MDD. The factors that were significantly associated with MDD were: positive coping style (protective against MDD), with those with medium scores having an aOR of 0.23 (95% CI: 0.18-0.30), those with high scores having an aOR of 0.11 (95% CI: 0.08-0.14) all compared with those with low scores; past psychiatric history 3.64 (95% CI: 2.62-5.05); increasing childhood trauma scores, those with a score of 3 having an aOR of 2.07 (95% CI: 1.66-2.59), those with a score of 4-5 having an aOR of 3.27 (95% CI: 2.04-5.23) all compared with those with a score of 1-2; and a family history of mental illness 1.65

Table 3 Psychological and clinical correlates of Major Depressive Disorder in the Wayo- Nero study population, Northern Uganda

	Number in study (N,%)	Major depressive disorder (n,%)	Unadjusted OR (95% CI)	Adjusted ORs(95% CI)	p-value
Positive Coping Style index					
Low	479 (19.9)	284 (59.3)	1	1	
Medium	705 (29.3)	164 (23.3)	0.21 (0.16-0.27)	0.23 (0.18-0.30)	<0.001
High	1222 (50.8)	151 (12.4)	0.10 (0.08-0.12)	0.11 (0.08-0.14)	<0.001
Past Psychiatric history					
No	2156 (90.1)	485 (22.5)	1	1	
Yes	173 (7.2)	89 (51.5)	3.65 (2.66-5.00)	3.64 (2.62-5.05)	<0.001
Don't Know	64 (2.7)	22 (34.4)	1.80 (1.07-3.05)	1.54 (0.89-2.68)	0.127
Childhood Trauma index					
1-2	1668 (70.7)	353 (21.2)	1	1	
3	608 (25.8)	193 (31.7)	1.73 (1.41-2.13)	2.07 (1.66-2.59)	<0.001
4-5	85 (3.6)	36 (42.4)	2.74 (1.75-4.28)	3.27 (2.04-5.23)	<0.001
Family history of psychiatric illness					
No	1800 (76.1)	410 (22.8)	1	1	
Yes	566 (23.9)	180 (31.8)	1.58 (1.28-1.95)	1.65 (1.33-2.06)	<0.001
Life-time attempted suicide					
No	2389 (99.3)	585 (24.5)	1	1	
Yes	17 (0.7)	14 (82.4)	14.39 (4.12-50.25)	13.52 (3.78-48.33)	<0.001
Posttraumatic Stress disorder					
No	2122 (88.2)	447 (21.1)	1	1	
Yes	284 (11.8)	152 (53.5)	4.31 (3.34-5.57)	4.79 (3.65-6.29)	<0.001
Generalized anxiety disorder					
No	2051 (85.2)	372 (18.1)	1	1	
Yes	355 (14.8)	227 (63.9)	8.00 (6.27-10.22)	7.48 (5.81-9.63)	<0.001
Alcohol dependency disorder					
No	2244 (93.7)	554 (24.7)	1	1	
Yes	162 (6.7)	45 (27.8)	1.17 (0.82-1.68)	1.51 (1.01-2.26)	0.046

(95% CI: 1.33-2.06). Additionally, the following psychiatric disorders/problems were significantly associated with MDD: life-time attempted suicide 13.52 (95% CI: 3.78-48.33); PTSD 4.79 (95% CI: 3.65-6.29); generalized anxiety disorder 7.48 (95% CI: 5.81-9.63) and alcohol dependency disorder 1.51 (1.01-2.26).

Multivariate analysis of correlates of major depressive disorder

According to Table 4 the following risk factors retained statistical significance with MDD at multivariate analysis: female gender, self-report of being HIV positive, low social support scores, increasing number of negative life events, increasing negative life events stress scores, decreasing positive coping style scores and the psychiatric comorbidities of life-time attempted suicide, PTSD and generalized anxiety disorder.

Association between clinical and behavioural outcomes with major depressive disorder

Table 5, shows the association between MDD and various clinical and behavioural outcomes. Not receiving anti-retroviral therapy for those with HIV was the only outcome that was significantly associated with MDD aOR3.22 (95% CI: 1.08-9.57).

Discussion

In this study we investigated the burden of major depressive disorder (MDD) and its association with risk factors and negative outcomes as seen in northern Uganda seven years after the cessation of one of the worst conflicts in the world. The principal finding of this study is that post-conflict northern Uganda still has high rates of MDD although these seem to have reduced in the post-conflict time period. There is still poor uptake of mental health services. This can largely be attributed to the general lack of health services and other factors that affect uptake of mental health services such as traditional beliefs and stigma [24]. The risk factors for MDD fell under the broad domains of socio-demographic factors (female gender, increasing age, being widowed and being separated/divorced); distal psychosocial vulnerability factors (being HIV positive, low social support, increasing war trauma events previously experienced, war trauma stress scores previously experienced, past psychiatric history, family history of mental illness, negative coping style, increasing childhood trauma scores, life-time attempted suicide, PTSD, generalized anxiety disorder and alcohol dependency disorder) and proximal psychosocial stressors (food insufficiency, increasing negative life event scores and increasing stress scores). 'Not receiving anti-retroviral therapy' for those who were HIV positive was the only

Table 4 Multivariate model of risk factors for major depressive disorder in the Wayo-Nero study population, Northern Uganda

	Final model		
	Odds ratio	(95% CI)	P-value
Sex (Baseline group – Male)			
Female	1.63	1.25-2.11	<0.001
Age-Group (Baseline group – 18-24)			
25-34	1.06	0.75-1.52	0.728
35-44	1.37	0.95-1.98	0.090
45+	1.33	0.94-1.87	0.102
Self-reported HIV Status			
Negative/Don't Know			
Positive	1.83	1.22-2.74	0.003
Social Support score High (Median and above)			
(Low - < median)	1.35	1.05-1.75	0.018
Life events stress index (Baseline group – 0)			
1-5	1.77	0.83-3.78	0.140
6-10	2.74	1.27-5.89	0.010
11-15	4.26	1.96-9.28	<0.001
16-48	6.63	2.80-15.66	<0.001
Negative life events (Baseline group: 0-5 events)			
6-9 events	1.61	1.05-2.44	0.027
10-18 events	1.60	0.91-2.82	0.099
Positive Coping Style index (Baseline –Low)			
Medium	0.28	0.21-0.39	<0.001
High	0.19	0.14-0.26	<0.001
Life-time attempted suicide			
Present	6.18	1.53-25.01	0.011
Post-Traumatic Stress Disorder (PTSD)			
Present	1.90	1.37-2.65	<0.001
Generalized anxiety disorder (GAD)			
Present	4.09	3.06-5.49	<0.001

negative clinical and behavioral outcome associated with MDD in the study.

In this study there were more females (62.5%) than males (37.5%) with MDD. Roberts and colleagues [8] in war conflict northern Uganda observed a similar gender ratio. A similar gender ratio is reported by Sheikh et al. in their study among internally displaced persons in Kaduna, north of Western Nigeria [36]. The majority of respondents were less than 45 years of age (71.6%), employed as peasant farmers (80%) and married (69.1%). There was

Table 5 Association between Major Depressive Disorder and clinical and behavioral outcomes as seen in the Wayo-Nero study population, Northern Uganda

	Number in study (N,%)	Major depressive disorder (n,%)	Unadjusted OR (95% CI)	Adjusted ORS (95% CI)	p-value
Whether currently receiving treatment for psychiatric illness (n = 173)					
No	159 (91.9)	81 (50.9)	1		
Yes	14 (8.1)	8 (57.1)	1.284 (0.426-3.870)	1.103 (0.349-3.480)	0.868
Whether receiving HIV treatment (n = 164)					
Yes	139 (84.8)	59 (42.5)	1		
No	25 (15.2)	15 (60.0)	2.034 (0.854-4.845)	3.218 (1.082-9.570)	0.036*
Whether receiving treatment for the epilepsy (n = 66)					
No	40 (61.6)	11 (27.5)	1		
Yes	26 (39.4)	9 (34.6)	1.396 (0.481-4.049)	1.789 (0.536-5.973)	0.344
High risk sexual behaviour (n = 139)					
No	2223 (94.2)	545 (24.5)	1		
Yes	139 (5.8)	37 (26.8)	1.128 (0.765-1.664)	1.374 (0.905-2.086)	0.136

Key: †Adjusted for sex, age and education.

however a considerable proportion who were widowed (12.3%) and separated/divorced (7.2%) all potential consequences of chronic war conflict. In this study 6.9% of the respondents self-reported that they were HIV positive with the majority (84.8%) reporting that they were receiving HIV/AIDS care. The self-reported prevalence of HIV in this study is close to the figure of 8.3% for HIV sero-prevalence for the mid-northern Uganda region (area where the survey was conducted) [37]. About a tenth (7.3%) of the respondents self-reported that they had ever suffered from a nervous or psychiatric problem of these only 8% reported that they had ever sought treatment for this condition from a health facility indicating that many of such cases are not reported to the modern health system. Kinyanda and colleagues in non-war affected south-western Uganda reported a prevalence of self-reported severe mental illness of 0.9% [27]. Other Ugandan studies [38,39] have noted that less than 20% of all persons with mental illness are treated at formal health services in the country. Closing the treatment gap for mental illness (as indicated in this study) is a challenge in poor communities such as northern Uganda [40].

The prevalence of major depressive disorder (MDD) in this study was 24.7%. This prevalence figure did not change much (increased to 29.1%) when a similar criteria used in previous studies in northern Uganda was applied to the data (where MDD was defined as meeting only the symptom criteria of the MDD module in the MINI excluding the function assessment dimension). In previous studies undertaken during the conflict in northern Uganda using non diagnostic assessment tools, prevalence figures for MDD of between 45% and 67% were reported which are about twice those observed in this study [5,8]. This

difference in the figures reported during the conflict and seven years after the conflict (period for the current study) suggest that the rates of MDD in northern Uganda may have reduced after the conflict. We obtained low prevalence rates largely due to two factors; a) we undertook MDD assessment using a diagnostic criteria and therefore we were more likely to get low prevalence rates. Kuwert et al. [41] urges that approaches that use symptom assessment normally provides presumptive rates of prevalence of mental health conditions [41] b) prevalence rates normally change over time.

On risk factors of MDD in this study, these will be discussed under socio-demographic factors, social factors and psychological and clinical factors. Most of the socio-demographic factors significantly associated with MDD in this study were indices of socio-economic disadvantage in this post-conflict situation in northern Uganda, these were: female gender, increasing age, being widowed, separated and divorced, and having no formal education, with female gender retaining statistical significance in the final multivariate analysis. Previous studies in conflict and post-conflict settings have reported the following socio-demographic factors to be associated with MDD: female gender [2-4,8,12,14]; increasing age [42]; being widowed, separated and divorced [2,8,43] and having no formal education [44].

The following social factors were significantly associated with MDD in this study: food insufficiency, being HIV positive, poor social support, increasing number of negative life events, increasing life events stress scores, increasing number of reported war trauma events and increasing war trauma events stress scores. As communities are working towards recovery, they still grapple with the long

term effects of the war on the individual, family and the entire community system. In the final multivariate model HIV positive status, poor social support, increasing number of negative life events and the associated stress scores retained statistical significance. Previous studies in conflict settings have reported the following social factors as significantly associated with MDD: degree of exposure to traumatic events [2-5,8,13,15,16]; negative life events [13]; and poor social support [45,46]. From studies undertaken in non-conflict settings HIV positive status [47] and food insufficiency [44] have been reported to be significantly associated with MDD.

The following psychological and clinical factors were significantly associated with MDD in this study: negative coping style, past psychiatric history, childhood trauma, family history of mental illness and the comorbid psychiatric problems of lifetime attempted suicide, PTSD, generalized anxiety disorder and marginally alcohol dependency disorder. At multivariate analysis the psychological and clinical factors that remained significantly associated with MDD were negative coping style and the comorbid psychiatric disorders of PTSD and generalized anxiety disorder. Previous studies in conflict settings have reported negative coping style [48] and psychiatric comorbidities [23,49,50] as risk factors for MDD. Sheikh et al. while focusing on a post-conflict setting in Kaduna Northwestern Nigeria reported the co-morbidity of PTSD and depression (those that had PTSD were more likely to have depression) [51].

Among the investigated negative outcomes, it was only poor uptake of antiretroviral therapy (ART) that was significantly associated with MDD. Gourlay and colleagues [50] in a systematic review observed that MDD was one of the barriers against the uptake of antiretroviral drugs in prevention of mother-to-child transmission of HIV programs in sub-Saharan Africa [51]. Similarly, Olisah et al. [52] reported that 42.3% of their respondents with depressive disorder did not use their medications regularly [52].

Study limitations

In this study, MDD was associated with poor uptake of antiretroviral therapy (ART). But given the cross sectional nature of this study, it was not possible to elucidate the direction of causality between these investigated factors and MDD. This could be better understood if this was a longitudinal study.

We were not able to develop a sampling frame because there were no updated lists of community members at community level in most villages. This implies that the total sample might be a true representation of the target population. However using the health centers as a central place, we could find out in one direction until all households in that direction were completed. We could then turn to another direction and the same process was

repeated throughout the study. This meant that households near and far from the roads were included into the sample.

We used Psychiatric nurses to administer the study tools. It is plausible that more accurate data on diagnosis assessment could have been obtained if we had used highly trained cadres in mental health. Though this was unavoidable given the fact that there are few mental health workers in the region of higher qualifications (by the time this study was conducted northern Uganda had only three psychiatrists and the out of three, two had retired and were only working with the University in Gulu). The region also had one clinical psychologist). We however, used a support team of psychologists and psychiatrists from Makerere University to supervise data collection and offer regular training from time to time.

Conclusion

This study showed that post-conflict northern Uganda still had high rates of MDD. The risk factors for MDD fell under the broad domains of socio-demographic factors (female gender, increasing age, being widowed and being separated/divorced); distal psychosocial vulnerability factors (being HIV positive, low social support, increasing war trauma events previously experienced, war trauma stress scores previously experienced, past psychiatric history, family history of mental illness, negative coping style, increasing childhood trauma scores, life-time attempted suicide, PTSD, generalized anxiety disorder and alcohol dependency disorder) and proximal psychosocial stressors (food insufficiency, increasing negative life event scores, increasing stress scores) in line with the stress-vulnerability model of depression [36].

Overall, given the high level of vulnerability in the study areas to MDD due to a number of factors (including psychiatric, psychological and social factors) there is the need for effective multi-sectoral programs tailored towards addressing the high rates of MDD in the region. Longitudinal studies are recommended to continuously assess the trend/rates of MDD in the region and remedial action taken.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Each author made substantive contribution towards the development of this paper. JM is a principle investigator for this study, coordinated all the data collection, conceptualized the paper and made substantive contribution in the drafting and writing up of the paper. HM is also a principle investigator for the study, supported data collection and writing up this paper. SM did all the statistical analysis and participated in writing the paper. EK contributed substantively in designing the study tools, training of research assistants, and in revising the manuscript for intellectual content. All authors read and approved the final manuscript.

Acknowledgements

This project was funded by Grand Challenges Canada and Bill & Melinda Gates Foundation (GMH 0094-04). We are grateful to all our respondents in the three study districts. We are also grateful to Drs. Nshemerirwe, S, Kizza D, Mutamba, B for supervising some of the fieldwork during data collection. We further appreciate all our research assistants.

Author details

¹Makerere University, Child Health and Development Center, School of Health Sciences, P. Box 6717, Makerere Hill, Kampala, Uganda. ²Butabika National Psychiatric Referral Hospital, P.o.Box 7017, Off Old Port Bell, Kampala, Uganda. ³Sør-Trøndelag University College, E. C. Dahls gate 2, 7012 Trondheim, Norway. ⁴Uganda Virus Research Institute (UVRI) HIV Reference Laboratory Program, 50-59 Nakiwogo Street, Entebbe, Uganda. ⁵MRC/UVRI Uganda Research Unit on AIDS, Uganda/MRC-DFID African Leadership Award, 50-59 Nakiwogo Street, Entebbe, Uganda. ⁶Department of Psychiatry, Makerere University College of Health Sciences, School of Health Sciences, Makerere Hill, Kampala, Uganda. ⁷London School of Hygiene & Tropical Medicine, Keppel Street, London WC1E, 7HT, UK.

Received: 6 August 2014 Accepted: 19 February 2015

Published online: 14 March 2015

References

- Roberts B, Patel P, McKee M. Noncommunicable diseases and post-conflict countries. *Bull World Health Organ.* 2012;90(1):2. 2A.
- Sabin M, Lopes Cardozo B, Nackerud L, Kaiser R, Varese L. Factors associated with poor mental health among Guatemalan refugees living in Mexico 20 years after civil conflict. *JAMA.* 2003;290(5):635–42.
- Cardozo BL, Bilukha OO, Crawford CA, Shaikh I, Wolfe MI, Gerber ML, et al. Mental health, social functioning, and disability in postwar Afghanistan. *JAMA.* 2004;292(5):575–84.
- Scholte WF, Olff M, Ventevogel P, de Vries GJ, Jansveld E, Cardozo BL, et al. Mental health symptoms following war and repression in eastern Afghanistan. *JAMA.* 2004;292(5):585–93.
- Vinck P, Pham PN, Stover E, Weinstein HM. Exposure to war crimes and implications for peace building in northern Uganda. *JAMA.* 2007;298(5):543–54.
- Rasekh Z, Bauer HM, Manos MM, Iacopino V. Women's health and human rights in Afghanistan. *JAMA.* 1998;280(5):449–55.
- Cardozo BL, Bilukha OO, Gotway CA, Wolfe MI, Gerber ML, Anderson M. Report from the CDC: mental health of women in postwar Afghanistan. *J Women's Health.* 2005;14(4):285–93.
- Roberts B, Ocaka KF, Browne J, Oyok T, Sondorp E. Factors associated with post-traumatic stress disorder and depression amongst internally displaced persons in northern Uganda. *BMC Psychiatry.* 2008;8:38.
- Bolton P, Neugebauer R, Ndogoni L. Prevalence of depression in rural Rwanda based on symptom and functional criteria. *J Nerv Ment Dis.* 2002;190(9):631–7.
- Wong EC, Marshall GN, Schell TL, Elliott MN, Babey SH, Hambarsoomians K. The unusually poor physical health status of Cambodian refugees two decades after resettlement. *J Immigr Minor Health.* 2011;13(5):876–82.
- Hovey JD, King CA. Acculturative stress, depression, and suicidal ideation among immigrant and second-generation Latino adolescents. *J Am Acad Child Adolesc Psychiatry.* 1996;35(9):1183–92.
- Ovuga E, Boardman J, Wasserman D. The prevalence of depression in two districts of Uganda. *Soc Psychiatry Psychiatr Epidemiol.* 2005;40(6):439–45.
- Kinyanda E, Woodburn P, Tugumisirize J, Kagugube J, Ndyabangi S, Patel V. Poverty, life events and the risk for depression in Uganda. *Soc Psychiatry Psychiatr Epidemiol.* 2011;46(1):35–44.
- Kizza D, Loa Knizek B, Kinyanda E, Hjelmeland H. An escape from agony: a qualitative psychological autopsy study of women's suicide in a post-conflict northern Uganda. *Int J Qual Stud Health Well-being.* 2012;7:1–13.
- Marshall C, Wolfe R, Kossman T, Wesselingh S, Harrington G, Spelman D. Risk factors for acquisition of methicillin-resistant *Staphylococcus aureus* (MRSA) by trauma patients in the intensive care unit. *J Hosp Infect.* 2004;57(3):245–52.
- Roberts B, Damundu EY, Lomoro O, Sondorp E. Post-conflict mental health needs: a cross-sectional survey of trauma, depression and associated factors in Juba Southern Sudan. *BMC Psychiatry.* 2009;9:7.
- Pfeiffer A, Elbert T. PTSD, depression and anxiety among former abductees in Northern Uganda. *Confl Heal.* 2011;5(1):14.
- Frontieres MS. Two decades of conflict take toll on northern Uganda. 2006. Two decades of conflict take toll on northern Uganda. <http://www.msf.org/article/two-decades-conflicttake-toll-northern-uganda>. Accessed, 17th June 2014. 1 www.isis-wicce.org/sites/default/.../Kitgum%20Medical%20Report.pdf.
- Kinyanda E, Musisi S, Biryabarema C, Ezati I, Oboke H, Ojiambo-Ochieng R, et al. War related sexual violence and its medical and psychological consequences as seen in Kitgum, Northern Uganda: A cross-sectional study. *BMC Int Health Hum Rights.* 2010;10:28.
- Liebling-Kalifani H, Ojiambo-Ochieng R, Marshall A, Were-Oguttu J, Musisi S, Kinyanda E. Violence against women in Northern Uganda: the neglected health consequences of war. *J Int Women's Stud.* 2008;9(3):174–92.
- Isis-WICCE. Medical Interventional Study of War Affected Kitgum District, Uganda: An Isis-WICCE Report. Kampala, Uganda: Isis-WICCE; 2006.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry.* 1998;22-33:34–57.
- Kinyanda E, Kizza R, Abbo C, Ndyabangi S, Levin J. Prevalence and risk factors of depression in childhood and adolescence as seen in four districts of North-Eastern Uganda. *BMC Int Health Hum Rights.* 2013;13:19.
- Muyinda Herbert. Utilizing the Wayo-Nero ("Aunt-Uncle") strategy as a vehicle for improving access to mental health services in post-conflict northern Uganda: Effectively bridging indigenous/traditional and modern/biomedical therapy systems, 2003. Grand Challenges Canada. <http://www.grandchallenges.ca/grantee-stars/0094-04/>.
- Kirkwood B, Sterne J. *Medical statistics.* Hoboken, NJ, USA: Blackwell Science; 2003. Edition Oxford edn.
- Kinyanda E, Hjelmeland H, Musisi S. Negative life events associated with deliberate self-harm in an African population in Uganda. *Crisis.* 2005;26(1):4–11.
- Kinyanda E, Waswa L, Baisley K, Maher D. Prevalence of severe mental distress and its correlates in a population-based study in rural south-west Uganda. *BMC Psychiatry.* 2011;11:97.
- Kerkhof A, Bernasco W, Bille-Brahe U, Platt S, Schmidtke A. WHO/EURO multicentre study on parasuicide: European parasuicide study interview schedule (EPSIS). Leiden: Department of Clinical Health and Personality Psychology, Leiden University; 1989.
- Musisi S, Kinyanda E, Leibling H, Mayengo K. Posttraumatic torture disorders in Uganda-A three year retrospective study of patient records as seen at a specialized torture treatment center in Kampala, Uganda. *Torture.* 2000;10(3):81–7.
- Dahlem NW, Zimet GD, Walker RR. The multidimensional scale of perceived social support: a confirmation study. *J Clin Psychol.* 1991;47(6):756–61.
- Bernstein DP, Fink L. *Childhood trauma questionnaire: a retrospective self-report: manual.* San Antonio, TX: Harcourt Brace & Company; 1998.
- Watson M, Greer S, Young J, Inayat Q, Burgess C, Robertson B. Development of a questionnaire measure of adjustment to cancer: the MAC scale. *Psychol Med.* 1988;18(1):203–9.
- Uganda AIDS Commission (UAC). *Rapid assessment of trends and drivers of the HIV epidemic and effectiveness of prevention interventions in Uganda.* Kampala, Uganda: UAC; 2005.
- Ministry of Health (MOH) [Uganda] and ORC Macro. *Uganda HIV/AIDS Sero-behavioural Survey 2004-2005.* Calverton, Maryland, USA: Ministry of Health and ORC Macro; 2006. <http://dhsprogram.com/pubs/pdf/AIS2/AIS2.pdf>.
- Uganda Bureau of Statistics (UBOS). *Uganda: demographic and health survey 2006: preliminary report.* Calverton, Maryland, USA: Ministry of Health (MOH) [Uganda] and ORC Macro International; 2006. Uganda Ministry of Finance, Planning and Economic Development, Kampala, Uganda.
- Sheikh L T, Mohammed S, Joseph Ike, Ebiti B, Oluwatosin A, et al. Correlates of depression among internally displaced persons after post-election violence in Kaduna, North Western Nigeria, *Affective Disorders.* 170, 46-55 DOI: <http://dx.doi.org/10.1016/j.jad.2014.08.050>
- Ministry of Health [Uganda]. *2011 Uganda AIDS Indicator Survey.* http://www.health.go.ug/docs/UAIS_2011_FACT_SHEET.pdf
- Okello ES. *Cultural explanatory models of depression in Uganda, Thesis, phd. School of Medicine (Sch. of Med.) Collections: Kalolinska University and Makerere University.* Stockholm, Sweden: Kalolinska University; 2006.
- Abbo C. *Profiles and outcome of traditional healing practices for severe mental illnesses in two districts of Eastern Uganda, PhD Thesis Karolinska Institute and Makerere University.* Stockholm, Sweden: Kalolinska University; 2009.
- Duggan MB. *Epilepsy and its effects on children and families in rural Uganda.* *Afr Health Sci.* 2013;13(3):613–23.

41. Kuwert P, Spitzer C, Tr"ader A, Freyberger H J, Ermann M: Six years later: Post-traumatic stress symptoms and current psychopathology in former German children of World War II. *Int Psychogeriatric Assoc.* 2006, p1-7. doi:10.1017/S104161020600442X.
42. Senarath U, Wickramage K, Peiris SL. Prevalence of depression and its associated factors among patients attending primary care settings in the post-conflict Northern Province in Sri Lanka: a cross-sectional study. *BMC Psychiatry.* 2014;14:85.
43. Roberts B, Felix Ocaka K, Browne J, Oyok T, Sondorp E. Factors associated with the health status of internally displaced persons in northern Uganda. *J Epidemiol Community Health.* 2009;63(3):227–32.
44. Kinyanda E, Hoskins S, Nakku J, Nawaz S, Patel V. Prevalence and risk factors of major depressive disorder in HIV/AIDS as seen in semi-urban Entebbe district Uganda. *BMC Psychiatry.* 2011;11:205.
45. Betancourt TS, Khan KT. The mental health of children affected by armed conflict: protective processes and pathways to resilience. *Int Rev Psychiatry.* 2008;20(3):317–28.
46. Murthy RS, Lakshminarayana R. Mental health consequences of war: a brief review of research findings. *World Psychiatry.* 2006;5(1):25–30.
47. Ciesla JA, Roberts JE. Meta-analysis of the relationship between HIV infection and risk for depressive disorders. *Am J Psychiatry.* 2001;158(5):725–30.
48. Hobfoll SE, Spielberger CD, Breznitz S, Figley C, Folkman S, Lepper-Green B, et al. War-related stress. Addressing the stress of war and other traumatic events. *Am Psychol.* 1991;46(8):848–55.
49. Ayazi T, Lien L, Eide AH, Ruom MM, Hauff E. What are the risk factors for the comorbidity of posttraumatic stress disorder and depression in a war-affected population? A cross-sectional community study in South Sudan. *BMC Psychiatry.* 2012;12:175.
50. Gourlay A, Birdthistle I, Mburu G, Iorpenda K, Wringe A. Barriers and facilitating factors to the uptake of antiretroviral drugs for prevention of mother-to-child transmission of HIV in sub-Saharan Africa: a systematic review. *J Int AIDS Soc.* 2013;16(1):18588.
51. Sheikh L T, Mohammed A, Agunbiade S, William NE, Oluwatosin A et al. Psycho-trauma, psychosocial adjustment, and symptomatic post- traumatic stress disorder among internally displaced persons I Kaduna, Northwestern Nigeria, *Frontiers in PSYCHIATRY* 5:127 doi: 10.3389/fpsy.2014.00127.
52. Olisah VO, Balyewu O, Sheikh TL. Adherence to highly active antiretroviral therapy in depressed patients with HIV/AIDS attending a Nigerian University teaching hospital clinic. *African J Psychiatry.* 2010;13:275–9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

