

Immune cells and preterm labour: do invariant NKT cells hold the key?

S.F. Rinaldi^{1,*}, A.G. Rossi², P.T.K. Saunders², and J.E. Norman¹

¹MRC Centre for Reproductive Health and Tommy's Centre for Maternal and Fetal Health, University of Edinburgh, Queen's Medical Research Institute, Edinburgh, UK ²MRC Centre for Inflammation Research, University of Edinburgh, Queen's Medical Research Institute, Edinburgh, UK

*Correspondence address. E-mail: S.Rinaldi@ed.ac.uk

Submitted on December 22, 2014; resubmitted on December 22, 2014; accepted on January 8, 2015

Whilst there is now considerable evidence that labour, both at term and preterm, is an inflammatory event associated with an immune cell influx into the utero-placental tissues (Thomson *et al.*, 1999; Young *et al.*, 2002; Osman *et al.*, 2003; Gomez-Lopez *et al.*, 2010; Rinaldi *et al.*, 2011; Hamilton *et al.*, 2012), the precise role that these immune cells play in the initiation of labour, either in the presence of an intrauterine infection preterm, or physiologically at term, is unclear. Previous work has shown that the infiltrating leukocytes are a major source of pro-inflammatory mediators, including interleukin (IL)-1 β , IL-8, IL-6, tumour necrosis factor (TNF)- α and MMP-9 (Roh *et al.*, 2000; Helmig *et al.*, 2002; Young *et al.*, 2002), suggesting that these immune cells are likely to contribute to the inflammatory response surrounding parturition. However, whether the infiltration of leukocytes is a key event in triggering the onset of parturition or merely a consequence of the heightened inflammatory signalling remains to be elucidated.

In vivo animal models have been an invaluable tool to study the role of immune cells in parturition onset, where in recent years several studies have utilized either antibody-mediated immune cell depletion or genetically modified mice to investigate the function of specific immune cell populations in normal and adverse pregnancy outcomes.

In this issue of *Molecular Human Reproduction*, Li *et al.* build upon their previous findings by identifying the molecular mechanisms underlying the role of activation of invariant natural killer T (iNKT) cells in inflammation-induced preterm birth in a mouse model (Li *et al.*, 2012, 2015).

iNKT cells are a subset of T cells which express both NK cell receptors and a rearranged T cell receptor (TCR) which has a semi-invariant TCR α chain (in mice this is a V α 14-J α 18 rearrangement, and in humans a V α 24-J α 18 rearrangement), that pairs with a restricted set of TCR β chains (Van Kaer *et al.*, 2013). This TCR rearrangement restricts iNKT cells to recognize self and foreign lipids presented by the MHC class I-related protein CD1d (Brennan *et al.*, 2013). Upon activation, iNKT cells rapidly secrete cytokines, such as interferon (IFN)- γ and IL-4, and these cytokines can activate other immune cells, such as NK cells, dendritic cells, macrophages, neutrophils, B cells and T cells, and can therefore participate in a wide range of immune responses (Brennan *et al.*, 2013). Due to their ability to produce a wide variety of cytokines and interact with other cells of the immune system, iNKT cells are often

considered as a 'bridge' between the innate and adaptive immune systems (Boyson *et al.*, 2008).

Several studies have now reported the presence of iNKT cells in the pregnant human and mouse decidua (Ito *et al.*, 2000; Tsuda *et al.*, 2001; Boyson *et al.*, 2002). iNKT cell stimulation with its specific ligand α -galactosylceramide (α -GalCer) results in both early pregnancy loss and preterm birth in mouse models (Ito *et al.*, 2000; Boyson *et al.*, 2006). Additionally, J α 18 KO mice, which have no iNKT cells, have a reduced rate of lipopolysaccharide (LPS)-induced early pregnancy loss and LPS-induced preterm birth (Li *et al.*, 2012, 2013). In this issue, Li *et al.* report that adoptive transfer of decidual iNKT cells to J α 18 KO mice promotes LPS-induced preterm birth and, by using various neutralizing antibodies, demonstrate that decidual iNKT cell activation is regulated by TLR-4-mediated signalling pathways, IL-12 and IL-18 secretion, and endogenous glycolipid antigens presented by CD1d.

In women, neutrophils are proposed to play a role in cervical ripening (Bokström *et al.*, 1997) and have been shown to infiltrate into the myometrium and cervix in association with term labour (Thomson *et al.*, 1999; Osman *et al.*, 2003). Furthermore, neutrophil infiltration has been specifically associated with infection-induced preterm labour. Hamilton *et al.* recently reported increased decidual neutrophil infiltration in women with infection-associated preterm labour, compared with women in either idiopathic or normal term labour (Hamilton *et al.*, 2012). A similar neutrophil influx has been reported in mouse models of inflammation-induced preterm labour (Shynlova *et al.*, 2013; Rinaldi *et al.*, 2014). However, in these models, neutrophil depletion in mice has been reported to have no effect on the timing of delivery either in normal term labour (Timmons and Mahendroo, 2006) or in models of CpG-oligodeoxynucleotide (CpG ODN; a TLR-9 agonist)-induced (Thaxton *et al.*, 2009; Sun *et al.*, 2013) and LPS-induced preterm birth (Rinaldi *et al.*, 2014), suggesting that neutrophil infiltration is not required for the onset of preterm labour.

The role of mast cells in the onset of parturition has also been investigated (Menzies *et al.*, 2011). Mast cells have been identified in the pregnant human and mouse uterus and cervix (Thomson *et al.*, 1999; Naik *et al.*, 2004; Garfield *et al.*, 2006; Menzies *et al.*, 2012); and there is evidence indicating that degranulation of mast cells can induce myometrial

contractility *in vitro* (Garfield *et al.*, 2000; Bytautiene *et al.*, 2004, 2008). However, mast-cell-deficient Kit^{W^{sh}} mice had no defects in the timing of normal term labour (Menzies *et al.*, 2012), suggesting that mast cells are not essential to the initiation of parturition in mice. Whether mast cells may be involved in the onset of inflammation-induced preterm birth has yet to be determined.

Conversely, macrophages appear to play a more crucial role in the onset of parturition. Macrophages account for around 20% of the decidual leukocyte population (Erlebacher, 2013) and there is an influx of macrophages into the myometrium, fetal membranes, decidua, placenta and cervix during spontaneous term labour (Thomson *et al.*, 1999; Osman *et al.*, 2003; Gomez-Lopez *et al.*, 2010) and in preterm labour (Hamilton *et al.*, 2012). In animal models of preterm birth, decidual macrophage numbers were elevated during labour (Hamilton *et al.*, 2012; Shynlova *et al.*, 2013). Importantly, macrophage depletion protected mice from CPG ODN-induced early pregnancy loss and preterm birth in IL-10 knockout (KO) (Thaxton *et al.*, 2009) and NK-cell deficient non-obese diabetic (NOD) mice (Sun *et al.*, 2013), and LPS-induced preterm delivery was prevented in wild-type mice (Gonzalez *et al.*, 2011).

Uterine natural killer (uNK) cells are another immune cell population that may be involved in the onset of parturition. During early pregnancy, uNK cells are the predominant immune cell population present in the human decidua, with their numbers gradually decreasing towards term (Bulmer *et al.*, 1991; Bartmann *et al.*, 2014). The uNK cells are proposed to play important roles in spiral arteriole remodelling and extravillous trophoblast invasion in early pregnancy (Erlebacher, 2013); however the role of uNKs in late pregnancy and parturition remains unclear. Depletion of uNK cells rescued LPS-induced fetal resorption and preterm birth in IL-10 KO mice (Murphy *et al.*, 2005, 2009), but not CpG ODN-induced fetal resorption or preterm birth in IL-10 KO mice (Thaxton *et al.*, 2009), suggesting that the role of uNK cells may depend on the nature of the inflammatory insult.

Several studies have also investigated the role of cells of the adaptive immune system, T and B cells, in pregnancy and parturition. Both T and B cells are found in the human myometrium, decidua and fetal membranes, with the number of T cells reported to increase in these tissues in association with the onset of labour at term (Thomson *et al.*, 1999; Gomez-Lopez *et al.*, 2013a, b). Regulatory T cells (Tregs), which are defined as CD4⁺CD25^{bright}FoxP3⁺ cells, appear to play an important, immunosuppressive role in the maintenance of pregnancy (Erlebacher, 2013), with animal studies demonstrating that depletion of FoxP3⁺ cells results in early pregnancy loss (Aluvihare *et al.*, 2004; Rowe *et al.*, 2012), and adoptive transfer of Tregs rescues CpG-ODN-induced pregnancy loss in NOD mice (Lin *et al.*, 2014). Other subpopulations of T cells, such as Th17 cells (CD3⁺CD4⁺IL-17A⁺), CD4⁺ cells and CD8⁺ cytotoxic T cells have also been identified at the maternal–fetal interface, but their precise role in the maintenance of pregnancy and/or onset of labour at term and preterm, remains unclear (Gomez-Lopez *et al.*, 2014). Interestingly, *Rag1* KO mice, which are deficient in both T and B cells, were not protected from LPS-induced preterm delivery, and were actually found to be more likely (than their wild-type counterparts) to deliver preterm following low-dose LPS (an effect which was partially reversed following adoptive transfer of CD4⁺ cells) (Bizargity *et al.*, 2009). Further work is needed to elucidate the precise functions of these adaptive immune cell populations during pregnancy and parturition.

Taken together, the above studies highlight the diverse immune cell populations, present at the maternal–fetal interface, which have the potential to be involved in the initiation of parturition. There will likely be differences in the function of each immune cell population in the presence of an intrauterine infection, compared with their role during spontaneous term labour, however the animal studies discussed here have provided promising evidence regarding the role of iNKT cells in the onset of inflammation-induced preterm birth, with some evidence suggesting macrophages and uNK cells may also play a role. For neutrophils, mast cells and T cells, although they have also been implicated in parturition, likely redundancy is demonstrated by studies where their depletion does not alter the timing of preterm parturition.

Whilst preterm birth remains the leading cause of neonatal mortality and morbidity worldwide, there is an urgent need for novel treatments (Blencowe *et al.*, 2013; Norman and Shennan, 2013). Therefore, studies identifying the underlying mechanisms leading to preterm birth, such as the work by Li *et al.*, published in this issue, are critical in highlighting new potential therapeutic targets. Although the precise role of iNKT cells in preterm birth in women remains to be elucidated, improved understanding of the role of individual immune cell populations in the onset of preterm birth may lead to the identification of novel therapeutic targets and ultimately the development of new treatments to delay preterm delivery and improve neonatal outcome.

Authors' roles

S.F.R., A.G.R., P.T.K.S. and J.E.N. all contributed to the preparation of the manuscript and gave final approval of the version to be published.

Funding

S.F.R. is supported by Medical Research Council (grant number MR/L002657/1) and Tommy's the baby charity. Funding to pay the Open Access publication charges for this article was provided by the Medical Research Council.

Conflict of interest

J.E.N., A.G.R. and P.T.K.S. hold a Medical Research Council grant investigating iNKT cells as drivers for preterm labour. S.F.R. is employed under this grant (Grant number MR/L002657/1).

References

- Aluvihare VR, Kallikourdis M, Betz AG. Regulatory T cells mediate maternal tolerance to the fetus. *Nat Immunol* 2004;**5**:266–271.
- Bartmann C, Segerer SE, Rieger L, Kapp M, Sutterlin M, Kammerer U. Quantification of the predominant immune cell populations in decidua throughout human pregnancy. *Am J Reprod Immunol* 2014;**71**:109–119.
- Bizargity P, Del Rio R, Phillippe M, Teuscher C, Bonney EA. Resistance to lipopolysaccharide-induced preterm delivery mediated by regulatory T cell function in mice. *Biol Reprod* 2009;**80**:874–881.
- Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller AB, Kinney M, Lawn J; Born Too Soon Preterm Birth Action Group. Born too soon: the global epidemiology of 15 million preterm births. *Reprod Health* 2013;**10**(Suppl 1):S2.

- Bokström H, Brännström M, Alexandersson M, Norström A. Leukocyte subpopulations in the human uterine cervical stroma at early and term pregnancy. *Hum Reprod* 1997;**12**:586–590.
- Boyson JE, Rybalov B, Koopman LA, Exley M, Balk SP, Racke FK, Schatz F, Masch R, Wilson SB, Strominger JL. CD1d and invariant NKT cells at the human maternal-fetal interface. *Proc Natl Acad Sci USA* 2002;**99**:13741–13746.
- Boyson JE, Nagarkatti N, Nizam L, Exley MA, Strominger JL. Gestation stage-dependent mechanisms of invariant natural killer T cell-mediated pregnancy loss. *Proc Natl Acad Sci USA* 2006;**103**:4580–4585.
- Boyson JE, Aktan I, Barkhoff DA, Chant A. NKT cells at the maternal-fetal interface. *Immunol Invest* 2008;**37**:565–582.
- Brennan PJ, Brigl M, Brenner MB. Invariant natural killer T cells: an innate activation scheme linked to diverse effector functions. *Nat Rev Immunol* 2013;**13**:101–117.
- Bulmer JN, Morrison L, Longfellow M, Ritson A, Pace D. Granulated lymphocytes in human endometrium: histochemical and immunohistochemical studies. *Hum Reprod* 1991;**6**:791–798.
- Bytautiene E, Vedernikov YP, Saade GR, Romero R, Garfield RE. Degranulation of uterine mast cell modifies contractility of isolated myometrium from pregnant women. *Am J Obstet Gynecol* 2004;**191**:1705–1710.
- Bytautiene E, Vedernikov YP, Saade GR, Romero R, Garfield RE. IgE-independent mast cell activation augments contractility of nonpregnant and pregnant guinea pig myometrium. *Int Arch Allergy Immunol* 2008;**147**:140–146.
- Erlbacher A. Immunology of the maternal-fetal interface. *Annu Rev Immunol* 2013;**31**:387–411.
- Garfield RE, Bytautiene E, Vedernikov YP, Marshall JS, Romero R. Modulation of rat uterine contractility by mast cells and their mediators. *Am J Obstet Gynecol* 2000;**183**:118–125.
- Garfield RE, Irani AM, Schwartz LB, Bytautiene E, Romero R. Structural and functional comparison of mast cells in the pregnant versus nonpregnant human uterus. *Am J Obstet Gynecol* 2006;**194**:261–267.
- Gomez-Lopez N, Guilbert LJ, Olson DM. Invasion of the leukocytes into the fetal-maternal interface during pregnancy. *J Leukoc Biol* 2010;**88**:625–633.
- Gomez-Lopez N, Hernandez-Santiago S, Lobb AP, Olson DM, Vadillo-Ortega F. Normal and premature rupture of fetal membranes at term delivery differ in regional chemotactic activity and related chemokine/cytokine production. *Reprod Sci* 2013a;**20**:276–284.
- Gomez-Lopez N, Vega-Sanchez R, Castillo-Castrejon M, Romero R, Cubeiro-Arreola K, Vadillo-Ortega F. Evidence for a role for the adaptive immune response in human term parturition. *Am J Reprod Immunol* 2013b;**69**:212–230.
- Gomez-Lopez N, StLouis D, Lehr MA, Sanchez-Rodriguez EN, Arenas-Hernandez M. Immune cells in term and preterm labor. *Cell Mol Immunol* 2014;**11**:571–581.
- Gonzalez JM, Franzke CW, Yang F, Romero R, Girardi G. Complement activation triggers metalloproteinases release inducing cervical remodeling and preterm birth in mice. *Am J Pathol* 2011;**179**:838–849.
- Hamilton S, Oomomian Y, Stephen G, Shynlova O, Tower CL, Garrod A, Lye SJ, Jones RL. Macrophages infiltrate the human and rat decidua during term and preterm labor: evidence that decidual inflammation precedes labor. *Biol Reprod* 2012;**86**:39.
- Helmig BR, Romero R, Espinoza J, Chaiworapongsa T, Bujold E, Gomez R, Ohlsson K, Uldbjerg N. Neutrophil elastase and secretory leukocyte protease inhibitor in prelabor rupture of membranes, parturition and intra-amniotic infection. *J Matern Fetal Neonatal Med* 2002;**12**:237–246.
- Ito K, Karasawa M, Kawano T, Akasaka T, Koseki H, Akutsu Y, Kondo E, Sekiya S, Sekikawa K, Harada M *et al*. Involvement of decidual Valpha14 NKT cells in abortion. *Proc Natl Acad Sci USA* 2000;**97**:740–744.
- Li LP, Fang YC, Dong GF, Lin Y, Saito S. Depletion of invariant NKT cells reduces inflammation-induced preterm delivery in mice. *J Immunol* 2012;**188**:4681–4689.
- Li L, Yang J, Ren L, Su N, Fang Y, Lin Y. Invariant NKT cells increase lipopolysaccharide-induced pregnancy loss by a mechanism involving Th1 and Th17 responses. *J Matern Fetal Neonatal Med* 2013;**26**:1212–1218.
- Li L, Yang J, Jiang Y, Tu J, Schust DJ. Activation of decidual invariant natural killer T cells promotes lipopolysaccharide-induced preterm birth. *Mol Hum Reprod* 2015;**21**:369–381.
- Lin Y, Liu X, Shan B, Wu J, Sharma S, Sun Y. Prevention of CpG-induced pregnancy disruption by adoptive transfer of *in vitro*-induced regulatory T cells. *PLoS One* 2014;**9**:e94702.
- Menzies FM, Shepherd MC, Nibbs RJ, Nelson SM. The role of mast cells and their mediators in reproduction, pregnancy and labour. *Hum Reprod Update* 2011;**17**:383–396.
- Menzies FM, Higgins CA, Shepherd MC, Nibbs RJ, Nelson SM. Mast cells reside in myometrium and cervix, but are dispensable in mice for successful pregnancy and labor. *Immunol Cell Biol* 2012;**90**:321–329.
- Murphy SP, Fast LD, Hanna NN, Sharma S. Uterine NK cells mediate inflammation-induced fetal demise in IL-10-null mice. *J Immunol* 2005;**175**:4084–4090.
- Murphy SP, Hanna NN, Fast LD, Shaw SK, Berg G, Padbury JF, Romero R, Sharma S. Evidence for participation of uterine natural killer cells in the mechanisms responsible for spontaneous preterm labor and delivery. *Am J Obstet Gynecol* 2009;**200**:e301–309.
- Naik R, Pai MR, Poornima Baliga B, Nayak KS, Shankararayanan ., Dighe P. Mast cell profile in uterine cervix. *Indian J Pathol Microbiol* 2004;**47**:178–180.
- Norman JE, Shennan AH. Prevention of preterm birth—why can't we do any better? *Lancet* 2013;**381**:184–185.
- Osman I, Young A, Ledingham MA, Thomson AJ, Jordan F, Greer IA, Norman JE. Leukocyte density and pro-inflammatory cytokine expression in human fetal membranes, decidua, cervix and myometrium before and during labour at term. *Mol Hum Reprod* 2003;**9**:41–45.
- Rinaldi SF, Hutchinson JL, Rossi AG, Norman JE. Anti-inflammatory mediators as physiological and pharmacological regulators of parturition. *Expert Rev Clin Immunol* 2011;**7**:675–696.
- Rinaldi SF, Catalano RD, Wade J, Rossi AG, Norman JE. Decidual neutrophil infiltration is not required for preterm birth in a mouse model of infection-induced preterm labor. *J Immunol* 2014;**192**:2315–2325.
- Roh CR, Oh WJ, Yoon BK, Lee JH. Up-regulation of matrix metalloproteinase-9 in human myometrium during labour: a cytokine-mediated process in uterine smooth muscle cells. *Mol Hum Reprod* 2000;**6**:96–102.
- Rowe JH, Ertelt JM, Xin L, Way SS. Pregnancy imprints regulatory memory that sustains energy to fetal antigen. *Nature* 2012;**490**:102–106.
- Shynlova O, Nedd-Roderique T, Li Y, Dorogin A, Nguyen T, Lye SJ. Infiltration of myeloid cells into decidua is a critical early event in the labour cascade and post-partum uterine remodelling. *J Cell Mol Med* 2013;**17**:311–324.
- Sun Y, Qin X, Shan B, Wang W, Zhu Q, Sharma S, Wu J, Lin Y. Differential effects of the CpG-TLR9 axis on pregnancy outcome in nonobese diabetic mice and wild-type controls. *Fertil Steril* 2013;**99**:1759–1767.
- Thaxton JE, Romero R, Sharma S. TLR9 activation coupled to IL-10 deficiency induces adverse pregnancy outcomes. *J Immunol* 2009;**183**:1144–1154.
- Thomson AJ, Telfer JF, Young A, Campbell S, Stewart CJ, Cameron IT, Greer IA, Norman JE. Leukocytes infiltrate the myometrium during human parturition: further evidence that labour is an inflammatory process. *Hum Reprod* 1999;**14**:229–236.

- Timmons BC, Mahendroo MS. Timing of neutrophil activation and expression of proinflammatory markers do not support a role for neutrophils in cervical ripening in the mouse. *Biol Reprod* 2006;**74**:236–245.
- Tsuda H, Sakai M, Michimata T, Tanebe K, Hayakawa S, Saito S. Characterization of NKT cells in human peripheral blood and decidual lymphocytes. *Am J Reprod Immunol* 2001;**45**:295–302.
- Van Kaer L, Parekh VV, Wu L. Invariant natural killer T cells as sensors and managers of inflammation. *Trends Immunol* 2013;**34**:50–58.
- Young A, Thomson AJ, Ledingham M, Jordan F, Greer IA, Norman JE. Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term. *Biol Reprod* 2002;**66**:445–449.