

# Role of Phosphatidylinositol Phosphate Signaling in the Regulation of the Filamentous-Growth Mitogen-Activated Protein Kinase Pathway

Hema Adhikari, Paul J. Cullen

Department of Biological Sciences at SUNY-Buffalo, Buffalo, New York, USA

**Reversible phosphorylation of the phospholipid phosphatidylinositol (PI) is a key event in the determination of organelle identity and an underlying regulatory feature in many biological processes. Here, we investigated the role of PI signaling in the regulation of the mitogen-activated protein kinase (MAPK) pathway that controls filamentous growth in yeast. Lipid kinases that generate phosphatidylinositol 4-phosphate [PI(4)P] at the Golgi (Pik1p) or PI(4,5)P<sub>2</sub> at the plasma membrane (PM) (Mss4p and Stt4p) were required for filamentous-growth MAPK pathway signaling. Introduction of a conditional allele of *PIK1* (*pik1-83*) into the filamentous ( $\Sigma$ 1278b) background reduced MAPK activity and caused defects in invasive growth and biofilm/mat formation. MAPK regulatory proteins that function at the PM, including Msb2p, Sho1p, and Cdc42p, were mislocalized in the *pik1-83* mutant, which may account for the signaling defects of the PI(4)P kinase mutants. Other PI kinases (Fab1p and Vps34p), and combinations of PIP (synaptojanin-type) phosphatases, also influenced the filamentous-growth MAPK pathway. Loss of these proteins caused defects in cell polarity, which may underlie the MAPK signaling defect seen in these mutants. In line with this possibility, disruption of the actin cytoskeleton by latrunculin A (LatA) dampened the filamentous-growth pathway. Various PIP signaling mutants were also defective for axial budding in haploid cells, cell wall construction, or proper regulation of the high-osmolarity glycerol response (HOG) pathway. Altogether, the study extends the roles of PI signaling to a differentiation MAPK pathway and other cellular processes.**

**M**APK (mitogen-activated protein kinase) pathways are evolutionarily conserved signal transduction modules (1, 2). MAPK cascades regulate the response to environmental challenges, such as changes in osmolarity, nutrient starvation, DNA damage, and damage to cell integrity. In the budding yeast *Saccharomyces cerevisiae*, MAPK pathways regulate cell wall integrity (3), pheromone response or mating (4), filamentous growth (5), and the response to high osmolarity (high-glycerol response [HOG] pathway [6]). Each MAPK pathway in yeast responds to a different stimulus. Under some circumstances, several MAPK pathways are required to mount an appropriate response (7–11).

The filamentous-growth MAPK pathway regulates differentiation to the filamentous cell type (12–14) and the development of biofilms or mats (15). During filamentous growth, the MAPK pathway, together with other pathways (16–18), induces a delay in the cell cycle (19), a reorganization of cell polarity, which leads to a distal-unipolar budding pattern (12, 13, 20, 21), and elevated expression of the cell adhesion molecule Flo1p (22). The developmental foraging responses that occur in *S. cerevisiae* are evolutionarily conserved across many fungal species. In pathogenic fungi, like *Candida albicans*, an orthologous differentiation MAPK pathway (called the Cek1p pathway) regulates filamentous/hyphal growth and biofilm formation (23–25). These behaviors are critical for virulence (24, 26–30). Studies of filamentous growth in a genetically tractable fungal system like *S. cerevisiae* provides information about the genetic basis of fungal behaviors that can be applied to other species, including pathogens.

In *S. cerevisiae*, the filamentous-growth MAPK pathway is regulated by Msb2p (31, 32), a member of the signaling mucin family of glycoprotein receptors (33). Msb2p, the tetraspan adaptor Sho1p (31, 34–36), and the cytosolic adaptor protein Bem4p (37) together regulate the Rho-type GTPase Cdc42p and effector p21-activated kinase Ste20p (38, 39) in the filamentous-growth MAPK pathway. Ste20p activates the MAPK cascade by phosphorylating

the MAPKKK Ste11p. Ste11p phosphorylates the MAPKK Ste7p, which in turn phosphorylates the MAPK Kss1p (13, 40). Opy2p is another transmembrane protein that recruits Ste11p to the PM through the adaptor protein Ste50p (35, 36, 41–43). The culmination of these events is the activation of transcription factors, Ste12p and Tec1p, which regulate target genes that control differentiation to the filamentous cell type (14, 44–47).

Phosphorylation of the key lipid phosphatidylinositol (PI) is a critical modification of membrane phospholipids in eukaryotes that is important for normal cellular function (48–50). PIPs are utilized as a mark for organelle identity and impact diverse cellular processes, including the reorganization of the actin cytoskeleton, protein trafficking through the endomembrane system, and protein secretion (51–56). A family of lipid kinases phosphorylates the inositol ring at different positions to designate organelles with specific PIP combinations (54). Two kinases generate PI(4)P in yeast: Pik1p regulates the level of PI(4)P at the Golgi (57), and Stt4p regulates PI(4)P at the PM (58). Mss4p regulates the distribution of PI(4,5)P<sub>2</sub> at the PM (59). In addition, Vps34p regulates the level of PI(3)P at the endosome (60–62), and Fab1p regulates PI(3,5)P<sub>2</sub> at the vacuole/lysosome (63). PIPs are recognized by

Received 29 January 2015 Accepted 23 February 2015

Accepted manuscript posted online 27 February 2015

Citation Adhikari H, Cullen PJ. 2015. Role of phosphatidylinositol phosphate signaling in the regulation of the filamentous-growth mitogen-activated protein kinase pathway. *Eukaryot Cell* 14:427–440. doi:10.1128/EC.00013-15.

Address correspondence to Paul J. Cullen, pjcullen@buffalo.edu.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/EC.00013-15>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved. doi:10.1128/EC.00013-15

TABLE 1 Yeast strains

Strain	Genotype <sup>a</sup>	Reference
PC313	<i>MATa ura3-52</i>	14
PC538	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52</i>	31
PC948	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 msb2::KanMX6</i>	31
PC1029	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 flo11::KanMX6</i>	156
PC1531	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sho1::HYG</i>	78
PC2053	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 pbs2::KanMX6</i>	106
PC2382	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 ste12::KanMX6</i>	78
PC776	<i>MATa his3 ura3-52 rsr1::HIS3</i>	20
PC677	<i>MATa his3 ura3-52 bud3::KIURA3</i>	20
PC2613	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 trp1::NAT</i>	This study
PC4305	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl1::KIURA3</i>	This study
PC4306	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl2::KIURA3</i>	This study
PC4307	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl3::KIURA3</i>	This study
PC4308	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl1::KIURA3 sjl2::NAT</i>	This study
PC4309	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl3::KIURA3 sjl2::NAT</i>	This study
PC4312	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl1::KIURA3 sjl3::NAT</i>	This study
PC4990	<i>SEY6210 MATα leu2-3,112 ura3-52 his3-Δ200 trp1-Δ901 lys2-801 suc2-Δ9<sup>b</sup></i>	94
PC4991	<i>SEY6210 MATα leu2-3,112 ura3-52 his3-Δ200 trp1-Δ901 lys2-801 suc2-Δ9 sjl3::TRP1 sjl2::HIS3 sjl1::HIS3 pRS314sjl2<sup>ts</sup>-8 (LEU2 CEN6 sjl2<sup>ts</sup>-8)<sup>b</sup></i>	68
PC4992	<i>SEY6210; stt4::HIS3 pRS415stt4-4 (LEU2 CEN6 stt4-4)<sup>b</sup></i>	57
PC4993	<i>SEY6210; mss4::HIS3MX6 Ycplacmss4-102 (LEU2 CEN6 mss4-102)<sup>b</sup></i>	68
PC4994	<i>SEY6210; pik1::HIS3 pRS314pik1-83 (TRP1 CEN6 pik1-83)<sup>b</sup></i>	57
PC5260	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl2::NAT sjl3::HYG</i>	This study
PC5276	<i>MATa ura3-52 sjl2::NAT sjl3::HYG</i>	This study
PC5294	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sjl2::NAT sjl3::HYG rga1::KIURA3</i>	This study
PC5303	<i>MATa ura3-52 sjl2::NAT sjl3::HYG ssk1::KIURA3</i>	This study
PC5319	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 trp1::NAT pik1::KLURA3 pRS314pik1-83 (TRP1 CEN6 pik1-83)</i>	This study
PC5433	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 fab1::HYG</i>	This study
PC5436	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 sac1::KIURA3</i>	This study
PC5438	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 trp1::NAT pik1::KLURA3 pRS314pik1-83 (TRP1 CEN6 pik1-83) URA3</i>	This study
PC5473	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 vps34::NAT</i>	This study
PC5659	<i>RSY255 MATa leu2-3,112 ura3-52</i>	157
PC5661	<i>MATα leu2-3,112 his4-519 ura3-52 suc2-Δ9 sec7-1</i>	116
PC5662	<i>RSY263 MATa leu2-3,112 ura3-52 sec12-4</i>	157
PC5663	<i>RSY271 MATa his4-619 ura3-52 sec18-1</i>	157
PC5664	<i>RSY529 MATa sec62-1 ura3-52, leu2-3,112, his4-619</i>	157
PC5712	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 msb2::KanMX6 sho1::NAT</i>	This study
PC6324	<i>MATa ste4 FUS1-lacZ FUS1-HIS3 ura3-52 inp54::KIURA3</i>	This study

<sup>a</sup> All strains were made in the  $\Sigma$ 1278b background unless otherwise indicated.

<sup>b</sup> Strain was made in the SEY6210 background.

specific domains on effector proteins (64–67). PIP modification is reversible by the action of phosphatases that dephosphorylate PIPs (54). Synaptojanin-type proteins Sjl1p, Sjl2p, and Sjl3p are the major PIP phosphatases in yeast. These proteins exhibit a high degree of functional redundancy (68).


Lipid signaling has established connections to the regulation of morphogenetic pathways (69). For example, activation of the yeast mating pathway requires recruitment of the mating-pathway specific scaffold Ste5p to the PM by recognition of PI(4,5)P<sub>2</sub> (70). The pheromone response and HOG pathways are regulated by Pik1p at the level of Ste11p (71). The guanine nucleotide exchange factor for the cell wall integrity pathway, Rom2p, interacts with PI(4,5)P<sub>2</sub> at the PM by its PH domain to regulate the PKC pathway activity (58). A clear link between PIP signaling and the regulation of the filamentous-growth MAPK pathway has yet to be defined. Intriguingly, in *C. albicans*, steep PI(4,5)P<sub>2</sub> gradients occur at hyphal tips and promote filamentous growth and invasion (72–74).

Here, we examined the impact of PI signaling on the regulation

of the filamentous-growth MAPK pathway in *S. cerevisiae*. We show that conditional PI kinase mutants exhibit defects in the filamentous-growth pathway. Membrane-associated regulators of the filamentous-growth MAPK pathway (including Msb2p, Sho1p, and Cdc42p) were mislocalized in PI(4)P kinase mutants, which may account for the reduction in MAPK activity in this context. Perhaps unexpectedly, other PI kinases, including Vps34p and to some degree Fab1p, were also involved. Polarity defects in these mutants might explain the MAPK signaling defects, as disruption of the actin cytoskeleton led to ablation of MAPK activity. Roles for PI kinases and PIP phosphatases in axial bud site selection in haploid cells, the maintenance of the yeast cell wall, and the HOG pathway were also uncovered. Therefore, PIP signaling plays a critical role in the regulation of a differentiation MAPK pathway and other aspects of cellular biogenesis and decision-making.

## MATERIALS AND METHODS

**Strains, media, and growth conditions.** Yeast strains are listed in Table 1.



**FIG 1** Role of PI(4)P kinases in regulating the filamentous-growth MAPK pathway. (A) Phosphorylation of Kss1p in wild-type cells and the *pik1-83*, *stt4-4*, and *mss4-102* mutants in the SEY6210 background. Cells were incubated at 37°C for 4 h and harvested for immunoblot analysis. Antibodies

Strains were grown under standard laboratory conditions (75). Strains were maintained at 30°C unless otherwise indicated. The medium used was YEPD (yeast extract-peptone-dextrose, 2% Glu) or YEP-Gal (2% Gal) for most experiments. For temperature shift experiments, cells were incubated at 37°C in prewarmed medium for 4 to 5 h. Plasmids were maintained on synthetic medium (generally on SD-Ura). Bacterial cultures of *Escherichia coli* were propagated in LB+CARB (carbenicillin) by standard methods (76).

Plasmid pAxl1p-HA was provided by the Boone lab (77). pMsb2p-HA (78), pMsb2p-GFP (78), pSho1p-GFP (79), *ppik1-83* (80), pGFP-Cdc42p (81), and Sec7p-DsRed (82) have been described previously.


Standard gene disruption techniques were used (83). Antibiotic resistance (84) and heterologous auxotrophic markers were used for gene disruption (85) and to create integrated fusion proteins (86). The *pik1-83* allele was introduced into the  $\Sigma$ 1278b strain background by allele replacement. Wild-type  $\Sigma$ 1278b cells (PC538) were transformed with a plasmid containing the *pik1-83* allele. The *PIK1* gene was subsequently disrupted in cells harboring the *pik1-83* plasmid. Gene disruptions were confirmed by PCR analysis.

**Biological assays for filamentous growth and biofilm/mat formation.** The plate-washing assay was performed as described previously (13). Biofilm/mat assays were performed as described previously (15). Cells were spotted onto low-agar medium (0.3% YEPD) for 3 days and photographed. Calcofluor white (CFW) staining was performed as described in reference 87. Cells were grown to saturation in YEPD medium at 30°C. Cells were fixed in 3.9% formaldehyde for 30 min at 30°C. Cells were harvested by centrifugation, washed with 1× PBS (phosphate-buffered saline, 137 mM NaCl, 2.7 mM KCl, 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.8 mM KH<sub>2</sub>PO<sub>4</sub> [pH 7.4]), and resuspended in 0.01% CFW for 5 min. Cells were washed 3 times in 1× PBS and observed by fluorescence microscopy using the DAPI (4',6-diamidino-2-phenylindole) channel (350 to 400 nm). Cells stained with CFW were also analyzed for budding pattern. Budding pattern determination was based on previous methods (88), except that cell position was also taken into consideration among adherent cells (20, 89).

**Protein immunoblot analysis.** Cells were harvested, and pellets were frozen at -80°C. Proteins were extracted from cell lysates using the trichloroacetic acid (TCA) precipitation method as described previously (90). Protein extracts were separated by 10% SDS-PAGE analysis and transferred onto nitrocellulose membrane. Phosphorylated Kss1p was detected using p42/p44 antibodies (1:4,000 dilution; no. 4370; Cell Signaling Technology). Antibodies to total Kss1p (1:5,000; no. sc6775; Santa Cruz), Hog1p (1:5,000; yc-20; no. sc6815; Santa Cruz), and Pgk1p (1:5,000 dilution; catalog no. 459250; Life Technologies) were purchased and used according to the manufacturers' specifications. Antibodies to green fluorescent protein (GFP; clones 7.1 and 13.1; no. 11814460001; Roche Diagnostics) and the hemagglutinin (HA) epitope (no. 11583816001; Roche) were used to detect epitope-tagged proteins. Secondary antibodies were used, including goat anti-mouse IgG conjugated to horseradish peroxidase (HRP; 170-6516; Bio-Rad) and goat anti-rabbit IgG-HRP (111-035-144; Jackson ImmunoResearch Laboratories, Inc.). Msb2p-HA secretion was evaluated as described previously (78). Protein bands were quantified by ImageJ by densitometry scanning analysis. The numbers indicate the ratio of phosphorylated protein to total protein, or the ratio of total protein to loading control where appropriate. To determine the change in phosphorylated proteins, total protein levels were normalized to the loading control (Pgk1p).

against Pgk1p were used to compare total protein levels between samples. (B) Evaluation of Msb2p-HA secretion in the *pik1-83* mutant. The wild type and the *pik1-83* mutant harboring pMsb2p-HA were grown to saturation at 37°C for 4 h and separated into supernatant (SUP) and pellet fractions by centrifugation. Cell extracts were prepared and examined by immunoblot analysis for HA-Msb2p levels and Pgk1p levels as a control for protein loading. (C) Phosphorylation of Kss1p in wild-type cells and the *pik1-83* mutant in the  $\Sigma$ 1278b background. Cells were incubated in YEPD and YEP-Gal medium at 30°C for 4 h. (D) Cells used for panel C were incubated at 37°C for 4 h.





**FIG 2** The role of Pik1p in regulating haploid invasive growth and biofilm/mat formation. (A) Plate-washing assay. Wild-type cells and the *pik1-83* and *ste12Δ* mutants were spotted onto YEPD medium and incubated at 30°C for 2 days. Plates were photographed (left), washed in a stream of water, and photographed again (right). (B) Cells scraped from the invasive scars on the washed YEPD plate were examined by differential interference contrast (DIC) microscopy at a magnification of  $\times 40$ . Bar, 20  $\mu\text{m}$ . (C) Biofilm/mat formation by the indicated strains on YEPD+0.3% agar medium. Wild-type cells and the *pik1-83*, *ste12Δ*, and *flo11Δ* mutants were spotted onto plates and incubated for 5 days at 30°C. Representative colonies are shown. Bar, 1 cm.

**Microscopy and protein localization experiments.** For protein localization experiments, cells were grown in SD-Ura for 16 h at 30°C, shifted to 37°C for 4 h, and examined at 37°C. Differential interference contrast (DIC) and fluorescence microscopy using fluorescein isothiocyanate (FITC) filter sets were performed using an Axioplan 2 fluorescence microscope (Zeiss) with a Plan-Apochromat 100 $\times$ /1.4 (oil) objective (numerical aperture [NA], 0.17). Digital images were obtained with the Axiocam MRm camera (Zeiss). Axiovision 4.4 software (Zeiss) was used for image acquisition and analysis. Cells were examined by oil immersion on glass slides (no. 2947-75; Corning, Inc., Corning, NY) with a glass coverslip (VWR 48366-227) using the 100 $\times$  objective. Cells were photographed at 37°C using a slide warmer (no. 0115.000; PeCon GmbH, Germany).

## RESULTS

**PI(4)P kinases regulate the filamentous-growth MAPK pathway.** The role of PI kinases in regulating the filamentous-growth pathway was examined. The filamentous-growth pathway was evaluated by measuring phosphorylation of the filamentous-growth MAPK Kss1p (91–93). Conditional (temperature-sensitive) alleles of PI kinases *pik1-83*, *stt4-4*, and *mss4-102* were tested, which showed reduced phosphorylation of Kss1p at 37°C (Fig. 1A). Thus, the generation of PI(4)P is necessary for activation of the filamentous-growth MAPK pathway. The above-described test was performed in a laboratory strain (SEY6210 background [94]). Many lab strains have lost the ability to undergo filamentous growth (95). Indeed, growth in the nonpreferred carbon source galactose (Gal) induces the filamentous-growth MAPK pathway (36, 78, 96) but led to a decrease in levels of phosphorylated Kss1p (P~Kss1p) in the SEY6210 background (Fig. 1A, compare GLU lanes to GAL lanes). To better evaluate the role of PI(4)P signaling in regulating the filamentous-growth pathway, the *pik1-83* allele was introduced into the filamentous ( $\Sigma 1287b$ ) background (12, 95). The *pik1-83* strain behaved as expected, based on temperature sensitivity at 37°C (97), aberrant Golgi morphology (see Fig. S1 in the supplemental material) (57), and a defect in protein secretion (98).


The extracellular domain of Msb2p is highly glycosylated and migrates as a smear at  $\sim 250$  kDa (78). The extracellular inhibitory domain is proteolytically processed by the aspartyl protease Yps1p

and is shed from cells (78). Consistent with the secretion defect of the *pik1-83* mutant, the large extracellular domain of HA-Msb2p was not shed (Fig. 1B, SUP) and accumulated in cell pellets (Fig. 1B, PELLET). The *pik1-83* mutant is also partly compromised for function at semipermissive temperatures (30°C) (70). The *pik1-83* mutant showed reduced levels of P~Kss1p at 30°C (Fig. 1C, 30°C) and complete loss of P~Kss1p at 37°C (Fig. 1D). At 37°C, P~Kss1p levels were equivalent in glucose and galactose in wild-type cells. These results establish a requirement for Pik1p and other PI(4)P kinases in regulating the filamentous-growth MAPK pathway.


**Generation of PI(4)P is required for filamentous growth and biofilm/mat formation.** The fact that Pik1p regulates the filamentous-growth MAPK pathway suggested a role for PI(4)P in the regulation of filamentous growth (12) and biofilm/mat formation (15). At 30°C, the *pik1-83* mutant showed a defect in invasive growth by the plate-washing assay (Fig. 2A). The *pik1-83* mutant was more defective for invasive growth than a mutant that completely lacks pathway activity (Fig. 2A, *ste12Δ*). This may indicate that Pik1p has roles in regulating filamentous growth that extend beyond the regulation of the MAPK pathway. Microscopic analysis of the *pik1-83* mutant showed that cells were defective for cell elongation and filament formation compared to wild-type cells (Fig. 2B). These phenotypes are controlled by the filamentous-growth MAPK pathway (Fig. 2B, *ste12Δ*).

Like many microbial species, including fungal pathogens (99, 100), budding yeast forms biofilms or mats (15). On low-agar medium, the *pik1-83* mutant was defective for biofilm/mat formation (Fig. 2C). Specifically, *pik1-83* mats were small and smooth in appearance. In contrast, wild-type mats showed a wrinkled pattern, which was dependent on the filamentous-growth MAPK pathway (Fig. 2C, *ste12Δ*) and the cell adhesion molecule Flo11p (Fig. 2C, *flo11Δ*). Thus, generation of PI(4)P is required for filamentous growth and biofilm/mat formation in yeast.

**PI(4)P is required for localization of membrane-associated regulators of the filamentous-growth MAPK pathway.** PI kinases are critical regulators of membrane trafficking (51). Modification of PI(4)P at the Golgi (98, 101) and modification of



**FIG 3** The localization and relative levels of PM regulators of the filamentous-growth MAPK pathway evaluated in the *pik1-83* mutant. (A) Localization of Msb2p-GFP, Sho1p-GFP, and GFP-Cdc42p in wild-type cells and the *pik1-83* mutant. Cells were incubated to mid-log phase at 30°C and shifted to 37°C for 4 h. Bar, 5  $\mu$ m. The localization patterns shown are representative of the patterns seen in most cells over multiple independent trials. More than 200 cells were examined. The panels on the right show higher magnifications of the areas marked by rectangles in the middle column. Arrows indicate differences between wild type and the *pik1-83* allele. (B) Relative levels of Msb2p-GFP in wild-type cells and the *pik1-83* mutant compared to a loading control, Pgk1p. Cells were grown in YEPD medium to mid-log phase at 30°C and shifted to 37°C for 4 h. (C and D) Relative levels of Sho1p-GFP (C) and GFP-Cdc42p (D). (E) Phosphorylation of Kss1p in wild-type cells and the indicated trafficking mutants at 37°C for 4 h.




**FIG 4** The role of PI kinases Vps34p and Fab1p in regulating the filamentous-growth MAPK pathway. (A) Kss1p phosphorylation in the *vps34Δ* mutant incubated at 30°C for 6 h in YEPD (GLU) and YEP-Gal (GAL) medium. (B) Biofilm/mat formation in the *vps34Δ* mutant and control strains incubated in YEPD+0.3% agar at 30°C for 3 days. (C) Plate-washing assay of the indicated strains incubated at 30°C after 3 days. The *pbs2Δ* mutant was used as a hyperinvasive growth mutant as a control. (D) Kss1p phosphorylation in the *fab1Δ* mutant incubated at 30°C for 6 h in YEPD (GLU) and YEP-Gal (GAL) medium.


PI(4,5)P<sub>2</sub> at the PM (102, 103) are required for delivery of vesicles and cargoes to the PM. The signaling defect of the *pik1-83* and other PI(4)P mutants might result from mislocalization of PM proteins that regulate the filamentous-growth MAPK pathway. A functional Msb2p-GFP fusion shows primarily vacuolar localization (78), which results from turnover of the protein from the PM (104). Msb2p-GFP failed to show precise vacuolar localization in the *pik1-83* mutant, which indicates that the protein is mislocalized (Fig. 3A). Msb2p may fail to be delivered to the vacuole in the *pik1-83* mutant because it does not reach the PM, although we cannot rule out the possibility that Msb2p is trafficked from the Golgi to the vacuole directly.

Sho1p is a transmembrane protein and adaptor for the filamentous-growth MAPK pathway (31, 32, 34) and the HOG MAPK pathway (105). Sho1p-GFP localizes to the PM and the mother-bud neck (Fig. 3A) (106–108). Sho1p-GFP was also mislocalized in the *pik1-83* mutant (Fig. 3A). Sho1p-GFP was specifically identified in internal sites in the *pik1-83* mutant. Sho1p-GFP was also mislocalized in the *stt4-4* and *mss4-102* mutants (see Fig. S2 in the supplemental material).

Cdc42p is a Rho-type GTPase that regulates (among other things) the filamentous-growth MAPK pathway (38, 39, 109). Cdc42p is localized to the PM membrane by a lipid modification, geranylgeranylation (110–114). GFP-Cdc42p was also mislocal-



**FIG 5** Regulation of the filamentous-growth MAPK pathway by synaptojanin-type PIP phosphatases. Phosphorylation of Kss1p in combinations of PI phosphatase mutants was assessed. Wild-type cells and the *sjl1Δ*, *sjl2Δ*, *sjl3Δ*, *sjl2Δ sjl3Δ*, *sjl1Δ sjl2Δ*, and *sjl1Δ sjl3Δ* mutants were incubated in YEPD (for wild-type [GLU]) or YEP-Gal (GAL) medium for 6 h at 30°C.



ized in the *pik1-83* mutant. Compared to wild-type cells, which show Cdc42p localization at the PM and vacuolar membrane (Fig. 3A), GFP-Cdc42p was seen primarily in internal sites in the *pik1-83* mutant (Fig. 3A). GFP-Cdc42p was also mislocalized in the *stt4-4* and *mss4-102* mutants (see Fig. S3 in the supplemental material).

It was previously reported that Cdc42p levels are reduced in the *mss4-102* mutant (103). The localization defects of Cdc42p and other filamentous-growth MAPK regulatory proteins might result from a decrease in protein stability. The processed form of Msb2p-GFP, Msb2<sup>p</sup>, migrates at 55 kDa, and a minor product migrates at 75 kDa (104). The level of Msb2<sup>p</sup>-GFP was not reduced in the *pik1-83*, *stt4-4*, or *mss4-102* mutants (Fig. 3B). The higher levels of Msb2p seen in these mutants might result from a delay in turnover of the protein. The level of Sho1p-GFP was reduced in the *pik1-83* mutant (Fig. 3C). The levels of GFP-Cdc42p were not reduced in the *pik1-83* mutant (Fig. 3D). Thus, defects in trafficking of pathway components to the PM in a PI(4)P-dependent manner, not loss of protein abundance, may account for the defect in activation of the filamentous-growth MAPK pathway in PI kinase mutants.


To further test this possibility, protein trafficking mutants that trap PM cargoes in the secretory pathway were tested for filamentous-growth MAPK pathway activity. Specifically, the *sec62-1*, *sec12-14*, *sec18-1*, and *sec7-1* mutants, which are defective for protein trafficking at various steps in the secretory pathway (115–120), were tested. Like PI kinase mutants, these mutants showed a decrease in MAPK activity (Fig. 3E) and mislocalization of PM regulators of the filamentation MAPK pathway (data not shown). These results are consistent with the idea that delivery of membrane-associated regulators of the filamentous-growth MAPK pathway to the PM is required for activation of the filamentous-growth MAPK pathway.

**Other PI kinases regulate the filamentous-growth MAPK pathway.** Other PI kinases may also regulate the filamentous-growth MAPK pathway. Vps34p regulates PI(3)P levels at the endosome/multivesicular body and is required for protein trafficking to the vacuole/lysosome (62, 121–123). The *vps34Δ* mutant showed reduced P~Kss1p levels (Fig. 4A). The levels of total Kss1p were also reduced, which may result from a positive-feedback loop, given that the *KSS1* gene is a transcriptional target of the filamentous-growth pathway (124).

The *vps34Δ* mutant was also defective for biofilm/mat formation (Fig. 4B). The *flo11Δ* mutant was used as a control in evaluating biofilm/mat formation. The *vps34Δ* mutant also had a growth defect; however, *vps34Δ* mats were smaller than wild-type mats and smooth in appearance. The unstructured appearance of these mats suggests that Vps34p plays a role in their development. Vps34p was also required for invasive growth (Fig. 4C), although its growth defect on YEPD (Glu) medium may contribute to its invasive growth defect. The PI(3,5)P<sub>2</sub> kinase Fab1p (63, 125) was also tested. The *fab1Δ* mutant showed a modest reduction in

**FIG 6** The role of LatA on the activity of the filamentous-growth MAPK pathway. (A) Wild-type cells were grown in YEPD (GLU) or YEP-Gal (GAL) for 6 h at 30°C and treated with LatA at the indicated concentrations (μM) for 3 h at 30°C. (B) Phosphorylation of Kss1p in wild-type cells and *msb2Δ*, *sho1Δ*, and *msb2Δ sho1Δ* mutants. Data are the ratios of the phosphorylated Kss1p levels to total Kss1p quantified by ImageJ. (C and D) Same experiment performed in the *msb2Δ* (C) and *sho1Δ* (D) mutants with 20 μM LatA for 3 h at 30°C.





**FIG 7** Role of the PI kinase Fab1p in regulating axial budding pattern and Axl1p protein levels. (A) CFW staining of the *fab1Δ* mutant and control strain grown to saturation in YEPD medium at 30°C. (B) Levels of Axl1p-HA in the wild-type cells and *fab1Δ* mutant at 30°C. Cells were incubated for 6 h in YEPD (GLU) and YEP-Gal (GAL) medium.

P~Kss1p levels in high-glucose (basal) conditions (Fig. 4D) and exhibited a defect in invasive growth (Fig. 4C). Thus, Fab1p may play a minor role in regulating the filamentous-growth MAPK pathway and may have several roles in regulating filamentous growth.

**Synaptojanin-type PIP phosphatases regulate the filamentous-growth pathway.** PIP phosphatases dephosphorylate PI(4,5)P<sub>2</sub> at different cellular locations to maintain PIP balance (54, 68, 126). The role of PIP phosphatases in regulating the filamentous-growth MAPK pathway was evaluated. The level of Kss1p~P was reduced in PIP phosphatase *sjl1Δ*, *sjl2Δ*, and *sjl3Δ* single mutants (Fig. 5). *sjl1Δ sjl2Δ*, *sjl2Δ sjl3Δ*, and *sjl1Δ sjl3Δ* double mutants also showed reduced MAPK activity (Fig. 5). The *sjl1Δ sjl2Δ* mutant showed higher P~Kss1p levels than the single mutants. The Sjl proteins are functionally redundant, and it is possible that Sjl3p alone carries out a new function in this genetic context. Therefore, PIP phosphatases are involved in regulating the filamentous-growth MAPK pathway. Two other phosphatases, Sac1p and Inp54p, were also tested but did not regulate the filamentous-growth MAPK pathway (data not shown). Thus, altering the balance of PIP signaling through perturbation of PI kinases or PIP phosphatases impacts the filamentous-growth MAPK pathway.

**An intact actin cytoskeleton is required for filamentous-growth pathway activity.** The trafficking defect of the PM regulators Msb2p, Sho1p, and Cdc42p in PI(4)P kinase mutants may explain the defect in filamentous-growth MAPK pathway activity. Other PI kinase mutants and PIP phosphatase mutants did not show dramatic localization defects of these proteins (data not shown). PI kinases and PIP phosphatases also regulate cell polarity and the actin cytoskeleton (58, 59, 68, 82, 127–129). Defects in the actin cytoskeleton may underlie the MAPK signaling defects of these mutants. To test this possibility, a pharmacological inhibitor of filamentous actin, latrunculin A (LatA) (130), was tested. Addition of a minimal concentration of LatA (10 μM) led to reduced P~Kss1p levels (Fig. 6A), despite the fact that total Kss1p levels are higher under this condition. Addition of higher concentration of LatA (30 μM) caused a more severe reduction in P~Kss1p levels (Fig. 6A). Thus, an intact cytoskeleton is required for filamentous-growth MAPK pathway signaling.

The actin cytoskeleton is required for many different cellu-

lar processes, such as delivery of vesicles and cargoes to the PM and turnover of proteins from the PM by endocytosis (131, 132). The individual contributions of Msb2p and Sho1p in regulating MAPK signaling were examined in response to treatment with LatA. Under nutrient-limiting conditions, Msb2p and Sho1p were partly redundant for activation of the filamentous-growth MAPK pathway (Fig. 6B), which is consistent with the invasive growth phenotypes of the *msb2Δ* and *sho1Δ* mutants (31). The *msb2Δ* mutant, which is somewhat defective for MAPK activity, showed a further reduction in P~Kss1p levels upon treatment with LatA (Fig. 6C). Thus, in the *msb2Δ* mutant, the Sho1p-dependent signal requires an intact actin cytoskeleton. The *sho1Δ* mutant showed a similar response (Fig. 6D), indicating that the Msb2p-dependent signal also requires an intact actin cytoskeleton. Therefore, the actin cytoskeleton is required to facilitate filamentous-growth MAPK pathway signaling by a mechanism that is dependent on Msb2p and Sho1p.

**PIP signaling contributes to axial budding in haploid yeast.** Cell polarity in yeast is also controlled by the actin cytoskeleton

**TABLE 2** Budding patterns of mutants defective for PI signaling and control strains in haploid yeast cells


Mutation(s) <sup>a</sup>	Percentage of cells with pattern		
	Distal unipolar	Random	Axial
None (wild type)	8	<1	92
<i>rsr1Δ</i> <sup>b</sup>	10	32	58
<i>bud3Δ</i> <sup>c</sup>	36	1	63
<i>fab1Δ</i>	20	<1	80
<i>sac1Δ</i>	19	1	80
<i>sjl1Δ</i>	11	1	88
<i>sjl2Δ</i>	7	3	90
<i>sjl3Δ</i>	6	<1	81
<i>sjl1Δ sjl2Δ</i>	21	<1	79
<i>sjl1Δ sjl3Δ</i>	18	1	81
<i>sjl2Δ sjl3Δ</i>	30	6	64

<sup>a</sup> Cells were grown to saturation in YEPD medium, fixed, and stained with CFW. More than 200 cells were counted for each mutant.

<sup>b</sup> The *rsr1Δ* mutant exhibits a random budding pattern (135) and was used as a control.

<sup>c</sup> The *bud3Δ* mutant exhibits a distal-pole budding pattern (135) and was used as a control.





**FIG 8** Role of the PI kinase and PIP phosphatase mutants in regulating the HOG pathway. (A) Hog1p phosphorylation in essential PI kinase mutants. Cells were grown for 37°C for 4 h. Cells were treated with 0.4 M KCl for the indicated times. (B) Hog1p phosphorylation in the indicated combinations of PIP phosphatase mutants. (C) CFW staining of control and PIP phosphatase mutants grown to saturation in YEPD medium at 30°C.

(133, 134). Depending on cell type and growth condition, yeast cells bud at the proximal or distal poles (88, 135–138). Many genes regulate bud site selection in haploid and diploid cells (135, 139, 140). PI kinase mutants showed a bud site selection defect. Specifically, at 30°C, haploid *pik1-83* and *fab1Δ* mutants failed to bud axially and showed an increase in distal-unipolar budding (Fig. 7A

shows data for *fab1Δ*; results for other mutants are shown in Fig. S4 in the supplemental material [arrows]). The increased distal-pole budding in the *pik1-83* mutant, which would be expected to promote filamentous growth, was not sufficient to restore agar invasion to that mutant (Fig. 2A). This phenotype was quantitated for the *fab1Δ* mutant (Table 2) but not for the essential kinase

mutants, because at 37°C the mutants fail to produce buds due to their growth defect. Combinations of PIP phosphatase mutants also showed axial budding defects, including the *sjl1Δ sjl2Δ*, *sjl2Δ sjl3Δ*, and *sjl1Δ sjl3Δ* double mutants (Table 2; also, see Fig. S4 [arrows]). The Golgi PI(4)P phosphatase mutant *sac1Δ* also showed an axial budding defect (Table 2; also, see Fig. S4). Thus, PIP signaling contributes to axial budding in haploid yeast.

The defect in axial budding might be related to the levels of Axl1p, an axial-specific protein expressed in haploid cells (141). Axl1p is not produced in diploid cells (77, 141–143) or haploid cells grown under nutrient-limiting conditions (20). One of the PI kinase mutants that showed an axial budding defect was tested for the levels of Axl1p. The *fab1Δ* mutant showed reduced Axl1p-HA levels by immunoblot analysis, as seen under glucose-rich and glucose-limited conditions (Fig. 7B). Therefore, changes in Axl1p levels may provide a connection between axial budding and PI signaling.

**Role of PI signaling in regulating the HOG pathway and yeast cell wall.** The filamentous-growth and HOG pathways share a subset of components (1, 144–147). PIP regulators may impact the activity of the HOG pathway, which is measured by phosphorylation of the MAPK Hog1p (148, 149). It was previously shown that Pik1p plays a positive role in regulating the HOG pathway (71). We found that the *stt4-4* and *mss4-102* mutants showed constitutive HOG pathway activity at the restrictive temperature (Fig. 8A), although not to the levels seen under activating conditions. Therefore, PI(4)P kinases that function at the PM play an inhibitory role in regulating the HOG pathway.

Several PI kinase and PIP phosphatase mutant combinations showed growth defects in high-osmolarity medium. The *vps34Δ* mutant had a growth defect on YEPD supplemented with 1 M KCl that was similar to the *pbs2Δ* mutant (data not shown). No significant reduction in P~Hog1p was observed in the *vps34Δ* mutant (data not shown). Some combinations of PIP phosphatase mutants also showed osmotic sensitivity, such as the *sjl2Δ sjl3Δ* mutant (128). The *sjl2Δ sjl3Δ* mutant showed normal HOG pathway activity compared to wild-type cells (Fig. 8B). This may indicate that PIP phosphatases play roles in regulating osmotic tolerance through a mechanism that is independent of the HOG pathway.

A connection between the cell wall integrity pathway and PIP signaling has been described (58). Calcofluor white (CFW), which stains chitin in the yeast cell wall (150, 151), showed an irregular pattern in the *sjl1Δ sjl2Δ* mutant (Fig. 8C). A triple PIP phosphatase mutant, the *sjl1Δ sjl2Δ<sup>ts</sup> sjl3Δ* mutant, also showed a defect in chitin deposition at 37°C (Fig. 8C). Single *sjl* mutants showed uniform distribution of CFW on the cell surface (see Fig. S4 in the supplemental material). Thus, combinations of Sjl-type phosphatases have a function in maintenance of the yeast cell wall.

## DISCUSSION

PIP signaling is an essential cellular process that is critical for the regulation of protein secretion, actin cytoskeleton reorganization, and organelle identity, biogenesis, transport, and inheritance. Here, we describe a role for PIP signaling in the regulation of an ERK-type MAPK pathway that controls filamentous growth in yeast. We specifically show that generation of PI(4)P is required for filamentous-growth MAPK pathway signaling. This may result from mislocalization of PM regulators of the MAPK pathway in PI(4)P kinase mutants. Failure of PM regulators to reach the PM may be expected to result in MAPK signaling defects. Previous

studies have implicated PI signaling in the regulation of filamentous growth, particularly in *C. albicans* (72–74). Here, we posit that this connection can be explained, at least in part, at the level of the MAPK pathway.

We also demonstrate that the filamentous-growth MAPK pathway requires other PI kinases that generate PI(3)P and PI(3,5)P<sub>2</sub> and Sjl-type PIP phosphatases. PM MAPK regulatory proteins are not mislocalized in these mutants (data not shown); thus, how the MAPK is functionally connected to these PIP species is not clear. We show that perturbation of PIP signaling influences cell polarity. Thus, the MAPK signaling defect in these mutants may result from problems in cell polarity. In support of this possibility, pharmacological disruption of filamentous actin results in a defect in filamentous-growth MAPK pathway activity. Previous reports have suggested a link between actin cytoskeleton and filamentous growth (152). Thus, the observations reported here extend this connection by linking the actin cytoskeleton to the activity of the filamentous-growth MAPK pathway.

We also show that PIP signaling is required for a specific aspect of cell polarity regulation, that of axial bud site selection in haploid cells. This may result from a general defect in the actin cytoskeleton. However, the levels of the axial-specific factor Axl1p are reduced in at least one PI kinase mutant and may reflect a specific connection between the two pathways. PIP signaling in yeast is required for other cellular processes, including proper regulation of the cell wall (58), and we identify cell wall defects in some *sjl* mutant combinations. Intriguingly, different PIP combinations influence each of these processes, which indicates a high degree of functional specialization of PIP regulators.

Different PIP species differentially regulate the HOG pathway. A Golgi PI(4)P kinase, Pik1p, positively regulates the HOG pathway (71), whereas PM PI(4)P kinases negatively regulate the HOG pathway. The mechanistic basis for the antagonistic roles of these PI(4)P kinases is not clear and underscores the importance of future studies of PI(4)P in regulation of the p38 MAPK pathway. The filamentous-growth (ERK-type) and HOG (p38-type) MAPK pathways have opposing functions in the cell (7, 106, 153–155). The fact that PI(4)P has different effects on ERK and p38 MAPK pathways could, in principle, influence the specificity of MAPK outputs. Future studies on how PIP signaling differentially activates MAPK pathways will shed light on the overall regulation of signaling pathways in this system.

## ACKNOWLEDGMENTS

We thank Scott Emr (Cornell University), Chris Stefan (University Medical College London), Alan Tartakoff (Case Western Reserve University), Doug Johnson (University of Vermont), and C. Boone (University of Toronto) for reagents. We thank Hay-Oak Park for discussions and assistance.

P.J.C. is supported by the U.S. Public Health Service (GM098629).

## REFERENCES

- Schwartz MA, Madhani HD. 2004. Principles of map kinase signaling specificity in *Saccharomyces cerevisiae*. *Annu Rev Genet* 38:725–748. <http://dx.doi.org/10.1146/annurev.genet.39.073003.112634>.
- Widmann C, Gibson S, Jarpe MB, Johnson GL. 1999. Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol Rev* 79:143–180.
- Levin DE. 2011. Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* 189:1145–1175. <http://dx.doi.org/10.1534/genetics.111.128264>.
- Dohlman HG, Thorner JW. 2001. Regulation of G protein-initiated signal transduction in yeast: paradigms and principles. *Annu Rev*

- Biochem 70:703–754. <http://dx.doi.org/10.1146/annurev.biochem.70.1.703>.
5. Cullen PJ, Sprague GF, Jr. 2012. The regulation of filamentous growth in yeast. *Genetics* 190:23–49. <http://dx.doi.org/10.1534/genetics.111.127456>.
  6. Saito H, Posas F. 2012. Response to hyperosmotic stress. *Genetics* 192:289–318. <http://dx.doi.org/10.1534/genetics.112.140863>.
  7. Adhikari H, Cullen PJ. 2014. Metabolic respiration induces AMPK- and Ire1p-dependent activation of the p38-type HOG MAPK pathway. *PLoS Genet* 10:e1004734. <http://dx.doi.org/10.1371/journal.pgen.1004734>.
  8. Baltanas R, Bush A, Couto A, Durrieu L, Hohmann S, Colman-Lerner A. 2013. Pheromone-induced morphogenesis improves osmoadaptation capacity by activating the HOG MAPK pathway. *Sci Signal* 6:ra26. <http://dx.doi.org/10.1126/scisignal.2003312>.
  9. Zarzov P, Mazzoni C, Mann C. 1996. The SLT2(MPK1) MAP kinase is activated during periods of polarized cell growth in yeast. *EMBO J* 15:83–91.
  10. Buehrer BM, Errede B. 1997. Coordination of the mating and cell integrity mitogen-activated protein kinase pathways in *Saccharomyces cerevisiae*. *Mol Cell Biol* 17:6517–6525.
  11. Errede B, Cade RM, Yashar BM, Kamada Y, Levin DE, Irie K, Matsumoto K. 1995. Dynamics and organization of MAP kinase signal pathways. *Mol Reprod Dev* 42:477–485. <http://dx.doi.org/10.1002/mrd.1080420416>.
  12. Gimeno CJ, Ljungdahl PO, Styles CA, Fink GR. 1992. Unipolar cell divisions in the yeast *S. cerevisiae* lead to filamentous growth: regulation by starvation and RAS. *Cell* 68:1077–1090. [http://dx.doi.org/10.1016/0092-8674\(92\)90079-R](http://dx.doi.org/10.1016/0092-8674(92)90079-R).
  13. Roberts RL, Fink GR. 1994. Elements of a single MAP kinase cascade in *Saccharomyces cerevisiae* mediate two developmental programs in the same cell type: mating and invasive growth. *Genes Dev* 8:2974–2985. <http://dx.doi.org/10.1101/gad.8.24.2974>.
  14. Liu H, Styles CA, Fink GR. 1993. Elements of the yeast pheromone response pathway required for filamentous growth of diploids. *Science* 262:1741–1744. <http://dx.doi.org/10.1126/science.8259520>.
  15. Reynolds TB, Fink GR. 2001. Bakers' yeast, a model for fungal biofilm formation. *Science* 291:878–881. <http://dx.doi.org/10.1126/science.291.5505.878>.
  16. Bharucha N, Ma J, Dobry CJ, Lawson SK, Yang Z, Kumar A. 2008. Analysis of the yeast kinome reveals a network of regulated protein localization during filamentous growth. *Mol Biol Cell* 19:2708–2717. <http://dx.doi.org/10.1091/mbc.E07-11-1199>.
  17. Rupp S, Summers E, Lo HJ, Madhani H, Fink G. 1999. MAP kinase and cAMP filamentation signaling pathways converge on the unusually large promoter of the yeast FLO11 gene. *EMBO J* 18:1257–1269. <http://dx.doi.org/10.1093/emboj/18.5.1257>.
  18. Ryan O, Shapiro RS, Kurat CF, Mayhew D, Baryshnikova A, Chin B, Lin ZY, Cox MJ, Vizeacoumar F, Cheung D, Bahr S, Tsui K, Tebbji F, Sellam A, Istel F, Schwarzmuller T, Reynolds TB, Kuchler K, Gifford DK, Whiteway M, Giaever G, Nislow C, Costanzo M, Gingras AC, Mitra RD, Andrews B, Fink GR, Cowen LE, Boone C. 2012. Global gene deletion analysis exploring yeast filamentous growth. *Science* 337:1353–1356. <http://dx.doi.org/10.1126/science.1224339>.
  19. Kron SJ, Styles CA, Fink GR. 1994. Symmetric cell division in pseudohyphae of the yeast *Saccharomyces cerevisiae*. *Mol Biol Cell* 5:1003–1022. <http://dx.doi.org/10.1091/mbc.5.9.1003>.
  20. Cullen PJ, Sprague GF, Jr. 2002. The roles of bud-site-selection proteins during haploid invasive growth in yeast. *Mol Biol Cell* 13:2990–3004. <http://dx.doi.org/10.1091/mbc.E02-03-0151>.
  21. Taheri N, Kohler T, Braus GH, Mosch HU. 2000. Asymmetrically localized Bud8p and Bud9p proteins control yeast cell polarity and development. *EMBO J* 19:6686–6696. <http://dx.doi.org/10.1093/emboj/19.24.6686>.
  22. Guo B, Styles CA, Feng Q, Fink GR. 2000. A *Saccharomyces* gene family involved in invasive growth, cell-cell adhesion, and mating. *Proc Natl Acad Sci U S A* 97:12158–12163. <http://dx.doi.org/10.1073/pnas.220420397>.
  23. Cote P, Sulea T, Dignard D, Wu C, Whiteway M. 2011. Evolutionary reshaping of fungal mating pathway scaffold proteins. *mBio* 2:e00230–10. <http://dx.doi.org/10.1128/mBio.00230-10>.
  24. Csank C, Schroppel K, Leberer E, Harcus D, Mohamed O, Meloche S, Thomas DY, Whiteway M. 1998. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infect Immun* 66:2713–2721.
  25. Liu H, Kohler J, Fink GR. 1994. Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. *Science* 266:1723–1726. <http://dx.doi.org/10.1126/science.7992058>.
  26. Ramage G, VanDeWalle K, Lopez-Ribot JL, Wickes BL. 2002. The filamentation pathway controlled by the Efg1 regulator protein is required for normal biofilm formation and development in *Candida albicans*. *FEMS Microbiol Lett* 214:95–100. <http://dx.doi.org/10.1111/j.1574-6968.2002.tb11330.x>.
  27. Paramonova E, Krom BP, van der Mei HC, Busscher HJ, Sharma PK. 2009. Hyphal content determines the compression strength of *Candida albicans* biofilms. *Microbiology* 155:1997–2003. <http://dx.doi.org/10.1099/mic.0.021568-0>.
  28. Lengeler KB, Davidson RC, D'Souza C, Harashima T, Shen WC, Wang P, Pan X, Vaughn M, Heitman J. 2000. Signal transduction cascades regulating fungal development and virulence. *Microbiol Mol Biol Rev* 64:746–785. <http://dx.doi.org/10.1128/MMBR.64.4.746-785.2000>.
  29. Lo HJ, Kohler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. 1997. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* 90:939–949. [http://dx.doi.org/10.1016/S0092-8674\(00\)80358-X](http://dx.doi.org/10.1016/S0092-8674(00)80358-X).
  30. Lorenz MC, Heitman J. 1998. Regulators of pseudohyphal differentiation in *Saccharomyces cerevisiae* identified through multicopy suppressor analysis in ammonium permease mutant strains. *Genetics* 150:1443–1457.
  31. Cullen PJ, Sabbagh W, Jr, Graham E, Erick MM, van Olden EK, Neal C, Delrow J, Bardwell L, Sprague GF, Jr. 2004. A signaling mucin at the head of the Cdc42- and MAPK-dependent filamentous growth pathway in yeast. *Genes Dev* 18:1695–1708. <http://dx.doi.org/10.1101/gad.1178604>.
  32. O'Rourke SM, Herskowitz I. 2002. A third osmosensing branch in *Saccharomyces cerevisiae* requires the Msb2 protein and functions in parallel with the Sho1 branch. *Mol Cell Biol* 22:4739–4749. <http://dx.doi.org/10.1128/MCB.22.13.4739-4749.2002>.
  33. Kufe DW. 2009. Mucins in cancer: function, prognosis and therapy. *Nat Rev Cancer* 9:874–885. <http://dx.doi.org/10.1038/nrc2761>.
  34. O'Rourke SM, Herskowitz I. 1998. The Hog1 MAPK prevents cross talk between the HOG and pheromone response MAPK pathways in *Saccharomyces cerevisiae*. *Genes Dev* 12:2874–2886. <http://dx.doi.org/10.1101/gad.12.18.2874>.
  35. Yamamoto K, Tatebayashi K, Tanaka K, Saito H. 2010. Dynamic control of yeast MAP kinase network by induced association and dissociation between the Ste50 scaffold and the Opy2 membrane anchor. *Mol Cell* 40:87–98. <http://dx.doi.org/10.1016/j.molcel.2010.09.011>.
  36. Karunanithi S, Cullen PJ. 2012. The filamentous growth MAPK pathway responds to glucose starvation through the Mig1/2 transcriptional repressors in *Saccharomyces cerevisiae*. *Genetics* 192:869–887. <http://dx.doi.org/10.1534/genetics.112.142661>.
  37. Pitoniak A, Chavel CA, Chow J, Smith J, Camara D, Karunanithi S, Li B, Wolfe KH, Cullen PJ. 2015. Cdc42p-interacting protein bem4p regulates the filamentous-growth mitogen-activated protein kinase pathway. *Mol Cell Biol* 35:417–436. <http://dx.doi.org/10.1128/MCB.00850-14>.
  38. Peter M, Neiman AM, Park HO, van Lohuizen M, Herskowitz I. 1996. Functional analysis of the interaction between the small GTP binding protein Cdc42 and the Ste20 protein kinase in yeast. *EMBO J* 15:7046–7059.
  39. Leberer E, Wu C, Leeuw T, Fourest-Lieuvin A, Segall JE, Thomas DY. 1997. Functional characterization of the Cdc42p binding domain of yeast Ste20p protein kinase. *EMBO J* 16:83–97. <http://dx.doi.org/10.1093/emboj/16.1.83>.
  40. Madhani HD, Styles CA, Fink GR. 1997. MAP kinases with distinct inhibitory functions impart signaling specificity during yeast differentiation. *Cell* 91:673–684. [http://dx.doi.org/10.1016/S0092-8674\(00\)80454-7](http://dx.doi.org/10.1016/S0092-8674(00)80454-7).
  41. Truckses DM, Bloomekatz JE, Thorner J. 2006. The RA domain of Ste50 adaptor protein is required for delivery of Ste11 to the plasma membrane in the filamentous growth signaling pathway of the yeast *Saccharomyces cerevisiae*. *Mol Cell Biol* 26:912–928. <http://dx.doi.org/10.1128/MCB.26.3.912-928.2006>.
  42. Kwan JJ, Warner N, Maini J, Chan Tung KW, Zakaria H, Pawson T, Donaldson LW. 2006. *Saccharomyces cerevisiae* Ste50 binds the MAPKKK Ste11 through a head-to-tail SAM domain interaction. *J Mol Biol* 356:142–154. <http://dx.doi.org/10.1016/j.jmb.2005.11.012>.



43. Ramezani-Rad M. 2003. The role of adaptor protein Ste50-dependent regulation of the MAPKKK Ste11 in multiple signalling pathways of yeast. *Curr Genet* 43:161–170. <http://dx.doi.org/10.1007/s00294-003-0383-6>.
44. Madhani HD, Fink GR. 1997. Combinatorial control required for the specificity of yeast MAPK signaling. *Science* 275:1314–1317. <http://dx.doi.org/10.1126/science.275.5304.1314>.
45. Bardwell L, Cook JG, Voora D, Baggott DM, Martinez AR, Thorner J. 1998. Repression of yeast Ste12 transcription factor by direct binding of unphosphorylated Kss1 MAPK and its regulation by the Ste7 MEK. *Genes Dev* 12:2887–2898. <http://dx.doi.org/10.1101/gad.12.18.2887>.
46. Bardwell L, Cook JG, Zhu-Shimoni JX, Voora D, Thorner J. 1998. Differential regulation of transcription: repression by unactivated mitogen-activated protein kinase Kss1 requires the Dig1 and Dig2 proteins. *Proc Natl Acad Sci U S A* 95:15400–15405. <http://dx.doi.org/10.1073/pnas.95.26.15400>.
47. Cook JG, Bardwell L, Kron SJ, Thorner J. 1996. Two novel targets of the MAP kinase Kss1 are negative regulators of invasive growth in the yeast *Saccharomyces cerevisiae*. *Genes Dev* 10:2831–2848. <http://dx.doi.org/10.1101/gad.10.22.2831>.
48. Majerus PW, York JD. 2009. Phosphoinositide phosphatases and disease. *J Lipid Res* 50(Suppl):S249–S254. <http://dx.doi.org/10.1194/jlr.R800072-JLR200>.
49. Dillon RL, White DE, Muller WJ. 2007. The phosphatidylinositol 3-kinase signaling network: implications for human breast cancer. *Oncogene* 26:1338–1345. <http://dx.doi.org/10.1038/sj.onc.1210202>.
50. Mellman I, Nelson WJ. 2008. Coordinated protein sorting, targeting and distribution in polarized cells. *Nat Rev Mol Cell Biol* 9:833–845. <http://dx.doi.org/10.1038/nrm2525>.
51. Fruman DA, Meyers RE, Cantley LC. 1998. Phosphoinositide kinases. *Annu Rev Biochem* 67:481–507. <http://dx.doi.org/10.1146/annurev.biochem.67.1.481>.
52. Di Paolo G, De Camilli P. 2006. Phosphoinositides in cell regulation and membrane dynamics. *Nature* 443:651–657. <http://dx.doi.org/10.1038/nature05185>.
53. Vicinanza M, D'Angelo G, Di Campli A, De Matteis MA. 2008. Function and dysfunction of the PI system in membrane trafficking. *EMBO J* 27:2457–2470. <http://dx.doi.org/10.1038/emboj.2008.169>.
54. Strahl T, Thorner J. 2007. Synthesis and function of membrane phosphoinositides in budding yeast, *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1771:353–404. <http://dx.doi.org/10.1016/j.bbali.2007.01.015>.
55. Saarikangas J, Zhao H, Lappalainen P. 2010. Regulation of the actin cytoskeleton-plasma membrane interplay by phosphoinositides. *Physiol Rev* 90:259–289. <http://dx.doi.org/10.1152/physrev.00036.2009>.
56. Shewan A, Eastburn DJ, Horvost K. 2011. Phosphoinositides in cell architecture. *Cold Spring Harbor Perspect Biol* 3:a004796. <http://dx.doi.org/10.1101/cshperspect.a004796>.
57. Audhya A, Foti M, Emr SD. 2000. Distinct roles for the yeast phosphatidylinositol 4-kinases, Stt4p and Pik1p, in secretion, cell growth, and organelle membrane dynamics. *Mol Biol Cell* 11:2673–2689. <http://dx.doi.org/10.1091/mbc.11.8.2673>.
58. Audhya A, Emr SD. 2002. Stt4 PI 4-kinase localizes to the plasma membrane and functions in the Pkc1-mediated MAP kinase cascade. *Dev Cell* 2:593–605. [http://dx.doi.org/10.1016/S1534-5807\(02\)00168-5](http://dx.doi.org/10.1016/S1534-5807(02)00168-5).
59. Desrivieres S, Cooke FT, Parker PJ, Hall MN. 1998. MSS4, a phosphatidylinositol-4-phosphate 5-kinase required for organization of the actin cytoskeleton in *Saccharomyces cerevisiae*. *J Biol Chem* 273:15787–15793. <http://dx.doi.org/10.1074/jbc.273.25.15787>.
60. Slessareva JE, Rott SM, Temple B, Bankaitis VA, Dohlman HG. 2006. Activation of the phosphatidylinositol 3-kinase Vps34 by a G protein alpha subunit at the endosome. *Cell* 126:191–203. <http://dx.doi.org/10.1016/j.cell.2006.04.045>.
61. Herman PK, Stack JH, DeModena JA, Emr SD. 1991. A novel protein kinase homolog essential for protein sorting to the yeast lysosome-like vacuole. *Cell* 64:425–437. [http://dx.doi.org/10.1016/0092-8674\(91\)90650-N](http://dx.doi.org/10.1016/0092-8674(91)90650-N).
62. Herman PK, Emr SD. 1990. Characterization of VPS34, a gene required for vacuolar protein sorting and vacuole segregation in *Saccharomyces cerevisiae*. *Mol Cell Biol* 10:6742–6754.
63. Yamamoto A, DeWald DB, Boronenkov IV, Anderson RA, Emr SD, Koshland D. 1995. Novel PI(4)P 5-kinase homologue, Fab1p, essential for normal vacuole function and morphology in yeast. *Mol Biol Cell* 6:525–539. <http://dx.doi.org/10.1091/mbc.6.5.525>.
64. Lemmon MA. 2003. Phosphoinositide recognition domains. *Traffic* 4:201–213. <http://dx.doi.org/10.1034/j.1600-0854.2004.00071.x>.
65. Bottomley MJ, Salim K, Panayotou G. 1998. Phospholipid-binding protein domains. *Biochim Biophys Acta* 1436:165–183. [http://dx.doi.org/10.1016/S0005-2760\(98\)00141-6](http://dx.doi.org/10.1016/S0005-2760(98)00141-6).
66. Itoh T, Takenawa T. 2002. Phosphoinositide-binding domains: Functional units for temporal and spatial regulation of intracellular signalling. *Cell Signal* 14:733–743. [http://dx.doi.org/10.1016/S0898-6568\(02\)00028-1](http://dx.doi.org/10.1016/S0898-6568(02)00028-1).
67. Hurley JH, Meyer T. 2001. Subcellular targeting by membrane lipids. *Curr Opin Cell Biol* 13:146–152. [http://dx.doi.org/10.1016/S0955-0674\(00\)00191-5](http://dx.doi.org/10.1016/S0955-0674(00)00191-5).
68. Stefan CJ, Audhya A, Emr SD. 2002. The yeast synaptojanin-like proteins control the cellular distribution of phosphatidylinositol (4,5)-bisphosphate. *Mol Biol Cell* 13:542–557. <http://dx.doi.org/10.1091/mbc.01-10-0476>.
69. Singh A, Del Poeta M. 2011. Lipid signalling in pathogenic fungi. *Cell Microbiol* 13:177–185. <http://dx.doi.org/10.1111/j.1462-5822.2010.01550.x>.
70. Garrenton LS, Stefan CJ, McMurray MA, Emr SD, Thorner J. 2010. Pheromone-induced anisotropy in yeast plasma membrane phosphatidylinositol-4,5-bisphosphate distribution is required for MAPK signaling. *Proc Natl Acad Sci U S A* 107:11805–11810. <http://dx.doi.org/10.1073/pnas.1005817107>.
71. Cappell SD, Dohlman HG. 2011. Selective regulation of MAP kinase signaling by an endomembrane phosphatidylinositol 4-kinase. *J Biol Chem* 286:14852–14860. <http://dx.doi.org/10.1074/jbc.M110.195073>.
72. Guillas I, Vernay A, Vitagliano JJ, Arkowitz RA. 2013. Phosphatidylinositol 4,5-bisphosphate is required for invasive growth in *Saccharomyces cerevisiae*. *J Cell Sci* 126:3602–3614. <http://dx.doi.org/10.1242/jcs.122606>.
73. Vernay A, Schaub S, Guillas I, Bassilana M, Arkowitz RA. 2012. A steep phosphoinositide bis-phosphate gradient forms during fungal filamentous growth. *J Cell Biol* 198:711–730. <http://dx.doi.org/10.1083/jcb.201203099>.
74. Hairfield ML, Westwater C, Dolan JW. 2002. Phosphatidylinositol-4-phosphate 5-kinase activity is stimulated during temperature-induced morphogenesis in *Candida albicans*. *Microbiology* 148:1737–1746.
75. Rose MD, Winston F, Hieter P. 1990. *Methods in yeast genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
76. Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
77. Adames N, Blundell K, Ashby MN, Boone C. 1995. Role of yeast insulin-degrading enzyme homologs in propheromone processing and bud site selection. *Science* 270:464–467. <http://dx.doi.org/10.1126/science.270.5235.464>.
78. Vadaie N, Dionne H, Akajagbor DS, Nickerson SR, Krysan DJ, Cullen PJ. 2008. Cleavage of the signaling mucin Msb2 by the aspartyl protease Yps1 is required for MAPK activation in yeast. *J Cell Biol* 181:1073–1081. <http://dx.doi.org/10.1083/jcb.200704079>.
79. Marles JA, Dahesh S, Haynes J, Andrews BJ, Davidson AR. 2004. Protein-protein interaction affinity plays a crucial role in controlling the Sho1p-mediated signal transduction pathway in yeast. *Mol Cell* 14:813–823. <http://dx.doi.org/10.1016/j.molcel.2004.05.024>.
80. Hendricks KB, Wang BQ, Schnieders EA, Thorner J. 1999. Yeast homologue of neuronal frequenin is a regulator of phosphatidylinositol-4-OH kinase. *Nature Cell Biol* 1:234–241. <http://dx.doi.org/10.1038/12058>.
81. Richman TJ, Johnson DI. 2000. *Saccharomyces cerevisiae* cdc42p GTPase is involved in preventing the recurrence of bud emergence during the cell cycle. *Mol Cell Biol* 20:8548–8559. <http://dx.doi.org/10.1128/MCB.20.22.8548-8559.2000>.
82. Foti M, Audhya A, Emr SD. 2001. SacI lipid phosphatase and Stt4 phosphatidylinositol 4-kinase regulate a pool of phosphatidylinositol 4-phosphate that functions in the control of the actin cytoskeleton and vacuole morphology. *Mol Biol Cell* 12:2396–2411. <http://dx.doi.org/10.1091/mbc.12.8.2396>.
83. Baudin A, Ozier-Kalogeropoulos O, Denouel A, Lacroute F, Cullin C. 1993. A simple and efficient method for direct gene deletion in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 21:3329–3330. <http://dx.doi.org/10.1093/nar/21.14.3329>.
84. Goldstein AL, McCusker JH. 1999. Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. *Yeast* 15:1541–



1553. [http://dx.doi.org/10.1002/\(SICI\)1097-0061\(199910\)15:14<1541::AID-YEA476>3.0.CO;2-K](http://dx.doi.org/10.1002/(SICI)1097-0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K).
85. Wach A, Brachat A, Alberti-Segui C, Rebischung C, Philippsen P. 1997. Heterologous HIS3 marker and GFP reporter modules for PCR-targeting in *Saccharomyces cerevisiae*. *Yeast* 13:1065–1075. [http://dx.doi.org/10.1002/\(SICI\)1097-0061\(19970915\)13:11<1065::AID-YEA159>3.0.CO;2-K](http://dx.doi.org/10.1002/(SICI)1097-0061(19970915)13:11<1065::AID-YEA159>3.0.CO;2-K).
  86. Longtine MS, McKenzie A, III, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR. 1998. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14:953–961. [http://dx.doi.org/10.1002/\(SICI\)1097-0061\(199807\)14:10<953::AID-YEA293>3.0.CO;2-U](http://dx.doi.org/10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U).
  87. Cabib E, Farkas V, Kosik O, Blanco N, Arroyo J, McPhie P. 2008. Assembly of the yeast cell wall. Crh1p and Crh2p act as transglycosylases in vivo and in vitro. *J Biol Chem* 283:29859–29872. <http://dx.doi.org/10.1074/jbc.M804274200>.
  88. Chant J, Pringle JR. 1995. Patterns of bud-site selection in the yeast *Saccharomyces cerevisiae*. *J Cell Biol* 129:751–765. <http://dx.doi.org/10.1083/jcb.129.3.751>.
  89. Chavel CA, Caccamise LM, Li B, Cullen PJ. 2014. Global regulation of a differentiation MAPK pathway in yeast. *Genetics* 198:1309–1328. <http://dx.doi.org/10.1534/genetics.114.168252>.
  90. Lee MJ, Dohlman HG. 2008. Coactivation of G protein signaling by cell-surface receptors and an intracellular exchange factor. *Curr Biol* 18:211–215. <http://dx.doi.org/10.1016/j.cub.2008.01.007>.
  91. Ma D, Cook JG, Thorner J. 1995. Phosphorylation and localization of Kss1, a MAP kinase of the *Saccharomyces cerevisiae* pheromone response pathway. *Mol Biol Cell* 6:889–909. <http://dx.doi.org/10.1091/mbc.6.7.889>.
  92. Sabbagh W, Jr, Flatauer LJ, Bardwell AJ, Bardwell L. 2001. Specificity of MAP kinase signaling in yeast differentiation involves transient versus sustained MAPK activation. *Mol Cell* 8:683–691. [http://dx.doi.org/10.1016/S1097-2765\(01\)00322-7](http://dx.doi.org/10.1016/S1097-2765(01)00322-7).
  93. Maleri S, Ge Q, Hackett EA, Wang Y, Dohlman HG, Errede B. 2004. Persistent activation by constitutive Ste7 promotes Kss1-mediated invasive growth but fails to support Fus3-dependent mating in yeast. *Mol Cell Biol* 24:9221–9238. <http://dx.doi.org/10.1128/MCB.24.20.9221-9238.2004>.
  94. Robinson JS, Klionsky DJ, Banta LM, Emr SD. 1988. Protein sorting in *Saccharomyces cerevisiae*: isolation of mutants defective in the delivery and processing of multiple vacuolar hydrolases. *Mol Cell Biol* 8:4936–4948.
  95. Liu H, Styles CA, Fink GR. 1996. *Saccharomyces cerevisiae* S288C has a mutation in FLO8, a gene required for filamentous growth. *Genetics* 144:967–978.
  96. Cullen PJ, Sprague GF, Jr. 2000. Glucose depletion causes haploid invasive growth in yeast. *Proc Natl Acad Sci U S A* 97:13619–13624. <http://dx.doi.org/10.1073/pnas.240345197>.
  97. Flanagan CA, Schnieders EA, Emerick AW, Kunisawa R, Admon A, Thorner J. 1993. Phosphatidylinositol 4-kinase: gene structure and requirement for yeast cell viability. *Science* 262:1444–1448. <http://dx.doi.org/10.1126/science.8248783>.
  98. Walch-Solimena C, Novick P. 1999. The yeast phosphatidylinositol-4-OH kinase pik1 regulates secretion at the Golgi. *Nat Cell Biol* 1:523–525. <http://dx.doi.org/10.1038/70319>.
  99. Finkel JS, Mitchell AP. 2011. Genetic control of *Candida albicans* biofilm development. *Nat Rev Microbiol* 9:109–118. <http://dx.doi.org/10.1038/nrmicro2475>.
  100. Ganguly S, Mitchell AP. 2011. Mucosal biofilms of *Candida albicans*. *Curr Opin Microbiol* 14:380–385. <http://dx.doi.org/10.1016/j.mib.2011.06.001>.
  101. Hama H, Schnieders EA, Thorner J, Takemoto JY, DeWald DB. 1999. Direct involvement of phosphatidylinositol 4-phosphate in secretion in the yeast *Saccharomyces cerevisiae*. *J Biol Chem* 274:34294–34300. <http://dx.doi.org/10.1074/jbc.274.48.34294>.
  102. Routt SM, Ryan MM, Tyeryar K, Rizzieri KE, Mousley C, Roumanie O, Brennhald PJ, Bankaitis VA. 2005. Nonclassical PITPs activate PLD via the Ste4p PtdIns-4-kinase and modulate function of late stages of exocytosis in vegetative yeast. *Traffic* 6:1157–1172. <http://dx.doi.org/10.1111/j.1600-0854.2005.00350.x>.
  103. Yakir-Tamang L, Gerst JE. 2009. A phosphatidylinositol-transfer protein and phosphatidylinositol-4-phosphate 5-kinase control Cdc42 to regulate the actin cytoskeleton and secretory pathway in yeast. *Mol Biol Cell* 20:3583–3597. <http://dx.doi.org/10.1091/mbc.E08-10-1073>.
  104. Adhikari H, Vadaie N, Chow J, Caccamise LM, Chavel CA, Li B, Bowitch A, Stefan CJ, Cullen PJ. 9 February 2015. Role of the unfolded protein response in regulating the mucin-dependent filamentous growth MAPK pathway. *Mol Cell Biol* <http://dx.doi.org/10.1128/MCB.01501-14>.
  105. Maeda T, Takekawa M, Saito H. 1995. Activation of yeast PBS2 MAPKK by MAPKKs or by binding of an SH3-containing osmosensor. *Science* 269:554–558. <http://dx.doi.org/10.1126/science.7624781>.
  106. Pitoniak A, Birkaya B, Dionne HM, Vadaie N, Cullen PJ. 2009. The signaling mucins Msb2 and Hkr1 differentially regulate the filamentation mitogen-activated protein kinase pathway and contribute to a multimodal response. *Mol Biol Cell* 20:3101–3114. <http://dx.doi.org/10.1091/mbc.E08-07-0760>.
  107. Raitt DC, Posas F, Saito H. 2000. Yeast Cdc42 GTPase and Ste20 PAK-like kinase regulate Sho1-dependent activation of the Hog1 MAPK pathway. *EMBO J* 19:4623–4631. <http://dx.doi.org/10.1093/emboj/19.17.4623>.
  108. Reiser V, Salah SM, Ammerer G. 2000. Polarized localization of yeast Pbs2 depends on osmotic stress, the membrane protein Sho1 and Cdc42. *Nat Cell Biol* 2:620–627. <http://dx.doi.org/10.1038/35023568>.
  109. Mosch HU, Roberts RL, Fink GR. 1996. Ras2 signals via the Cdc42/Ste20/mitogen-activated protein kinase module to induce filamentous growth in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* 93:5352–5356. <http://dx.doi.org/10.1073/pnas.93.11.5352>.
  110. Johnson JL, Erickson JW, Cerione RA. 2009. New insights into how the Rho guanine nucleotide dissociation inhibitor regulates the interaction of Cdc42 with membranes. *J Biol Chem* 284:23860–23871. <http://dx.doi.org/10.1074/jbc.M109.031815>.
  111. Ohya Y, Goebel M, Goodman LE, Petersen-Bjorn S, Friesen JD, Tamañoi F, Anraku Y. 1991. Yeast CAL1 is a structural and functional homologue to the DPR1 (RAM) gene involved in ras processing. *J Biol Chem* 266:12356–12360.
  112. Ohya Y, Qadota H, Anraku Y, Pringle JR, Botstein D. 1993. Suppression of yeast geranylgeranyl transferase I defect by alternative prenylation of two target GTPases, Rho1p and Cdc42p. *Mol Biol Cell* 4:1017–1025. <http://dx.doi.org/10.1091/mbc.4.10.1017>.
  113. Ziman M, Preuss D, Mulholland J, O'Brien JM, Botstein D, Johnson DI. 1993. Subcellular localization of Cdc42p, a *Saccharomyces cerevisiae* GTP-binding protein involved in the control of cell polarity. *Mol Biol Cell* 4:1307–1316. <http://dx.doi.org/10.1091/mbc.4.12.1307>.
  114. Ziman M, O'Brien JM, Ouellette LA, Church WR, Johnson DI. 1991. Mutational analysis of CDC42Sc, a *Saccharomyces cerevisiae* gene that encodes a putative GTP-binding protein involved in the control of cell polarity. *Mol Cell Biol* 11:3537–3544.
  115. Nanduri J, Tartakoff AM. 2001. The arrest of secretion response in yeast: signaling from the secretory path to the nucleus via Wsc proteins and Pkc1p. *Mol Cell* 8:281–289. [http://dx.doi.org/10.1016/S1097-2765\(01\)00312-4](http://dx.doi.org/10.1016/S1097-2765(01)00312-4).
  116. Achstetter T, Franzusoff A, Field C, Schekman R. 1988. SEC7 encodes an unusual, high molecular weight protein required for membrane traffic from the yeast Golgi apparatus. *J Biol Chem* 263:11711–11717.
  117. Barlowe C, Schekman R. 1993. SEC12 encodes a guanine-nucleotide-exchange factor essential for transport vesicle budding from the ER. *Nature* 365:347–349. <http://dx.doi.org/10.1038/365347a0>.
  118. Newman AP, Ferro-Novick S. 1990. Defining components required for transport from the ER to the Golgi complex in yeast. *Bioessays* 12:485–491. <http://dx.doi.org/10.1002/bies.950121006>.
  119. Novick P, Field C, Schekman R. 1980. Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. *Cell* 21:205–215. [http://dx.doi.org/10.1016/0092-8674\(80\)90128-2](http://dx.doi.org/10.1016/0092-8674(80)90128-2).
  120. Deshaies RJ, Schekman R. 1989. SEC62 encodes a putative membrane protein required for protein translocation into the yeast endoplasmic reticulum. *J Cell Biol* 109:2653–2664. <http://dx.doi.org/10.1083/jcb.109.6.2653>.
  121. Kihara A, Noda T, Ishihara N, Ohsumi Y. 2001. Two distinct Vps34 phosphatidylinositol 3-kinase complexes function in autophagy and carboxypeptidase Y sorting in *Saccharomyces cerevisiae*. *J Cell Biol* 152:519–530. <http://dx.doi.org/10.1083/jcb.152.3.519>.
  122. Burda P, Padilla SM, Sarkar S, Emr SD. 2002. Retromer function in endosome-to-Golgi retrograde transport is regulated by the yeast Vps34

- PtdIns 3-kinase. *J Cell Sci* 115:3889–3900. <http://dx.doi.org/10.1242/jcs.00090>.
123. Schu PV, Takegawa K, Fry MJ, Stack JH, Waterfield MD, Emr SD. 1993. Phosphatidylinositol 3-kinase encoded by yeast VPS34 gene essential for protein sorting. *Science* 260:88–91. <http://dx.doi.org/10.1126/science.8385367>.
  124. Roberts CJ, Nelson B, Marton MJ, Stoughton R, Meyer MR, Bennett HA, He YD, Dai H, Walker WL, Hughes TR, Tyers M, Boone C, Friend SH. 2000. Signaling and circuitry of multiple MAPK pathways revealed by a matrix of global gene expression profiles. *Science* 287:873–880. <http://dx.doi.org/10.1126/science.287.5454.873>.
  125. Odorizzi G, Babst M, Emr SD. 1998. Fab1p PtdIns(3)P 5-kinase function essential for protein sorting in the multivesicular body. *Cell* 95:847–858. [http://dx.doi.org/10.1016/S0092-8674\(00\)81707-9](http://dx.doi.org/10.1016/S0092-8674(00)81707-9).
  126. Stefan CJ, Padilla SM, Audhya A, Emr SD. 2005. The phosphoinositide phosphatase Sjl2 is recruited to cortical actin patches in the control of vesicle formation and fission during endocytosis. *Mol Cell Biol* 25:2910–2923. <http://dx.doi.org/10.1128/MCB.25.8.2910-2923.2005>.
  127. Garcia-Bustos JF, Marini F, Stevenson I, Frei C, Hall MN. 1994. PIK1, an essential phosphatidylinositol 4-kinase associated with the yeast nucleus. *EMBO J* 13:2352–2361.
  128. Ooms LM, McColl BK, Wiradajaja F, Wijayaratnam AP, Gleeson P, Gething MJ, Sambrook J, Mitchell CA. 2000. The yeast inositol polyphosphate 5-phosphatases inp52p and inp53p translocate to actin patches following hyperosmotic stress: mechanism for regulating phosphatidylinositol 4,5-bisphosphate at plasma membrane invaginations. *Mol Cell Biol* 20:9376–9390. <http://dx.doi.org/10.1128/MCB.20.24.9376-9390.2000>.
  129. Yin HL, Janmey PA. 2003. Phosphoinositide regulation of the actin cytoskeleton. *Annu Rev Physiol* 65:761–789. <http://dx.doi.org/10.1146/annurev.physiol.65.092101.142517>.
  130. Coue M, Brenner SL, Spector I, Korn ED. 1987. Inhibition of actin polymerization by latrunculin A. *FEBS Lett* 213:316–318. [http://dx.doi.org/10.1016/0014-5793\(87\)81513-2](http://dx.doi.org/10.1016/0014-5793(87)81513-2).
  131. Pruyne D, Bretscher A. 2000. Polarization of cell growth in yeast. I. Establishment and maintenance of polarity states. *J Cell Sci* 113:365–375.
  132. Karpova TS, Reck-Peterson SL, Elkind NB, Mooseker MS, Novick PJ, Cooper JA. 2000. Role of actin and Myo2p in polarized secretion and growth of *Saccharomyces cerevisiae*. *Mol Biol Cell* 11:1727–1737. <http://dx.doi.org/10.1091/mbc.11.5.1727>.
  133. Irazoqui JE, Lew DJ. 2004. Polarity establishment in yeast. *J Cell Sci* 117:2169–2171. <http://dx.doi.org/10.1242/jcs.00953>.
  134. Bi E, Park HO. 2012. Cell polarization and cytokinesis in budding yeast. *Genetics* 191:347–387. <http://dx.doi.org/10.1534/genetics.111.132886>.
  135. Chant J, Herskowitz I. 1991. Genetic control of bud site selection in yeast by a set of gene products that constitute a morphogenetic pathway. *Cell* 65:1203–1212. [http://dx.doi.org/10.1016/0092-8674\(91\)90015-Q](http://dx.doi.org/10.1016/0092-8674(91)90015-Q).
  136. Chant J, Pringle JR. 1991. Budding and cell polarity in *Saccharomyces cerevisiae*. *Curr Opin Genet Dev* 1:342–350. [http://dx.doi.org/10.1016/S0959-437X\(05\)80298-9](http://dx.doi.org/10.1016/S0959-437X(05)80298-9).
  137. Freifelder D. 1960. Bud position in *Saccharomyces cerevisiae*. *J Bacteriol* 80:567–568.
  138. Casamayor A, Snyder M. 2002. Bud-site selection and cell polarity in budding yeast. *Curr Opin Microbiol* 5:179–186. [http://dx.doi.org/10.1016/S1369-5274\(02\)00300-4](http://dx.doi.org/10.1016/S1369-5274(02)00300-4).
  139. Ni L, Snyder M. 2001. A genomic study of the bipolar bud site selection pattern in *Saccharomyces cerevisiae*. *Mol Biol Cell* 12:2147–2170. <http://dx.doi.org/10.1091/mbc.12.7.2147>.
  140. Pringle JR, Bi E, Harkins HA, Zahner JE, De Virgilio C, Chant J, Corrado K, Fares H. 1995. Establishment of cell polarity in yeast. *Cold Spring Harbor Symp Quant Biol* 60:729–744. <http://dx.doi.org/10.1101/SQB.1995.060.01.079>.
  141. Fujita A, Oka C, Arikawa Y, Katagai T, Tonouchi A, Kuhara S, Misumi Y. 1994. A yeast gene necessary for bud-site selection encodes a protein similar to insulin-degrading enzymes. *Nature* 372:567–570. <http://dx.doi.org/10.1038/372567a0>.
  142. Lord M, Inose F, Hiroko T, Hata T, Fujita A, Chant J. 2002. Subcellular localization of Axl1, the cell type-specific regulator of polarity. *Curr Biol* 12:1347–1352. [http://dx.doi.org/10.1016/S0960-9822\(02\)01042-4](http://dx.doi.org/10.1016/S0960-9822(02)01042-4).
  143. Strathern J, Hicks J, Herskowitz I. 1981. Control of cell type in yeast by the mating type locus. The alpha 1-alpha 2 hypothesis. *J Mol Biol* 147:357–372.
  144. Saito H. 2010. Regulation of cross-talk in yeast MAPK signaling pathways. *Curr Opin Microbiol* 13:677–683. <http://dx.doi.org/10.1016/j.mib.2010.09.001>.
  145. Qi M, Elion EA. 2005. MAP kinase pathways. *J Cell Sci* 118:3569–3572. <http://dx.doi.org/10.1242/jcs.02470>.
  146. Murphy LO, Blenis J. 2006. MAPK signal specificity: the right place at the right time. *Trends Biochem Sci* 31:268–275. <http://dx.doi.org/10.1016/j.tibs.2006.03.009>.
  147. Bardwell L. 2006. Mechanisms of MAPK signalling specificity. *Biochem Soc Trans* 34:837–841. <http://dx.doi.org/10.1042/BST0340837>.
  148. Maeda T, Wurgler-Murphy SM, Saito H. 1994. A two-component system that regulates an osmosensing MAP kinase cascade in yeast. *Nature* 369:242–245. <http://dx.doi.org/10.1038/369242a0>.
  149. Brewster JL, de Valoir T, Dwyer ND, Winter E, Gustin MC. 1993. An osmosensing signal transduction pathway in yeast. *Science* 259:1760–1763. <http://dx.doi.org/10.1126/science.7681220>.
  150. Elorza MV, Rico H, Sentandreu R. 1983. Calcofluor white alters the assembly of chitin fibrils in *Saccharomyces cerevisiae* and *Candida albicans* cells. *J Gen Microbiol* 129:1577–1582.
  151. Wood PJ. 1980. Specificity in the interaction of direct dyes with polysaccharides. *Carbohydr Res* 85:271–287. [http://dx.doi.org/10.1016/S0008-6215\(00\)84676-5](http://dx.doi.org/10.1016/S0008-6215(00)84676-5).
  152. Cali BM, Doyle TC, Botstein D, Fink GR. 1998. Multiple functions for actin during filamentous growth of *Saccharomyces cerevisiae*. *Mol Biol Cell* 9:1873–1889. <http://dx.doi.org/10.1091/mbc.9.7.1873>.
  153. Shock TR, Thompson J, Yates JR, III, Madhani HD. 2009. Hog1 mitogen-activated protein kinase (MAPK) interrupts signal transduction between the Kss1 MAPK and the Tec1 transcription factor to maintain pathway specificity. *Eukaryot Cell* 8:606–616. <http://dx.doi.org/10.1128/EC.00005-09>.
  154. Davenport KD, Williams KE, Ullmann BD, Gustin MC. 1999. Activation of the *Saccharomyces cerevisiae* filamentation/invasion pathway by osmotic stress in high-osmolarity glycogen pathway mutants. *Genetics* 153:1091–1103.
  155. Westfall PJ, Thorner J. 2006. Analysis of mitogen-activated protein kinase signaling specificity in response to hyperosmotic stress: use of an analog-sensitive HOG1 allele. *Eukaryot Cell* 5:1215–1228. <http://dx.doi.org/10.1128/EC.00037-06>.
  156. Karunanithi S, Vadaie N, Chavel CA, Birkaya B, Joshi J, Grell L, Cullen PJ. 2010. Shedding of the mucin-like flocculin Flo11p reveals a new aspect of fungal adhesion regulation. *Curr Biol* 20:1389–1395. <http://dx.doi.org/10.1016/j.cub.2010.06.033>.
  157. Kaiser CA, Schekman R. 1990. Distinct sets of SEC genes govern transport vesicle formation and fusion early in the secretory pathway. *Cell* 61:723–733. [http://dx.doi.org/10.1016/0092-8674\(90\)90483-U](http://dx.doi.org/10.1016/0092-8674(90)90483-U).