

SHORT REPORT

Open Access

Validity of calendar day-based definitions for community-onset bloodstream infections

Kevin B Laupland^{1,2,3*}, Daniel B Gregson^{2,3,4} and Deirdre L Church^{2,3,4}

Abstract

Background: Community-onset (CO) bloodstream infections (BSI) are those BSI where the blood culture is drawn <48 hours from hospital admission. However, exact times of culture draw or hospital admission are not always available. We evaluated the validity of using 2- or 3- calendar day based definitions for CO-BSI by comparing to a “gold standard” 48-hour definition.

Findings: Among the population-based cohort of 14,106 episodes of BSI studied, 10,543 were classified as CO based on “gold standard” 48-hour criteria. When 2-day and 3-day definitions were applied, 10,396 and 10,707 CO-BSI episodes were ascertained, respectively. All but 147 (1.4%) true CO-BSI cases were included by using the 2-day definition. When the 3-day definition was applied, all cases of CO-BSI were identified but an additional 164 (1.5%) cases of hospital-onset HO-BSI were also included. Thus the sensitivity and specificity of the 2-day definition was 98.6% and 100% and for the 3-day definition was 100% and 98.5%, respectively. Overall, only 311 (2.2%) cases were potentially miss-classifiable using either the 2- or 3-calendar day based definitions.

Conclusions: Use of either a 2- or 3-day definition is highly accurate for classifying CO-BSI.

Keywords: Nosocomial, Bacteremia, Community-acquired, Incidence, Epidemiology

Introduction

Community onset (CO) bloodstream infections (BSI) are those that occur in outpatients or among those with first positive blood culture drawn less than 48 hours following admission to hospital [1]. These may be further categorized as healthcare-associated or community-acquired CO-BSI, and are mutually exclusive of hospital-onset (HO) BSI that are first identified 48 hours or more following hospital admission [1-3]. While dates of admission and of culture draw date are typically routinely recorded, exact times of these occurrences may not be readily available. Use of a calendar day-based definition that utilizes the difference in dates of first positive culture draw and admission may be used to classify CO-BSI. However, because a 48-hour time difference between admission and culture draw may occur over a 2 or 3 calendar day period, it is not known whether use of such calendar day-based definitions accurately ascertains

the presence of a CO-BSI. The objective of this study was to evaluate the validity of calendar day- versus exact time-based definitions for CO-BSI.

Methods

The base study population consisted of all Calgary, Canada, area residents with incident BSI occurring during 2000–2009 as previously described [4,5]. True cases of CO-BSI were established by applying the “gold standard” or reference CO-BSI definition based on the exact times listed for each of culture draw and hospital admission registration. Two CO-BSI study cohorts were then developed by applying 2 and 3-calendar day definitions to the overall cohort. In these cases, only dates, and not times, of culture draw and admission were considered. The 2-day definition included those where the date of culture was before, on the same day, or one day after admission, and the 3-day definition included those where the culture draw was before, on the same day, or within the next two days following admission to hospital. This study was approved by the ethics review board at the University of Calgary.

* Correspondence: klaupland@gmail.com

¹Department of Medicine, Royal Inland Hospital, 311 Columbia Street, Kamloops, BC V2C 2T1, Canada

²Department of Medicine, University of Calgary, Calgary, AB, Canada
Full list of author information is available at the end of the article

Findings

During the study, there were 14,233 incident episodes of BSI; 10,543 were classified as CO and 3,680 as HO. Among this cohort, 127 episodes classified as HO-BSI had first positive cultures less than 48 hours following admission to one of the study hospitals as a result of either a recent hospital discharge or transfer from another institution and were excluded from further analysis. Among the 10,543 CO-BSI, 8,730 were admitted and 1,813 were not associated with an admission to hospital. The overall cohort admitted to hospital included 12,283 BSI cases of which 8,730 were CO and 3,553 were HO.

Among the 8,730 true CO-BSI cases that were admitted to hospital, the median culture draw minus hospital admission time was -1 (IQR, -5 to 2) hours. The time of culture draw for these cases was distributed tightly around the time of admission as shown in Figure 1. Only 339 (3.9%) true CO-BSI cases were cultured between 24 and 47 hours following admission. In comparison, among the HO-BSI cases, the median time to culture draw was 12 (IQR, 6-24) days following admission. The distribution of these cases showed a progressive decrease from day 2. Of the HO-BSI cases, 290 (8%) were first cultured between 48-71 hours and 217 (6%) between 72 and 95 hours following hospital admission.

When the 2-day definition was applied, all but 147/10,543 (1.4%) true CO-BSI cases were included. When the 3-day definition was applied, all cases of CO-BSI were identified but an additional 164 (1.5%) cases of HO-BSI were also included. In other words, the sensitivity and

specificity of the 2-day definition was 98.6% and 100% and for the 3-day definition was 100% and 98.5%, respectively. Overall, only 311 (2.2%) cases were potentially misclassifiable using either the 2- or 3-calendar day based definitions. Among this cohort of 311 cases there were no evident features as regards gender, age, poly-microbial, antibiotic resistance, and species distribution that would allow their further classification as either CO- or HO-BSI.

Discussion

In this study we show that use of either a 2 or 3 calendar-day based definition is highly accurate (>98%) to classify cases as to true CO-BSI cases based on a 48-hour definition. The ability to accurately classify CO-BSI arises due to the observation that the majority of CO cases are identified within a few hours of hospital admission (Figure 1). Use of the 2 calendar day definition results in the inclusion of only true cases, although it does fail to include a very small (1.4%) number of these cases. On the other hand, the 3-day definition results in the inclusion of all true CO-BSI cases albeit with the inclusion of a small number (1.5%) of misclassified HO-BSI cases. These data indicate that both the 2- or 3-day definitions are highly accurate for classification of CO-BSI with the former having slightly higher specificity and the latter sensitivity.

Accurate determination of CO-BSI is important for epidemiology and surveillance purposes. Traditionally BSI's were classified based on attempts at defining location of acquisition infection with application of complicated




Figure 1 Timing of blood culture draw following admission to hospital for community-onset bloodstream infections.

definitions on a case-by-case review [6]. While standardized criteria are used in these cases, they still require a degree of subjective interpretation. As a result, reproducibility and inter-observer variation are considerable [7]. In contrast, CO-BSI are strictly defined by easily applied and objective time based criteria. These attributes allow objective inter-facility and regional comparisons and facilitate the use of electronic surveillance systems [8-10].

Strengths of this study include that we studied a large cohort of patients and that data were consistently and systematically recorded. Our system comprehensively identifies all patients with BSI in our population including those who are delayed in being admitted to hospital or who are not admitted to hospital for management of their BSI. We included all incident BSIs occurring in a well-defined population such that selection bias was minimized [4]. However, a potential limitation is that we defined the admission time as that registered within our hospital administrative database. This time is entered once an order has been written to admit the patient to hospital. The possibility exists that there could be delays from presentation/decision to admit the patient to hospital and the actual entry of the admission order as patients may have tests performed and assessments by consultants prior to establishing the patients admission disposition and order. However, this time is unlikely exceed a few hours at most and since the vast majority of CO-BSI occur with hours of admission, it is unlikely to have any significant effect on our study conclusions.

In summary, this study validates the practice of using calendar day based definitions for CO-BSI. Furthermore, these data indicate that 2- and 3-day based definitions are highly specific and sensitive and should be adequate for most surveillance purposes where exact times of admission and culture draw are not readily available.

Abbreviations

BSI: Bloodstream infection; CO: Community-onset; HO: Hospital-onset.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KBL conceived and designed the study, collected data, conducted the primary analysis, and drafted the manuscript. DBG and DLC contributed to data collection and revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors take full responsibility for the conduct of the study and no other individuals were involved in the preparation of this manuscript. This was an unfunded study and no external funds were utilized.

Author details

¹Department of Medicine, Royal Inland Hospital, 311 Columbia Street, Kamloops, BC V2C 2T1, Canada. ²Department of Medicine, University of Calgary, Calgary, AB, Canada. ³Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, AB, Canada. ⁴Division of Microbiology, Calgary Laboratory Services, #9, 3535 Research Road NW, Calgary, AB T2A 2K8, Canada.

Received: 17 December 2014 Accepted: 11 March 2015

Published online: 02 April 2015

References

1. Laupland KB, Church DL. Population-Based Epidemiology and Microbiology of Community-Onset Bloodstream Infections. *Clin Microbiol Rev.* 2014;27(4):647–64.
2. Friedman ND, Kaye KS, Stout JE, McGarry SA, Trivette SL, Briggs JP, et al. Health care-associated bloodstream infections in adults: a reason to change the accepted definition of community-acquired infections. *Ann Intern Med.* 2002;137(10):791–7.
3. Morin CA, Hadler JL. Population-based incidence and characteristics of community-onset *Staphylococcus aureus* infections with bacteremia in 4 metropolitan Connecticut areas, 1998. *J Infect Dis.* 2001;184(8):1029–34.
4. Laupland KB. Defining the epidemiology of bloodstream infections: the 'gold standard' of population-based assessment. *Epidemiol Infect.* 2013;141(10):2149–57.
5. Lenz R, Leal JR, Church DL, Gregson DB, Ross T, Laupland KB. The distinct category of healthcare associated bloodstream infections. *BMC Infect Dis.* 2012;12:85.
6. Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control.* 1988;16(3):128–40.
7. Mayer J, Greene T, Howell J, Ying J, Rubin MA, Trick WE. Agreement in classifying bloodstream infections among multiple reviewers conducting surveillance. *Clin Infect Dis.* 2012;55(3):364–70.
8. Leal J, Gregson DB, Ross T, Flemons WW, Church DL, Laupland KB. Development of a novel electronic surveillance system for monitoring of bloodstream infections. *Infect Control Hosp Epidemiol.* 2010;31(7):740–7.
9. Leal J, Laupland KB. Validity of electronic surveillance systems: a systematic review. *J Hosp Infect.* 2008;69(3):220–9.
10. Laupland KB, Lyytikäinen O, Sogaard M, Kennedy KJ, Knudsen JD, Ostergaard C, et al. The changing epidemiology of *Staphylococcus aureus* bloodstream infection: a multinational population-based surveillance study. *Clin Microbiol Infect.* 2013;19(5):465–71.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

