

# Heteroresistance to Colistin in *Klebsiella pneumoniae* Associated with Alterations in the PhoPQ Regulatory System

Aurélie Jayol,<sup>a</sup> Patrice Nordmann,<sup>a,b</sup> Adrian Brink,<sup>c</sup> Laurent Poirel<sup>a</sup>

Medical and Molecular Microbiology “Emerging Antibiotic Resistance” Unit, Department of Medicine, Faculty of Science, University of Fribourg,<sup>a</sup> and HFR–Hôpital Cantonal de Fribourg,<sup>b</sup> Fribourg, Switzerland; Department of Clinical Microbiology, Ampath National Laboratory Services, Milpark Hospital, Johannesburg, South Africa<sup>c</sup>

**A multidrug-resistant *Klebsiella pneumoniae* isolate exhibiting heteroresistance to colistin was investigated. The colistin-resistant subpopulation harbored a single amino acid change (Asp191Tyr) in protein PhoP, which is part of the PhoPQ two-component system that activates *pmrHFIJKLM* expression responsible for L-aminoarabinose synthesis and polymyxin resistance. Complementation assays with a wild-type *phoP* gene restored full susceptibility to colistin. Then, analysis of the colistin-susceptible subpopulation showed a partial deletion (25 bp) in the *phoP* gene compared to that in the colistin-resistant subpopulation. That deletion disrupted the reading frame of *phoP*, leading to a longer and inactive protein (255 versus 223 amino acids long). This is the first report showing the involvement of mutation(s) in PhoP in colistin resistance. Furthermore, this is the first study to decipher the mechanisms leading to colistin heteroresistance in *K. pneumoniae*.**

*Klebsiella pneumoniae* is a Gram-negative bacterium often associated with nosocomial infections, including urinary tract infection, pneumonia, and bloodstream infection (1). While multidrug resistance is increasingly reported in that species due to acquisition of numerous resistance traits, including extended-spectrum  $\beta$ -lactamase and carbapenemase genes, colistin is increasingly used for treating infections due to multidrug-resistant isolates. Colistin and polymyxin B correspond to important therapeutic options for treating infections caused by multidrug-resistant *K. pneumoniae*, particularly in countries with a high prevalence of carbapenemase producers (2). Those drugs are bactericidal for Gram-negative bacteria, interacting with the lipid A moiety of lipopolysaccharide (LPS) and subsequently causing disorganization of the outer membrane (3).

In *K. pneumoniae*, colistin resistance is being increasingly reported, and some cases of heteroresistant *K. pneumoniae* isolates have also been observed (4, 5). Colistin heteroresistance is defined as “the emergence of resistance to colistin by a subpopulation from an otherwise susceptible (MIC of  $\leq 2$  mg/liter) population” (6) that may be related to exposure to a suboptimal polymyxin concentration (7). A comparison of the *in vitro* susceptibility testing methods for colistin showed that the Etest and agar dilution methods are reliable for detecting resistant subpopulations in contrast to some automatic methods; nonetheless, the broth microdilution method remains the reference test for determination of colistin MICs (8).

Several studies have shown that mutations in the PmrAB or PhoPQ regulatory systems may confer colistin resistance in *K. pneumoniae* (9–12). In addition, inactivation of the *mgrB* gene, which encodes the MgrB protein known to negatively regulate the PhoPQ signaling system, may also be the source of acquired resistance to colistin (13–16).

Acquired resistance to polymyxins is mediated by the addition of 4-deoxyaminoarabinose (LAra4N) and/or phosphoethanolamine (pEtN) to lipid A. LAra4N synthesis requires the products of the *pmrE* gene and *pmrHFIJKLM* operon, and pEtN synthesis is encoded by the *pmrC* gene. These modifications create a more positively charged lipopolysaccharide and thus reduce the affinity

of LPS to positively charged polymyxins (7). Of note, mechanisms leading to heteroresistance in *K. pneumoniae* remain unknown.

PmrB and PhoQ are sensor cytoplasmic membrane-bound kinases activated by high concentrations of iron ( $\text{Fe}^{3+}$ ) and an acidic pH (pH 5.5) for the PmrAB system (17) and by low extracellular concentrations of magnesium ( $\text{Mg}^{2+}$ ) for the PhoPQ system (18). Upon activation, PmrB activates the response regulator PmrA, which in turn upregulates *pmrC* and *pmrHFIJKLM*. Likewise, PhoQ activates PhoP, which in turn activates the expression of the *pmrHFIJKLM* operon, directly by binding to the *pmrHFIJKLM* promoter and indirectly via PmrD-dependent activation of the PmrA protein also binding to the *pmrHFIJKLM* promoter (19).

The aim of this study was to determine the mechanism(s) responsible for heteroresistance to colistin in *K. pneumoniae*.

## MATERIALS AND METHODS

**Bacterial strains.** A multidrug-resistant *K. pneumoniae* clinical isolate named Kp75 was recovered in 2012 in South Africa. It produced the carbapenemase OXA-48, which has been reported elsewhere (20). The wild-type and colistin-susceptible *K. pneumoniae* ATCC 53153 strain was used as a control throughout the study.

**Antimicrobial susceptibility assays.** MICs were determined using Etest strips (AB bioMérieux, La Balme-les-Grottes, France) on Mueller-Hinton agar plates (Bio-Rad, Marnes-la-Coquette, France) with a 0.5 McFarland inoculum. The MICs of colistin were also determined by a broth

Received 15 December 2014 Returned for modification 11 January 2015

Accepted 23 February 2015

Accepted manuscript posted online 2 March 2015

Citation Jayol A, Nordmann P, Brink A, Poirel L. 2015. Heteroresistance to colistin in *Klebsiella pneumoniae* associated with alterations in the PhoPQ regulatory system. *Antimicrob Agents Chemother* 59:2780–2784. doi:10.1128/AAC.05055-14.

Address correspondence to Laurent Poirel, laurent.poirel@unifr.ch.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.05055-14>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved. doi:10.1128/AAC.05055-14

TABLE 1 Oligonucleotides used as primers in this study

Primer	Sequence (5' to 3')	Gene	Reference or study
pmrA ext F	CAT TTC CGC GCA CTG TCT GC	<i>pmrA</i>	12
pmrA ext R	CAG GTT TCA GTT GCA AAC AG	<i>pmrA</i>	12
pmrB ext F	ACC TAC GCG AAA AGA TTG GC	<i>pmrB</i>	12
pmrB ext R	GAT GAG GAT AGC GCC CAT GC	<i>pmrB</i>	12
phoP ext F	GAG CTT CAG ACT ACT ATC GA	<i>phoP</i>	12
phoP ext R	GGG AAG ATA TGC CGC AAC AG	<i>phoP</i>	12
phoQ ext F	ATA CCC ACA GGA CGT CAT CA	<i>phoQ</i>	12
phoQ ext R	CAG GTG TCT GAC AGG GAT TA	<i>phoQ</i>	12
mgrB ext F	TTA AGA AGG CCG TGC TAT CC	<i>mgrB</i>	13
mgrB ext R	AAG GCG TTC ATT CTA CCA CC	<i>mgrB</i>	13
mdh F	CCC AAC TCG CTT CAG GTT CAG	<i>mdh</i>	26
mdh R	CCG TTT TTC CCC AGC AGC AG	<i>mdh</i>	26
pmrC int F	GCG TGA TGA ATA TCC TCA CCA	<i>pmrC</i>	12
pmrC int R	CAC GCC AAA GTT CCA GAT GA	<i>pmrC</i>	12
pmrA int F	GAT GAA GAC GGG CTG CAT TT	<i>pmrA</i>	12
pmrA int R	ACC GCT AAT GCG ATC CTC AA	<i>pmrA</i>	12
pmrB int F	TGC CAG CTG ATA AGC GTC TT	<i>pmrB</i>	12
pmrB int R	TTC TGG TTG TTG TGC CCT TC	<i>pmrB</i>	12
pmrD int F	GAT CGC AGA GAT TGA AGC CT	<i>pmrD</i>	12
pmrD int R	GCG TTG CGG ATC TTC AAA GT	<i>pmrD</i>	12
pmrE int F	GCA TAC CGT AAT GCC GAC TA	<i>pmrE</i>	12
pmrE int R	GGG TTG ATC TCT GTG ACA TC	<i>pmrE</i>	12
pmrK int F	AGT ATC GGT CAG TGG CTG TT	<i>pmrK</i>	12
pmrK int R	CCG CTT ATC ACG AAA GAT CC	<i>pmrK</i>	12
phoP int F	GCG TCA CCA CCT CAA AGT TC	<i>phoP</i>	This study
phoP int R	GGC GAT ATC CGG GAG ATG TT	<i>phoP</i>	This study
phoQ int F	CTC AAG CGC AGC TAT ATG GT	<i>phoQ</i>	This study
phoQ int R	TCT TTG GCC AGC GAC TCA AT	<i>phoQ</i>	This study
rpsL int F	CCG TGG CGG TCG TGT TAA AGA	<i>rpsL</i>	13
rpsL int R	GCC GTA CTT GGA GCG AGC CTG	<i>rpsL</i>	13

culture microdilution method as recommended by the CLSI (21), with Tween 80 being added at 0.002% as suggested in several studies (22, 23). According to the EUCAST guidelines, isolates with MICs of  $\leq 2$   $\mu\text{g/ml}$  are categorized as susceptible, and those with MICs of  $> 2$   $\mu\text{g/ml}$  are categorized as resistant (24).

**PCR amplification and sequencing.** Whole-cell DNA was extracted using a commercially available kit (QIAquick; Qiagen, Valencia, CA) according to the manufacturer's instructions. The *pmrA*, *pmrB*, *phoP*, *phoQ*, and *mgrB* genes were amplified with specific oligonucleotide primers (the primers used in this study are listed in Table 1). Molecular detection of the  $\beta$ -lactam resistance genes was performed as reported previously (25). The amplified DNA fragments were purified with the QIAquick PCR purification kit (Qiagen, Courtaboeuf, France). Both strands of the amplification products obtained were sequenced with an ABI 3100 sequencer (Applied Biosystems, Foster City, CA). The nucleotide and deduced protein sequences were analyzed at the National Center for Biotechnology Information website ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) by the Basic Local Alignment Search Tool (BLAST) program.

**Analysis of the primary and secondary structures of the PhoP protein.** The primary structure of PhoP was analyzed by using the Pfam website (<http://pfam.xfam.org/>). The predicted secondary structures of the proteins were obtained by the GOR method using the EMBOSS 6.3.1 software available on the Mobylye@Pasteur portal (<http://mobylye.pasteur.fr/cgi-bin/portal.py>).

**Complementation experiments.** The entire *phoP* and *phoQ* genes from the colistin-susceptible *K. pneumoniae* ATCC 53153 reference strain (MIC of colistin at 0.125  $\mu\text{g/ml}$ ) were amplified by PCR using 2 $\times$  Phusion HF master mix (Thermo Scientific Finnzymes, Illkirch, France) and primers sets phoP ext F/phoP ext R and phoQ ext F/phoQ ext R, respectively.

The noncoding sequence *mdh* was also amplified with primer set mdh F/mdh R as described previously (26) (the primers used are listed in Table 1). The amplified fragments were cloned into plasmid pCR-BluntII-TOPO (Applied Biosystems by Life Technologies, Carouge, Switzerland) and the resulting plasmids pTOPO-phoP, pTOPO-phoQ, and pTOPO-mdh were, respectively, introduced into the colistin-resistant isolate Kp75 by electroporation. Transformants were selected by overnight incubation at 37°C on Mueller-Hinton agar supplemented with 100  $\mu\text{g/ml}$  of zeocin. The MICs of colistin were determined by Etest for all *K. pneumoniae* transformants.

**Growth curves.** Growth kinetics were determined with subpopulations of isolate Kp75 in the absence of any selective pressure. Two hundred-milliliter volumes of Luria-Bertani broth were inoculated independently with  $10^8$  CFU of each strain, and cultures were grown for 24 h at 100 rpm and 37°C. Absorbance at 600 nm was measured every hour for the first 10 h and after 24 h. The colony counting was performed by serial dilution and final plating on solid medium.

**Transcriptional analysis by qRT-PCR.** Quantitative real-time PCR (qRT-PCR) was used to measure the expression of the *phoP*, *phoQ*, *pmrD*, *pmrC*, *pmrA*, *pmrB*, and *pmrK* genes, using specific oligonucleotide primers (Table 1). A culture volume of 0.5 ml was taken during the mid-log phase of growth (optical density at 600 nm [ $\text{OD}_{600 \text{ nm}}$ ] = 1.1) and combined with 1 ml of RNeasy Protect (Qiagen, Courtaboeuf, France). Total RNA was extracted from cell lysates using the RNeasy minikit (Qiagen), according to the manufacturer's instructions. qRT-PCR was carried out using a Rotor-Gene Q instrument (Qiagen), and QuantiFast SYBR green (Qiagen) was used as a signal reporter. In each run, a blank sample (distilled water) and a no reverse transcriptase control were included to exclude DNA contamination. Relative gene expression differences were cal-




FIG 1 MIC determination using a colistin Etest strip for *K. pneumoniae* Kp75. The arrow indicates the colistin-resistant subpopulation (Kp75b).

culated by first normalizing the  $C_T$  values by subtracting the 12S rRNA control (*rpsL* gene) (13) and then comparing to the values obtained for the susceptible isolate.

**Nucleotide sequence accession numbers.** The nucleotide and protein sequences of the PhoP proteins of isolates Kp75a and Kp75b were registered in GenBank under accession numbers [KP742843](#) and [KP742844](#), respectively.

## RESULTS AND DISCUSSION

**Isolate Kp75 exhibits multidrug resistance, including resistance to colistin.** Isolate Kp75 was resistant to all  $\beta$ -lactams, including carbapenems (the MICs of imipenem, meropenem, and ertapenem were 3, 8, and  $>32$   $\mu\text{g/ml}$ , respectively). In addition, it was resistant to chloramphenicol, fluoroquinolones, gentamicin, tetracycline, and tobramycin. The MIC of rifampin was 16  $\mu\text{g/ml}$ , and the MIC of nitrofurantoin was  $>128$   $\mu\text{g/ml}$ . It remained susceptible only to sulfonamides, tigecycline, and amikacin. Determination of the MIC of colistin by using the broth culture microdilution method showed that isolate Kp75 was resistant to colistin (MIC of 128  $\mu\text{g/ml}$ ). Molecular detection of the  $\beta$ -lactamase resistance genes identified the narrow-spectrum  $\beta$ -lactamase *bla*<sub>TEM-1</sub> gene, the expanded-spectrum  $\beta$ -lactamase *bla*<sub>CTX-M-15</sub> gene, and the carbapenem-hydrolyzing class D  $\beta$ -lactamase *bla*<sub>OXA-48</sub> gene.

**Susceptibility testing by Etest revealed a heteroresistant phenotype.** When the MIC of colistin for isolate Kp75 was measured by using Etest strips, two subpopulations were identified close to the Etest strip (Fig. 1). Bacterial colonies located in the resistance and susceptible areas were picked up, and the two subpopulations were separated. The MICs for the two distinct subpopulations were again measured by Etest. The colistin-susceptible subpopulation (then termed Kp75a) and the colistin-resistant subpopulation (then termed Kp75b) had MICs of colistin at 0.12 and 12  $\mu\text{g/ml}$ , respectively. With use of the microdilution method, the MICs were found to be 0.12 and 128  $\mu\text{g/ml}$ , respectively. These data further highlight the discrepancies between values obtained by the two methods and reveal that the MICs obtained by the Etest

are significantly lower than those obtained by the microdilution method for such a resistant isolate.

**A single amino acid substitution in PhoP leads to colistin resistance in isolate Kp75b.** Total DNAs from isolate Kp75b and colistin-susceptible *K. pneumoniae* reference strain ATCC 53153 were used as the templates for PCRs to amplify the *pmrA*, *pmrB*, *phoP*, *phoQ*, and *mgrB* genes possibly involved in colistin resistance. A single base pair substitution was identified in the *phoP* gene of isolate Kp75b, leading to a single amino acid substitution (Asp191Tyr) compared to that of the PhoP proteins available in the GenBank databases and also compared to that of strain ATCC 53153 (Fig. 2). No substitution was identified in the *pmrA*, *pmrB*, *phoQ*, and *mgrB* genes.

The amino acid substitution was identified at position 191 of the PhoP protein, which is located in its C-terminal domain. We therefore speculate that the replacement of a polar negative amino acid by an aromatic amino acid partially hydrophobic in the C-terminal domain of PhoP might induce a constitutive activation of PhoP, deregulating the two-component regulatory system PhoPQ. The predicted secondary structures of two PhoP proteins, namely that of a wild-type strain and that of isolate Kp75b, were compared, and it was revealed that the Asp191Tyr amino acid substitution significantly modified the secondary structure of the protein, with an interruption of the  $\alpha$ -helix (see Fig. S1 in the supplemental material). It was therefore likely that resistance to colistin in isolate Kp75b was related to that single substitution in PhoP. To further assess this hypothesis, an experiment was conducted in order to complement isolate Kp75b with the entire *phoP* gene obtained from the wild-type strain ATCC 53153. The entire *phoP* and *phoQ* genes, together with a fragment of the *mdh* gene (a noncoding sequence as control), were amplified from the reference strain ATCC 53153 and cloned in plasmid pTOPO-BluntII, giving rise to, respectively, pTOPO-*phoP*, pTOPO-*phoQ*, and pTOPO-*mdh* and then transformed into the resistant-colistin clinical isolate Kp75b. The MIC determinations showed a complete reversion to susceptibility only for the transformant complemented with pTOPO-*phoP* (MIC at 0.125  $\mu\text{g/ml}$ ), whereas the MICs of transformants complemented with recombinant plasmids pTOPO-*phoQ* or pTOPO-*mdh* remained unchanged. This result strongly suggested that the amino acid substitution identified in PhoP was indeed responsible for the colistin resistance in Kp75b (Fig. 3).

**A partial deletion of the *phoP* gene is responsible for a reversion in colistin susceptibility.** Similarly to what was performed with isolate Kp75b, PCRs followed by sequencing were performed to amplify the *pmrA*, *pmrB*, *phoP*, *phoQ*, and *mgrB* genes of the colistin-susceptible isolate Kp75a. All sequences were found to be identical with those obtained with Kp75b, except for the *phoP* gene. Indeed, the *phoP* gene harbored not only a substitution leading to the Asp191Tyr amino acid substitution in the corresponding protein sequence but also a partial deletion (25 bp long) in the *phoP* gene of Kp75a, disrupting its reading frame and consequently giving rise to a longer protein of 255 amino acids instead of 223 amino acids as found in Kp75b and all other PhoP sequences in the databases (Fig. 2). This deletion was therefore considered to be responsible for the reversion into a colistin-susceptible phenotype. Figure 3 provides a summary of the evolution process observed in Kp75. This mechanism of partial gene deletion involved in reversion to colistin susceptibility was previously





In addition, we want to emphasize that the phenomenon of colistin heteroresistance remains very likely underestimated, considering that it may not be seen when the method (broth microdilution) recommended by the CLSI is used for the evaluation of susceptibility to polymyxins. We showed here that heteroresistance may be well demonstrated when only Etest strips are used. Notably, the clinical significance of colistin heteroresistance in *K. pneumoniae* is still unknown and remains to be evaluated.

## ACKNOWLEDGMENTS

This work was funded by the University of Fribourg, Switzerland, and by a grant from the European Community (R-GNOSIS, FP7/HEALTH-F3-2011-282512).

## REFERENCES

- Podschun R, Ullmann U. 1998. *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clin Microbiol Rev* 11:589–603.
- Falagas ME, Lourida P, Poulidakos P, Rafailidis PI, Tansarli GS. 2014. Antibiotic treatment of infections due to carbapenem-resistant *Enterobacteriaceae*: systematic evaluation of the available evidence. *Antimicrob Agents Chemother* 58:654–663. <http://dx.doi.org/10.1128/AAC.01222-13>.
- Hancock RE. 1997. Peptide antibiotics. *Lancet* 349:418–422. [http://dx.doi.org/10.1016/S0140-6736\(97\)80051-7](http://dx.doi.org/10.1016/S0140-6736(97)80051-7).
- Meletis G, Tzampaz E, Sianou E, Tzavaras I, Sofianou D. 2011. Colistin heteroresistance in carbapenemase-producing *Klebsiella pneumoniae*. *J Antimicrob Chemother* 66:946–947. <http://dx.doi.org/10.1093/jac/dkr007>.
- Poudyal A, Howden BP, Bell JM, Gao W, Owen RJ, Turnidge JD, Nation RL, Li J. 2008. In vitro pharmacodynamics of colistin against multidrug-resistant *Klebsiella pneumoniae*. *J Antimicrob Chemother* 62:1311–1318. <http://dx.doi.org/10.1093/jac/dkn425>.
- Li J, Nation RL, Turnidge JD, Milne RW, Coulthard K, Rayner CR, Paterson DL. 2006. Colistin: the re-emerging antibiotic for multidrug-resistant Gram-negative bacterial infections. *Lancet Infect Dis* 6:589–601. [http://dx.doi.org/10.1016/S1473-3099\(06\)70580-1](http://dx.doi.org/10.1016/S1473-3099(06)70580-1).
- Falagas ME, Rafailidis PI, Matthaïou DK. 2010. Resistance to polymyxins: mechanisms, frequency and treatment options. *Drug Resist Updat* 13:132–138. <http://dx.doi.org/10.1016/j.drup.2010.05.002>.
- Humphries RM. 2015. Susceptibility testing of the polymyxins: where are we now? *Pharmacotherapy* 35:22–27. <http://dx.doi.org/10.1002/phar.1505>.
- Cannatelli A, Di Pilato V, Giani T, Arena F, Ambretti S, Gaibani P, D'Andrea MM, Rossolini GM. 2014. In vivo evolution to colistin resistance by PmrB sensor kinase mutation in KPC-producing *Klebsiella pneumoniae* is associated with low-dosage colistin treatment. *Antimicrob Agents Chemother* 58:4399–4403. <http://dx.doi.org/10.1128/AAC.02555-14>.
- Cheng HY, Chen YF, Peng HL. 2010. Molecular characterization of the PhoPQ-PmrD-PmrAB mediated pathway regulating polymyxin B resistance in *Klebsiella pneumoniae* CG43. *J Biomed Sci* 17:60. <http://dx.doi.org/10.1186/1423-0127-17-60>.
- Choi MJ, Ko KS. 2014. Mutant prevention concentrations of colistin for *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* clinical isolates. *J Antimicrob Chemother* 69:275–277. <http://dx.doi.org/10.1093/jac/dkt315>.
- Jayol A, Poirel L, Brink A, Villegas MV, Yilmaz M, Nordmann P. 2014. Resistance to colistin associated with a single amino acid change in protein PmrB among *Klebsiella pneumoniae* isolates of worldwide origin. *Antimicrob Agents Chemother* 58:4762–4766. <http://dx.doi.org/10.1128/AAC.00084-14>.
- Cannatelli A, D'Andrea MM, Giani T, Di Pilato V, Arena F, Ambretti S, Gaibani P, Rossolini GM. 2013. In vivo emergence of colistin resistance in *Klebsiella pneumoniae* producing KPC-type carbapenemase mediated by insertional inactivation of the PhoQ/PhoP *mgrB* regulator. *Antimicrob Agents Chemother* 57:5521–5526. <http://dx.doi.org/10.1128/AAC.01480-13>.
- Cannatelli A, Giani T, D'Andrea MM, Di Pilato V, Arena F, Conte V, Tryfinopoulou K, Vatopoulos A, Rossolini GM. 2014. MgrB inactivation is a common mechanism of colistin resistance in KPC-producing *Klebsiella pneumoniae* of clinical origin. *Antimicrob Agents Chemother* 58:5696–5703. <http://dx.doi.org/10.1128/AAC.03110-14>.
- Poirel L, Jayol A, Bontron S, Villegas MV, Özdamar M, Turkoglu S, Nordmann P. 2015. The *mgrB* gene as a key target for acquired resistance to colistin in *Klebsiella pneumoniae*. *J Antimicrob Chemother* 70:75–80. <http://dx.doi.org/10.1093/jac/dku323>.
- Wright MS, Suzuki Y, Jones MB, Marshall SH, Rudin SD, van Duin D, Kaye K, Jacobs MR, Bonomo RA, Adams MD. 2015. Genomic and transcriptomic analyses of colistin-resistant clinical isolates of *Klebsiella pneumoniae* reveal multiple pathways of resistance. *Antimicrob Agents Chemother* 59:536–543. <http://dx.doi.org/10.1128/AAC.04037-14>.
- Gunn JS. 2008. The *Salmonella* PmrAB regulon: lipopolysaccharide modifications, antimicrobial peptide resistance and more. *Trends Microbiol* 16:284–290. <http://dx.doi.org/10.1016/j.tim.2008.03.007>.
- Groisman EA. 2001. The pleiotropic two-component regulatory system PhoP-PhoQ. *J Bacteriol* 183:1835–1842. <http://dx.doi.org/10.1128/JB.183.6.1835-1842.2001>.
- Mitrophanov AY, Jewett MW, Hadley TJ, Groisman EA. 2008. Evolution and dynamics of regulatory architectures controlling polymyxin B resistance in enteric bacteria. *PLoS Genet* 4:e1000233. <http://dx.doi.org/10.1371/journal.pgen.1000233>.
- Potron A, Poirel L, Rondinaud E, Nordmann P. 2013. Intercontinental spread of OXA-48  $\beta$ -lactamase-producing *Enterobacteriaceae* over a 11-year period, 2001 to 2011. *Euro Surveill* 18(31):pii=20549. <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=20549>.
- Clinical and Laboratory Standards Institute. 2014. Performance standards for antimicrobial susceptibility testing; 24th informational supplement. CLSI document M100-S24. Clinical and Laboratory Standards Institute, Wayne, PA.
- Hindler JA, Humphries RM. 2013. Colistin MIC variability by method for contemporary clinical isolates of multidrug-resistant Gram-negative bacilli. *J Clin Microbiol* 51:1678–1684. <http://dx.doi.org/10.1128/JCM.03385-12>.
- Sader HS, Rhomberg PR, Flamm RK, Jones RN. 2012. Use of a surfactant (polysorbate 80) to improve MIC susceptibility testing results for polymyxin B and colistin. *Diagn Microbiol Infect Dis* 74:412–414. <http://dx.doi.org/10.1016/j.diagmicrobio.2012.08.025>.
- European Committee on Antimicrobial Susceptibility Testing. 2015. Breakpoint tables for interpretation of MICs and zone diameters. Version 2.0. [http://www.eucast.org/clinical\\_breakpoints/](http://www.eucast.org/clinical_breakpoints/).
- Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis* 17:1791–1798. <http://dx.doi.org/10.3201/eid1710.110655>.
- Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. 2005. Multi-locus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* 43:4178–4182. <http://dx.doi.org/10.1128/JCM.43.8.4178-4182.2005>.
- Adams MD, Nickel GC, Bajaksouzian S, Lavender H, Murthy AR, Jacobs MR, Bonomo RA. 2009. Resistance to colistin in *Acinetobacter baumannii* associated with mutations in the PmrAB two-component system. *Antimicrob Agents Chemother* 53:3628–3634. <http://dx.doi.org/10.1128/AAC.00284-09>.