



# HHS Public Access

Author manuscript

*Ticks Tick Borne Dis.* Author manuscript; available in PMC 2015 June 01.

Published in final edited form as:

*Ticks Tick Borne Dis.* 2014 June ; 5(4): 449–452. doi:10.1016/j.ttbdis.2014.02.004.

## Detection of a *Borrelia* species in questing Gulf Coast ticks, *Amblyomma maculatum*

Jung Keun Lee<sup>a</sup>, Whitney Crow Smith<sup>b</sup>, Chelsea McIntosh<sup>b</sup>, Flavia Girao Ferrari<sup>c</sup>, Brittany Moore-Henderson<sup>d</sup>, and Andrea Varela-Stokes<sup>b,\*</sup>

<sup>a</sup>Department of Pathobiology and Population Medicine, Mississippi State University, Mississippi State, MS, USA

<sup>b</sup>Department of Basic Sciences, Mississippi State University, Mississippi State, MS, USA

<sup>c</sup>Merck Animal Health, Desoto, KS, USA

<sup>d</sup>College of Veterinary Medicine, Mississippi State University, Mississippi State, MS, USA

### Abstract

*Borrelia* spp. are agents of Lyme disease and relapsing fever, diseases which use *Ixodes* hard ticks and *Ornithodoros* soft ticks, respectively, as primary vectors. Some relapsing fever spirochetes, such as *B. miyamotoi*, are also found in hard ticks. To date, no *Borrelia* sp. is known to use the hard tick, *Amblyomma maculatum*, as a vector. However, both *B. burgdorferi* and *B. lonestari* were recently detected in *A. maculatum* removed from hosts. In our study, DNA extracts from 306 questing adult *A. maculatum* collected in Mississippi in 2009 and 2010 were tested for *Borrelia* spp. DNA by PCR amplification of *flaB* and 16S rRNA gene targets. An additional 97 *A. maculatum* collected in 2013 were tested by amplification of 16S rRNA gene target. Two ticks, one collected in 2009 and the other in 2010, were positive by PCR of the *flaB* and 16S rRNA gene targets; both were collected from the same location in central Mississippi. Interestingly, 16S rRNA gene amplicons from these two tick extracts were 98% identical to twelve *Borrelia* spp. including the reptile-associated spirochete *B. turcica* and *Borrelia* sp. “tAG158M”; *flaB* amplicons from these two ticks shared closest identity (89%) to the reptile-associated spirochete, *B. turcica*. These results demonstrate a *Borrelia* sp. in unfed *A. maculatum* ticks that is unique from other species in the NCBI database and in a clade with reptile-associated *Borrelia* species. Detection of a previously unrecognized *Borrelia* in a hard tick species generates additional questions regarding the bacterial fauna in these arthropods and warrants further studies to better understand this fauna.

### Keywords

*Borrelia*; Gulf Coast ticks; *Amblyomma maculatum*; Spirochetes

© 2014 Elsevier GmbH. All rights reserved

\*Corresponding author at: Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, 240 Wise Center Dr., Mississippi State, MS39762, USA. Tel.: +1 662 325 1345; fax: +1 662 325 1031. stokes@cvm.msstate.edu (A. Varela-Stokes).

## Introduction

Lyme disease is the most commonly reported vector-borne disease and perhaps the most socio-politically charged infectious disease in the United States, where the etiologic agent, *Borrelia burgdorferi* sensu stricto, is transmitted by the hard tick, *Ixodes scapularis* Say (1821) (Burgdorfer et al., 1982). Relapsing fever, the other endemic borreliosis in the United States, is most commonly caused by *B. hermsii* (Schwan and Piesman, 2002) and transmitted by soft ticks in the genus *Ornithodoros*. *Borrelia* species within the relapsing fever group have also been identified in hard ticks. For example, *B. miyamotoi*, first discovered in *I. persulcatus* ticks from Japan (Fukunaga et al., 1995), has been found in *I. ricinus* in Europe (Fraenkel et al., 2002; Richter et al., 2003), *I. scapularis* in the eastern United States (Scoles et al., 2001; Ullmann et al., 2005), and *I. pacificus* in the western U.S. (Mun et al., 2006). Interestingly, *B. miyamotoi* was recently associated with human disease in both the U.S. and Russia (Gugliotta et al., 2013; Krause et al., 2013; Platonov et al., 2011). *Borrelia lonestari*, another hard-tick relapsing fever group spirochete, was first detected in *Amblyomma americanum*, ticks common in the southeastern and south-central U.S. (Barbour et al., 1996). While this spirochete was associated with one case of human disease, “southern tick-associated rash illness,” over a decade ago (James et al., 2001), additional evidence implicating *B. lonestari* as a human disease agent is lacking (Feder et al., 2011). The Gulf Coast tick, *Amblyomma maculatum* Koch (1844) is a hard tick found primarily along the Gulf and Atlantic Coasts of the U.S., with a host range that includes rodents and ground-dwelling birds for immature stages, and large mammals, such as cattle, for adult stages (Teel et al., 2010). While *A. maculatum* is a known vector for the human spotted fever rickettsial pathogen, *Rickettsia parkeri* (Paddock et al., 2004), and the canine protozoal pathogen *Hepatozoon americanum* (Ewing and Panciera, 2003), *A. maculatum* is not known to harbor or transmit a *Borrelia* species. However, a recent study of *A. maculatum* in Arkansas reported the presence of a *Borrelia flaB* gene amplicon (*B. lonestari* and *B. burgdorferi*) in approximately 28% of *A. maculatum* collected from canines and white-tailed deer (Fryxell et al., 2012). To our knowledge, the presence of a *Borrelia* sp. in unfed *A. maculatum* has not been shown.

The objective of this study was to evaluate questing (unfed) adult *A. maculatum* for the presence of *Borrelia* species. Based on previous evidence of *B. burgdorferi* and *B. lonestari* in *A. maculatum* removed from hosts, the possibility of acquiring these spirochetes during the blood meal and the lack of evidence for *A. maculatum* as a vector for either *Borrelia* sp., we hypothesized that *B. burgdorferi* and *B. lonestari* would not be present in questing *A. maculatum*. We tested *A. maculatum* collected in Mississippi (MS) in 2009 and 2010 as part of another study (Ferrari et al., 2012), and an additional 97 samples collected in 2013, using PCR assays and amplicon sequencing.

## Materials and methods

### *Amblyomma maculatum* collection

A subset of 306 (177 in 2009 and 129 in 2010) questing adult *A. maculatum* that were collected as part of a previous study was used in the present study (Ferrari et al., 2012). An additional 97 questing *A. maculatum* that was collected in May and June of 2013 from 2

sites in northern MS and one site in central MS were also tested. The three 2013 sites were among those included in the previous study (Ferrari et al., 2012).

### DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted using an Illustra Tissue and Cells GenomicPrep Mini Spin Kit (GE Healthcare Life Sciences, Piscataway, NJ, USA) from the 306 whole-individual *A. maculatum* collected previously and from the 97 half tick bodies collected in 2013. *Amblyomma maculatum* collected in 2013 were first cleaned with 20% sodium hypochlorite (NaClO), 0.5% benzalkonium chloride, and 70% ethyl alcohol, rinsed in sterile PBS and then bisected sagittally with the half tick used for DNA extraction and the other half used as part of another study.

DNA extracts from 2009 and 2010 were used in 2 nested PCR assays to amplify approximately 330-base-pair (bp) and 650-bp regions of the *flaB* and 16S rRNA genes, respectively, for *Borrelia* spp. based on previously published assays (Barbour et al., 1996; Richter et al., 2003). Samples from 2013 were tested by *Borrelia* spp. 16S rRNA gene PCR only (Richter et al., 2003). All ticks were also tested in a PCR amplifying a portion of the tick mitochondrial 16S rRNA gene to ensure presence of amplifiable DNA (Black and Piesman, 1994; Ferrari et al., 2012). DNA extracted from cultured *B. burgdorferi* strain B31 and *B. hermsii* strain DAH were used as positive controls; molecular grade water was used as a non-template negative control. PCR products were detected by electrophoresis in a 1.5% agarose gel and visualized with ethidium bromide. PCR amplicons were purified using a DNA Clean and Concentrator Kit (Zymo Research, Irvine, CA, USA), bidirectionally sequenced (Eurofins MWG Operon, Huntsville, AL, USA), aligned using ClustalX2 (Larkin et al., 2007), and compared to available sequences in the National Center for Biotechnology Information (NCBI) database. Alignments were also used in MEGA 5 (Molecular Evolutionary Genetics Analysis) (Tamura et al., 2011) to generate phylogenetic trees.

## Results

### *Amblyomma maculatum* collection and amplicon sequence

A total of 177 adult *A. maculatum* collected in 2009, comprised of 84 females (48%) and 93 males (53%), were tested by PCR assay. Of these ticks, 78 (44%), 26 (15%), and 73 (41%) were collected from northern, central, and southern Mississippi, respectively. In 2010, 129 ticks were tested of which 81 were females (63%) and 48 were males (37%). The majority ( $n = 123$ ; 95%) of these ticks was collected from sites in northern Mississippi, while 6 (5%) were collected from central Mississippi. In 2013, 97 ticks were tested, of which 55 were females (57%) and 42 were males (43%). The majority ( $n = 95$ ; 98%) of these ticks were collected from northern Mississippi, while 2 (2%) ticks were collected from the central Mississippi site.

Two *A. maculatum* tick extracts out of 306 (0.65%) that were collected in 2009 and 2010 generated bands of appropriate size on PCR amplification of both the *flaB* and 16SrRNA gene targets. The positive tick samples were collected from the same location in central Mississippi (Byram, MS, USA). No tick extracts collected in 2013 were positive by PCR

assay of the *Borrelia* 16S rRNA gene target. 16S rRNA gene sequences amplified from the 2 positive *A. maculatum* samples were 98% identical to 12 *Borrelia* spp. including *B. turcica* (AB473539.1), *Borrelia* sp. tAG158 M (AB529425.1) and *Borrelia* sp. Tortoise 14M1 (AB473533.1). *FlaB* amplicon sequences from these same 2 tick samples were both 89% identical to *B. turcica flaB*, isolate Tortoise 7S2 and 7S1 (Accession: AB473520.1 and AB473519.1). Phylogenetic analyses using partial 16S rRNA gene sequences (Fig. 1) and *flaB* (Fig. 2) sequences from this study with selected sequences available in GenBank showed that sequences detected from the 2 ticks in this study were not closely related to Lyme disease group or relapsing fever *Borrelia* group. Rather, they were more closely related to reptile-associated borreliae. Two additional tick extracts demonstrated *flaB* amplicons on PCR that shared 100% identity to multiple strains of *B. hermsii* including DAH. A *flaB* amplicon from a third sample shared 100% identity to *B. burgdorferi* sensu stricto including strain N40 and 99% identity to *B. burgdorferi* strain B31. However, these 3 tick extracts were negative using the 16S rRNA gene target and on subsequent PCR assays to reamplify the *flaB* gene. Given that we could not amplify a 16S rRNA gene product from these samples and both *B. hermsii* DAH and *B. burgdorferi* B31 were included as PCR controls, we suspect that the *flaB* PCR results were false-positive and likely a result of PCR contamination.

GenBank accession numbers for *Borrelia* sp. partial 16S rRNA gene sequences are KF395228 and KF395229. Accession numbers for *Borrelia* sp. partial *flaB* gene sequences are KF395230 and KF395231.

## Discussion

Of *A. maculatum* extracts tested in this study, 2 out of 403 (0.5%) of the extracts were positive by PCR for *Borrelia* species. Both ticks were collected from the same central Mississippi site (Byram, MS, USA) during 2009 and 2010. Sequences of the 16S rRNA gene amplicons from the 2 samples were 100% identical to each other; *flaB* amplicon sequences for the 2 samples were also identical to each other. The identification of a *Borrelia* sp. in unfed *A. maculatum* is intriguing considering *Borrelia* spp. have been detected by PCR from only fed *A. maculatum*, and were limited to *B. lonestari* and *B. burgdorferi* (Fryxell et al., 2012). The absence of *B. burgdorferi* and *B. lonestari* amplicons in over 400 questing adult *A. maculatum* tested here suggests that the spirochetes detected previously in *A. maculatum* attached to canids and deer may have been acquired from the blood meal (Fryxell et al., 2012). Alternatively, the geographic areas sampled here differed from the previous report and there may be differences in infection rates with *Borrelia* spp. between *A. maculatum* in these areas.

Phylogenetic analyses have revealed that spirochetes in the genus *Borrelia* fall into 3 groups (Parola and Raoult, 2001; Takano et al., 2010, 2011). The Lyme disease group of *Borrelia* spp. contains the most common tick-borne disease agents in the northern Hemisphere, including *B. burgdorferi* sensu stricto, *B. afzelii*, and *B. garinii*. The relapsing fever group of *Borrelia* spp. includes *B. hermsii*, *B. duttonii*, and *B. turcicatae*, all transmitted by soft ticks (*Ornithodoros* spp.). There is also a sister group of hard tick-associated relapsing fever spirochetes, including the species *B. miyamotoi* and *B. lonestari*. The third recently proposed

group of borreliae consists of reptile-associated *Borrelia* spp. found in hard ticks of the genera *Amblyomma* and *Hyalomma* (Güner et al., 2004; Takano et al., 2011). Phylogenetic analyses of the *flaB* and 16S rRNA partial gene sequences identified in this study placed these *A. maculatum* associated *Borrelia* sequences in a clade with or as a sister group to the reptile-associated borreliae, and distinct from the relapsing fever group and Lyme disease groups. These results are consistent with the GenBank query results which showed closest *flaB* sequence identity (89%) to the *B. turcica* isolates Tortoise 7S1 and 7S2 from the hard tick, *Hyalomma aegyptium*, removed from a tortoise in Turkey (Güner et al., 2004), and closest 16S rRNA gene sequence identity (98%) to species including *B. turcica*. That the sequences identified here are closest to reptile-associated *Borrelia* species is curious considering *A. maculatum* are not known to feed on reptiles, and opens additional questions.

In this study, both 16S rRNA and *flaB* gene sequences amplified from *A. maculatum*-associated *Borrelia* sp. were unique compared to available *Borrelia* spp. sequences in the NCBI database. The presence of this previously uncharacterized *Borrelia* sp. and absence of other borreliae suggests that a distinct *Borrelia* sp. is present and supports absence of *B. burgdorferi* and *B. lonestari* in unfed *A. maculatum* tested in Mississippi. While infected *A. maculatum* may be rare, the existence of an additional *Borrelia* sp. in a southern tick species known to bite humans (Goddard, 2002) is significant. Additional studies that include continued testing to determine prevalence in *A. maculatum* throughout their range and more in depth characterization of the *Borrelia* species are warranted.


## Acknowledgments

The authors would like to thank Jerome Goddard and Christopher Paddock for intellectual contributions to this study and the Mississippi State University College of Veterinary Medicine for funding support. It is specifically administered through the Office of Research and Graduate Studies MSU-CVM.

## References


- Barbour AG, Maupin GO, Teltow GJ, Carter CJ, Piesman J. Identification of an uncultivable *Borrelia* species in the hard tick *Amblyomma americanum*: possible agent of a Lyme disease-like illness. *J. Infect. Dis.* 1996; 173:403–409. [PubMed: 8568302]
- Black WC, Piesman J. Phylogeny of hard- and soft-tick taxa (Acari: Ixodida) based on mitochondrial 16S rDNA sequences. *Proc. Natl. Acad. Sci. U. S. A.* 1994; 91:10034–10038. [PubMed: 7937832]
- Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. Lyme disease - a tick-borne spirochetosis? *Science (New York, NY)*. 1982; 216:1317–1319.
- Ewing SA, Panciera RJ. American canine hepatozoonosis. *Clin. Microbiol. Rev.* 2003; 16:688–697. [PubMed: 14557294]
- Feder HM Jr, Hoss DM, Zemel L, Telford SR 3rd, Dias F, Wormser GP. Southern tick-associated rash illness (STARI) in the North: STARI following a tick bite in Long Island, New York. *Clin. Infect. Dis.* 2011; 53(10):e142–e146. [PubMed: 21940418]
- Ferrari FA, Goddard J, Paddock CD, Varela-Stokes AS. *Rickettsia parkeri* and *Candidatus Rickettsia andeanae* in Gulf Coast ticks, Mississippi, USA. *Emerg. Infect. Dis.* 2012; 18:1705–1707. [PubMed: 23018026]
- Fraenkel CJ, Garpmo U, Berglund J. Determination of novel *Borrelia* genospecies in Swedish *Ixodes ricinus* ticks. *J. Clin. Microbiol.* 2002; 40:3308–3312. [PubMed: 12202571]
- Fryxell RT, Steelman CD, Szalanski AL, Kvamme KL, Billingsley PM, Williamson PC. Survey of Borreliae in ticks, canines, and white-tailed deer from Arkansas, USA. *Parasit. Vectors.* 2012; 5:139. [PubMed: 22781030]

- Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, Nakao M. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov. isolated from the ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int. J. Syst. Bacteriol.* 1995; 45:804–810. [PubMed: 7547303]
- Goddard J. A ten-year study of tick biting in Mississippi: implications for human disease transmission. *J. Agromed.* 2002; 8:25–32.
- Gugliotta JL, Goethert HK, Berardi VP, Telford SR. Meningoencephalitis from *Borrelia miyamotoi* in an immunocompromised patient. *N. Engl. J. Med.* 2013; 368:240–245. [PubMed: 23323900]
- Güner ES, Watanabe M, Hashimoto N, Kadosaka T, Kawamura Y, Ezaki T, Kawabata H, Imai Y, Kaneda K, Masuzawa T. *Borrelia turcica* sp. nov. isolated from the hard tick *Hyalomma aegyptium* in Turkey. *Int. J. Syst. Evol. Microbiol.* 2004; 54:1649–1652. [PubMed: 15388724]
- James AM, Liveris D, Wormser GP, Schwartz I, Montecalvo MA, Johnson BJ. *Borrelia lonestari* infection after a bite by an *Amblyomma americanum* tick. *J. Infect. Dis.* 2001; 183:1810–1814. [PubMed: 11372036]
- Krause PJ, Narasimhan S, Wormser GP, Rollend L, Fikrig E, Lepore T, Barbour A, Fish D. Human *Borrelia miyamotoi* infection in the United States. *N. Engl. J. Med.* 2013; 368:291–293. [PubMed: 23323920]
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. Clustal W and Clustal X version 2.0. *Bioinformatics.* 2007; 23:2947–2948. [PubMed: 17846036]
- Mun J, Eisen RJ, Eisen L, Lane RS. Detection of a *Borrelia miyamotoi* sensu lato relapsing-fever group spirochete from *Ixodes pacificus* in California. *J. Med. Entomol.* 2006; 43:120–123. [PubMed: 16506458]
- Paddock CD, Sumner JW, Comer JA, Zaki SR, Goldsmith CS, Goddard J, McLellan SLF, Tamminga CL, Ohl CA. *Rickettsia parkeri*: a newly recognized cause of spotted fever rickettsiosis in the United States. *Clin. Infect. Dis.* 2004; 38:805–811. [PubMed: 14999622]
- Parola P, Raoult D. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. *Clin. Infect. Dis.* 2001; 32:897–928. [PubMed: 11247714]
- Platonov AE, Karan LS, Kolyasnikova NM, Makhneva NA, Toporkova MG, Maleev VV, Fish D, Krause PJ. Humans infected with relapsing fever spirochete *Borrelia miyamotoi*, Russia. *Emerg. Infect. Dis.* 2011; 17:1816–1823. [PubMed: 22000350]
- Richter D, Schlee DB, Matuschka FR. Relapsing fever-like spirochetes infecting European vector tick of Lyme disease agent. *Emerg. Infect. Dis.* 2003; 9:697–701. [PubMed: 12781009]
- Schwan TG, Piesman J. Vector interactions and molecular adaptations of Lyme disease and relapsing fever spirochetes associated with transmission by ticks. *Emerg. Infect. Dis.* 2002; 8:115–121. [PubMed: 11897061]
- Scoles GA, Papero M, Beati L, Fish D. A relapsing fever group spirochete transmitted by *Ixodes scapularis* ticks. *Vector Borne Zoonotic Dis.* 2001; 1:21–34. [PubMed: 12653133]
- Takano A, Fujita H, Kadosaka T, Konnai S, Tajima T, Watanabe H, Ohnishi M, Kawabata H. Characterization of reptile-associated *Borrelia* sp. in the vector tick *Amblyomma geoemydae*, and its association with Lyme disease and relapsing fever *Borrelia* spp. *Environ. Microbiol. Rep.* 2011; 3:632–637. [PubMed: 23761345]
- Takano A, Goka K, Une Y, Shimada Y, Fujita H, Shiino T, Watanabe H, Kawabata H. Isolation and characterization of a novel *Borrelia* group of tick-borne borreliae from imported reptiles and their associated ticks. *Environ. Microbiol.* 2010; 12:134–146. [PubMed: 19758349]
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 2011; 28:2731–2739. [PubMed: 21546353]
- Teel PD, Ketchum HR, Mock DE, Wright RE, Strey OF. The Gulf Coast tick: a review of the life history, ecology, distribution, and emergence as an arthropod of medical and veterinary importance. *J. Med. Entomol.* 2010; 47:707–722. [PubMed: 20939363]
- Ullmann AJ, Gabitzsch ES, Schulze TL, Zeidner NS, Piesman J. Three multiplex assays for detection of *Borrelia burgdorferi* sensu lato and *Borrelia miyamotoi* sensu lato in field-collected *Ixodes* nymphs in North America. *J. Med. Entomol.* 2005; 42:1057–1062. [PubMed: 16465748]



**Fig. 1.**

Unrooted maximum likelihood phylogenetic tree based on 568-bp partial 16S rRNA gene sequences from *Borrelia* spp. F3S6 and F17S6. Numbers at the branch notes denote bootstrap values as a percentage of 1000 replications. GenBank accession numbers for sequences used in tree construction are included for each species.



**Fig. 2.** Unrooted maximum likelihood phylogenetic tree based on 267 and 226 bp partial *flaB* gene sequences from *Borrelia* spp. F3S6 and F17S6, respectively. Numbers at the branch notes denote bootstrap values as a percentage of 1000 replications. GenBank accession numbers for sequences used in tree construction are included for each species. The scale bar demonstrates degree of divergence and corresponds to a distance of 0.05.