

# Draft Genome Sequence of *Enterobacter aerogenes*, a DDE-Degrading and Plant Growth-Promoting Strain Isolated from *Cucurbita pepo*

Nele Eevers,<sup>a</sup> Jonathan D. Van Hamme,<sup>b</sup> Eric M. Bottos,<sup>b</sup> Nele Weyens,<sup>a</sup> Jaco Vangronsveld<sup>a</sup>

Centre for Environmental Sciences, Hasselt University, Agoralaan Building D, Diepenbeek, Belgium<sup>a</sup>; Department of Biological Sciences, Thompson Rivers University, Kamloops, British Columbia, Canada<sup>b</sup>

**We report here the draft genome of *Enterobacter aerogenes*, a Gram-negative bacterium of the *Enterobacteriaceae* isolated from *Cucurbita pepo* root tissue. This bacterium shows 2,2-bis(*p*-chlorophenyl)-1,1-dichloroethylene (DDE)-degrading potential and plant growth-promoting capacity. An analysis of its 4.5-Mb draft genome will enhance the understanding of DDE degradation pathways and phytoremediation applications for DDE-contaminated soils.**

Received 3 March 2015 Accepted 9 March 2015 Published 16 April 2015

**Citation** Eevers N, Van Hamme JD, Bottos EM, Weyens N, Vangronsveld J. 2015. Draft genome sequence of *Enterobacter aerogenes*, a DDE-degrading and plant growth-promoting strain isolated from *Cucurbita pepo*. *Genome Announc* 3(2):e00317-15. doi:10.1128/genomeA.00317-15.

**Copyright** © 2015 Eevers et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jaco Vangronsveld, jaco.vangronsveld@uhasselt.be.

DDT [2,2-bis(*p*-chlorophenyl)-1,1,1-trichloroethane] (1) is a pesticide that has been used in agriculture and gardening since 1943 (2). In soils, DDT degrades to 2,2-bis(*p*-chlorophenyl)-1,1-dichloroethylene (DDE). Both DDT and DDE are classified as persistent organic pollutants (3) and are of concern to human and animal health because of their toxicity and hormone-disrupting properties (4). Endophyte-enhanced phytoremediation using *Cucurbita pepo*, a DDE-accumulating plant (5), is being explored for a role in the remediation of DDE-contaminated soils.

A DDE-degrading bacterial strain was isolated from the roots of *C. pepo* plants exposed to 100 mg·liter<sup>-1</sup> DDE. Identified as *Enterobacter aerogenes* by partial 16S rRNA gene sequencing and phenotypic profiling, the closest related 16S rRNA sequence (87%) was from *E. aerogenes* strain EA1509E (GenBank accession no. FO203355.1) (6).

Genomic DNA was isolated using a DNeasy blood and tissue kit (Qiagen, Venlo, The Netherlands). An Ion Torrent PGM (Life Technologies, Inc., Carlsbad, CA) was used for sequencing. The DNA was digested, and sequencing adapters were ligated using an Ion Xpress Plus fragment library kit (Life Technologies, Inc., Burlington, Ontario, Canada), according to the manufacturer's instructions. Adapter-ligated DNA was size selected to a target length of 480 bp on a 2% E-Gel SizeSelect agarose gel, and Agencourt AMPure XP beads (Beckman Coulter, Mississauga, Ontario, Canada) were used for the purification steps. An Ion library quantitation kit was used to calculate the library dilution factor prior to amplification and enrichment with an Ion PGM Template OT2 400 kit on an Ion OneTouch 2 system. An Ion Sphere quality control kit was used to quantify the percent enriched Ion Sphere Particles prior to sequencing with an Ion PGM 400 sequencing kit.

In total, 1.67 million reads (mean length, 298 bases) generated 499 Mb of data in Torrent Suite 4.2.1. These were assembled using MIRA 3.9.9 (7) into 69 contigs > 500 bp, giving a consensus length of 4,474,344 bp at 51 × coverage (largest contig, 602,746 bp; *N*<sub>50</sub>, 139,424 bp). Mauve (8) was used to order contigs using the genome of *E. aerogenes* EA1509E (GenBank accession no.

FO203355.1) (6), and the annotation was completed using the PGAP (NCBI) pipeline (9). The genome of *E. aerogenes* consists of a single circular chromosome (53.8% G+C content), which includes 4,191 coding genes that were arranged into 309 pathways using Pathway Tools (10, 11), 260 pseudogenes, 40 rRNAs (5S, 16S, 23S), 77 tRNAs, and 7 noncoding RNAs (ncRNAs).

This *E. aerogenes* strain showed increased growth when exposed to DDE in comparison to that under control conditions. Analyses of the draft genome showed the presence of dioxygenases and hydrolases that have been associated with DDE degradation (12–15). Genes for plant growth-promoting capacities, 1-aminocyclopropane-1-carboxylate deaminase activity, siderophore production, auxin biosynthesis, and phosphorous solubilization, are present, corroborating the results from phenotypic tests. These characteristics make *E. aerogenes* a promising strain for DDE phytoremediation.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/Genbank under the accession no. [JXTQ000000000](https://www.ncbi.nlm.nih.gov/nuccore/JXTQ000000000). The version described in this paper is version [JXTQ010000000](https://www.ncbi.nlm.nih.gov/nuccore/JXTQ010000000).

## ACKNOWLEDGMENTS

This work was supported by Ph.D. grant 121243 from the Agency for Innovation by Science and Technology (IWT) and a postdoc grant to N.W. from the Fund for Scientific Research-Flanders, Belgium (FWO Vlaanderen) and the Methusalem project 08M03VGRJ.

## REFERENCES

1. Thomas JE, Ou LT, All-Agely A. 2008. DDE remediation and degradation. *Rev Environ Contam Toxicol* 194:55–70. [http://dx.doi.org/10.1007/978-0-387-74816-0\\_3](http://dx.doi.org/10.1007/978-0-387-74816-0_3).
2. Turusov V, Rakitsky V, Tomatis L. 2002. Dichlorodiphenyltrichloroethane (DDT): ubiquity, persistence, and risks. *Environ Health Perspect* 110:125–128.
3. White JC. 2010. Inheritance of *p,p'*-DDE phytoextraction ability in hybridized *Cucurbita pepo* cultivars. *Environ Sci Technol* 44:5165–5169. <http://dx.doi.org/10.1021/es100706t>.
4. Colles A, Koppen G. 2007. DDE serumwaarden in landelijke regio's en de

- Albertkanaalzone boven het Vlaamse referentiegemiddelde: enquête naar de oorzaak. De Biomonitor, digitale nieuwsbrief van het medisch milieukundig netwerk (VITO).
5. White JC, Parrish ZD, Isleyen M, Gent MPN, Iannucci-Berger W, Eitzer BD, Mattina MJI. 2005. Uptake of weathered *p,p'*-DDE by plant species effective at accumulating soil elements *Microchem J* 81:148–155. <http://dx.doi.org/10.1016/j.microc.2005.01.010>.
  6. Diene SM, Merhej V, Henry M, El Filali A, Roux V, Robert C, Azza S, Gavory F, Barbe V, La Scola B, Raoult D, Rolain J-. 2013. The rhizome of the multidrug-resistant *Enterobacter aerogenes* genome reveals how new “killer bugs” are created because of a sympatric lifestyle. *Mol Biol Evol* 30:369–383. <http://dx.doi.org/10.1093/molbev/mss236>.
  7. Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* Computer science and biology. Proceedings of the German Conference on Bioinformatics, GCB '99. GCB, Hannover, Germany.
  8. Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of draft genomes using the Mauve aligner. *Bioinformatics* 25:2071–2073. <http://dx.doi.org/10.1093/bioinformatics/btp356>.
  9. Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of Standard Operating Procedures (SOPs) for (meta)genomic annotation. *12:137–141*. <http://dx.doi.org/10.1089/omi.2008.0017>.
  10. Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, Holland TA, Keseler IM, Kothari A, Kubo A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P, Weaver DS, Weerasinghe D, Zhang P, Karp PD. 2014. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res* 42:D459–D471. <http://dx.doi.org/10.1093/nar/gkt1103>.
  11. Karp PD, Paley S, Romero P. 2002. The pathway tools software. *Bioinformatics* 18(Suppl 1):S225–S232. [http://dx.doi.org/10.1093/bioinformatics/18.suppl\\_1.S225](http://dx.doi.org/10.1093/bioinformatics/18.suppl_1.S225).
  12. Quensen JF, Mueller SA, Jain MK, Tiedje JM. 1998. Reductive dechlorination of DDE to DDMU in marine sediment microcosms. *Science* 280:722–724. <http://dx.doi.org/10.1126/science.280.5364.722>.
  13. Hay AG, Focht DD. 1998. Cometabolism of 1,1-dichloro-2,2-bis(4-chlorophenyl)ethylene by *Pseudomonas acidovorans* M3GY grown on biphenyl. *Appl Environ Microbiol* 64:2141–2146.
  14. Nadeau LJ, Menn FM, Breen A, Saylor GS. 1994. Aerobic degradation of 1,1,1-trichloro-2,2-bis(4-chlorophenyl)ethane (DDT) by *Alcaligenes eutrophus* A5. *Appl Environ Microbiol* 60:51–55.
  15. Hay AG, Focht DD. 2000. Transformation of 1,1-dichloro-2,2-(4-chlorophenyl)ethane (DDD) by *Ralstonia eutropha* strain A5. *FEMS Microbiol Ecol* 31:249–253. <http://dx.doi.org/10.1111/j.1574-6941.2000.tb00690.x>.