

Evaluation of Emerging *Fusarium* mycotoxins beauvericin, Enniatins, Fusaproliferin and Moniliformin in Domestic Rice in Iran

Firouzeh Nazari^{a,b}, Michael Sulyok^b, Farzad Kobarfard^c, Hassan Yazdanpanah^{d*} and Rudolf Krška^b

^aFood and Drug Organization, Iran University of Medical Sciences, Tehran, Iran. ^bCenter for Analytical Chemistry, Department for Agrobiotechnology (IFA-Tulln), University of Natural Resources and Life Sciences Vienna (BOKU), Konrad-Lorenz-Str. 20, A-3430 Tulln, Austria. ^cDepartment of Medicinal Chemistry, School of Pharmacy, Phytochemistry Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^dDepartment of Toxicology and Pharmacology, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

The occurrence of emerging *Fusarium* mycotoxins beauvericin (BEA), enniatins (ENNs) (A, A1, B, B1), Fusaproliferin and moniliformin was evaluated by a liquid chromatography/electrospray ionization-tandem mass spectrometric (LC/ESI-MS/MS) technique in 65 domestic rice samples produced in Gilan and Mazandaran Provinces in Iran. The results showed that 46% of the samples were contaminated with at least one of the emerging mycotoxins. BEA was the most prevalent mycotoxin, which was found in 26 out of 65 rice samples at the concentrations up to 0.47 µg/Kg. Enniatin A1 which was the only member of ENNs was detected in the samples, occurred in 7.7% of samples with an average level of 0.06 µg/Kg. No detectable level of Fusaproliferin and moniliformin was found. This is the first report concerning the contamination of Iranian domestic rice samples with the emerging *Fusarium* mycotoxins.

Keywords: Beauvericin; Enniatins; Fusaproliferin; Moniliformin; Rice; Iran.

Introduction

Mycotoxins are natural toxic secondary metabolites produced by various filamentous fungi *Aspergillus*, *Penicillium* and *Fusarium* species in a broad range of foodstuff (1). The existence of mycotoxins in food commodities is of major concern for public health because of their toxic properties including carcinogenicity, genotoxicity, immunotoxicity, mutagenicity and reproductive and developmental toxicity (2). *Fusarium* species produce the most common mycotoxins such as trichothecenes, fumonisins, moniliformin and zearalenone, as

well as emerging mycotoxins like beauvericin (BEA) enniatins (ENNs), fusaproliferin (FUS) and moniliformin (MON) (3). There is only limited data available about these metabolites. This is not only due to their late recognition but especially because of the late understanding of their role as mycotoxins (4). Because of presence of high concentrations of emerging *Fusarium* mycotoxins in many food products, there is a great interest on these toxins. In addition, the European Food Safety Authority (EFSA) is collecting occurrence data for risk assessment at the moment.

BEA and ENNs are cyclic hexa depsipeptide mycotoxins (5) produced by different *Fusarium* species such as *F. avenaceum*, *F. subglutinans*, *F. oxysporum*, *F. proliferatum*, *F. poae* and *F.*

* Corresponding author:

E-mail: yazdanpanah@sbm.ac.ir




Figure 1. The chemical structures of beavericin (BEA), enniatins (A, A1, B and B1), moniliformin and fusaproliferin.

tricinctum (6-8). *Fusaproliferinis* are produced by *F. antophilum*, *F. begoniae*, *F. bulbicola*, *F. circinatum*, *F. concentricum*, *F. succisae*, and *F. udum* (9). MON is sodium or potassium salt of 3-hydroxycyclobut-3-ene-1, 2-dione (10) and produced by several species of *Fusarium* including *F. avenaceum*, *F. tricinctum*, and *F. proliferatum* (11, 12). Chemical structures of BEA, enniatins, fusaproliferin and moniliformin have been shown in Figure 1.

All of them are bioactive compounds and elicit a wide range of different toxicological effects. BEA has shown genotoxic and cytotoxic effects in human lymphocytes and animal species, respectively (13, 14). It induces chromosomal aberrations, sister-chromatid exchanges and micronuclei formation (13). It also induces apoptosis (15, 16), mitochondrial dysfunction (17) and erythrocyte cell membrane scrambling (18). Finally, BEA has shown to be a specific cholesterol acyltransferase inhibitor (19). ENNs possess a wide range of biological activities. They have shown cytotoxic and apoptotic activity (20, 21). They are cationophoric compounds and specific inhibitors of acyl-coenzyme A cholesterol acyltransferase (ACAT) (22). The most well-known ENNs reported as natural contaminants are ENN A, A1, B and B1. Several other analogues (B2, B3, B4, D, E, F and G) have also been identified (23). ENNs have been found to have synergistic toxicity effects (15). ENN B is the most bioactive ENNs among B1, A and A1 and showed mitochondrial toxicity

in isolated rat liver mitochondria (17).

FUS is a sesterterpene mycotoxin with a variety of toxic properties. This compound has been found to be toxic on *Artemia salina* (9, 24), human B lymphocytes (25), brine shrimp larvae and insect cells (26), and Caco-2 cells (27). It also has teratogenic and pathogenic effects (25).

MON is a potent inhibitor of the pyruvate dehydrogenase complex (28). MON induces acidosis and causes muscular weakness (29-30), cardiotoxicity (31), immunosuppression (32) and intestinal problems (33).

The incidence of emerging *Fusarium* mycotoxins has been reported in different foods such as maize (34, 35), wheat (36, 37), rice (38), grain-based products (39), cereal-based foodstuffs (40), breakfast, infant cereals (41), and barley (36). Although the majority of the studies are related to cereal and cereal products samples, there are some studies indicating the emergence of mycotoxin contamination in other food commodities. Abia *et al.* (2013) found BEA and ENNs in groundnuts, soybean, and groundnut soup in Cameroon (42). Sebastià *et al.* (2012) detected BEA and ENNs in tiger-nuts collected in Spain (43). Ezekiel *et al.* (2012) found BEA in peanut cake at concentrations up to 3.36 mg/Kg (44).

Rice (*Oryza sativa* L.) is one of the most important food crops worldwide especially for Iranian population. It can be ideal substrates for the growth of mycotoxin producing fungi during inappropriate storage conditions, moisture and

insect infestation (45).

Too little information is available regarding the incidence of BEA and EENS in rice samples (38, 46). Therefore, the aim of this study was to evaluate the presence of BEA, ENNs, MON and FUS in Iranian domestic rice using a highly sensitive and selective LC-MS/MS multi-mycotoxin method.

Experimental

Reagents and equipment

Methanol (MeOH) and acetonitrile (ACN) (both LC gradient grade) were purchased from Merck, Darmstadt, Germany and VWR, International Leuven, respectively; ammonium acetate (MS grade) and glacial acetic acid (p.a.) were obtained from Sigma Aldrich (Vienna, Austria). Water was purified successively by reverse osmosis and a Milli-Q plus system from Millipore (Molsheim, France).

Mycotoxin standards were dissolved in acetonitrile (ACN) predominantly to yield a multi-analyte working solution. This stock solution was subsequently used to prepare different working solutions for calibration. All standard solutions were stored at -20 °C and were brought to room temperature before use. Extraction (ACN/water/glacial acetic acid 79:20:1, v/v/v) and dilution (ACN/ water/ acetic acid 20:79:1, v/v/v) solvents were freshly prepared and kept at room temperature before use. Two eluents that contained 5 mM ammonium acetate each were prepared using MeOH/water/ glacial acetic acid (10:89:1, v/v/v) (eluent A) and (97:2:1, v/v/v) (eluent B).

Sampling

A total of 65 rice samples were collected randomly from local markets in Tehran, originally from Gilan Province and Mazandaran Province in northern Iran during April 2010 to April 2011. The minimum amount of samples was 1000 g. All samples were finely grounded and homogenized by milling (Romer Series II® Mill) and subsamples were stored at -20 °C until analysis.

Methodology

Rice samples were analysed for the incidence

and quantification of emerging *Fusarium* mycotoxins by LC-MS/MS according to Sulyok *et al.* (2006) (47).

Extraction

Five grams of milled rice samples were extracted with 20 mL extraction solvent for 90 min at 180 rpm using a GFL 3017 rotary shaker (GFL 3017, Burgwedel, Germany). Aliquots of 500 µL extracts were transferred into glass vials containing an equal volume of the dilution solvent. After appropriate mixing, 5 µL of the diluted extracts were injected into the LC-MS/MS system.

Chromatographic and mass spectrometric parameters

Detection and quantification were performed using a QTRAP 5500 LC-MS/MS System (AB SCIEX, Foster City, CA) equipped with Turbo Ion Spray electrospray ionization (ESI) source and a 1290 Series HPLC System (Agilent Technologies 1290 Infinity).

Chromatographic separation was performed at 25 °C on a Gemini® C18 column, 150×4.6-mm *i.d.*, 5-µm particle size, equipped with a C18 4×3-mm-*i.d.* security guard cartridge (all from Phenomenex, Torrance, CA, US). After an initial time of 2 min at 100% eluent A, the proportion of eluent B was increased linearly to 50% within 2-5 min and to 100% within 5-14 min, followed by a holding-time of 4 min at 100% eluent B and 2.5 min column re-equilibration at 100% eluent A pumped at a flow rate of 1 mL/min.

The declustering potential (DP), entrance potential collision energy (CE) and collision cell exit potential (CXP) were optimized for each toxin (Table 1). ESI-MS/MS was performed in the scheduled multiple reaction monitoring (sMRM) mode both in positive and negative polarities in two separated chromatography runs per sample by scanning two fragmentation reaction per analyte with the following setting: source temperature 550 °C; curtain gas 30 psi; ion source gas 1 (sheath gas) 80 psi, ion source gas 2 (drying gas) 80 psi, ion spray voltage -4500 V, and +5500 V, respectively, collision gas (nitrogen) medium. The MRM detection windows were 54 and 96 s in the positive and negative ionization mode, respectively, and the

Table 1. Optimized ESI-MS/MS parameters.

Analyte	Precursor ion	Product ion	DP	EP	CE	CXP	Rt (min)
Beauvericin	801.5[M+NH ₄] ⁺	243.2	116	10	47	12	14.3
	806.5[M+Na] ⁺	384.4	191	10	73	10	
Enniatin A	699.4 [M+NH ₄] ⁺	210.1/228	106	10	43/47	12/18	14.8
Enniatin A1	685 [M+NH ₄] ⁺	210.1/228.2	96	10	41/49	8/20	14.5
Enniatin B	657.5 [M+NH ₄] ⁺	196.3/214.1	81	10	45/47	18/18	14
Enniatin B1	671.4 [M+NH ₄] ⁺	196/210	111	10	43/41	12/12	14.2
Fusaproliferin	445.2[M+H] ⁺	368.1/350.1	96	10	25/29	10/10	13.3
Moniliformin	96.9 [M-H] ⁻	41.2	-100	-10	-24	-5	3.2

DP: declustering potential, EP: entrance potential, CE: collision energy, collision CXP: cell exit potential, Rt: retention time

cycle time was set to 1 s.

Mycotoxins were quantified using external calibration (1/x weighted) and levels were later adjusted based on recoveries determined by spiking four samples. Limits of detection (LOD) and quantification (LOQ) were estimated at the lowest concentration in spiked samples corresponding to a signal-to-noise ratio (S/N) of 3:1 and 10:1, respectively 10:1.

Spiking experiment

To determine the apparent recovery, spiking experiments were performed. In this regard, four blank (least contaminated) samples (each 0.25 g) were spiked with multi-analyte standard, thoroughly mixed and stored at room temperature in the dark cupboard about four hours to establish equilibration between the analytes and the matrix. Subsequently, one mL of extraction solvent was added and placed on a rotary shaker for 90 min at 180 rpm. After diluting 300 µL of extracts with an equal volume of the dilution solvent and mixing it for 30 s by vortex, 5 µL of the diluted extracts were injected into the LC-MS/MS system.

Results and Discussion

MS/MS parameters such as precursors and product ion, declustering potentials, collision energies and cell exit potentials were optimized for all analytes. All of the compounds exhibited precursor ions and product ions with reasonably high signal intensities in positive mode with the exception of MON which gave no signal in the positive ion mode. For each analyte, the polarity

giving the most abundant product ions was selected; a second product ion was monitored for confirmation of the identity (with the exception of MON, which showed only one product ion). Table 1 summarizes the parameters of the optimized MRM transitions.

As shown in Table 1, among the seven emerging *Fusarium* mycotoxins, five mycotoxins (ENNB, ENNB1, ENNA1, ENNA and BEA) appeared as their ammonium adduct in mass spectra. Examined the chemical structures of these mycotoxins reveals that all these mycotoxins which appear as ammonium adduct, contain a 1,7,13-trioxa-4,10,16-triazacyclooctadecane ring as the backbone of their molecules. It could then be concluded that the presence of 18 membered ring of trioxa triazacyclooctadecane is responsible for the formation of ammonium adducts in mass spectra.

The recoveries of all mycotoxins ranged from 94 % to 111% (Table 2). It was observed that correlation coefficients (r²) were > 0.99 for all the analytes.

Occurrence of emerging Fusarium mycotoxins in rice samples

Despite the great concern about the toxicity of these toxins and their natural incidence in different food stuff, no maximum limits and tolerable daily intakes (EDI) have been established for them.

In this study, the occurrence of BEA, ENNs, MON and FUS was evaluated in rice samples and the results have been shown in Table 3. The results indicated that 46% of samples were contaminated with at least one of the emerging

Table 2. Method Performance Characteristics for Beauvericin, and Enniatins, moniliformin and fusaproliferin rice samples.

Analyte	LOD ^a (µg/Kg)	LOQ ^b (µg/Kg)	Recovery (%) ^c
Beauvericin	0.02	0.06	101.9 ± 5.1
Enniatin A	0.02	0.06	98.1 ± 3.8
Enniatin A1	0.02	0.06	94.1 ± 2.6
Enniatin B	0.02	0.06	101.8 ± 5.1
Enniatin B1	0.02	0.06	100.5 ± 3.9
Moniliformin	0.08	0.25	105.3 ± 2.1
Fusaproliferin	0.02	0.06	110.8 ± 23.1

^aLOD, (S/N = 3:1) expressed as µg/Kg sample

^bLOQ (S/N = 10:1) expressed as µg/Kg sample

^cMean and standard deviation of four samples

mycotoxins.

Beauvericin

The results showed that the most prevalent mycotoxin was BEA, which was found in 26 out of 65 rice samples at the concentrations up to 0.47 µg/Kg.

There are limited data available on the occurrence of emerging mycotoxins in rice. Meca *et al.* (2010) found BEA contamination in rice samples from Spain at a level of 11.78 mg/Kg (46). In Morocco, Sifou *et al.* (2011) found that BEA was present in 75.7% of rice samples, and contamination level up to 26.3 mg/Kg was observed (38). Nazari *et al.* found that 88% of imported rice samples to Iran were contaminated with low level of BEA (range: 0.01 to 1.56 µg/Kg) (unpublished data). Natural occurrence of BEA in other food commodities has been reported in different countries. Logrieco *et al.* (2002) reported that BEA was found in all 13 wheat samples from Finland in the concentration range of 0.64 to 3.5 µg/g (6). In another study in

Finland, BEA was detected in 95% of 38 grain samples (39). Mahnine *et al.* (2011) reported the presence of BEA in 5.8% of 68 cereal derivatives products (breakfast cereals and infant cereals) from Morocco with a maximum level of 10.6 mg/Kg (41).

Enniatins

In our study, ENNA1 was the only ENN among ENNB, ENNB1, ENNA1, and EENA, which was found in the samples. ENNA1 contamination occurred in 7.7% of rice samples with an average level of 0.06 µg/Kg. Sifou *et al.* (2011) reported that total ENNs was found in 50% of rice samples (38). ENNB was the most prevalent ENN which found in 30% of the samples, while ENNB1, ENNA and ENNA1 were found in 24.6% (ranged: 4.4 to 26.2 mg/Kg), 22.8% (ranged: 8.4 to 119.5 mg/Kg) and 5.7% (ranged: 56.2 to 448.7 mg/Kg) of total samples, respectively (38). Nazari *et al.* reported that ENNs (ENNB, ENNB1, ENNB2, and ENNA1) were detected in 9% of imported rice

Table 3. Occurrence of emerging *Fusarium* mycotoxins in Iranian rice samples.

Analyte	Incidence (%)	Mean(µg/Kg)	Max(µg/Kg)	Daily intake(ng/Kgbw/day)	
				Mean	Max
Beauvericin	26 (40)	0.11	0.47	0.2	0.86
Enniatin A1	5 (7.7%)	0.06	0.08	0.11	0.15
Enniatin A	nd	-	-	-	-
Enniatin B	nd	-	-	-	-
Enniatin B1	nd	-	-	-	-
Moniliformin	nd	-	-	-	-
Fusaproliferin	nd	-	-	-	-

nd: not detected

samples to Iran with contamination frequencies of 3%, 3%, 1%, and 2% for ENNB, ENNB1, ENNB2, and ENNA1, respectively (unpublished data). Meca *et al.* (2010) detected the occurrence of ENA1 (814.42 mg/Kg) and ENB (7.95 mg/Kg) in one sample of rice from Spain (46).

Fusaproliferin

As for FUS, no detectable amount of FUS was identified in the Iranian rice samples. The low incidence of FUS in rice has been previously reported by others (38, 41, and 46). In Morocco, 4.3% of rice samples were contaminated with FUS at the concentration range of 0.2 to 19.6 mg/Kg (38). In another study in Morocco, Mahnine *et al.* (2011) reported that one rice-based infant cereal was contaminated to FUS at the concentration level of 7.4 mg/Kg (41). In Spain, Meca *et al.* (2010) reported that FUS was found in one rice sample at the level of 3.17 mg/Kg (46).

Moniliformin

In our study, no detectable level of MON was found in samples. Data on the occurrence of MON in rice is limited. Nazari *et al.* (unpublished data) reported that only one out of 100 imported rice samples was contaminated to MON at the level of 8.25 µg/Kg. Yu *et al.* (1995) analyzed 123 rice samples and found MON was present in 8 samples ranging from 73.6 to 265.3 µg/Kg (48).

Estimated daily intake of emerging fusarium mycotoxins

In Iran, two studies regarding estimation of daily intake of AFB1 and DON in foods have been done (49-50). The results obtained in the present study were used to estimate the daily intake of emerging fusarium mycotoxins from rice by the Iranian population. Exposure to mycotoxins for each type of food depends on the mycotoxins level in food and the amount of food consumed. In this study, the consumption rate of rice (110 g per day per person) was based on a consumption survey performed in Iran during 2001-2003 (51) and body weight (b.w) for adults is assumed 60 Kg.

In this study, the EDI of BEA and ENN A1 were 0.2 and 0.11 ng/Kg b.w/day, respectively,

and maximum daily intake of BEA and ENN A1 were 0.8 and 0.15 ng/Kg b.w/day, respectively.

Since JECFA has not set a provisional maximum tolerable daily intake (PMTDI), it is not possible to confirm that these contamination levels of ENs and BEA represent a risk for Iranian consumers.

Conclusion

For the first time, the presence of emerging *Fusarium* mycotoxins especially BEA in Iranian rice was reported. Although the contamination level of BEA was low, considering the toxic effects of BEA, high incidence of the toxin, and high consumption of rice in Iran, BEA may pose health effects to Iranian population. The results of present study indicate the potential health threat due to the contamination of rice with some non-classic mycotoxins which might be neglected in conventional mycotoxin monitoring. Therefore, in Iran, more comprehensive mycotoxin monitoring programs may be needed in rice and other food commodities, in which emerging *Fusarium* mycotoxins, particularly BEA, should be included. In addition, the effect of different methods of food processing on the level of these new emerging mycotoxins in foods, and environmental conditions which influence the occurrence of them should be studied.

References

- (1) Zöllner P and Mayer-Helm B. Trace mycotoxin analysis in complex biological and food matrices by liquid chromatography-atmospheric pressure ionization mass spectrometry. *J. Chromatogr. A* (2006) 1136: 123-169.
- (2) Bennett JW and Klich M. Mycotoxins. *Clin. Microbiol. Rev.* (2003) 16: 497-516.
- (3) Uhlig S, Jestoi M and Parikka P. *Fusarium avenaceum*-the North European situation. *Int. J. Food Microbiol.* (2007) 119: 17-24.
- (4) Jestoi M. Emerging fusarium-mycotoxins fusaproliferin, beauvericin, enniatins, and moniliformin: a review. *Crit. Rev. Food Sci. Nutr.* (2008) 48: 21-49.
- (5) Xu L, Wang J, Zhao J, Li P, Shan T, Wang J, Li X and Zhou L. Beauvericin from the endophytic fungus, *Fusarium redolens*, isolated from *Dioscorea zingiberensis* and its antibacterial activity. *Nat. Prod. Commun.* (2010) 5: 811-814.
- (6) Logrieco A, Rizzo A, Ferracane R and Ritieni A. Occurrence of beauvericin and enniatins in wheat

- affected by *Fusarium avenaceum* head blight. *Appl. Environ. Microbiol.* (2002) 68: 82-85.
- (7) Jestoi M, Rokka M, Rizzo A and Peltonen K. Determination of *Fusarium* mycotoxins beauvericin and enniatins with liquid chromatography-tandem mass spectrometry (LC-MS/MS). *J. Liq. Chromatogr. Related Technol.* (2005) 28: 369-381.
 - (8) Thrane U, Adler A, Clasen PE, Galvano F, Langseth W, Lew H, Logrieco A, Nielsen KF and Ritieni A. Diversity in metabolite production by *Fusarium langsethiae*, *Fusarium poae*, and *Fusarium sporotrichioides*. *Int. J. Food Microbiol.* (2004) 95: 257-266.
 - (9) Moretti A, Mule G, Ritieni A and Logrieco A. Further data on the production of beauvericin, enniatins and fusaproliferin and toxicity to *Artemia salina* by *Fusarium* species of *Gibberella fujikuroi* species complex. *Int. J. Food Microbiol.* (2007) 118: 158-163.
 - (10) Springer J P, Clardy J, Cole R J, Kirksey JW, Hill RK, Carlson RM and Isidor JL. Structure and synthesis of moniliformin, a novel cyclobutane microbial toxin. *J. Am. Chem. Soc.* (1974) 96: 2267-2268.
 - (11) Cole RJ, Kirksey JW, Cutler HG, Doupnik BL and Peckham JC. Toxin from *Fusarium moniliforme*: effects on plants and animals. *Sci.* (1973) 179: 1324-1326.
 - (12) Schutt F, Nirenberg HI and Deml G. Moniliformin production in the genus *Fusarium*. *Mycotoxin Res.* (1998) 14: 35-40.
 - (13) Celik M, Aksoy H and Yilmaz S. Evaluation of beauvericin genotoxicity with the chromosomal aberrations, sister-chromatid exchanges and micronucleus assays. *Ecotoxicol. Environ. Saf.* (2010) 73: 1553-1557.
 - (14) Tan DC, Flematti GR, Ghisalberti EL, Sivasithamparam K and Barbetti MJ. Toxicity of enniatins from Western Australian *Fusarium* species to brine shrimp (*Artemia franciscana*). *Toxicon.* (2011) 57: 817-825.
 - (15) Jow G, Chou C, Chen B and Tsai J. Beauvericin induces cytotoxic effects in human acute lymphoblastic leukemia cells through cytochrome c release, caspase 3 activation: the causative role of calcium. *Cancer Lett.* (2004) 216: 165-173.
 - (16) Lin HI, Lee YJ, Chen BF, Tsai, MC, Lu JL, Chou CJ and Jow GM. Involvement of Bcl-2 family, cytochrome c and caspase 3 in induction of apoptosis by beauvericin in human non-small cell lung cancer cells. *Cancer Lett.* (2005) 230: 248-259.
 - (17) Tonshin AA, Teplova VV, Andersson MA and Salkinoja Salonen MS. The *Fusarium* mycotoxins enniatins and beauvericin cause mitochondrial dysfunction by affecting the mitochondrial volume regulation, oxidative phosphorylation and ion homeostasis. *Toxicol.* (2010) 276: 49-57.
 - (18) Qadri SM, Kucherenko Y and Lang F. Beauvericin induced erythrocyte cell membrane scrambling. *Toxicol.* (2011) 283: 24-31.
 - (19) Tomoda H, Huang XH, Cao J, Nishida H, Nagao R, Okuda S, Tanaka H, Omura S, Arai H and Inoue K. Inhibition of acyl-CoA-cholesterol acyltransferase activity by cyclodepsipeptide antibiotics. *J. Antibiot.* (1992) 45: 1626-1632.
 - (20) Juan-García A, Manyes L, Ruiz MJ and Font G. Involvement of enniatins-induced cytotoxicity in human HepG2 cells. *Toxicol. Lett.* (2013) 218: 166-173.
 - (21) Hyun U, Lee DH, Lee C and Shin CG. Apoptosis induced by enniatins H and MK1688 isolated from *Fusarium oxysporum* FB1501. *Toxicon.* (2009) 53: 723-728.
 - (22) Dornetshuber R, Heffeter P, Kamyar MR, Peterbauer T, Berger W and Lemmens-Gruber R. Enniatin exerts p53-dependent cytostatic and p53-independent cytotoxic activities against human cancer cells. *Chem. Res. Toxicol.* (2007) 20: 465-473.
 - (23) Feifel SC, Schmiederer T, Hornbogen T, Berg H, Süßmuth RD and Zocher R. *In-vitro* synthesis of new enniatins: Probing the alpha-D-hydroxy carboxylic acid binding pocket of the multienzyme enniatin synthetase. *Chem. Biochem.* (2007) 8: 1767-1770.
 - (24) Steenkamp ET, Wingfield BD, Desjardins AE, Marasas WFO and Wingfield MJ. Cryptic speciation in *Fusarium* subglutinans. *Mycologia.* (2002) 94: 1032-1043.
 - (25) Logrieco A, Moretti A, Fornelli F, Fogliano V, Ritieni A, Caiaffa MF, Randazzo G, Bottalico A and Macchia L. Fusaproliferin production by *Fusarium* subglutinans and its toxicity to *Artemia salina* SF-9 insect cells and IARC/LCL 171 human B lymphocytes. *Appl. Environ. Microbiol.* (1996) 62: 3378-3384.
 - (26) Ritieni A, Monti SM, Randazzo G, Logrieco A, Moretti A, Peluso G, Ferracane R and Fogliano V. Teratogenic effects of fusaproliferin on chicken embryos. *J. Agric. Food Chem.* (1997) 45: 3039-3043.
 - (27) Prosperini A, Meca G, Font G and Ruiz MJ. Study of the cytotoxic activity of beauvericin and fusaproliferin and bioavailability *in-vitro* on Caco-2 cells. *Food Chem. Toxicol.* (2012) 50: 2356-2361.
 - (28) Gathercole PS, Thiel PG and Hofmeyr JH. Inhibition of pyruvate dehydrogenase complex by moniliformin. *Biochem. J.* (1986) 233: 719-723.
 - (29) Nagaraj RY, Wu W, Will JA, Valdivia HH and Vesonder RF. Is lactic acid responsible for the electrocardiographic changes induced by intravenous injection of moniliformin. *Poultry Sci.* (1997) 76: 69.
 - (30) Wu W and Vesonder RF. Moniliformin but not fumonisin B1 induced cardiomegaly and elevated serum lactate in broilers. *Poultry Sci.* (1997) 76: 126.
 - (31) Nagaraj RY, Wu W, Will J and Vesonder RF. Acute cardiotoxicity of moniliformin in broiler chickens as measured by electrocardiography. *Avian. Dis.* (1996) 40: 223-227.
 - (32) Li YC, Ledoux DR, Bermudez AJ, Fritsche KL and Rottinghaus GE. The individual and combined effects of fumonisin B1 and moniliformin on performance and selected immune parameters in turkey poults. *Poultry Sci.* (2000) 79: 871-878.
 - (33) Abbas HK, Mirocha CJ, Vesonder RF and Gunther R. Acute toxic effects of an isolate of moniliformin-

- producing *Fusarium oxysporum* and purified moniliformin in rats. *Arch. Environ. Contam. Toxicol.* (1990) 19: 433-436.
- (34) Sulyok M, Krska R and Schuhmacher R. Application of an LC-MS/MS based multi-mycotoxin method for the semi-quantitative determination of mycotoxins occurring in different types of food infected by moulds. *Food Chem.* (2010) 119: 408-416.
- (35) Sørensen JL, Nielsen KF, Rasmussen PH and Thrane U. Development of a LC-MS/MS method for the analysis of enniatins and beauvericin in whole fresh and ensiled maize. *J. Agric. Food Chem.* (2008) 56: 10439-10443.
- (36) Zinedine A, Meca G, Manes J and Font G. Further data on the occurrence of *Fusarium* emerging mycotoxins enniatins (A, A1, B, B1), fusaproliferin and beauvericin in raw cereals commercialized in Morocco. *Food Control* (2011) 22: 1-5.
- (37) Adler A, Lew H, Broddacz W, Edinger W and Oberforster M. Occurrence of oniliformin, deoxynivalenol, and zearalenone in durum wheat. *Mycotoxin Res.* (1995) 11: 9-15.
- (38) Sifou A, Meca G, Serrano AB, Mahnine N, El Abidi A, Mañes J, El Azzouzi M and Zinedine A. First report on the presence of emerging *Fusarium* mycotoxins enniatins (A, A1, B, B1), beauvericin and fusaproliferin in rice on the Moroccan retail markets. *Food Control* (2011) 22: 1826-1830.
- (39) Jestoi M, Rokka M, Yli-Mattila T, Parikka P, Rizzo A and Peltonen K. Presence and concentrations of the *Fusarium* related mycotoxins beauvericin, enniatins and moniliformin in Finnish grain samples. *Food Addit. Contam.* (2004) 21: 794-802.
- (40) Blesa J, Marín R, Lino CM and Mañes J. Evaluation of enniatins A, A1, B, B1 and beauvericin in Portuguese cereal-based foods. *Food Addit. Contam. Part A Chem. Anal. Control Expo. Risk Assess.* (2012) 29: 1727-1735.
- (41) Mahnine N, Meca G, El Abidi A, Fekhaoui M, Saoiabi A, Font G, Manes J and Zinedine A. Further data on the levels of emerging *Fusarium* mycotoxins enniatins (A, A1, B, B1), beauvericin and fusaproliferin in breakfast and infant cereals from Morocco. *Food Chem.* (2011) 124: 481-485.
- (42) Abia WA, Warth B, Sulyok M, Krska R, Tchana AN, Njobeh PB, Dutton MF and Moundipa PF. Determination of multi-mycotoxin occurrence in cereals, nuts and their products in Cameroon by liquid chromatography tandem mass spectrometry (LC-MS/MS). *Food Control* (2013) 31: 438-453.
- (43) Sebastián N, Meca G, Soriano JM and Mañes J. Presence of *Fusarium* emerging mycotoxins in tiger-nuts commercialized in Spain. *Food Control* (2012) 25: 631-635.
- (44) Ezekiel CN, Sulyok M, Warth B, Odebode AC and Krska R. Natural occurrence of mycotoxins in peanut cake from Nigeria. *Food Control* (2012) 27: 338-342.
- (45) Cotty PJ and Jaime-García R. Influences of climate on aflatoxin producing fungi and aflatoxin contamination. *Int. J. Food Microbiol.* (2007) 119: 109-115.
- (46) Meca G, Zinedine A, Blesa J, Font G and Manes J. Further data on the presence of *Fusarium* emerging mycotoxins enniatins, fusaproliferin and beauvericin in cereals available on the Spanish markets. *Food Chem. Toxicol.* (2010) 48: 1412-1416.
- (47) Sulyok M, Berthiller F, Krska R and Schuhmacher R. Development and validation of a liquid chromatography/tandem mass spectrometric method for the determination of 39 mycotoxins in wheat and maize. *Rapid. Commun. Mass Spectrom.* (2006) 20: 2649-2659.
- (48) Yu SR, Liu XJ, Wang YH and Liu J. A survey of moniliformin contamination in rice and corn from Keshan disease endemic and non-KSD areas in China. *Biomed. Environ. Sci.* (1995) 8: 330-334.
- (49) Yazdanpanah H, Zarghi A, Shafaati A, Foroutan SM, Aboul-Fathi F, Khoddam A, Nazari F and Shaki F. Analysis of Aflatoxin B1 in Iranian foods using HPLC and a monolithic column and estimation of its dietary intake. *Iran. J. Pharm. Res.* (2013) 12: 83-89.
- (50) Yazdanpanah H, Shafaati A, Foroutan SM, Zarghi A, Aboul-Fathi F, Khoddam A, Shaki F and Nazari F. Occurrence of deoxynivalenol in foods for human consumption from Tehran, Iran. *Iran. J. Pharm. Res.* (2014) 13: 87-92.
- (51) National Nutrition and Food Technology Research Institute. *Comprehensive Study of Food Basket Pattern, and Nutrition Status in Iran during 2000-2002*. Shahid Beheshti University of Medical Sciences, Tehran (2004).